



Title	MIDPOINTS FOR THOMPSON' S METRIC ON SYMMETRIC CONES
Author(s)	Lemmens, Bas; Roelands, Mark
Citation	Osaka Journal of Mathematics. 2017, 54(1), p. 197-208
Version Type	VoR
URL	https://doi.org/10.18910/61900
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

MIDPOINTS FOR THOMPSON'S METRIC ON SYMMETRIC CONES

BAS LEMMENS and MARK ROELANDS

(Received July 9, 2015, revised February 5, 2016)

Abstract

We characterise the affine span of the midpoints sets, $\mathcal{M}(x, y)$, for Thompson's metric on symmetric cones in terms of a translation of the zero-component of the Peirce decomposition of an idempotent. As a consequence we derive an explicit formula for the dimension of the affine span of $\mathcal{M}(x, y)$ in case the associated Euclidean Jordan algebra is simple. In particular, we find for A and B in the cone positive definite Hermitian matrices that

$$\dim(\text{aff } \mathcal{M}(A, B)) = q^2,$$

where q is the number of eigenvalues μ of $A^{-1}B$, counting multiplicities, such that

$$\mu \neq \max\{\lambda_+(A^{-1}B), \lambda_-(A^{-1}B)^{-1}\},$$

where $\lambda_+(A^{-1}B) := \max\{\lambda : \lambda \in \sigma(A^{-1}B)\}$ and $\lambda_-(A^{-1}B) := \min\{\lambda : \lambda \in \sigma(A^{-1}B)\}$. These results extend work by Y. Lim [18].

1. Introduction

The space of $n \times n$ Hermitian matrices contains a cone $\Pi_n(\mathbb{C})$ of all positive-semidefinite matrices. Its interior, $\Pi_n(\mathbb{C})^\circ$, consists of all invertible elements, and is a prime example of a symmetric cone. It is well known, see for example [4], that $\Pi_n(\mathbb{C})^\circ$ can be equipped with a Riemannian metric

$$\delta_2(A, B) := \|\log(A^{-1}B)\|_2 = \left(\sum_{i=1}^n (\log \lambda_i(A^{-1}B))^2 \right)^{1/2},$$

where the $\lambda_i(A^{-1}B)$'s are the eigenvalues of $A^{-1}B$. The metric space $(\Pi_n(\mathbb{C})^\circ, \delta_2)$ is geodesic, i.e., any two points are connected by a geodesic. In fact, in this case the geodesic is unique and given by the path,

$$t \mapsto A^{1/2}(A^{-1/2}BA^{-1/2})^t A^{1/2},$$

where $t \in [0, 1]$, and the *geometric mean*

$$A \sharp B := A^{1/2}(A^{-1/2}BA^{-1/2})^{1/2}A^{1/2}$$

is the unique midpoint of A and B .

Another natural metric on $\Pi_n(\mathbb{C})^\circ$ is *Thompson's metric*,

$$d_T(A, B) := \|\log(A^{-1}B)\|_\infty = \max_i |\log \lambda_i(A^{-1}B)|.$$

The space $(\Pi_n(\mathbb{C})^\circ, d_T)$ is a geodesic Finsler metric space, see [17, 21], in which the path $t \mapsto A^{1/2}(A^{-1/2}BA^{-1/2})^t A^{1/2}$ is also a geodesic, but in general not unique.

Thompson's metric, which was introduced in [24], is a useful metric that can be defined on the interior of any closed cone in a normed space. It is widely applied in the spectral theory of linear and nonlinear operators on cones [1, 8, 9, 14, 15, 22, 24].

It is also studied in the geometry of spaces of positive operators, where it provides a useful alternative for the usual Riemannian metric, see [3, 5, 6, 10, 20]. Unlike the Riemannian case, Thompson's metric is not uniquely geodesic. Understanding the geodesic structure of these metric spaces is not only of geometric interest, but also plays an important role in the study of operator means [11, 12, 13, 19, 23]. Motivated by understanding the properties of various matrix means, Lim studied in [18] the geometry of the *midpoints set*

$$\mathcal{M}(A, B) := \left\{ C \in \Pi_n(\mathbb{C})^\circ : d_T(A, C) = \frac{1}{2}d_T(A, B) = d_T(C, B) \right\}$$

for $A, B \in \Pi_n(\mathbb{C})^\circ$. Among other results Lim [18, Theorem 5.2] showed that $\mathcal{M}(A, B)$ is a singleton if, and only if, the spectrum of $A^{-1}B$ satisfies $\sigma(A^{-1}B) \subseteq \{\alpha, \alpha^{-1}\}$ for some $\alpha > 0$. This result has been generalised by the authors in [16] to the cone of positive self-adjoint elements in a unital C^* algebra and symmetric cones.

In general cones the midpoints set $\mathcal{M}(x, y) := \{z \in K^\circ : d_T(x, z) = \frac{1}{2}d_T(x, y) = d_T(z, y)\}$ is convex, as it is the intersection of the Thompson metric balls $B(x, 1/2)$ and $B(y, 1/2)$, which are both convex, see [14, Lemma 2.6.2]. The main goal of this paper is to characterise the affine span of the midpoints set $\mathcal{M}(x, y)$ for x and y in a symmetric cone in terms of a translation of the zero-component of the Peirce decomposition of an idempotent. As a corollary we obtain an explicit formula for the dimension of $\mathcal{M}(x, y)$, which is equal to the dimension of its affine span ($\mathcal{M}(x, y)$ is convex), in case the associated Euclidean Jordan algebra is simple. In the special case where $A, B \in \Pi_n(\mathbb{C})^\circ$ we find that

$$\dim(\mathcal{M}(A, B)) = q^2,$$

where q is the number of eigenvalues μ of $A^{-1}B$, counting multiplicities, such that

$$\mu \neq \max\{\lambda_+(A^{-1}B), \lambda_-(A^{-1}B)^{-1}\}$$

and $\lambda_+(A^{-1}B) := \max\{\lambda : \lambda \in \sigma(A^{-1}B)\}$ and $\lambda_-(A^{-1}B) := \min\{\lambda : \lambda \in \sigma(A^{-1}B)\}$.

To obtain the results we first prove a characterisation of the midpoints set in a general cone in terms of its faces, see Theorem 3.2. This result is subsequently used in Section 4 to find the affine span of the midpoints set, and its dimension, in symmetric cones. We begin by recalling some basic definitions.

2. Preliminaries

A cone K in a vector space V is a convex subset such that $K \cap (-K) = \{0\}$ and $\lambda K \subseteq K$ for all $\lambda \geq 0$. It induces a partial ordering $\leq_K$ on V by putting $x \leq_K y$ if $y - x \in K$. Given $x \leq_K y$ in V we denote the *order interval* by $[x, y]_K := \{z \in V : x \leq_K z \leq_K y\}$. A non-empty convex subset $F \subseteq K$ is said to be a *face* of K if $x, y \in K$ such that $\lambda x + (1 - \lambda)y \in F$ for some $0 < \lambda < 1$ implies that $x, y \in F$. The face generated by $z \in K$ is denoted F_z , i.e.,

$$F_z := \{y \in K : \lambda y + (1 - \lambda)z \in K \text{ for some } \lambda < 0\}.$$

The faces can be characterised as follows.

Lemma 2.1. *If (V, K) is a partially ordered vector space and $z \in K$, then*

$$F_z = \bigcup_{n \geq 1} [0, nz]_K.$$

Proof. Note that $\lambda y + (1 - \lambda)z \in K$ for some $\lambda < 0$ if and only if $-\mu y + (1 + \mu)z \in K$ for some $\mu > 0$, which is equivalent to $y \leq_K \alpha z \in K$ for some $\alpha > 0$. Thus, $F_z = \cup_{\alpha \geq 1} [0, \alpha z]_K = \cup_{n \geq 1} [0, nz]_K$. $\square$

We say that a cone K is *Archimedean* if for all $x \in V$ and $y \in K$ with $nx \leq_K y$ for $n \geq 1$ we have that $x \leq_K 0$. An element $u \in K$ is called an *order unit* if for each $x \in V$ there exists $\lambda > 0$ such that $x \leq_K \lambda u$. The triple (V, K, u) is called an *order unit space* if K is Archimedean and u is an order unit.

An order unit space (V, K, u) can be equipped with the *order unit norm* $\|\cdot\|_u$, which is defined by

$$\|x\|_u := \inf\{\lambda > 0 : -\lambda u \leq_K x \leq_K \lambda u\}.$$

With respect to this norm, the cone K is closed by [2, Theorem 2.55(2)]. Furthermore, the norm $\|\cdot\|_u$ is *monotone*, that is, $\|x\|_u \leq \|y\|_u$ for all $0 \leq_K x \leq_K y$. In particular, K is a *normal* cone with respect to $\|\cdot\|_u$, i.e., there exists a constant $\kappa > 0$ such that $\|x\|_u \leq \kappa\|y\|_u$ whenever $x \leq_K y$ in V . It is known, see [2, Lemma 2.5], that each interior point of K is an order unit of K . On the other hand, if $x \in K$ is an order unit of K , then there exists $M > 0$ such that $u \leq_K Mx$. So, for $y \in V$ with $\|y\|_u \leq 1/M$ we have that $0 \leq_K x - u/M \leq_K x - y$, which show that $x \in K^\circ$. Thus, the interior K° coincides with the set of order units of K .

We see that given an order unit space (V, K, u) and $x, y \in K^\circ$, there are constants $0 < \beta$ such that $x \leq_K \beta y$, and hence we can define

$$M(x/y) := \inf\{\beta > 0 : x \leq_K \beta y\} < \infty.$$

Now *Thompson's metric* on K° is given by

$$d_T(x, y) := \log(\max\{M(x/y), M(y/x)\})$$

and was introduced in [24]. The set of midpoints is denoted

$$\mathcal{M}(x, y) := \left\{ z \in K^\circ : d_T(x, z) = \frac{1}{2}d_T(x, y) = d_T(z, y) \right\}.$$

Thompson's metric spaces (K°, d_T) are geodesic spaces, see [14, Section 2.6], that is to say, any two points in K° are connected by a geodesic segment. Recall that a map γ from an (open, closed, bounded, or, unbounded) interval $I \subseteq \mathbb{R}$ into a metric space (X, d) is called a *geodesic path* if $d(\gamma(s), \gamma(t)) = |s - t|$ for all $s, t \in I$. The image of γ is called a *geodesic segment* in (X, d) , and for $x, y \in (X, d)$.

3. Midpoints in general cones

Before we give the characterisation of the midpoints set in (K°, d_T) , where (V, K, u) is an order unit space, we make some preliminary observations. To begin, we note that if $x, y \in K^\circ$ are linearly dependent, then the straight-line segment connecting x and y is a unique geodesic segment for Thompson's metric, see [16, Lemma 3.3], and hence the midpoints set is a singleton in that case. So, in the sequel we only need to consider the midpoints sets of linearly independent elements of K° .

If $x, y \in K^\circ$ are linearly independent, then we write $V(x, y) = \text{span}(x, y)$ and we let $K(x, y) := V(x, y) \cap K$ be the 2-dimensional cone containing x and y , which has relative interior $K(x, y)^\circ$ in $V(x, y)$. Note that for $w, z \in K(x, y)^\circ$ the distance $d_T(w, z)$ with respect to $K(x, y)$ is the same as $d_T(w, z)$ with respect to K . As $K(x, y)$ is a closed cone in $V(x, y)$ and $K(x, y)^\circ$ is non-empty, we know [14, Theorem A.5.1] that there exist linearly independent linear functionals ψ_1 and ψ_2 on $V(x, y)$ such that

$$K(x, y) = \{z \in V(x, y) : \psi_1(z) \geq 0 \text{ and } \psi_2(z) \geq 0\}.$$

The linear map $\Psi : V(x, y) \rightarrow \mathbb{R}^2$ given by, $\Psi(z) = (\psi_1(z), \psi_2(z))$ for $z \in V(x, y)$, maps $K(x, y)$ onto the standard positive cone $\mathbb{R}_+^2 := \{(w_1, w_2) : w_1 \geq 0 \text{ and } w_2 \geq 0\}$. Furthermore for $u, v \in K(x, y)^\circ$ we have that $M(u/v) = M(\Psi(u)/\Psi(v))$, and hence Ψ is a d_T -isometry. One can verify that in $((\mathbb{R}_+^2)^\circ, d_T)$ the path $t \mapsto \Psi(x)^{1-t}\Psi(y)^t$, for $t \in [0, 1]$, where $\Psi(x)^{1-t}\Psi(y)^t := (\Psi(x)_1^{1-t}\Psi(y)_1^t, \Psi(x)_2^{1-t}\Psi(y)_2^t) \in (\mathbb{R}_+^2)^\circ$, is a geodesic path from $\Psi(x)$ to $\Psi(y)$. The pull-back of this geodesic path under the isometry Ψ is an geodesic path connecting x and y in $(K(x, y)^\circ, d_T)$. We will call it the *canonical geodesic* connecting x and y and denote it by γ_{xy} . Moreover, the midpoint

$$m_{xy} := \gamma_{xy}(1/2)$$

is said to be the *canonical midpoint* of x and y . Note that

$$M(x/m_{xy}) = M(\Psi(x)/\Psi(x)^{1/2}\Psi(y)^{1/2}) = M(\Psi(x)^{1/2}/\Psi(y)^{1/2}) = M(x/y)^{1/2},$$

so that $M(x/y) = M(x/m_{xy})^2$. Likewise it can be shown that

$$(3.1) \quad M(m_{xy}/y)^2 = M(x/y) \quad \text{and} \quad M(m_{xy}/x)^2 = M(y/x) = M(y/m_{xy})^2.$$

Given $x, y \in K^\circ$, we let $\ell_{xy}^+ := \{\lambda y + (1 - \lambda)x : \lambda \geq 0\}$ be the half-line emanating from x through y . The following basic observation will be useful.

Lemma 3.1. *Let (V, K, u) be an order unit space and $x, y \in K^\circ$. If $M := M(x/y) > 1$, then ℓ_{xy}^+ intersects ∂K in*

$$(3.2) \quad y' := \frac{M}{M-1}y + \frac{1}{1-M}x.$$

Proof. Note that, as K is closed in (V, K, u) , we have that $y - M^{-1}x \in \partial K$. This implies that $y' := \frac{M}{M-1}y + \frac{1}{1-M}x \in \partial K$. As y' is also on ℓ_{xy}^+ the result follows. $\square$

Similarly, if $M = M(y/x) > 1$, we get that ℓ_{yx}^+ intersects ∂K in the point

$$(3.3) \quad x' := \frac{M}{M-1}x + \frac{1}{1-M}y.$$

For a non-empty subset $W \subseteq V$ we will denote the affine span of W in V by $\text{aff } W$.

Theorem 3.2. *Let (V, K, u) be an order unit space and $x, y \in K^\circ$ be linearly independent. Then the affine span of $\mathcal{M}(x, y)$ satisfies:*

- (i) $\text{aff } \mathcal{M}(x, y) = m_{xy} + \text{span } F_{y'}$, if $M(x/y) > M(y/x)$;
- (ii) $\text{aff } \mathcal{M}(x, y) = m_{xy} + \text{span } F_{x'}$, if $M(y/x) > M(x/y)$;
- (iii) $\text{aff } \mathcal{M}(x, y) = m_{xy} + \text{span } F_{x'} \cap \text{span } F_{y'}$, if $M(x/y) = M(y/x)$,

where x' and y' are as in (3.3) and (3.2), respectively.

Proof. Note that case (ii) follows from case (i) by symmetry. So, suppose that $M(x/y) > M(y/x)$ and write $M := M(x/y)$. As $d_T(x, y) = \log M$, we see that $M > 1$. It now follows from Lemma 3.1 that ℓ_{xy}^+ intersects ∂K in

$$y' := \frac{M}{M-1}y + \frac{1}{1-M}x.$$

Let $z \in \mathcal{M}(x, y)$. We deduce from

$$d_T(x, y) \leq \log M(x/z) + \log M(z/y) \leq d_T(x, z) + d_T(z, y) = d_T(x, y)$$

that $d_T(x, z) = \log M(x/z) = \log M(z/y) = d_T(z, y)$, so that $M(x/z)^2 = M(z/y)^2 = M$. Write $N := M(x/z) > 1$. Again using Lemma 3.1 the half-line ℓ_{xz}^+ intersects ∂K in

$$z_1 := \frac{N}{N-1}z + \frac{1}{1-N}x,$$

and ℓ_{zy}^+ intersects ∂K in

$$z_2 := \frac{N}{N-1}y + \frac{1}{1-N}z,$$

see Figure 1.




FIG. 1. Endpoints

Working out the following convex combination

$$\begin{aligned} \frac{1}{N+1}z_1 + \frac{N}{N+1}z_2 &= \left(\frac{N}{N^2-1}z + \frac{1}{1-N^2}x \right) + \left(\frac{N^2}{N^2-1}y + \frac{N}{1-N^2}z \right) \\ &= \frac{N^2}{N^2-1}y + \frac{1}{1-N^2}x = \frac{M}{M-1}y + \frac{1}{1-M}x \\ &= y' \end{aligned}$$

shows that z_1 and z_2 both belong to the face $F_{y'}$.

Write $z := m_{xy} + v$ for some $v \in V$. Let $\bar{m}_{xy} \in F_{y'}$ be the points of intersection of $\ell_{m_{xy}y}^+$ and ∂K . So,

$$\bar{m}_{xy} = \frac{N}{N-1}y + \frac{1}{1-N}m_{xy}$$

by Lemma 3.1. It follows that $v = (N-1)(\bar{m}_{xy} - z_2) \in \text{span } F_{y'}$ which yields the inclusion

$$\text{aff } \mathcal{M}(x, y) \subseteq m_{xy} + \text{span } F_{y'}.$$

Conversely, suppose $v \in \text{span } F_{y'}$, with $v \neq 0$. Define $z := m_{xy} + v$. By Lemma 3.1 the point

$$\bar{m}_{xy} := \frac{N}{N-1}y + \frac{1}{1-N}m_{xy}$$

lies in ∂K . As γ_{xy} lies in $K(x, y)$, we see that $\bar{m}_{xy}$ is a positive scalar multiple of y' , and hence $\bar{m}_{xy}$ lies in the relative interior of $F_{y'}$. Let $t = (1-N)^{-1}$ and note that

$$\bar{m}_{xy} + tv = \frac{N}{N-1}y + \frac{1}{1-N}(m_{xy} + v).$$

As $\bar{m}_{xy}$ is in the relative interior of $F_{y'}$, we can replace v by ϵv for some $\epsilon > 0$ sufficiently small, and assume that $\bar{m}_{xy} + tv \in F_{y'}$ and $m_{xy} + v \in K^\circ$. We know from [14, pp. 28–29] that

$$M(m_{xy}/y) = \frac{|m_{xy}\bar{m}_{xy}|}{|y\bar{m}_{xy}|} \quad \text{and} \quad M(z/y) = \frac{|z(\bar{m}_{xy} + tv)|}{|y(\bar{m}_{xy} + tv)|},$$

where $|uw'|/|ww'|$ denotes the ratio of the lengths of the straight-line segments $[u, w']$ and $[w, w']$, see Figure 2. Using similarity of triangles we conclude that $M(m_{xy}/y) = M(z/y)$.




FIG. 2. The endpoints in $F_{y'}$

Similarly, let

$$\hat{m}_{xy} = \frac{N}{N-1}m_{xy} + \frac{1}{1-N}x.$$

Note that $\hat{m}_{xy} \in \partial K$ by Lemma 3.1 and is a positive scalar multiple of y' , as γ_{xy} is contained in $K(x, y)$. By possibly further reducing $\epsilon > 0$ we may assume for $s := N/(N-1)$ that

$$\hat{m}_{xy} + sv = \frac{N}{N-1}(m_{xy} + v) + \frac{1}{1-N}x \in F_{y'}$$

and $m_{xy} + v \in K^\circ$. Again using similarity of triangles, see Figure 3, we get that

$$M(x/m_{xy}) = \frac{|x\hat{m}_{xy}|}{|m_{xy}\hat{m}_{xy}|} = \frac{|x(\hat{m}_{xy} + sv)|}{|z(\hat{m}_{xy} + sv)|} = M(x/z).$$




FIG. 3. The endpoints in $F_{y'}$

It now follows from (3.1) that

$$M(x/z)^2 = M(x/m_{xy})^2 = M(x/y) > M(y/x) = M(m_{xy}/x)^2.$$

As the map $(u, w) \mapsto M(u/w)$ is continuous on $K^\circ \times K^\circ$, see [15, Lemma 2.2], we can assume, after possibly further reducing $\epsilon > 0$, that $M(z/x)^2 < M(x/y)$. It now follows that $d_T(x, z) = d_T(x, m_{xy}) = \frac{1}{2}d_T(x, y)$. In the same way it can be shown that $d_T(z, y) = d_T(m_{xy}, y) = \frac{1}{2}d_T(x, y)$. We conclude that $z \in \mathcal{M}(x, y)$ and hence $m_{xy} + \text{span } F_{y'} \subseteq \text{aff } \mathcal{M}(x, y)$.

Finally, suppose that $M(x/y) = M(y/x)$. We have already shown that the inclusions $\text{aff } \mathcal{M}(x, y) \subseteq m_{xy} + \text{span } F_{x'}$ and $\text{aff } \mathcal{M}(x, y) \subseteq m_{xy} + \text{span } F_{y'}$ hold, which immediately implies that

$$\text{aff } \mathcal{M}(x, y) \subseteq m_{xy} + \text{span } F_{x'} \cap \text{span } F_{y'}.$$

Moreover, if $v \in \text{span } F_{x'} \cap \text{span } F_{y'}$ and $z := m_{xy} + \epsilon v$, then we have also shown that for small enough $\epsilon > 0$, the equalities $\log M(x/z) = \log M(z/y) = \frac{1}{2}d_T(x, y)$ hold. Now since $M(x/y) = M(y/x)$, we can apply the same argument to show that $\log M(z/x) = \log M(y/z) = \frac{1}{2}d_T(x, y)$, and hence

$$m_{xy} + \text{span } F_{x'} \cap \text{span } F_{y'} \subseteq \text{aff } \mathcal{M}(x, y),$$

which proves the last assertion. $\square$

4. Midpoints in symmetric cones

The interior K° of a closed cone K in a finite-dimensional inner-product space $(V, \langle \cdot, \cdot \rangle)$ is called a *symmetric cone* if the *dual cone*, $K^* := \{y \in V : \langle y, x \rangle \geq 0 \text{ for all } x \in K\}$ satisfies $K^* = K$, and the automorphism group $\text{Aut}(K) := \{A \in \text{GL}(V) : A(K) = K\}$ acts transitively on K° . A prime example is the cone of positive definite Hermitian matrices.

It is well known that the symmetric cones in finite dimensions are precisely the interiors of the cones of squares of Euclidean Jordan algebras. We will follow the notation and terminology from [7], which gives detailed account of the theory of symmetric cones.

A *Euclidean Jordan algebra* is a finite-dimensional real inner-product space $(V, \langle \cdot, \cdot \rangle)$ equipped with a bilinear product $(x, y) \mapsto x \bullet y$ from $V \times V$ into V such that for each $x, y \in V$:

- (i) $x \bullet y = y \bullet x$,
- (ii) $x \bullet (x^2 \bullet y) = x^2 \bullet (x \bullet y)$, and
- (iii) for each $x \in V$, the linear map $L(x) : V \rightarrow V$ given by $L(x)y := x \bullet y$ satisfies

$$\langle L(x)y, z \rangle = \langle y, L(x)z \rangle \quad \text{for all } y, z \in V.$$

A Euclidean Jordan algebra is not associative in general, but it is commutative. The *unit* in a Euclidean Jordan algebra is denoted by e . An element $c \in V$ is called an *idempotent* if $c^2 = c$. A set $\{c_1, \dots, c_k\}$ is called a *complete system of orthogonal idempotents* if

- (i) $c_i^2 = c_i$ for all i ,
- (ii) $c_i \bullet c_j = 0$ for all $i \neq j$, and
- (iii) $c_1 + \dots + c_k = e$.

The spectral theorem [7, Theorem III.1.1] says that for each $x \in V$ there exist unique real numbers $\lambda_1, \dots, \lambda_k$, all distinct, and a complete system of orthogonal idempotents $c_1, \dots, c_k$ such that $x = \lambda_1 c_1 + \dots + \lambda_k c_k$. The numbers λ_i are called the *eigenvalues* of x . The *spectrum* of x is denoted by $\sigma(x) = \{\lambda : \lambda \text{ eigenvalue of } x\}$, and we write

$$\lambda_+(x) = \max\{\lambda : \lambda \in \sigma(x)\} \quad \text{and} \quad \lambda_-(x) = \min\{\lambda : \lambda \in \sigma(x)\}.$$

The spectral decomposition gives rise to a functional calculus on V . For example, for $x = \lambda_1 c_1 + \dots + \lambda_k c_k$ with $\lambda_i > 0$ for all $i = 1, \dots, k$, we can define $x^{-1/2} := \lambda_1^{-1/2} c_1 + \dots + \lambda_k^{-1/2} c_k$.

For $x \in V$ the linear mapping, $P(x) = 2L(x)^2 - L(x^2)$, is called the *quadratic representation* of x . Note that $P(x^{-1/2})x = e$ for all $x \in K^\circ$. It is known that $P(x^{-1}) = P(x)^{-1}$ for all $x \in K^\circ$ and $P(x) \in \text{Aut}(K)$ whenever $x \in K^\circ$, see [7, Proposition III.2.2]. So, $P(x)$ is an isometry of (K°, d_T) if $x \in K^\circ$ by [14, Corollary 2.1.4]. For $x, y \in K^\circ$ we write

$$\lambda_+(x, y) = \lambda_+(P(y^{-1/2})x) \quad \text{and} \quad \lambda_-(x, y) = \lambda_-(P(y^{-1/2})x).$$

Note that for $x, y \in K^\circ$, $x \leq_K \beta y$ if and only if $0 \leq_K \beta e - P(y^{-1/2})x$, and hence

$$M(x/y) = \lambda_+(x, y).$$

Similarly, $\alpha y \leq_K x$ is equivalent with $0 \leq_K P(y^{-1/2})x - \alpha e$, and hence

$$M(y/x)^{-1} = \lambda_-(x, y).$$

So, for $x, y \in K^\circ$ the Thompson metric distance is given by

$$d_T(x, y) = \log \left(\max \{ \lambda_+(x, y), \lambda_-(x, y)^{-1} \} \right).$$

For $A, B \in \Pi_n(\mathbb{C})^\circ$ we have that $P(B^{-\frac{1}{2}})A = B^{-\frac{1}{2}}AB^{-\frac{1}{2}}$; so, in that case

$$\begin{aligned} d_T(A, B) &= \max \left\{ \max_i \log \lambda_i(B^{-\frac{1}{2}}AB^{-\frac{1}{2}}), \max_i -\log \lambda_i(B^{-\frac{1}{2}}AB^{-\frac{1}{2}}) \right\} \\ &= \max_i |\log \lambda_i(B^{-1}A)|. \end{aligned}$$

The quadratic representation $P(y^{-1/2})$ of $y \in K^\circ$ is an isometry with respect to Thompson's metric, and hence $z \in \mathcal{M}(x, y)$ if and only if $P(y^{-\frac{1}{2}})z \in \mathcal{M}(P(y^{-\frac{1}{2}})x, e)$. Thus, without loss of generality, we may consider midpoint sets of the form $\mathcal{M}(x, e)$ where $x \in K^\circ$.

The following lemma, which is Exercise III.3 in [7] will be useful in the sequel. A proof can be found in [16, Lemma 6.1].

Lemma 4.1. *Let K° be a symmetric cone. For $x, y \in K$ we have $\langle x, y \rangle = 0$ if and only if $x \bullet y = 0$.*

Given an idempotent $c \in K$ we have the Peirce decomposition

$$V = V(c, 0) \oplus V(c, \frac{1}{2}) \oplus V(c, 1)$$

where $V(c, \lambda)$ are the corresponding eigenspaces of the only possible eigenvalues λ that the linear operator $L(c)$ can have, see [7, Proposition III.1.3]. Although this is a direct sum of vector spaces, both components $V(c, 0)$ and $V(c, 1)$ are Jordan subalgebras [7, Proposition IV.1.1], and for $\lambda = 0, 1$ we will denote the cone of squares in $V(c, \lambda)$ by $K(c, \lambda)$. Regarding the midpoint sets, we are particularly interested in $V(c, 0)$. Note that this subalgebra has $e - c$ as a unit.

For $x \in K^\circ$ with spectral decomposition $x = \sum_{i=1}^k \lambda_i c_i$ we let

$$C_x := \{c_i \in \{c_1, \dots, c_k\} : \max\{\lambda_i, \lambda_i^{-1}\} = \max\{\lambda_+(x), \lambda_-(x)^{-1}\}\}.$$

Note that after reordering the eigenvalues $\lambda_1 < \lambda_2 < \dots < \lambda_k$ we have that $C_x \subseteq \{c_1, c_n\}$. Before we characterise the affine span of the midpoints set $\mathcal{M}(x, e)$ for $x \in K^\circ$, we first prove the following lemma.

Lemma 4.2. *Let V be a Euclidean Jordan algebra with cone of squares K and let $c \in K$ be an idempotent. Then $K(c, 0)$ is a face of K with relative interior*

$$K(c, 0)^\circ = \text{Inv}(V(c, 0)) \cap K(c, 0),$$

where $\text{Inv}(V(c, 0))$ denotes the set of invertible elements in the subalgebra $V(c, 0)$.

Proof. Let $z \in K(c, 0)$. If $\xi_1, \xi_2 \in K$ and $0 < t < 1$ are such that $z = t\xi_1 + (1 - t)\xi_2$, then

$$0 \leq t \langle c, \xi_1 \rangle + (1 - t) \langle c, \xi_2 \rangle = \langle c, z \rangle = 0,$$

so $\xi_1, \xi_2 \in K(c, 0)$ by Lemma 4.1, and hence $K(c, 0)$ is a face of K . Note that the Jordan subalgebra $V(c, 0)$ has unit $e - c \in K(c, 0)$, since $(e - c)^2 = e - c$. The fact that $K(c, 0)^\circ =$

$\text{Inv}(V(c, 0)) \cap K(c, 0)$ now follows from [7, Theorem III.2.1]. $\square$

Note that $K(c, 0) = F_{e-c}$. Indeed, if $x \in K(c, 0)$, then $x \leq n(e - c)$ for some $n \geq 1$, as $e - c$ is an order unit in $V(c, 0)$. So, $K(c, 0) \subseteq F_{e-c}$ by Lemma 2.1. Conversely, if $y \in F_{e-c}$, then $0 \leq_K y \leq_K n(e - c)$ for some $n \geq 1$. It now follows that

$$0 \leq \langle c, y \rangle = \langle c, y \rangle - \langle c, n(e - c) \rangle = \langle c, y - n(e - c) \rangle \leq 0,$$

and hence $y \in K(c, 0)$ by Lemma 4.1.

We can now prove the characterisation of the midpoints set in symmetric cones.

Theorem 4.3. *Let K° be a symmetric cone. For $x \in K^\circ \setminus \{e\}$ let C_x be defined as above and put $c := \sum_{c_i \in C_x} c_i$. The affine span of $\mathcal{M}(x, e)$ satisfies*

$$\text{aff } \mathcal{M}(x, e) = m_{xe} + V(c, 0).$$

Proof. Let $x = \lambda_1 c_1 + \cdots + \lambda_k c_k$ be the spectral decomposition of x with $\lambda_1 < \cdots < \lambda_k$. First suppose that $C_x = \{c_k\}$. Note that $\lambda_k > 1$, as $d_T(x, e) = \log \lambda_k > 0$. The endpoint

$$x' = \frac{\lambda_k}{\lambda_k - 1} e - \frac{1}{\lambda_k - 1} x = \sum_{i=1}^{k-1} \frac{\lambda_k - \lambda_i}{\lambda_k - 1} c_i,$$

where e is between x' and x , is in the relative interior of $K(c, 0)$ by Lemma 4.2, as it is invertible in $V(c, 0)$ with respect to $e - c_k = c_1 + \cdots + c_{k-1}$. The desired equality now follows from Theorem 3.2 since $K(c, 0)$ is generating in $V(c, 0)$. In the same way it can be shown that the assertion holds if $C_x = \{c_1\}$. Finally, if $C_x = \{c_1, c_k\}$, then $c = c_1 + c_k$ and it follows from Theorem 3.2 that

$$\text{aff } \mathcal{M}(x, e) = m_{xe} + V(c_1, 0) \cap V(c_k, 0).$$

Clearly $V(c, 0) \supseteq V(c_1, 0) \cap V(c_k, 0)$. As $V(c_i, 0) = \ker L(c_i)$ for $i = 1, k$ and $K(c, 0) \subseteq \ker L(c_1) \cap \ker L(c_k)$, we must have that $V(c, 0) = V(c_1, 0) \cap V(c_k, 0)$, since $\text{span } K(c, 0) = V(c, 0)$. $\square$

It follows from Theorem 4.3 that $\dim \text{aff}(\mathcal{M}(x, e)) = \dim V(c, 0)$. If V is a simple n -dimensional Euclidean Jordan algebra, i.e., V has no non-trivial ideals, then for any two orthogonal primitive idempotents c_1 and c_2 in V the dimension

$$d := \dim V(c_1, \tfrac{1}{2}) \cap V(c_2, \tfrac{1}{2})$$

is independent of c_1 and c_2 . In fact, if $\text{rank}(V) = r$, then

$$n = r + \frac{d}{2}r(r - 1)$$

by [7, Corollary IV.2.6].

Now let us decompose x with respect to a Jordan frame, for details see [7, Theorem III.1.2], where we might have different primitive idempotents corresponding to the same eigenvalues, so

$$x = \lambda_1 c_1 + \cdots + \lambda_r c_r.$$

We can rearrange the eigenvalues in such a way that

$$(4.1) \quad x = \sum_{j \in C_x^c} \lambda_j c_j + \sum_{c_i \in C_x} \lambda_i c_i.$$

It follows that for $q := |C_x^c|$ the dimensional formula

$$\dim V(c, 0) = \dim V(e - c, 1) = q + \frac{d}{2}q(q - 1)$$

holds by [7, Proposition IV.3.1]. As $\dim \mathcal{M}(x, e) = \dim \text{aff}(\mathcal{M}(x, e))$, this immediately gives the following corollary.

Corollary 4.4. *Let V be a simple n -dimensional Euclidean Jordan algebra with cone of squares K and rank $r > 1$. If $x \in K^\circ$, then the affine dimension of the midpoints set satisfies*

$$\dim \mathcal{M}(x, e) = q + \frac{d}{2}q(q - 1) = q + \frac{n - r}{r(r - 1)}q(q - 1).$$

As an example, let us consider the Hermitian matrices $\mathbb{H}_n(\mathbb{C})$ and compute $\dim(\text{aff } \mathcal{M}(A, I_n))$ for $A \in \Pi_n(\mathbb{C})^\circ$ and I_n the $n \times n$ identity matrix. There exists a unitary matrix U such that $UAU^* = D$ where D is a diagonal matrix with the eigenvalues on the diagonal arranged as in (4.1). Conjugating with U is a linear automorphism of $\Pi_n(\mathbb{C})$, and $B \in \mathcal{M}(A, I_n)$ if and only if $U^*BU \in \mathcal{M}(D, I_n)$. So, to compute the dimension of $\text{aff } \mathcal{M}(A, I_n)$ we may assume without loss of generality that A is a diagonal matrix as described above. In that case the projection C is of the form

$$C = \sum_{i=q+1}^n E_{ii} = \begin{pmatrix} 0 & 0 \\ 0 & I_{n-q} \end{pmatrix},$$

where $q = |C_A^c|$. It is easily checked that $V(C, 0)$ equals

$$V(C, 0) = \left\{ \begin{pmatrix} A_q & 0 \\ 0 & 0 \end{pmatrix} : A_q \in \mathbb{H}_q(\mathbb{C}) \right\},$$

see [7, p. 63], and $\dim(V(C, 0)) = q^2$. Since $\text{rank}(V) = n$ and $\dim(V) = n^2$, it follows from Corollary 4.4 that

$$\dim \mathcal{M}(A, I_n) = q + q(q - 1) = q^2.$$

References

- [1] M. Akian, S. Gaubert, B. Lemmens, and R.D. Nussbaum: *Iteration of order preserving subhomogeneous maps on a cone*, Math. Proc. Cambridge Philos. Soc. **140** (2006), 157–176.
- [2] C.D. Aliprantis and R. Tourky: *Cones and duality*, Graduate Studies in Mathematics, 84. American Mathematical Society, Providence, RI, 2007.
- [3] E. Andruchow, G. Corach and D. Stojanoff: *Geometrical significance of Löwner-Heinz inequality*, Proc. Amer. Math. Soc. **128** (2000), 1031–1037.
- [4] R. Bhatia: *Positive definite matrices*. Princeton Series in Applied Mathematics. Princeton University Press, Princeton, NJ, 2007.

- [5] G. Corach and A.L. Maestri: *Differential and metrical structure of positive operators*, Positivity **3** (1999), 297–315.
- [6] G. Corach, H. Porta, and L. Recht: *Convexity of the geodesic distance on spaces of positive operators*, Illinois J. Math. **38** (1994), 87–94.
- [7] J. Faraut and A. Korányi: *Analysis on Symmetric Cones*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1994.
- [8] S. Gaubert and Z. Qu: *The contraction rate in Thompson’s part metric of order-preserving flows on a cone: application to generalized Riccati equations*, J. Differential Equations **256** (2014), 2902–2948.
- [9] D.H. Hyers, G. Isac, and T.M. Rassias: *Topics in nonlinear analysis & applications*. World Scientific Publishing Co., Inc., River Edge, NJ, 1997.
- [10] J. Lawson and Y. Lim: *Metric convexity of symmetric cones*, Osaka J. Math. **44** (2007), 795–816.
- [11] J. Lawson and Y. Lim: *Weighted means and Karcher equations of positive operators*, Proc. Natl. Acad. Sci. USA **110** (2013), 5626–5632.
- [12] J. Lawson and Y. Lim: *Karcher means and Karcher equations of positive definite operators*, Trans. Amer. Math. Soc. Ser. B **1**, (2012), 1–22.
- [13] H. Lee and Y. Lim: *Carlson’s iterative mean algorithm of positive definite matrices*, Linear Algebra Appl. **439** (2013), 1183–1201.
- [14] B. Lemmens and R. Nussbaum: *Nonlinear Perron-Frobenius theory*. Cambridge Tracts in Mathematics **189**, Cambridge Univ. Press, Cambridge, 2012.
- [15] B. Lemmens, B. Lins, R. Nussbaum, and M. Wortel: *Denjoy-Wolff theorems for Hilbert’s and Thompson’s metric spaces*, J. Anal. Math., to appear.
- [16] B. Lemmens and M. Roelands: *Unique geodesics for Thompson’s metric*, Ann. Inst. Fourier (Grenoble) **65** (2015), 315–348.
- [17] Y. Lim: *Finsler metrics on symmetric cones*, Math. Ann. **316**, (2000), 379–389.
- [18] Y. Lim: *Geometry of midpoint sets for Thompson’s metric*, Linear Algebra Appl., **439** (2013), 211–227.
- [19] Y. Lim and M. Pálfi: *The matrix power means and the Karcher mean*, J. Funct. Anal. **262**, (2012), 1498–1514.
- [20] L. Molnár: *Thompson isometries of the space of invertible positive operators*, Proc. Amer. Math. Soc. **137** (2009), 3849–3859.
- [21] R.D. Nussbaum: *Finsler structures for the part metric and Hilbert’s projective metric and applications to ordinary differential equations*, Differential Integral Equations **7** (1994), 1649–1707.
- [22] R.D. Nussbaum: *Hilbert’s projective metric and iterated nonlinear maps*, Mem. Amer. Math. Soc. **391**, (1988), 1–137.
- [23] M. Pálfi and D. Petz: *Weighted multivariable operator means of positive definite operators*, Linear Algebra Appl. **463** (2014), 134–153.
- [24] A.C. Thompson: *On certain contraction mappings in a partially ordered vector space*, Proc. Amer. Math. Soc. **14** (1963), 438–443.

Bas Lemmens
 School of Mathematics
 Statistics & Actuarial Sciences
 University of Kent
 CT27NF, Canterbury
 United Kingdom
 e-mail: B.Lemmens@kent.ac.uk

Mark Roelands
 School of Mathematics
 Statistics & Actuarial Sciences
 University of Kent
 United Kingdom
 e-mail: mark.roelands@gmail.com