



Title	Disseminated metastatic tissue calcification after orthotopic liver transplantation: A case report
Author(s)	Deguchi, K.; Ueno, T.; Matsuura, R. et al.
Citation	Transplantation Proceedings. 2016, 48(1), p. 251–254
Version Type	AM
URL	<a href="https://hdl.handle.net/11094/100146">https://hdl.handle.net/11094/100146</a>
rights	© 2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Table 1. Laboratory data related to common causes of hypercalcemia on post-retransplantation day 24.

BAP ( 3.7-20.9 µg/L)	83.7 µg/L
TRACP-5b (170-590 mIU/dL)	>1500 mIU/dL
intact PTH (10-65 pg/ml)	<10.0 pg/ml
PTHrP (<1.1 pmol/L)	<1.1 pmol/L
1 alpha ,25(OH) <sub>2</sub> D <sub>3</sub> (20-60 pg/mL)	6 pg/ml
Corrected Calcium (8.6-10.2 mg/dL)	13.5 mg/dl
Phosphate (8.5-10.2 mg/dL)	3.7 mg/dl
Mgnesium (1.8-2.4 mg/dL)	2.4 mg/dL

BAP:Bone alkaline phosphatase , TRACP-5b: Tartrate-resistant acid phosphatase type 5b ,

PTH: Parathyroid hormone, PTHrP: Parathyroid hormone related protein

Normal range are shown in parentheses.