

Title	Littlewood-Paley inequality for a diffusion satisfying the logarithmic Sobolev inequality and for the Brownian motion on a Riemannian manifold with boundary
Author(s)	Shigekawa, Ichiro
Citation	Osaka Journal of Mathematics. 2002, 39(4), p. 897-930
Version Type	VoR
URL	<a href="https://doi.org/10.18910/10092">https://doi.org/10.18910/10092</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

# LITTLEWOOD-PALEY INEQUALITY FOR A DIFFUSION SATISFYING THE LOGARITHMIC SOBOLEV INEQUALITY AND FOR THE BROWNIAN MOTION ON A RIEMANNIAN MANIFOLD WITH BOUNDARY

ICHIRO SHIGEKAWA

(Received April 20, 2001)

## 1. Introduction

In this paper, we discuss the Littlewood-Paley inequality. Typical example is the Brownian motion on the Euclidean space and it leads to the following inequality: for any  $p > 1$  there exist a positive constant  $C$  such that

$$(1.1) \quad C^{-1} \|\nabla u\|_p \leq \|\sqrt{-\Delta}u\|_p \leq C \|\nabla u\|_p.$$

$\sqrt{-\Delta}$ , the square root of the minus Laplacian, is called the Cauchy operator. (1.1) is equivalent to the  $L^p$ -boundedness of the Riesz transformation.

This kind of inequality also holds for the Ornstein-Uhlenbeck process on an abstract Wiener space, which was proved by P.A. Meyer [11] in a probabilistic approach.

In this paper, we attempt to extend this inequality for a diffusion process associated with a Dirichlet form that admits a square field operator. There have been several related works, e.g., Bakry [3, 4], Shigekawa-Yoshida [16]. In these papers, they assumed that  $\Gamma_2$  is positive or bounded from below. We replace this boundedness assumption with the exponential integrability of negative part of  $\Gamma_2$ . To handle this case, we assume that the logarithmic Sobolev inequality holds. Moreover our square field operator is of the gradient form, i.e., the Dirichlet form  $\mathcal{E}$  is given as follows;

$$(1.2) \quad \mathcal{E}(u, v) = \int_M (\nabla u, \nabla v) \mu(dx).$$

We adopt a probabilistic approach which was developed by Meyer and Bakry. We will show the inequality for the Littlewood-Paley  $G$ -function. Since the square field operator is given as a gradient, we consider another semigroup that acts on vector valued functions and use the semigroup domination to estimate vector valued functions. Using this method, the estimate for vector valued functions can be reduced to the scalar case. But the unboundedness of  $\Gamma_2$  causes some troubles and so we could not prove the exact inequality. We only show that the  $L^p$ -norm is dominated by  $L^q$ -norm for  $1 < p < q$  (see the precise statement in §2).

We also discuss the Brownian motion on a Riemannian manifold with boundary. We impose the Neumann boundary condition on the Brownian motion. In this case, the quantity corresponding to  $\Gamma_2$  is singular (i.e., it is not a function but a smooth measure). We deal with it by way of an associated additive functional. The additive functional belongs to the Kato class and we can show the exact inequality (i.e., no loss of exponent).

The organization of the paper is as follows. We give a formulation and a main result in §2. We define  $\Gamma_2$  in our formulation. It is a generalization of Ricci curvature and is based on a square field operator for vector valued functions. In §3, the maximal ergodic inequality for a semigroup with a potential is given. Here the logarithmic Sobolev inequality is essential. We give a proof of the main theorem in §5. To do this, we prepare fundamental inequalities for the Littlewood-Paley  $G$ -function in §4. A proof for the Littlewood-Paley inequality is given in §5. Combining this with the intertwining property of semigroups, we can get the main result. The Brownian motion on a Riemannian manifold with boundary is dealt with in §6.

## 2. Symmetric diffusion

Let us introduce a diffusion process that we use in the paper. Let  $M$  be a topological space. We assume  $M$  to be Souslinian. Suppose we are given a Borel probability measure  $\mu$  on  $M$  and a Dirichlet form  $\mathcal{E}$  in  $L^2(\mu)$ . We assume that there exists a Hunt diffusion process  $(X_t, P_x)_{x \in M}$  associated with  $\mathcal{E}$ . We denote the generator and the semigroup by  $L$  and  $\{T_t\}$ , respectively. We assume that  $1 \in \text{Dom}(L)$  and  $L1 = 0$  where 1 denotes the function that is identically equal to 1. Hence the diffusion  $(X_t)$  is conservative. We also assume that the Dirichlet form satisfies the following defective logarithmic Sobolev inequality: there exist  $\alpha > 0$  and  $\beta \geq 0$  such that

$$(2.1) \quad \int_M u^2 \log \left( \frac{u}{\|u\|_2} \right) \mu(dx) \leq \alpha \mathcal{E}(u, u) + \beta(u, u).$$

Here  $(\cdot, \cdot)$  denotes the inner product in  $L^2$ .

Further we assume that the square field operator  $\Gamma$  is well-defined. Here  $\Gamma: \text{Dom}(\mathcal{E}) \times \text{Dom}(\mathcal{E}) \rightarrow L^1(\mu)$  is a continuous bilinear map which is characterized as follows:

$$(2.2) \quad 2(\Gamma(v, w), u) = \mathcal{E}(vw, u) - \mathcal{E}(v, wu) - \mathcal{E}(vu, w), \quad \forall u, v, w \in \mathcal{E} \cap L^\infty.$$

A crucial assumption is as follows; there exists a ‘gradient operator’  $\nabla$  such that  $\nabla$  is a closed operator from  $L^2(\mu)$  to  $L^2(\mu; K)$  and it satisfies  $\Gamma(u, v) = (\nabla u, \nabla v)$ . Here  $K$  is a (separable) Hilbert space.  $L^2(\mu; K)$  may be possibly the set of all square integrable section of a vector bundle over  $M$ . But we use  $L^2(\mu; K)$  for notational convention. We need another semigroup  $\{\hat{T}_t\}$  in  $L^2(\mu; K)$ . Let  $\{\hat{T}_t\}$  be a contraction symmetric semigroup associated with a bilinear form  $\hat{\mathcal{E}}$ . We also need a square field operator

for  $\{\hat{T}_t\}$  and so we assume that

(A.1) For  $\theta \in \text{Dom}(\hat{L})$ , it holds that  $|\theta|_K^2 \in \text{Dom}(L_1)$ .

Here  $L_1$  is the generator in  $L^1(\mu)$ . Under this condition, we define a square field operator  $\hat{\Gamma}$  as

$$(2.3) \quad 2\hat{\Gamma}(\theta, \eta) = L(\theta, \eta)_K - (\hat{L}\theta, \eta)_K - (\theta, \hat{L}\eta)_K.$$

We assume the following two properties: the positivity and the derivation property.

(A.2)  $\hat{\Gamma}(\theta, \theta) \geq 0$  for  $\theta \in \text{Dom}(\hat{\mathcal{E}})$ .

(A.3) For  $\theta, \eta \in \text{Dom}(\hat{\mathcal{E}}) \cap L^\infty$  and  $u \in \text{Dom}(\mathcal{E}) \cap L^\infty$ , it holds that

$$(2.4) \quad 2u\hat{\Gamma}(\theta, \eta) = -(\nabla u, \nabla(\theta, \eta)) + \hat{\Gamma}(\theta, u\eta) + \hat{\Gamma}(u\theta, \eta).$$

Then, by the semigroup domination theorem (see [14]), we have

$$(2.5) \quad |\hat{T}_t\theta| \leq T_t|\theta|.$$

Let  $S_b(K)$  be the space of all self-adjoint operator on  $K$  that is bounded from below. Let  $R$  be a function on  $M$  taking values in  $S_b(K)$ . Define a bilinear form  $\hat{\mathcal{E}}^R$  by

$$(2.6) \quad \hat{\mathcal{E}}^R(\theta, \eta) = \hat{\mathcal{E}}(\theta, \eta) + \int_M (R(x)\theta(x), \eta(x))_K \mu(dx).$$

The associated semigroup will be denoted by  $\hat{T}_t^R$ . We assume the following intertwining property, which is crucial in the paper.

$$(2.7) \quad \nabla T_t u = \hat{T}_t^R \nabla u, \quad \text{for } u \in \text{Dom}(\nabla).$$

$R$  plays the role of so called  $\Gamma_2$ .

We take a scalar function  $V$  such that

$$(2.8) \quad (R(x)k, k)_K \geq V(x)(k, k)_K.$$

The semigroup generated by  $L - V$  is denoted by  $\{T_t^V\}$ . The generator of  $\hat{T}_t^R$  is  $\hat{L} - R$ . Again by the domination theorem, it holds that

$$(2.9) \quad |\hat{T}_t^R \theta| \leq T_t^V |\theta|.$$

$V$  can be decomposed as  $V = V_+ - V_-$  where  $V_+ = V \vee 0$  and  $V_- = (-V) \vee 0$ . The last assumption is that

(A.4)  $e^{V_-} \in L^{\infty-} = \bigcap_{p \geq 1} L^p$ .

For scalar functions, we can define two kinds of norms:  $\|\nabla u\|_p$  and  $\|\sqrt{1 - Lu}\|_p$ . It is a fundamental question whether these norms are equivalent or not. For example,

if the generator  $L$  is the Ornstein-Uhlenbeck operator on an abstract Wiener space, then the equivalence of two norms are known as the Meyer equivalence.

Under our conditions, we can get the following result.

**Theorem 2.1.** *For any  $1 < p < q < \infty$ , we have*

$$(2.10) \quad \|\nabla u\|_p \lesssim \|\sqrt{1 - Lu}\|_q,$$

$$(2.11) \quad \|\sqrt{1 - Lu}\|_p \lesssim \|\nabla u\|_q + \|u\|_q.$$

In the above theorem, the notation  $A \lesssim B$  stands for  $A \leq kB$  for a positive constant  $k$ . Further, in (2.10) for example, the constant depends only on  $p$  but is independent of  $u$ . We use this convention in the sequel without mentioning.

To prove the theorem, we use the Littlewood-Paley  $G$ -function. We introduce it in §4 and give a proof of the theorem in §5.

### 3. Maximal ergodic inequality

In this section, we discuss the maximal ergodic inequality. This inequality is known for a symmetric Markov semigroup (see e.g., Stein [17]). Here we consider a semigroup with a potential. To show the inequality, we adopt a probabilistic method due to Rota [13].

We consider an additive functional  $A_t$  associated to a smooth signed measure  $\rho$  under the Revuz correspondence. We define a Dirichlet form by

$$(3.1) \quad \mathcal{E}^\rho(u, v) = \mathcal{E}(u, v) + \int_M \tilde{u}\tilde{v}\rho(dx)$$

where  $\tilde{u}$  denotes the quasi-continuous modification of  $u$ . The associated semigroup is denoted by  $\{T_t^\rho\}$ , which is expressed as

$$(3.2) \quad T_t^\rho u(x) = E_x[u(X_t)e^{-A_t}]$$

where  $E_x$  denote the expectation under the measure  $P_x$ .

**Theorem 3.1.** *Assume that for any  $q \geq 1$ , there exist constants  $c_q, \beta_q$  such that*

$$(3.3) \quad E_x[e^{-qA_t}]^{1/q} \leq c_q e^{\beta_q t} \quad \forall t \geq 0, \text{ q.e.-}x.$$

Here “q.e.” means that it holds except for a set of capacity 0. Then for any  $p > 1$  there exist constants  $\lambda, c$  such that

$$(3.4) \quad \left\| \sup_{t \geq 0} |e^{-\lambda t} T_t^\rho u| \right\|_p \leq c \|u\|_p, \quad \forall u \in L^p.$$

In particular, if  $\rho$  is non-negative (i.e.,  $A_t$  is non-negative), we can take  $\lambda = 0$ .

Proof. We note that  $|T_t^\rho u| \leq T_t^{-\rho_-} |u|$ , where  $\rho = \rho_+ - \rho_-$  is the Hahn decomposition of  $\rho$ . Without loss of generality, we may assume that  $\rho$  is non-positive.

Set

$$(3.5) \quad M_t = T_{T-t}^\rho u(X_t) e^{-A_t}.$$

Here  $\theta_t$  is the shift operator. We show first that  $\{M_t\}$  is a martingale under  $P_\mu := \int_M P_x \mu(dx)$ . In fact,

$$\begin{aligned} E_\mu[u(X_T) e^{-A_T} \mid \mathcal{F}_t] &= E_\mu[u(X_{T-t} \circ \theta_t) e^{-A_{T-t} \circ \theta_t - A_t} \mid \mathcal{F}_t] \\ &= e^{-A_t} E_\mu[u(X_{T-t} \circ \theta_t) e^{-A_{T-t} \circ \theta_t} \mid \mathcal{F}_t] \\ &= e^{-A_t} E_{X_t}[u(X_{T-t}) e^{-A_{T-t}}] \quad (\text{Markov property}) \\ &= e^{-A_t} T_{T-t}^\rho u(X_t). \end{aligned}$$

We note, by the Markov property,

$$\begin{aligned} T_{T-t}^\rho u(X_T) &= E_{X_T}[u(X_{T-t}) e^{-A_{T-t}}] \\ &= E_\mu[u(X_{T-t} \circ \theta_T) e^{-A_{T-t} \circ \theta_T} \mid X_T] \\ &= E_\mu[u(X_{2T-t}) e^{-A_{2T-t} + A_T} \mid X_T]. \end{aligned}$$

Now, using the reversibility of  $(X_t)$ , i.e.,  $(X_{2T-t})_{0 \leq t \leq 2T}$  has the same law as  $(X_t)_{0 \leq t \leq 2T}$ , we have

$$T_{T-t}^\rho u(X_T) = E_\mu[u(X_t) e^{-A_T + A_t} \mid X_T].$$

Hence

$$\begin{aligned} T_{2(T-t)}^\rho u(X_T) &= T_{T-t}^\rho T_{T-t}^\rho u(X_T) \\ &= E_\mu[T_{T-t}^\rho u(X_t) e^{-A_T + A_t} \mid X_T] \\ &= E_\mu[M_t e^{-A_T + 2A_t} \mid X_T]. \end{aligned}$$

Noting that we have taken  $A$  to be non-positive, we have

$$\begin{aligned} \sup_{0 \leq t \leq T} |T_{2(T-t)}^\rho u(X_T)| &\leq E_\mu \left[ \sup_{0 \leq t \leq T} |M_t| e^{-A_T} \mid X_T \right] \\ &\leq E_\mu \left[ \sup_{0 \leq t \leq T} |M_t|^p \mid X_T \right]^{1/p} E_\mu [e^{-qA_T} \mid X_T]^{1/q}. \quad \left( \frac{1}{p} + \frac{1}{q} = 1 \right) \end{aligned}$$

On the other hand,

$$(3.6) \quad E_\mu [e^{-qA_T} \mid X_T]^{1/q} = E_\mu [e^{-q(A_{2T} - A_T)} \mid X_T]^{1/q} \quad (\text{reversibility})$$

$$\begin{aligned}
 &= E_\mu [e^{-qA_T \circ \theta_T} \mid X_T]^{1/q} \quad (\text{additivity}) \\
 &= E_{X_T} [e^{-qA_T}]^{1/q} \quad (\text{Markovian property}) \\
 &\leq c_q e^{\beta_q T}. \quad (\because (3.3))
 \end{aligned}$$

Thus we have

$$\sup_{0 \leq t \leq T} |T_{2(T-t)}^\rho u(X_T)| \leq c_q e^{\beta_q T} E_\mu \left[ \sup_{0 \leq t \leq T} |M_t|^p \mid X_T \right]^{1/p}$$

Hence, by the Doob inequality

$$\begin{aligned}
 &\left\| \sup_{0 \leq t \leq T} |T_{2(T-t)}^\rho u| \right\|_p \\
 &\leq c_q e^{\beta_q T} E_\mu \left[ E_\mu \left[ \sup_{0 \leq t \leq T} |M_t|^p \mid X_T \right] \right]^{1/p} \\
 &= c_q e^{\beta_q T} E_\mu \left[ \sup_{0 \leq t \leq T} |M_t|^p \right]^{1/p} \\
 &\leq C' c_q e^{\beta_q T} E_\mu [|M_T|^p]^{1/p} \quad (\text{Doob's inequality}) \\
 &= C' c_q e^{\beta_q T} E_\mu [|u(X_T)|^p e^{-pA_T}]^{1/p} \\
 &= C' c_q e^{\beta_q T} E_\mu [E_\mu [|u(X_T)|^p e^{-pA_T} \mid X_T]]^{1/p} \\
 &= C' c_q e^{\beta_q T} E_\mu [|u(X_T)|^p E_\mu [e^{-pA_T} \mid X_T]]^{1/p} \\
 &\leq C' c_q e^{\beta_q T} c_p e^{\beta_p T} E_\mu [|u(X_T)|^p]^{1/p} \quad (\because (3.6)) \\
 &= C' c_p c_q e^{(\beta_p + \beta_q)T} \|u\|_p.
 \end{aligned}$$

Thus we can find constants  $k > 0$  and  $C > 0$  which are independent of  $T$  and  $u$  such that

$$\left\| \sup_{0 \leq t \leq 2T} |T_t^\rho u| \right\|_p \leq C e^{2kT} \|u\|_p$$

We take  $\lambda > k$ . Note that for any integer  $n$ ,

$$\begin{aligned}
 \left\| \sup_{n \leq t \leq n+1} e^{-\lambda t} |T_t^\rho u| \right\|_p &\leq e^{-\lambda n} \left\| \sup_{0 \leq t \leq n+1} |T_t^\rho u| \right\|_p \\
 &\leq C e^{-\lambda n} e^{k(n+1)} \|u\|_p.
 \end{aligned}$$

Summing up in  $n$ ,

$$\sum_{n=0}^{\infty} \left\| \sup_{n \leq t \leq n+1} e^{-\lambda t} |T_t^\rho u| \right\|_p \leq C e^k \sum_{n=0}^{\infty} e^{-(\lambda-k)n} \|u\|_p$$

$$\leq \frac{C}{e^{-k} - e^{-\lambda}} \|u\|_p.$$

Clearly this leads us to

$$\begin{aligned} \left\| \sup_{0 \leq t < \infty} e^{-\lambda t} |T_t^\rho u| \right\|_p &\leq \left\| \sum_{n=0}^{\infty} \sup_{n \leq t \leq n+1} e^{-\lambda t} |T_t^\rho u| \right\|_p \\ &\leq \sum_{n=0}^{\infty} \left\| \sup_{n \leq t \leq n+1} e^{-\lambda t} |T_t^\rho u| \right\|_p \\ &\leq \frac{C}{e^{-k} - e^{-\lambda}} \|u\|_p. \end{aligned}$$

This completes the proof. □

The assumption (3.3) is rather strong. We replace it with the assumption (A.4). In this case, we set  $\rho = Vm$ . Hence, the associated additive functional is given by

$$(3.7) \quad A_t = \int_0^t V(X_s) ds.$$

Here we denote the semigroup  $\{T_t^\rho\}$  by  $\{T_t^V\}$ . Since  $\mathcal{E}$  satisfies the logarithmic Sobolev inequality (2.1), we have (see, e.g., [14]),

$$(3.8) \quad \|T_t^V u\|_p \leq \|e^{-V}\|_{\alpha p^2/4(p-1)}^t e^{4\beta t/\alpha} \|u\|_p.$$

This means that there exists a constant  $\gamma_p$  such that

$$(3.9) \quad \|T_t^V\|_{p \rightarrow p} \leq e^{\gamma_p t}.$$

E.g., set  $\gamma_p = (4\beta/\alpha) \log \|e^{-V}\|_{\alpha p^2/4(p-1)}$ .

In particular, when  $p = 2$ ,

$$(3.10) \quad \|T_t^V u\|_2 \leq \|e^{-V}\|_{\alpha}^t e^{4\beta t/\alpha} \|u\|_2$$

In this case, taking  $u = 1$ , we have

$$\begin{aligned} E_\mu[e^{-A_t}] &= \int_X E_x[1e^{-A_t}] \mu(dx) \\ &= \|T_t^V 1\|_1 \\ &\leq \|T_t^V 1\|_2 \\ &\leq \|e^{-V}\|_{\alpha}^t e^{4\beta t/\alpha} \|1\|_2 \\ &\leq \|e^{-V}\|_{\alpha}^t e^{4\beta t/\alpha}. \end{aligned}$$



Hence, for any  $\gamma > 0$ , it holds that

$$(3.11) \quad E_\mu[e^{-\gamma A_T}] \leq \|e^{-\gamma V}\|_\alpha^t e^{4\beta t/\alpha}$$

Noticing this inequality, we can get the following maximal ergodic inequality.

**Theorem 3.2.** *Take any  $1 < p < r < \infty$ . If we take  $\lambda > 0$  to be sufficiently large, then there exists a constant  $c > 0$  such that*

$$(3.12) \quad \left\| \sup_{t \geq 0} |e^{-\lambda t} T_t^V u| \right\|_p \leq c \|u\|_r.$$

Proof. By the same proof as in Theorem 3.1, we have

$$\sup_{0 \leq t \leq T} |T_{2(T-t)}^V u(X_T)| \leq E_\mu \left[ \sup_{0 \leq t \leq T} |M_t| e^{-A_T} \mid X_T \right].$$

Hence,

$$\begin{aligned} & \left\| \sup_{0 \leq t \leq T} |T_{2(T-t)}^V u| \right\|_p \\ & \leq E_\mu \left[ E_\mu \left[ \sup_{0 \leq t \leq T} |M_t| e^{-A_T} \mid X_T \right]^p \right]^{1/p} \\ & \leq E_\mu \left[ \sup_{0 \leq t \leq T} |M_t|^p e^{-pA_T} \right]^{1/p} \\ & \leq E_\mu \left[ \sup_{0 \leq t \leq T} |M_t|^{pq/p} \right]^{(p/q) \cdot (1/p)} E_\mu [e^{-puA_T}]^{(q-p)/pq} \left( \frac{1}{q/p} + \frac{1}{u} = 1 \right) \\ & \leq C E_\mu [|M_T|^q]^{1/q} E_\mu [e^{-puA_T}]^{(q-p)/pq} \quad (\text{Doob's inequality}) \\ & = C E_\mu [|u(X_T) e^{-A_T}|^q]^{1/q} E_\mu [e^{-puA_T}]^{(q-p)/pq} \\ & \leq C E_\mu [|u(X_T)|^{rq/q}]^{(q/r) \cdot (1/q)} E_\mu [e^{-qvA_T}]^{(r-q)/rq} E_\mu [e^{-puA_T}]^{(q-p)/pq} \\ & \quad \left( \frac{1}{r/q} + \frac{1}{v} = 1 \right) \\ & \leq C \|u\|_r \|e^{-qvV}\|_\alpha^{(r-q)T/rq} e^{4\beta(r-q)T/\alpha r q} \|e^{-puV}\|_\alpha^{(q-p)T/pq} e^{4\beta(q-p)T/\alpha pq}. \end{aligned}$$

Thus we can find a constant  $k > 0$  which is independent of  $T$  and  $u$  such that

$$\left\| \sup_{0 \leq t \leq 2T} |T_t^V u| \right\|_p \leq C e^{2kT} \|u\|_r.$$

The rest is the same as Theorem 3.1. This completes the proof. □

**4. Littlewood-Paley  $G$ -functions**

Let us introduce the Littlewood-Paley  $G$ -functions. To do this, we recall the subordination of a semigroup. Set  $T_t^\lambda = e^{-\lambda t} T_t$  ( $\lambda \geq 0$ ). We take  $\lambda$  to be large enough. For any  $t \geq 0$ , define a measure  $\mu_t$  on  $[0, \infty)$  by

$$(4.1) \quad \mu_t(ds) = \frac{t}{2\sqrt{\pi}} e^{-t^2/4s} s^{-3/2} ds.$$

In terms of the Laplace transform, this measure is characterized as

$$\int_0^\infty e^{-\alpha s} \mu_t(ds) = e^{-\sqrt{\alpha}t} \quad \text{for } \alpha > 0.$$

Then the subordination  $\{Q_t^\lambda\}$  of  $\{T_t^\lambda\}$  is defined by

$$(4.2) \quad Q_t^\lambda = \int_0^\infty T_s^\lambda \mu_t(ds).$$

The generator of  $\{Q_t^\lambda\}$  in  $L^2(\mu)$  is  $-\sqrt{\lambda - L}$ .

We recall that  $\{T_t^V\}$  is the semigroup with the potential  $V$ . We set  $T_t^{\lambda+V} = e^{-\lambda t} T_t^V$  and we also define the subordination of  $\{T_t^V\}$  as

$$(4.3) \quad Q_t^{\lambda+V} = \int_0^\infty T_s^{\lambda+V} \mu_t(ds).$$

The operator norm of  $\{Q_t^{\lambda+V}\}$  in  $L^p$  is estimated as

$$\begin{aligned} \|Q_t^{\lambda+V}\|_{p \rightarrow p} &\leq \int_0^\infty \|T_s^{\lambda+V}\|_{p \rightarrow p} \mu_t(ds) \\ &\leq \int_0^\infty e^{-\lambda s + \gamma_p s} \mu_t(ds) \\ &= e^{-\sqrt{\lambda - \gamma_p}t}. \end{aligned}$$

Here  $\gamma_p$  is the constant in (3.9). Moreover, by the semigroup domination  $|\hat{T}_t^{\lambda+R}\theta| \leq T_t^{\lambda+V}|\theta|$ , we have

$$(4.4) \quad \|\hat{T}_t^{\lambda+R}\|_{p \rightarrow p} \leq \|T_t^{\lambda+V}\|_{p \rightarrow p} \leq e^{-(\lambda - \gamma_p)t}.$$

Similarly we have

$$(4.5) \quad \|\hat{Q}_t^{\lambda+R}\|_{p \rightarrow p} \leq e^{-\sqrt{\lambda - \gamma_p}t}.$$

For any real valued function  $u$ , define

$$(4.6) \quad g^-(x, t) = |\partial_t Q_t^\lambda u(x)|^2,$$

$$(4.7) \quad g^\uparrow(x, t) = |\nabla Q_t^\lambda u(x)|_K^2,$$

$$(4.8) \quad g(x, t) = g^\rightarrow(x, t) + g^\uparrow(x, t).$$

Here  $\partial_t = \partial/\partial t$ . Then, the Littlewood-Paley  $G$ -function is defined by

$$(4.9) \quad G^\rightarrow u(x) = \left\{ \int_0^\infty t g^\rightarrow(x, t) dt \right\}^{1/2},$$

$$(4.10) \quad G^\uparrow u(x) = \left\{ \int_0^\infty t g^\uparrow(x, t) dt \right\}^{1/2},$$

$$(4.11) \quad Gu(x) = \left\{ \int_0^\infty t g(x, t) dt \right\}^{1/2}.$$

Moreover, we define the  $H$ -functions by

$$(4.12) \quad H^\rightarrow u(x) = \left\{ \int_0^\infty t Q_t g^\rightarrow(x, t) dt \right\}^{1/2},$$

$$(4.13) \quad H^\uparrow u(x) = \left\{ \int_0^\infty t Q_t g^\uparrow(x, t) dt \right\}^{1/2},$$

$$(4.14) \quad Hu(x) = \left\{ \int_0^\infty t Q_t g(x, t) dt \right\}^{1/2}.$$

For vector valued function  $\theta$ , we define  $G$ -function and  $H$ -function, similarly. That is, e.g.,

$$(4.15) \quad \hat{g}^\rightarrow(x, t) = |\partial_t \hat{Q}_t^{\lambda+R} \theta(x)|^2,$$

$$(4.16) \quad \hat{G}^\rightarrow \theta(x) = \left\{ \int_0^\infty t \hat{g}^\rightarrow(x, t) dt \right\}^{1/2}$$

$$(4.17) \quad \hat{H}^\rightarrow \theta(x) = \left\{ \int_0^\infty t Q_t \hat{g}^\rightarrow(x, t) dt \right\}^{1/2}.$$

Notice that, in this case, we use the semigroup  $\{\hat{Q}_t^{\lambda+R}\}$  that is the subordination of  $\{\hat{T}_t^{\lambda+R}\}$ .  $\hat{G}^\uparrow \theta$ ,  $\hat{H}^\uparrow \theta$ ,  $\hat{G}\theta$ , and  $\hat{H}\theta$  are defined similarly. For example,

$$\hat{g}^\uparrow(x, t) = \hat{\Gamma}(\hat{Q}_t^{\lambda+R} \theta, \hat{Q}_t^{\lambda+R} \theta)(x)$$

(see (2.3) for the definition of  $\hat{\Gamma}$ ).

The following proposition is easily obtained by the spectral decomposition:

**Proposition 4.1.** *It holds that*

$$(4.18) \quad \|G^\rightarrow u\|_2 = \frac{1}{2} \|u\|_2,$$

and

$$(4.19) \quad \|\hat{G} \rightarrow \theta\|_2 = \frac{1}{2} \|\theta\|_2.$$

Later we need the interrelationship between  $G$  and  $H$  functions and so we first prepare the following.

**Lemma 4.2.** *We have the following estimate:*

$$(4.20) \quad |T_t^{\lambda+V} u(x)|^2 \leq \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} T_t |u|^2(x),$$

$$(4.21) \quad |Q_t^{\lambda+V} u(x)|^2 \leq \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} Q_t |u|^2(x).$$

*Proof.* By the Feynman-Kac formula, we have

$$\begin{aligned} |T_t^{\lambda+V} u(x)|^2 &= \left| E_x \left[ \exp \left\{ -\lambda t - \int_0^t V(X_s) ds \right\} u(X_t) \right] \right|^2 \\ &\leq E_x \left[ \exp \left\{ -2\lambda t - 2 \int_0^t V(X_s) ds \right\} \right] E_x [ |u(X_t)|^2 ] \\ &= T_t^{2(\lambda+V)} 1(x) \cdot T_t |u|^2(x) \\ &\leq \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} \cdot T_t |u|^2(x). \end{aligned}$$

Further we have,

$$\begin{aligned} |Q^{\lambda+V} u(x)|^2 &= \left| \int_0^\infty T_s^{\lambda+V} u(x) \lambda_t(ds) \right|^2 \\ &\leq \int_0^\infty |T_s^{\lambda+V} u(x)|^2 \lambda_t(ds) \\ &\leq \int_0^\infty \left\{ \sup_{r \geq 0} T_r^{2(\lambda+V)} 1(x) \right\} \cdot T_s |u|^2(x) \lambda_t(ds) \\ &= \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} \cdot Q_t |u|^2(x). \end{aligned}$$

This completes the proof. □

Now we can show the following estimate between  $G$ -functions and  $H$ -functions.

**Proposition 4.3.** *We have that*

$$(4.22) \quad \hat{G}^{-}\theta \leq 2 \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\}^{1/2} \cdot \hat{H}^{-}\theta.$$

For scalar function, we have

$$(4.23) \quad G^{-}u \leq 2H^{-}u,$$

$$(4.24) \quad G^{+}u \leq 2 \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\}^{1/2} \cdot H^{+}u.$$

Proof. We have,

$$|\hat{Q}_t^{\lambda+R}\theta(x)| \leq \int_0^\infty |\hat{T}_s^{\lambda+R}\theta(x)|\mu_t(ds) \leq \int_0^\infty T_s^{\lambda+V}|\theta|(x)\mu_t(ds) = Q_t^{\lambda+V}|\theta|(x).$$

Using Lemma 4.2, we have

$$|\hat{Q}_t^{\lambda+R}\theta(x)|^2 \leq \{Q_t^{\lambda+V}|\theta|(x)\}^2 \leq \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} \cdot Q_t|\theta|^2(x).$$

Therefore

$$\begin{aligned} \hat{g}^{-}(x, 2t) &= |\partial_s \hat{Q}_s^{\lambda+R}\theta(x)|^2 \Big|_{s=2t} \\ &= \left| \sqrt{\lambda - \hat{L} + R} \hat{Q}_{2t}^{\lambda+R}\theta(x) \right|^2 \\ &= \left| \hat{Q}_t^{\lambda+R} \sqrt{\lambda - \hat{L} + R} \hat{Q}_t^{\lambda+R}\theta(x) \right|^2 \\ &\leq \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} Q_t \left| \sqrt{\lambda - \hat{L} + R} \hat{Q}_t^{\lambda+R}\theta \right|^2(x) \\ &= \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} Q_t \hat{g}^{-}(x, t). \end{aligned}$$

From this,

$$\begin{aligned} \hat{G}^{-}\theta(x) &= \left\{ \int_0^\infty t \hat{g}^{-}(x, t) dt \right\}^{1/2} = \left\{ 4 \int_0^\infty t \hat{g}^{-}(x, 2t) dt \right\}^{1/2} \\ &\leq 2 \left\{ \int_0^\infty t \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} Q_t \hat{g}^{-}(x, t) dt \right\}^{1/2} \\ &\leq 2 \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\}^{1/2} \left\{ \int_0^\infty t Q_t \hat{g}^{-}(x, t) dt \right\}^{1/2} \end{aligned}$$

$$= 2 \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\}^{1/2} \hat{H}^{-\rightarrow} \theta(x).$$

For the scalar function, it holds that  $G^{-\rightarrow} u \leq 2H^{-\rightarrow} u$  since we have  $|Q_t u(x)|^2 \leq Q_t |u|^2(x)$ .

Let us next estimate  $G^\uparrow u$ .

$$\begin{aligned} G^\uparrow u(x) &= \left\{ \int_0^\infty t |\nabla Q_t^\lambda u(x)|^2 dt \right\}^{1/2} \\ &= \left\{ 4 \int_0^\infty t |\nabla Q_{2t}^\lambda u(x)|^2 dt \right\}^{1/2} \\ &= 2 \left\{ \int_0^\infty t |\hat{Q}_t^{\lambda+R} \nabla Q_t^\lambda u(x)|^2 dt \right\}^{1/2} \\ &\leq 2 \left\{ \int_0^\infty t \{ Q_t^{\lambda+V} |\nabla Q_t^\lambda u(x)| \}^2 dt \right\}^{1/2} \\ &\leq 2 \left\{ \int_0^\infty t \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\} Q_t |\nabla Q_t^\lambda u|^2(x) dt \right\}^{1/2} \\ &= 2 \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\}^{1/2} \left\{ \int_0^\infty t Q_t g^\uparrow(x, t) dt \right\}^{1/2} \\ &= 2 \left\{ \sup_{s \geq 0} T_s^{2(\lambda+V)} 1(x) \right\}^{1/2} H^\uparrow u(x). \end{aligned}$$

Thus we have (4.24). This completes the proof. □

In the next section, we use the diffusion process generated by  $L + \partial_a^2$ . So we will do some calculation on  $L + \partial_a^2$ .

**Lemma 4.4.** *For any  $\theta$ , set  $\hat{f}(x, a) = |\hat{Q}_a^{\lambda+R} \theta(x)|$  and for  $\varepsilon > 0$ ,  $\hat{f}_\varepsilon(x, a) = \sqrt{\hat{f}(x, a)^2 + \varepsilon}$ . Then we have*

$$(4.25) \quad (L + \partial_a^2) \hat{f}^2 \geq 2(\lambda + V) \hat{f}^2 + 2\hat{g}.$$

and for  $1 < p \leq 2$ , it holds that

$$(4.26) \quad (L + \partial_a^2) \hat{f}_\varepsilon^p \geq p(\lambda + V) \hat{f}_\varepsilon^2 \hat{f}_\varepsilon^{p-2} + p(p - 1) \hat{f}_\varepsilon^{p-2} \hat{g}$$

where  $\hat{g} = \hat{g}(x, a)$  was defined by

$$\hat{g}(x, a) = |\partial_a \hat{Q}_a^{\lambda+R} \theta(x)|^2 + \hat{\Gamma}(\hat{Q}_a^{\lambda+R} \theta, \hat{Q}_a^{\lambda+R} \theta)(x).$$

For the scalar case, we define  $f(x, a) = |Q_a^\lambda u(x)|$ ,  $f_\varepsilon(x, a) = \sqrt{f(x, a)^2 + \varepsilon}$ . Then

we have

$$(4.27) \quad (L + \partial_a^2)f^2 \geq 2\lambda f^2 + 2g$$

and for  $1 < p \leq 2$ ,

$$(4.28) \quad (L + \partial_a^2)f_\varepsilon^p \geq p\lambda f^2 f_\varepsilon^{p-2} + p(p-1)f_\varepsilon^{p-2}g.$$

Proof. We first show (4.25). To show this, we note that  $(\hat{L} - \lambda - R + \partial_a^2) \times \hat{Q}_a^{\lambda+R}\theta(x) = 0$ . Moreover, using the identity  $2\hat{\Gamma}(\theta, \theta) = L|\theta|^2 - 2(\hat{L}\theta, \theta)$ , it holds that

$$L|\hat{Q}_a^{\lambda+R}\theta|^2 = 2(\hat{L}\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta) + 2\hat{\Gamma}(\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta).$$

Hence

$$\begin{aligned} (L + \partial_a^2)\hat{f}^2 &= (L + \partial_a^2)|\hat{Q}_a^{\lambda+R}\theta|^2 \\ &= 2(\partial_a^2\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta) + 2(\partial_a\hat{Q}_a^{\lambda+R}\theta, \partial_a\hat{Q}_a^{\lambda+R}\theta) \\ &\quad + 2(\hat{L}\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta) + 2\hat{\Gamma}(\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta) \\ &= -2((\hat{L} - \lambda - R)\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta) + 2|\partial_a\hat{Q}_a^{\lambda+R}\theta|^2 \\ &\quad + 2(\hat{L}\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta) + 2\hat{\Gamma}(\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta) \\ &\geq 2(\lambda + V)|\hat{Q}_a^{\lambda+R}\theta|^2 + 2\hat{g}(x, a). \end{aligned}$$

Secondly we show (4.26). To show this we recall the following fundamental relationship between  $L$  and  $\nabla$ : for  $F(\xi^1, \xi^2, \dots, \xi^n) \in C^\infty(\mathbb{R}^n)$  and  $f^1, f^2, \dots, f^n \in \text{Dom}(L)$ ,

$$LF(f^1, f^2, \dots, f^n) = \sum_{i=1}^n \frac{\partial F}{\partial \xi^i} Lf^i + \sum_{i,j=1}^n \frac{\partial^2 F}{\partial \xi^i \partial \xi^j} (\nabla f^i, \nabla f^j)$$

(see [5, Lemma 1]). Hence we have, for  $1 < p \leq 2$ ,

$$\begin{aligned} (L + \partial_a^2)\hat{f}_\varepsilon^p &= (L + \partial_a^2)(\hat{f}_\varepsilon^2)^{p/2} \\ &= \frac{p}{2}(\hat{f}_\varepsilon^2)^{p/2-1}(L + \partial_a^2)\hat{f}_\varepsilon^2 \\ &\quad + \frac{p}{2}\left(\frac{p}{2} - 1\right)(\hat{f}_\varepsilon^2)^{p/2-2}\{(\partial_a\hat{f}_\varepsilon^2)^2 + |\nabla\hat{f}_\varepsilon^2|^2\} \\ &= \frac{p}{2}\hat{f}_\varepsilon^{p-2}(L + \partial_a^2)\hat{f}_\varepsilon^2 + \frac{p}{4}(p-2)\hat{f}_\varepsilon^{p-4}\{(\partial_a\hat{f}_\varepsilon^2)^2 + |\nabla\hat{f}_\varepsilon^2|^2\}. \end{aligned}$$

Let us recall that (see, e.g., [14, (3.11)])

$$|\nabla\hat{f}_\varepsilon^2|^2 = |\nabla(\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta)|^2 \leq 4\hat{\Gamma}(\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta)|\hat{Q}_a^{\lambda+R}\theta|^2.$$

Taking this into account, we have

$$\begin{aligned}
 (L + \partial_a^2)\hat{f}_\varepsilon^p &\geq \frac{p}{2}\hat{f}_\varepsilon^{p-2}\{2(\lambda + V)|\hat{Q}_a^{\lambda+R}\theta|^2 + 2\hat{g}\} \\
 &\quad + \frac{p}{4}(p - 2)\hat{f}_\varepsilon^{p-4}\{4(\partial_a\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta)^2 + 4\hat{\Gamma}(\hat{Q}_a^{\lambda+R}\theta, \hat{Q}_a^{\lambda+R}\theta)|\hat{Q}_a^{\lambda+R}\theta|^2\} \\
 &\geq \frac{p}{2}\hat{f}_\varepsilon^{p-2}\{2(\lambda + V)|\hat{Q}_a^{\lambda+R}\theta|^2 + 2\hat{g}\} + p(p - 2)\hat{f}_\varepsilon^{p-4}\hat{f}^2\hat{g} \\
 &\geq p\hat{f}_\varepsilon^{p-2}(\lambda + V)|\hat{Q}_a^{\lambda+R}\theta|^2 + p\hat{f}_\varepsilon^{p-2}\hat{g} + p(p - 2)\hat{f}_\varepsilon^{p-2}\hat{g} \\
 &\geq p(\lambda + V)\hat{f}_\varepsilon^{p-2}\hat{f}^2 + p(p - 1)\hat{f}_\varepsilon^{p-2}\hat{g}.
 \end{aligned}$$

The scalar case can be proved similarly. This completes the proof. □

### 5. Equivalence of $L^p$ -norms

In this section, we give estimates of  $G$  and  $H$  functions by a probabilistic method and then show the domination of norms. The original idea is due to P.A. Meyer [9] but we mainly follow Bakry [4].

Let  $(X_t, P_x)$  be the diffusion process on  $M$  associated with  $\mathcal{E}$  as before. We need an additional 1-dimensional Brownian motion  $(B_t)_{t \geq 0}$  and we regard  $M$  as a vertical space. We write  $P_x^\uparrow$  in place of  $P_x$ . Let  $(B_t, P_a^\rightarrow)$  be a 1-dimensional Brownian motion starting at  $a \in \mathbb{R}$  with the generator  $d^2/da^2$ . Note that this Brownian motion is different from the standard one up to constant. Let  $\tau$  be the hitting time of  $(B_t)$  to 0, i.e.,

$$\tau = \inf\{t \geq 0; B_t = 0\}.$$

We consider the following diffusion  $(Y_t, \mathbf{P}_{(x,a)})$  on the state space  $M \times \mathbb{R}$ ;

$$(5.1) \quad Y_t := (X_t, B_t), \quad \mathbf{P}_{(x,a)} := P_x^\uparrow \otimes P_a^\rightarrow.$$

So the generator of  $(Y_t)$  is  $L + \partial_a^2$ . We denote the integration with respect to  $\mathbf{P}_{(x,a)}$  and  $\int_M \mathbf{P}_{(x,a)}\mu(dx)$  by  $\mathbf{E}_{(x,a)}$  and  $\mathbf{E}_{\mu \times \delta_a}$ , respectively.

We use the following identities (see Meyer [9] for the proof): Let  $\eta: M \times \mathbb{R}_+ \rightarrow [0, \infty)$  be measurable. Then, for  $a > 0$ ,

$$(5.2) \quad \mathbf{E}_{\mu \times \delta_a} \left[ \int_0^\tau \eta(X_t, B_t) dt \right] = \int_M \mu(dx) \int_0^\infty (a \wedge t)\eta(x, t) dt$$

and

$$(5.3) \quad \mathbf{E}_{\mu \times \delta_a} \left[ \int_0^\tau \eta(X_t, B_t) dt \mid X_\tau = x \right] = \int_0^\infty (a \wedge t)Q_t\eta(x, t) dt.$$

We need an inequality for submartingales. Let  $(Z_t)$  be a *non-negative* continuous



submartingale with the following Doob-Meyer decomposition;

$$Z_t = M_t + A_t$$

where  $(M_t)$  is a continuous martingale and  $(A_t)$  is a continuous increasing process with  $A_0 = 0$ . Then, for  $p \geq 1$ , it holds that

$$(5.4) \quad E[A_\infty^p] \leq C_p E[Z_\infty^p].$$

For the proof, see Lenglart-Lépingle-Pratelli [8].

Before going to estimate  $G$ -function we prepare the following;

**Proposition 5.1.** *For any  $p \geq 1$ , we have*

$$(5.5) \quad \sup_{\substack{\alpha \geq 0 \\ N \geq 0}} \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau \alpha e^{-\sqrt{\alpha} B_s} ds \right\}^p \right] < \infty.$$

*Proof.* By the Itô formula, we have

$$e^{-\sqrt{\alpha} B_t} = e^{-\sqrt{\alpha} B_0} - \sqrt{\alpha} \int_0^t e^{-\sqrt{\alpha} B_s} dB_s + \int_0^t \alpha e^{-\sqrt{\alpha} B_s} ds.$$

Hence

$$\int_0^{t \wedge \tau} \alpha e^{-\sqrt{\alpha} B_s} ds = e^{-\sqrt{\alpha} B_{t \wedge \tau}} - e^{-\sqrt{\alpha} B_0} + M_t$$

where  $(M_t)$  is a martingale defined by

$$M_t = \sqrt{\alpha} \int_0^{t \wedge \tau} e^{-\sqrt{\alpha} B_s} dB_s,$$

which satisfies

$$\langle M \rangle_t = 2\alpha \int_0^{t \wedge \tau} e^{-2\sqrt{\alpha} B_s} ds.$$

Now, by the Burkholder inequality

$$\begin{aligned} & \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau \alpha e^{-\sqrt{\alpha} B_s} ds \right\}^p \right] \\ & \leq C_p \mathbf{E}_{\mu \times \delta_N} \left[ (e^{-\sqrt{\alpha} B_\tau} - e^{-\sqrt{\alpha} B_0})^p \right] + C_p \mathbf{E}_{\mu \times \delta_N} [\langle M \rangle_\tau^{p/2}] \\ & \leq C_p + C_p \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau 4\alpha e^{-\sqrt{4\alpha} B_s} ds \right\}^{p/2} \right]. \end{aligned}$$

Thus it is enough to show (5.5) when  $p = 1$ .

$$\mathbf{E}_{\mu \times \delta_N} \left[ \int_0^\tau \alpha e^{-\sqrt{\alpha} B_s} ds \right] = \int_0^\infty (N \wedge a) \alpha e^{-\sqrt{\alpha} a} da \leq \int_0^\infty a \alpha e^{-\sqrt{\alpha} a} da = 1.$$

This completes the proof. □

$G$ -functions are now estimated as follows.

**Proposition 5.2.** *For any  $1 < p < q < 2$ , we have*

$$(5.6) \quad \|\hat{G}\theta\|_p \lesssim \|\theta\|_q$$

and

$$(5.7) \quad \|\theta\|_{q'} \lesssim \|\hat{G}^{-1}\theta\|_{p'}$$

where  $p'$  and  $q'$  are the conjugate exponent of  $p$  and  $q$ , respectively.

For scalar functions, we have

$$(5.8) \quad \|Gu\|_p \lesssim \|u\|_p.$$

Proof. Set  $\hat{f}(x, a) = |\hat{Q}_a^{\lambda+R}\theta(x)|$  and for  $\varepsilon > 0$ ,  $\hat{f}_\varepsilon(x, a) = \sqrt{\hat{f}(x, a)^2 + \varepsilon}$ . Define

$$Z_t^{(\varepsilon)} = \hat{f}_\varepsilon(X_{t \wedge \tau}, B_{t \wedge \tau})^p$$

and

$$Z_t = \hat{f}(X_{t \wedge \tau}, B_{t \wedge \tau})^p.$$

Then

$$M_t^{(\varepsilon)} = Z_t^{(\varepsilon)} - \int_0^{t \wedge \tau} (L + \partial_a^2) \hat{f}_\varepsilon(X_s, B_s)^p ds$$

is a martingale.

By Lemma 4.4, we have

$$(5.9) \quad \begin{aligned} (L + \partial_a^2) \hat{f}_\varepsilon^p &\geq p(\lambda + V) \hat{f}^2 \hat{f}_\varepsilon^{p-2} + p(p-1) \hat{g} \hat{f}_\varepsilon^{p-2} \\ &\geq -p(\lambda + V)_- \hat{f}^2 \hat{f}_\varepsilon^{p-2} + p(p-1) \hat{g} \hat{f}_\varepsilon^{p-2}. \end{aligned}$$

Hence

$$Z_t^{(\varepsilon)} + \int_0^\tau p(\lambda + V)_-(X_s) \hat{f}_\varepsilon(X_s, B_s)^p ds$$

$$= M_t^{(\varepsilon)} + \int_0^{t \wedge \tau} (L + \partial_a^2) \hat{f}_\varepsilon(X_s, B_s)^p ds + \int_0^\tau p(\lambda + V)_-(X_s) \hat{f}_\varepsilon(X_s, B_s)^p ds$$

is a non-negative submartingale. By letting  $\varepsilon \rightarrow 0$  in (5.9), we have

$$\liminf_{\varepsilon \rightarrow 0} (L + \partial_a^2) \hat{f}_\varepsilon^p \geq -p(\lambda + V)_- \hat{f}^p + p(p - 1) \hat{g} \hat{f}^{p-2}$$

which implies

$$(5.10) \quad \hat{g} \leq \frac{1}{p(p - 1)} \liminf_{\varepsilon \rightarrow 0} (L + \partial_a^2) \hat{f}_\varepsilon^p \cdot \hat{f}^{2-p} + \frac{1}{p - 1} V_- \hat{f}^2.$$

Now we can estimate  $\hat{G}\theta$ .

$$\begin{aligned} \|\hat{G}\theta\|_p^p &= \left\| \left\{ \int_0^\infty a \hat{g}(x, a) da \right\}^{p/2} \right\|_1 \\ &\lesssim \left\| \left\{ \int_0^\infty \left\{ a \liminf_{\varepsilon \rightarrow 0} (L + \partial_a^2) \hat{f}_\varepsilon^p + ap(\lambda + V)_- \hat{f}^p \right\} \hat{f}^{2-p} da \right\}^{p/2} \right\|_1 \\ &\lesssim \left\| \left\{ \sup_{t \geq 0} T_t^{\lambda+V} |\theta| \right\}^{p(2-p)/2} \left\{ \int_0^\infty a \left\{ \liminf_{\varepsilon \rightarrow 0} (L + \partial_a^2) \hat{f}_\varepsilon^p + p(\lambda + V)_- \hat{f}^p \right\} da \right\}^{p/2} \right\|_1 \\ &\leq \left\| \left\{ \sup_{t \geq 0} T_t^{\lambda+V} |\theta| \right\}^p \right\|_1^{(2-p)/2} \times \left\| \int_0^\infty a \left\{ \liminf_{\varepsilon \rightarrow 0} (L + \partial_a^2) \hat{f}_\varepsilon^p + p(\lambda + V)_- \hat{f}^p \right\} da \right\|_1^{p/2}. \end{aligned}$$

The first factor of the right hand side can be estimated as follows. By Theorem 3.2, we have

$$(5.11) \quad \left\| \left\{ \sup_{t \geq 0} T_t^{\lambda+V} |\theta| \right\}^p \right\|_1^{(2-p)/2} \leq \left\| \sup_{t \geq 0} T_t^{\lambda+V} |\theta| \right\|_p^{p(2-p)/2} \lesssim \|\theta\|_q^{p(2-p)/2}.$$

For the second factor, we have

$$\begin{aligned} &\left\| \int_0^\infty a \left\{ \liminf_{\varepsilon \rightarrow 0} (L + \partial_a^2) \hat{f}_\varepsilon^p + p(\lambda + V)_- \hat{f}^p \right\} da \right\|_1 \\ &= \lim_{N \rightarrow \infty} \mathbf{E}_{\mu \times \delta_N} \left[ \int_0^\tau \liminf_{\varepsilon \rightarrow 0} \left\{ (L + \partial_a^2) \hat{f}_\varepsilon^p(X_t, B_t) + p(\lambda + V)_-(X_t) \hat{f}^p(X_t) \right\} dt \right] \\ &\leq \lim_{N \rightarrow \infty} \liminf_{\varepsilon \rightarrow 0} \mathbf{E}_{\mu \times \delta_N} \left[ \int_0^\tau \left\{ (L + \partial_a^2) \hat{f}_\varepsilon^p(X_t, B_t) + p(\lambda + V)_-(X_t) \hat{f}^p(X_t) \right\} dt \right] \\ &\quad (\because \text{the Fatou lemma}) \\ &= \lim_{N \rightarrow \infty} \liminf_{\varepsilon \rightarrow 0} \mathbf{E}_{\mu \times \delta_N} \left[ Z_\infty^{(\varepsilon)} - Z_0^{(\varepsilon)} + \int_0^\tau p(\lambda + V)_-(X_t) \hat{f}^p(X_t) dt \right] \\ &= \lim_{N \rightarrow \infty} \mathbf{E}_{\mu \times \delta_N} \left[ Z_\infty - Z_0 + \int_0^\tau p(\lambda + V)_-(X_t) \hat{f}^p(X_t) dt \right] \end{aligned}$$

$$\begin{aligned} &\leq \lim_{N \rightarrow \infty} \mathbf{E}_{\mu \times \delta_N} [|\theta(X_\tau)|^p] + \lim_{N \rightarrow \infty} \mathbf{E}_{\mu \times \delta_N} \left[ \int_0^\tau p(\lambda + V)_-(X_t) \hat{f}^p(X_t) dt \right] \\ &= \|\theta\|_p^p + \left\| \int_0^\infty a(\lambda + V)_- \hat{f}^p da \right\|_1. \end{aligned}$$

The second term can be estimated as follows:

$$\begin{aligned} \left\| \int_0^\infty a(\lambda + V)_- \hat{f}^p da \right\|_1 &\leq \int_0^\infty a \|V_- \hat{f}^p\|_1 da \\ &\leq \int_0^\infty a \|V_- \|_r \| \hat{f}^p \|_{q/p} da \quad \left( \frac{1}{q/p} + \frac{1}{r} = 1 \right) \\ &\leq \|V_- \|_r \int_0^\infty a \| \hat{Q}_a^{\lambda+R} \theta \|_q^p da \\ &\leq \|V_- \|_r \int_0^\infty a \| \hat{Q}_a^{\lambda+R} \|_{q \rightarrow q}^p \| \theta \|_q^p da \\ &\leq \|V_- \|_r \int_0^\infty a e^{-\sqrt{\lambda - \gamma_q} a p} \| \theta \|_q^p da \quad (\because (4.5)) \\ &= \frac{\|V_- \|_r \| \theta \|_q^p}{(\lambda - \gamma_q) p^2}. \end{aligned}$$

Thus we have

$$\| \hat{G} \theta \|_p^p \lesssim \| \theta \|_q^{p(2-p)/2} (\| \theta \|_p^p + \| \theta \|_q^p)^{p/2} \lesssim \| \theta \|_q^p$$

which shows (5.6).

(5.7) is obtained by the duality argument. In fact, using Proposition 4.1, we have

$$\begin{aligned} \int_M (\theta(x), \eta(x))_K \mu(dx) &= 4 \int_M \mu(dx) \int_0^\infty a (\partial_a \hat{Q}_a^{\lambda+R} \theta(x), \partial_a \hat{Q}_a^{\lambda+R} \eta(x))_K da \\ &\leq 4 \int_M \hat{G}^\rightarrow \theta(x) \hat{G}^\rightarrow \eta(x) \mu(dx) \\ &\leq 4 \| \hat{G}^\rightarrow \theta \|_p \| \hat{G}^\rightarrow \eta \|_{p'} \\ &\lesssim \| \theta \|_q \| \hat{G}^\rightarrow \eta \|_{p'}. \end{aligned}$$

Now (5.7) follows easily.

(5.8) for scalar functions can be shown much easily. □

When  $p \geq 2$ , we estimate  $\hat{H}u$  and  $Hu$ .

**Proposition 5.3.** *For any  $2 < p < r$ , we have*

$$(5.12) \quad \| \hat{H} \theta \|_p \lesssim \| \theta \|_r.$$

For scalar functions, we have

$$(5.13) \quad \|Hu\|_p \lesssim \|u\|_p.$$

Proof. We set  $\hat{f}(x, a) = |\hat{Q}_a^{\lambda+R}\theta(x)|$  and define

$$Z_t = \hat{f}(X_{t \wedge \tau}, B_{t \wedge \tau})^2.$$

Then,

$$M_t = Z_t - \int_0^{t \wedge \tau} (L + \partial_a^2) \hat{f}^2(X_s, B_s) ds$$

is a martingale. By Lemma 4.4, we have

$$(5.14) \quad 2\hat{g} \leq (L + \partial_a^2) \hat{f}^2 + 2V_- \hat{f}^2.$$

Then, setting

$$(5.15) \quad A_t = \int_0^{t \wedge \tau} \{(L + \partial_a^2) \hat{f}^2(X_s, B_s) + 2V_- \hat{f}^2(X_s, B_s)\} ds,$$

we can see that  $(A_t)$  is an increasing process and have that

$$(5.16) \quad Z_t + \int_0^{t \wedge \tau} 2V_- \hat{f}^2(X_s, B_s) ds = M_t + A_t.$$

Hence  $Z_t + \int_0^{t \wedge \tau} 2V_- \hat{f}^2 ds$  is a non-negative submartingale and its increasing part  $(A_t)$  satisfies

$$(5.17) \quad A_t \geq \int_0^{t \wedge \tau} 2\hat{g}(X_s, B_s) ds.$$

Therefore, by (5.4), the following inequality hold.

$$\begin{aligned} \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau 2\hat{g}(X_s, B_s) ds \right\}^{p/2} \right] &\leq \mathbf{E}_{\mu \times \delta_N} [A_\infty^{p/2}] \\ &\lesssim \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ Z_\infty + \int_0^\tau 2V_- \hat{f}^2 ds \right\}^{p/2} \right] \\ &\lesssim \mathbf{E}_{\mu \times \delta_N} [Z_\infty^{p/2}] + \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau V_- \hat{f}^2 ds \right\}^{p/2} \right] \\ &= \|\theta\|_p^p + \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau V_- \hat{f}^2 ds \right\}^{p/2} \right]. \end{aligned}$$

The second term can be estimated as follows. We take any  $p < q < r$ .

$$\begin{aligned}
& \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau V_- \hat{f}^2 ds \right\}^{p/2} \right] \\
&= \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau e^{-\eta B_s} e^{\eta B_s} V_- \hat{f}^2 ds \right\}^{p/2} \right] \\
&\leq \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau e^{-\eta B_s q/(q-2)} ds \right\}^{(q-2)p/2q} \left\{ \int_0^\tau e^{\eta B_s q/2} V_-^{q/2} \hat{f}^q ds \right\}^{2p/2q} \right] \\
&\quad (\because \text{the Hölder inequality for the exponents } q/(q-2) \text{ and } q/2) \\
&\leq \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau e^{-\eta B_s q/(q-2)} ds \right\}^{u(q-2)p/2q} \right]^{1/u} \mathbf{E}_{\mu \times \delta_N} \left[ \int_0^\tau e^{\eta B_s q/2} V_-^{q/2} \hat{f}^q ds \right]^{p/q} \\
&\quad \left( \because \text{the Hölder inequality for the exponents } u \text{ and } q/p \text{ where } \frac{1}{u} + \frac{1}{q/p} = 1 \right) \\
&\lesssim \mathbf{E}_{\mu \times \delta_N} \left[ \int_0^\tau e^{\eta q B_s/2} V_-^{q/2} \hat{f}^q ds \right]^{p/q} \quad (\because \text{Proposition 5.1}) \\
&= \left\{ \int_M \mu(dx) \int_0^\infty (N \wedge a) e^{\eta q a/2} V_-^{q/2}(x) |\hat{Q}_a^{\lambda+R} \theta(x)|^q da \right\}^{p/q}.
\end{aligned}$$

To estimate the integral above, we recall that  $\|\hat{Q}_t^{\lambda+R} \theta\|_r \leq e^{-\sqrt{\lambda-\gamma_r} t} \|\theta\|_r$ . Therefore,

$$\begin{aligned}
& \int_M \mu(dx) \int_0^\infty (N \wedge a) e^{\eta q a/2} V_-^{q/2}(x) |\hat{Q}_a^{\lambda+R} \theta(x)|^q da \\
&\leq \int_0^\infty a e^{\eta q a/2} da \int_M V_-^{q/2}(x) |\hat{Q}_a^{\lambda+R} \theta(x)|^q \mu(dx) \\
&\leq \int_0^\infty a e^{\eta q a/2} da \left\{ \int_M V_-^{vq/2}(x) \mu(dx) \right\}^{1/v} \left\{ \int_M |\hat{Q}_a^{\lambda+R} \theta(x)|^{qr/q} \mu(dx) \right\}^{q/r} \\
&\quad \left( \frac{1}{v} + \frac{1}{r/q} = 1 \right) \\
&\lesssim \int_0^\infty a e^{\eta q a/2} \|\hat{Q}_a^{\lambda+R} \theta\|_r^q da \\
&\lesssim \int_0^\infty a e^{\eta q a/2} e^{-\sqrt{\lambda-\gamma_r} q a} \|\theta\|_r^q da \\
&\lesssim \|\theta\|_r^q \quad (\because \sqrt{\lambda-\gamma_r} q > \eta q/2).
\end{aligned}$$

Thus we have obtained

$$(5.18) \quad \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau \hat{g}(X_s, B_s) ds \right\}^{p/2} \right] \lesssim \|\theta\|_r^p.$$

Now we can estimate  $\hat{H}\theta$ .

$$\begin{aligned} \|\hat{H}\theta\|_p^p &= \left\| \left\{ \int_0^\infty a Q_a \hat{g}(x, a) da \right\}^{p/2} \right\|_1 \\ &= \lim_{N \rightarrow \infty} \int_M \mu(dx) \left\{ \int_0^\infty (a \wedge N) Q_a \hat{g}(x, a) da \right\}^{p/2} \\ &= \lim_{N \rightarrow \infty} \int_M \mu(dx) \mathbf{E}_{\mu \times \delta_N} \left[ \int_0^\tau \hat{g}(X_s, B_s) ds \mid X_\tau = x \right]^{p/2} \\ &\leq \lim_{N \rightarrow \infty} \int_M \mu(dx) \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau \hat{g}(X_s, B_s) ds \right\}^{p/2} \mid X_\tau = x \right] \\ &= \lim_{N \rightarrow \infty} \mathbf{E}_{\mu \times \delta_N} \left[ \left\{ \int_0^\tau \hat{g}(X_s, B_s) ds \right\}^{p/2} \right] \\ &\lesssim \|\theta\|_r^p \quad (\because (5.18)) \end{aligned}$$

The scalar case is easier. □

Combining Propositions 4.3, 5.3, we can get

**Proposition 5.4.** *For any  $2 \leq p < q < \infty$ , we have*

$$(5.19) \quad \|\hat{G}^{-1}\theta\|_p \lesssim \|\theta\|_q$$

and

$$(5.20) \quad \|\theta\|_{q'} \lesssim \|\hat{G}^{-1}\theta\|_{p'}$$

where  $p'$  and  $q'$  are the conjugate exponents of  $p$  and  $q$ , respectively.

For scalar functions, we have

$$(5.21) \quad \|Gu\|_p \lesssim \|u\|_p.$$

We are now ready to prove Theorem 2.1.

Proof of Theorem 2.1. We take  $\lambda$  to be large enough. Recall that  $\{Q_t^\lambda\}$  is the subordination of  $\{T_t^\lambda\}$ . Then, by the intertwining property (2.7), we have

$$\nabla Q_t^\lambda = \hat{Q}_t^{\lambda+R} \nabla.$$

Now take any  $1 < p < q < \infty$ . Then we have

$$\|\nabla u\|_p \lesssim \|\hat{G}^{-1} \nabla u\|_q = \left\| \left\{ \int_0^\infty a |\partial_a \hat{Q}_a^{\lambda+R} \nabla u(x)|^2 da \right\}^{1/2} \right\|_q$$

$$\begin{aligned} &= \left\| \left\{ \int_0^\infty a \left| \nabla Q_a^\lambda \sqrt{\lambda - Lu} \right|^2 da \right\}^{1/2} \right\|_q \\ &= \|G^\dagger \sqrt{\lambda - Lu}\|_q \\ &\lesssim \|\sqrt{\lambda - Lu}\|_q \end{aligned}$$

which proves (2.10).

The reversed inequality (2.11) is obtained by the duality argument. This completes the proof.  $\square$

**6. Riemannian manifold with boundary**

In this section, we discuss the reflected Brownian motion on a Riemannian manifold with boundary. Let  $M$  be a compact Riemannian manifold with boundary  $\partial M$ . Let  $(X_t, P_x)_{x \in M}$  be the Brownian motion on  $M$  with the Neumann boundary condition. We denote the Riemannian volume by  $m$ . In this section, the semigroup  $\{T_t\}$  is generated by  $L = \Delta$  with the Neumann boundary condition.  $\{T_t\}$  is a symmetric and strongly continuous contraction semigroup in  $L^2(m)$ . Further  $\{\hat{T}_t\}$  is the semigroup generated by the Hodge-Kodaira Laplacian  $\hat{L} = -dd^* - d^*d$  with absolute boundary condition. The associated bilinear forms with  $L$  and  $\hat{L}$  are denoted by  $\mathcal{E}$  and  $\hat{\mathcal{E}}$ . We can see that the following intertwining property holds for  $\{T_t\}$  and  $\{\hat{T}_t\}$ :

$$(6.1) \quad \nabla T_t = \hat{T}_t \nabla.$$

As in §5, we use an additional 1-dimensional Brownian motion  $(B_t, P_a^-)$  generated by  $d^2/da^2$ . Let  $\tau$  be the hitting time of  $(B_t)$  to 0, and  $(Y_t, \mathbf{P}_{(x,a)})$  be the product diffusion process on the state space  $M \times \mathbb{R}$ .

$$(6.2) \quad Y_t := (X_t, B_t), \quad \mathbf{P}_{(x,a)} := P_x \otimes P_a^-.$$

So the generator of  $(Y_t)$  is  $L + \partial_a^2$ .

We use the notation  $\mathbf{E}_{m \times \delta_a} = \int_M \mathbf{P}_{(x,a)} m(dx)$  in the same way as in §5. For any  $f \in C^\infty(M)$ , we have

$$f(X_t, B_t) - f(X_0, B_0) = \text{a martingale} + \int_0^t (L + \partial_a^2) f(X_s, B_s) ds + \int_0^t \nabla_N f(X_s, B_s) dl_s.$$

Here  $\{l_t\}$  is an additive functional corresponding to the smooth measure  $\sigma$  ( $\sigma$  is the surface measure of  $\partial M$ ),  $N$  is the inner normal vector and  $\nabla$  denotes the covariant differentiation. In particular we take 1-form  $\theta$  with absolute boundary condition and set  $f(x, a) = |Q_a \theta(x)|^2$ . Then,

$$f(X_{t \wedge \tau}, B_{t \wedge \tau}) - f(X_0, B_0)$$



$$\begin{aligned}
 &= M_t + \int_0^{t \wedge \tau} (L + \partial_a^2) f(X_s, B_s) ds + \int_0^{t \wedge \tau} \nabla_N f(X_s, B_s) dl_s \\
 &= M_t + \int_0^{t \wedge \tau} (L + \partial_a^2) f(X_s, B_s) ds + \int_0^{t \wedge \tau} \alpha(Q_{B_s} \theta(X_s), Q_{B_s} \theta(X_s)) dl_s.
 \end{aligned}$$

Here  $\alpha$  is the second fundamental form of  $\partial M$  (see [15] for this identity.) The quadratic variation of  $(M_t)$  is given by

$$\langle M \rangle_t = 2 \int_0^{t \wedge \tau} |\nabla Q_{B_s} \theta(X_s)|^2 + |\partial_a Q_{B_s} \theta(X_s)|^2 ds.$$

Hence we can do the same argument as in the previous section. But we have to tackle the additional term  $\int_0^{t \wedge \tau} \alpha(Q_{B_s} \theta(X_s), Q_{B_s} \theta(X_s)) dl_s$ .

Next we see the semigroup domination. We note that for 1-forms  $\theta, \eta$  and  $f \in C^\infty(M)$ ,

$$\begin{aligned}
 &-\mathcal{E}((\theta, \eta), f) + \hat{\mathcal{E}}(f\theta, \eta) + \hat{\mathcal{E}}(\theta, f\eta) \\
 (6.3) \quad &= 2 \int_M (\nabla\theta, \nabla\eta) f m(dx) + 2 \int_M \text{Ric}(\theta, \eta) f m(dx) + 2 \int_{\partial M} \alpha(\theta, \eta) f \sigma(dx)
 \end{aligned}$$

where Ric is the Ricci curvature (refer to [15] for this identity.)

We take  $\gamma \geq 0$  and  $\beta \geq 0$  so that  $\text{Ric}(\theta, \theta) \geq -\gamma|\theta|^2$  and  $\alpha(\theta, \theta) \geq -\beta|\theta|^2$ . Then  $\alpha(\theta, \theta)\sigma \geq -\beta|\theta|^2\sigma$  as measures. It is easy to see that  $\sigma$  is a smooth measure. We also note that in the interior of  $M$ , it holds that

$$(6.4) \quad L(\theta, \eta) - (\hat{L}\theta, \eta) - (\theta, \hat{L}\eta) = 2(\nabla\theta, \nabla\eta) - 2\text{Ric}(\theta, \eta).$$

By (6.3) and (6.4), the semigroup domination theorem implies (see [14, 15])

$$(6.5) \quad |\hat{T}_t \theta| \leq T_t^{-\gamma-\beta\sigma} |\theta|.$$

Here  $T_t^{-\gamma-\beta\sigma}$  is the semigroup which has  $-\gamma - \beta\sigma$  as a potential. It can be represented as

$$(6.6) \quad T_t^{-\gamma-\beta\sigma} u(x) = E_x [u(X_t) e^{\gamma t + \beta l_t}]$$

We can also show that  $(-l_t)$  satisfies the assumption of Theorem 3.1. To see this, take any function  $h \in C^\infty(M)$  such that  $\nabla_N h = 1$  on  $\partial M$ . Such a function can be constructed as follows. Take any local coordinate  $(x_1, \dots, x_{n-1}, r)$  such that  $\partial M = \{r = 0\}$  and  $r \mapsto (x_1, \dots, x_{n-1}, r)$  is a geodesic with unit velocity perpendicular to  $\partial M$ . Then  $h(x_1, \dots, x_{n-1}, r) = r$  satisfies the property above. Global existence of  $h$  can be obtained by using the partition of unity. Then

$$h(X_t) - h(X_0) = M_t + \int_0^t \Delta h(X_s) ds + l_t$$

where  $(M_t)$  is a martingale with  $d\langle M \rangle_t \leq C dt$  for a constant  $C > 0$ . Hence

$$E_x[e^{qh_t}] = E_x \left[ \exp \left\{ qh(X_t) - qh(X_0) - qM_t - q \int_0^t \Delta h(X_s) ds \right\} \right].$$

The right hand side is bounded in  $x$  because  $h$ ,  $\Delta h$  and  $\langle M \rangle$  is bounded (this implies that  $\sigma$  is a Kato class potential; for Kato class potentials, see Albeverio-Ma [2]). Therefore, there exist constant  $c_q > 0$  and  $\beta_q > 0$  such that, for q.e.- $x$ ,

$$(6.7) \quad E_x[e^{qh_t}]^{1/q} \leq c_q e^{\beta_q t}, \quad \forall t \geq 0.$$

Now we can apply Theorem 3.1 to  $T_t^{\lambda-\gamma-\beta\sigma}$ . For simplicity, we introduce the following notation:

$$M^{\lambda-\gamma-\beta\sigma}u(x) = \sup_{t \geq 0} |T_t^{\lambda-\gamma-\beta\sigma}u(x)|.$$

When  $\lambda - \gamma = 0$  and  $\beta = 0$ , we simply denote  $Mu$  in place of  $M^{\lambda-\gamma-\beta\sigma}u$ . Then, if  $\lambda$  is large enough, we have for any  $p > 1$ ,

$$(6.8) \quad \|M^{\lambda-\gamma-\beta\sigma}u\|_p \lesssim \|u\|_p.$$

We can also obtain an estimate for the subordination. Let  $\{Q_t^{\lambda-\gamma-\beta\sigma}\}$  be the subordination of  $\{T_t^{\lambda-\gamma-\beta\sigma}\}$ . Then

$$\begin{aligned} |Q_t^{\lambda-\gamma-\beta\sigma}u| &= \left| \int_0^\infty T_s^{\lambda-\gamma-\beta\sigma}u \mu_t(ds) \right| \\ &\leq \int_0^\infty e^{-\alpha^2 s} \sup_{r \geq 0} |T_r^{\lambda-\alpha^2-\gamma-\beta\sigma}u| \mu_t(ds) \\ &= e^{-\alpha t} M^{\lambda-\alpha^2-\gamma-\beta\sigma}u. \end{aligned}$$

Thus we have

$$(6.9) \quad \sup_{t \geq 0} \{e^{\alpha t} |Q_t^{\lambda-\gamma-\beta\sigma}u|\} \leq M^{\lambda-\alpha^2-\gamma-\beta\sigma}u.$$

We also note that  $\{\hat{T}_t\}$  is a bounded operator in  $L^p$  by virtue of (6.5) and there exist constants  $c_p > 0$  and  $\gamma_p > 0$  so that

$$(6.10) \quad \|\hat{T}_t\|_{p \rightarrow p} \leq c_p e^{\gamma_p t}.$$

Let  $\{\hat{Q}_t^\lambda\}$  be the subordination of  $\{\hat{T}_t^\lambda = e^{-\lambda t} \hat{T}_t\}$ . (6.5) implies  $|\hat{Q}_t^\lambda \theta| \leq Q_t^{\lambda-\gamma-\beta\sigma}|\theta|$ . We define  $\hat{G}$  and  $\hat{H}$  in terms of  $\{\hat{Q}_t^\lambda\}$ . Now we can easily see that Proposition 4.3 holds in this case. We have more. In fact, by virtue of (6.7), we can and do take  $\lambda$

large enough so that  $\sup_{t \geq 0} T_t^{\lambda - \gamma - \beta \sigma} 1(x)$  is bounded in q.e.- $x$  and thereby we have

$$(6.11) \quad \hat{G} \rightarrow \theta \lesssim \hat{H} \rightarrow \theta.$$

Similar estimate holds for  $G^\uparrow u$  and  $H^\uparrow u$ .

Lastly we note that, by combining the domination and (6.9),

$$(6.12) \quad \sup_{t \geq 0} \{e^{\alpha t} |\hat{Q}_t^\lambda \theta|\} \leq M^{\lambda - \alpha^2 - \gamma - \beta \sigma} |\theta|.$$

Next we extend (5.2) to additive functionals. Take any smooth measure  $\rho$  and let  $A_t$  be the additive functional associated with  $\rho$ . Then we have the following identity.

**Proposition 6.1.** *For any non-negative function  $f$  on  $M \times [0, \infty)$  and  $k$  on  $M$ , the following identity holds:*

$$(6.13) \quad \mathbf{E}_{m \times \delta_a} \left[ \int_0^\tau f(X_t, B_t) dA_t \right] = \int_M \rho(dx) \int_0^\infty (a \wedge t) f(x, t) dt$$

$$(6.14) \quad \mathbf{E}_{m \times \delta_a} \left[ k(X_\tau) \int_0^\tau f(X_t, B_t) dA_t \right] = \int_M \rho(dx) \int_0^\infty (a \wedge t) Q_t k(x) f(x, t) dt$$

Proof. Let us first recall the resolvent kernel for the absorbing Brownian motion on  $(0, \infty)$ . Here, the generator is  $d^2/da^2$ . For  $\alpha > 0$ , set

$$(6.15) \quad g_\alpha(a, b) = \begin{cases} \frac{1}{2\sqrt{\alpha}}(e^{\sqrt{\alpha}a} - e^{-\sqrt{\alpha}a})e^{-\sqrt{\alpha}b}, & a \leq b, \\ \frac{1}{2\sqrt{\alpha}}e^{-\sqrt{\alpha}a}(e^{\sqrt{\alpha}b} - e^{-\sqrt{\alpha}b}), & a \geq b. \end{cases}$$

Then the resolvent  $G_\alpha = (\alpha - d/(da^2))^{-1}$  is given by

$$(6.16) \quad G_\alpha h(a) = \int_0^\infty g_\alpha(a, b) h(b) db.$$

Moreover we note that  $\lim_{\alpha \rightarrow 0} g_\alpha(a, b) = a \wedge b$ .

By the Revuz correspondence, (see [6, the equation (5.1.14)]) we have

$$\begin{aligned} & \int_0^\infty h(a) da \mathbf{E}_{m \times \delta_a} \left[ \int_0^\tau e^{-\alpha s} f(X_s, B_s) dA_s \right] \\ &= \int_0^\infty G_\alpha h(a) da \int_M f(x, a) \rho(dx) \\ &= \int_0^\infty h(a) da \int_M \rho(dx) \int_0^\infty g_\alpha(a, b) f(x, b) db \quad (\because g_\alpha \text{ is symmetric}) \end{aligned}$$

Hence we have, for a.e.- $a$ ,

$$\mathbf{E}_{m \times \delta_a} \left[ \int_0^\tau e^{-\alpha s} f(X_s, B_s) dA_s \right] = \int_M \rho(dx) \int_0^\infty g_\alpha(a, b) f(x, b) db.$$

But both hands are quasi-continuous in  $a$  and one point has positive capacity, the above identity holds for all  $a \geq 0$ . By letting  $\alpha \rightarrow 0$ , we can get (6.13).

To show (6.14), set

$$H_t = \int_0^{t \wedge \tau} f(X_s, B_s) dA_s.$$

Then  $H_t$  is a process of bounded variation. Hence, by the Itô formula,

$$\begin{aligned} Q_{B_{t \wedge \tau}} k(X_{t \wedge \tau}) H_t &= \text{a martingale} + \int_0^{t \wedge \tau} (L + \partial_a^2) Q_{B_s} k(X_s) H_s ds + \int_0^{t \wedge \tau} Q_{B_s} k(X_s) dH_s \\ &= \text{a martingale} + \int_0^{t \wedge \tau} Q_{B_s} k(X_s) f(X_s, B_s) dA_s. \end{aligned}$$

Here we used that  $\nabla_N Q_a k = 0$  on  $\partial M$  because  $Q_a k$  belongs to the domain of the Neumann Laplacian. By taking expectation and letting  $t \rightarrow \infty$ , we have

$$\begin{aligned} \mathbf{E}_{m \times \delta_a} \left[ Q_{B_\tau} k(X_\tau) \int_0^\tau f(X_s, B_s) dA_s \right] &= \mathbf{E}_{m \times \delta_a} \left[ \int_0^\tau Q_{B_s} k(X_s) f(X_s, B_s) dA_s \right] \\ &= \int_M \rho(dx) \int_0^\infty (a \wedge t) Q_t k(x) f(x, t) dt. \quad (\because (6.13)) \end{aligned}$$

This completes the proof. □

Recall that  $\{Q_t^\lambda\}$ ,  $\{\hat{Q}_t^\lambda\}$  are subordinations of  $\{T_t^\lambda\}$ ,  $\{\hat{T}_t^\lambda\}$ , respectively and  $G$  and  $H$  functions are defined in terms of  $\{\hat{Q}_t^\lambda\}$ . Then we have the following estimate.

**Proposition 6.2.** *For  $1 < p \leq 2$ , we have*

$$(6.17) \quad \|\hat{G}\theta\|_p \lesssim \|\theta\|_p$$

and

$$(6.18) \quad \|\theta\|_{p'} \lesssim \|G^{-\rightarrow}\theta\|_{p'}$$

where  $p'$  is the conjugate exponent of  $p$ .

For scalar functions, we have

$$(6.19) \quad \|Gu\|_p \lesssim \|u\|_p.$$

**Proof.** We only show the 1-form case. We set  $\hat{f}(x, a) = |\hat{Q}_a^\lambda \theta(x)|$  and  $\hat{f}_\varepsilon = \sqrt{\hat{f}^2 + \varepsilon}$  ( $\varepsilon > 0$ ). Define

$$Z_t^{(\varepsilon)} = \hat{f}_\varepsilon(X_{t \wedge \tau}, B_{t \wedge \tau})^p$$

and

$$Z_t = \hat{f}(X_{t \wedge \tau}, B_{t \wedge \tau})^p.$$

Then,

$$M_t^{(\varepsilon)} = Z_t^{(\varepsilon)} - \int_0^{t \wedge \tau} (L + \partial_a^2) \hat{f}_\varepsilon(X_s, B_s)^p ds - \int_0^{t \wedge \tau} \nabla_N \hat{f}_\varepsilon(X_s, B_s)^p dl_s.$$

is a martingale. Note that

$$\nabla_N \hat{f}_\varepsilon^p = \nabla_N (\hat{f}^2 + \varepsilon)^{p/2} = \frac{p}{2} (\hat{f}^2 + \varepsilon)^{(p-2)/2} \alpha(\hat{Q}_a^\lambda \theta, \hat{Q}_a^\lambda \theta).$$

Therefore,

$$\begin{aligned} & \mathbf{E}_{m \times \delta_N} \left[ \int_0^\tau (L + \partial_a^2) \hat{f}_\varepsilon(X_s, B_s)^p ds \right] \\ &= \mathbf{E}_{m \times \delta_N} [Z_\infty^{(\varepsilon)} - Z_0^{(\varepsilon)}] - \frac{p}{2} \mathbf{E}_{m \times \delta_N} \left[ \int_0^\tau \hat{f}_\varepsilon^{p-2} \alpha(\hat{Q}_{B_s}^\lambda \theta, \hat{Q}_{B_s}^\lambda \theta) dl_s \right] \\ &\leq \mathbf{E}_{m \times \delta_N} [Z_\infty^{(\varepsilon)} - Z_0^{(\varepsilon)}] + \frac{p\beta}{2} \mathbf{E}_{m \times \delta_N} \left[ \int_0^\tau \hat{f}_\varepsilon^{p-2} |\hat{Q}_{B_s}^\lambda \theta|^2 dl_s \right]. \end{aligned}$$

By taking limit, we have

$$\begin{aligned} & \mathbf{E}_{m \times \delta_N} \left[ \liminf_{\varepsilon \rightarrow 0} \int_0^\tau (L + \partial_a^2) \hat{f}_\varepsilon(X_s, B_s)^p ds \right] \\ &\leq \liminf_{\varepsilon \rightarrow 0} \mathbf{E}_{m \times \delta_N} [Z_\infty^{(\varepsilon)} - Z_0^{(\varepsilon)}] + \frac{p\beta}{2} \liminf_{\varepsilon \rightarrow 0} \mathbf{E}_{m \times \delta_N} \left[ \int_0^\tau \hat{f}_\varepsilon^{p-2} |\hat{Q}_{B_s}^\lambda \theta(X_s)|^2 dl_s \right] \\ &\leq \|u\|_p^p + \frac{p\beta}{2} \liminf_{\varepsilon \rightarrow 0} \int_0^\infty (N \wedge a) da \int_M \hat{f}_\varepsilon^{p-2} |\hat{Q}_a^\lambda \theta|^2 \sigma(dx) \\ &\leq \|u\|_p^p + \frac{p\beta}{2} \liminf_{\varepsilon \rightarrow 0} \int_0^\infty (N \wedge a) da \int_M \hat{f}_\varepsilon^p \sigma(dx). \end{aligned}$$

We estimate the second term. We use the interpolation space. Taking  $\xi = 1 - (1/p)$ , we introduce the interpolation norm  $\|\cdot\|_{\xi,p}$  of  $\|\cdot\|_{0,p}$  and  $\|\cdot\|_{1,p}$ . Here  $\|\cdot\|_{0,p}$  is the  $L^p$  norm in  $L^p(M, dx)$  and  $\|\cdot\|_{1,p}$  is the Sobolev norm:

$$\|u\|_{1,p}^p = \int_M |u|^p m(dx) + \int_M |\nabla u|^p m(dx).$$

Then the following inequality holds (see e.g., [1, Chapter VII]):

$$\int_{\partial M} |u|^p d\sigma(x) \lesssim \|u\|_{\xi,p}^p.$$

Moreover, the general theory of interpolation implies (see [1, LEMMA 7.16])

$$\|u\|_{\xi,p}^p \lesssim \|u\|_{0,p}^{(1-\xi)p} \|u\|_{1,p}^{\xi p}.$$

Thus we have

$$(6.20) \quad \int_{\partial M} |u|^p \sigma(dx) \lesssim \|u\|_{0,p}^{(1-\xi)p} \|u\|_{1,p}^{\xi p}.$$

On the other hand

$$|\nabla \hat{f}_\varepsilon| = \left| \nabla \sqrt{\hat{f}^2 + \varepsilon} \right| = \frac{1}{2}(\hat{f}^2 + \varepsilon)^{-1/2} |\nabla \hat{f}^2| \leq \frac{1}{2}(\hat{f}^2 + \varepsilon)^{-1/2} 2\hat{f} |\nabla \hat{Q}_a^\lambda \theta| \leq |\nabla \hat{Q}_a^\lambda \theta|.$$

Using these inequalities, we have

$$\begin{aligned} \int_0^\infty (N \wedge a) da \int_M \hat{f}_\varepsilon^p \sigma(dx) &\lesssim \int_0^\infty (N \wedge a) \|\hat{f}_\varepsilon\|_{0,p}^{(1-\xi)p} \|\hat{f}_\varepsilon\|_{1,p}^{\xi p} da \\ &\lesssim \int_0^\infty (N \wedge a) \{ \|\hat{f}_\varepsilon\|_p^p + \|\hat{f}_\varepsilon\|_p^{(1-\xi)p} \|\nabla \hat{f}_\varepsilon\|_p^{\xi p} \} da \\ &\lesssim \int_0^\infty (N \wedge a) \{ \|\hat{f}_\varepsilon\|_p^p + \|\hat{f}_\varepsilon\|_p^{(1-\xi)p} \|\nabla \hat{Q}_a^\lambda \theta\|_p^{\xi p} \} da. \end{aligned}$$

By taking limit, we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^\infty (N \wedge a) da \int_M \hat{f}_\varepsilon^p \sigma(dx) &\lesssim \int_0^\infty (N \wedge a) \{ \|\hat{f}\|_p^p + \|\hat{f}\|_p^{(1-\xi)p} \|\nabla \hat{Q}_a^\lambda \theta\|_p^{\xi p} \} da \\ &\lesssim \int_0^\infty a e^{-\sqrt{\lambda-\gamma_p}pa} \|\theta\|_p^p da + \int_0^\infty a e^{-\sqrt{\lambda-\gamma_p}(1-\xi)pa} \|\theta\|_p^{(1-\xi)p} \|\nabla \hat{Q}_a^\lambda \theta\|_p^{\xi p} da \\ &\lesssim \|\theta\|_p^p + \|\theta\|_p^{(1-\xi)p} \int_0^\infty a e^{-\sqrt{\lambda-\gamma_p}(1-\xi)pa} da \left\{ \int_M |\nabla \hat{Q}_a^\lambda \theta(x)|^p m(dx) \right\}^\xi \\ &\leq \|\theta\|_p^p + \|\theta\|_p^{(1-\xi)p} \left\{ \int_0^\infty a e^{-\sqrt{\lambda-\gamma_p}(1-\xi)pa} a^{1/(1-\xi)} da \right\}^{1-\xi} \\ &\quad \times \left\{ \int_0^\infty a e^{-\sqrt{\lambda-\gamma_p}(1-\xi)pa} da \int_M |\nabla \hat{Q}_a^\lambda \theta(x)|^p m(dx) \right\}^\xi \\ &\quad \left( \frac{1}{1/(1-\xi)} + \frac{1}{1/\xi} = 1 \right) \end{aligned}$$

$$\begin{aligned}
 &\lesssim \|\theta\|_p^p + \|\theta\|_p^{(1-\xi)p} \left\{ \int_M m(dx) \int_0^\infty a e^{-\sqrt{\lambda-\gamma_p}(1-\xi)pa} |\nabla \hat{Q}_a^\lambda \theta(x)|^p da \right\}^\xi \\
 &\lesssim \|\theta\|_p^p + \|\theta\|_p^{(1-\xi)p} \left[ \int_M m(dx) \left\{ \int_0^\infty a e^{-\sqrt{\lambda-\gamma_p}(1-\xi)pva} da \right\}^{1/\nu} \right. \\
 &\quad \times \left. \left\{ \int_0^\infty a |\nabla \hat{Q}_a^\lambda \theta(x)|^{p \cdot 2/p} da \right\}^{p/2} \right]^\xi \quad \left( \frac{1}{\nu} + \frac{1}{2/p} = 1 \right) \\
 &\lesssim \|\theta\|_p^p + \|\theta\|_p^{(1-\xi)p} \left[ \int_M \left\{ \int_0^\infty a |\nabla \hat{Q}_a^\lambda \theta(x)|^2 da \right\}^{p/2} m(dx) \right]^\xi \\
 &\lesssim \|\theta\|_p^p + \|\theta\|_p^{(1-\xi)p} \|\hat{G}^\dagger \theta\|_p^{\xi p}.
 \end{aligned}$$

Further, as in the proof of Proposition 5.2, we can show that

$$\|\hat{G}\theta\|_p^p \lesssim \|\theta\|_p^{p(2-p)/2} \left\| \int_0^\infty a \liminf_{\varepsilon \rightarrow 0} (L + \partial_a^2) \hat{f}_\varepsilon^p da \right\|^{p/2}.$$

In fact,  $(\lambda + V)_-$  in the proof of Proposition 5.2 vanishes in this case. Combining these inequalities, we have

$$\begin{aligned}
 \|\hat{G}\theta\|_p^p &\lesssim \|\theta\|_p^{p(2-p)/2} \{ \|\theta\|_p^{p \cdot p/2} + \|\theta\|_p^{(1-\xi)p \cdot p/2} \|\hat{G}^\dagger \theta\|_p^{\xi p \cdot p/2} \} \\
 &\leq \|\theta\|_p^p + \|\theta\|_p^{(2-\xi)p/2} \|\hat{G}^\dagger \theta\|_p^{\xi p^2/2} \\
 &= \|\theta\|_p^p + \|\theta\|_p^{(3-p)p/2} \|\hat{G}^\dagger \theta\|_p^{(p-1)p/2} \quad \left( \xi = 1 - \frac{1}{p} \right) \\
 &\leq \|\theta\|_p^p + \frac{3-p}{2} \delta^{-(p-1)/(3-p)} \|\theta\|_p^p + \frac{p-1}{2} \delta \|\hat{G}^\dagger \theta\|_p^p \quad \left( \frac{3-p}{2} + \frac{p-1}{2} = 1 \right) \\
 &\lesssim \|\theta\|_p^p + \delta^{-(p-1)/(3-p)} \|\theta\|_p^p + \delta \|\hat{G}\theta\|_p^p.
 \end{aligned}$$

Since  $\delta$  is arbitrary, we can get

$$\|\hat{G}\theta\|_p^p \lesssim \|\theta\|_p^p$$

which is (6.17). Now the rest is easy. □

When  $p \geq 2$ , we estimate  $\hat{H}\theta$  and  $Hu$ .

**Proposition 6.3.** *We further assume that the second fundamental form  $\alpha$  is non-negative definite. Then, for any  $p \geq 2$ , we have*

$$(6.21) \quad \|\hat{H}\theta\|_p \lesssim \|\theta\|_p.$$

For scalar functions, we have

$$(6.22) \quad \|Hu\|_p \lesssim \|u\|_p.$$

Proof. Set  $\hat{f}(x, a) = |\hat{Q}_a^\lambda \theta(x)|$  for 1-form  $\theta$  and define

$$Z_t = \hat{f}(X_{t \wedge \tau}, B_{t \wedge \tau})^2.$$

Then

$$Z_t = Z_0 + M_t - \int_0^{t \wedge \tau} (L + \partial_a^2) \hat{f}^2(X_s, B_s) ds - \int_0^{t \wedge \tau} \alpha(\hat{Q}_{B_s}^\lambda \theta(X_s), \hat{Q}_{B_s}^\lambda \theta(X_s)) dl_s$$

where  $(M_t)$  is a martingale with the quadratic variation

$$\langle M \rangle_t = 2 \int_0^{t \wedge \tau} \{ |\nabla \hat{f}^2(X_s, B_s)|^2 + |\partial_a \hat{f}^2(X_s, B_s)|^2 \} ds.$$

By the assumption that  $\alpha$  is non-negative definite,  $(Z_t)$  is a submartingale and the increasing part is given as

$$A_t := \int_0^{t \wedge \tau} (L + \partial_a^2) \hat{f}^2(X_s, B_s) ds + \int_0^{t \wedge \tau} \alpha(\hat{Q}_{B_s}^\lambda \theta(X_s), \hat{Q}_{B_s}^\lambda \theta(X_s)) dl_s.$$

Now, recalling that (see Lemma 4.4)

$$(L + \partial_a^2) \hat{f}^2 \geq 2\hat{g},$$

we have

$$A_t \geq \int_0^{t \wedge \tau} 2\hat{g}(X_s, B_s) ds.$$

By virtue of the submartingale inequality (5.4), we obtain

$$\begin{aligned} \mathbf{E}_{m \times \delta_N} \left[ \left\{ \int_0^\tau 2\hat{g}(X_s, B_s) ds \right\}^{p/2} \right] &\leq \mathbf{E}_{m \times \delta_N} [A_\infty^{p/2}] \\ &\lesssim \mathbf{E}_{m \times \delta_N} [Z_\infty^{p/2}] \\ &= \|\theta\|_p^p. \end{aligned}$$

Thus we have

$$\begin{aligned} \|\hat{H}\theta\|_p^p &= \left\| \left\{ \int_0^\infty a Q_a \hat{g}(x, a) da \right\}^{p/2} \right\|_1 \\ &= \lim_{N \rightarrow \infty} \int_M \mu(dx) \left\{ \int_0^\infty (a \wedge N) Q_a \hat{g}(x, a) da \right\}^{p/2} \\ &= \lim_{N \rightarrow \infty} \int_M \mu(dx) \mathbf{E}_{m \times \delta_N} \left[ \int_0^\tau \hat{g}(X_s, B_s) ds \mid X_\tau = x \right]^{p/2} \end{aligned}$$



$$\begin{aligned} &\leq \lim_{N \rightarrow \infty} \int_M \mu(dx) \mathbf{E}_{m \times \delta_N} \left[ \left\{ \int_0^\tau \hat{g}(X_s, B_s) ds \right\}^{p/2} \middle| X_\tau = x \right] \\ &= \lim_{N \rightarrow \infty} \mathbf{E}_{m \times \delta_N} \left[ \left\{ \int_0^\tau \hat{g}(X_s, B_s) ds \right\}^{p/2} \right] \\ &\lesssim \|\theta\|_p^p. \end{aligned}$$

Scalar case is easier. □

By combining Propositions 4.3 and 6.3, we easily obtain the following estimates for  $G$ -functions:

**Proposition 6.4.** *Assume that  $\alpha$  is non-negative definite. Then, for any  $p \geq 2$ , we have*

$$(6.23) \quad \|\hat{G} \rightarrow \theta\|_p \lesssim \|\theta\|_p$$

and

$$(6.24) \quad \|\theta\|_{p'} \lesssim \|\hat{G} \rightarrow \theta\|_{p'}$$

where  $p'$  is the conjugate exponent of  $p$ .

For scalar functions, we have

$$(6.25) \quad \|Gu\|_p \lesssim \|u\|_p.$$

Now the following theorem can be proved in the same way as Theorem 2.1.

**Theorem 6.5.** *For any  $1 < p \leq 2$ , it holds that*

$$(6.26) \quad \|u\|_p + \|\nabla u\|_p \lesssim \|\sqrt{1 - \Delta}u\|_p,$$

$$(6.27) \quad \|\sqrt{1 - \Delta}u\|_{p'} \lesssim \|u\|_{p'} + \|\nabla u\|_{p'}.$$

where  $p'$  is the conjugate exponent of  $p$ .

If we further assume that the second fundamental form  $\alpha$  is non-negative definite, then the inequalities above hold for  $p \geq 2$ .

ACKNOWLEDGEMENT. The author is grateful to the referee for his careful reading. He pointed out a defect of the proof of Proposition 5.2 and corrected it.

## References

- [1] R.A. Adams: Sobolev spaces, Academic press, New York, 1975.
- [2] S. Albeverio and Z.-M. Ma: *Perturbation of Dirichlet forms—lower semiboundedness, closability, and form cores*, J. Funct. Anal. **99** (1991), 332–356.
- [3] D. Bakry: *Transformations de Riesz pour les semigroupes symétriques*, Séminaire de Prob. XIX, Lecture Notes in Math. **1123**, 130–174, Springer-Verlag, Berlin-Heidelberg-New York, 1985.
- [4] D. Bakry: *Etude des transformations de Riesz dans les variétés riemanniennes à courbure de Ricci minorée*, Séminaire de Prob. XXI, Lecture Notes in Math. **1247**, 137–172, Springer-Verlag, Berlin-Heidelberg-New York, 1987.
- [5] D. Bakry and M. Emery: *Diffusions hypercontractives*, Séminaire de Prob. XIX, Lecture Notes in Math. **1123**, 177–206, Springer-Verlag, Berlin-Heidelberg-New York 1985.
- [6] M. Fukushima, T. Oshima and M. Takeda: Dirichlet forms and Markov processes, Walter de Gruyter, Berlin-New York, 1994.
- [7] L. Gross: *Logarithmic Sobolev inequalities*, Amer. J. Math. **97** (1975), 1061–1083.
- [8] E. Lenglart, D. Lépingle and M. Pratelli: *Présentation unifiée de certaines inégalités de théorie des martingales*, Séminaire de Prob. XIV, Lecture Notes in Math. **784**, 26–48. Springer-Verlag, Berlin-Heidelberg-New York, 1980.
- [9] P.A. Meyer: *Démonstration probabiliste de certaines inégalité de Littlewood-Paley*, Séminaire de Prob. X, ed. par P.A. Meyer, Lecture Notes in Math. **511**, 125–183, Springer-Verlag, Berlin, 1976.
- [10] P.A. Meyer: *Notes sur les processus d’Ornstein-Uhlenbeck*, Séminaire de Prob. XVI, ed. par J. Azema et M. Yor, Lecture Notes in Math. **920**, 95–133, Springer-Verlag, Berlin-Heidelberg-New York, 1982.
- [11] P.A. Meyer: *Quelques results analytiques sur le semigroupe d’Ornstein-Uhlenbeck en dimension infinie*, Theory and application of random fields, Proceedings of IFIP-WG 7/1 Working conf. at Bangalore, ed. by G. Kallianpur, Lecture Notes in Cont. and Inform. Sci. **49**, 201–214, Springer-Verlag, Berlin-Heidelberg-New York. 1983.
- [12] E. Ouhabaz: *Invariance of closed convex sets and domination criteria for semigroups Potential Analysis*, **5** (1996), 611–625.
- [13] G.C. Rota: *An “Alternierende Verfahren” for general positive operators*, Bull. Amer. Math. Soc. **68** (1962), 95–102.
- [14] I. Shigekawa:  *$L^p$  contraction semigroups for vector valued functions*, J. Funct. Anal. **147** (1997), 69–108.
- [15] I. Shigekawa: *Semigroup domination on a Riemannian manifold with boundary*, Acta Applicandae Math, **63** (2000), 385–410.
- [16] I. Shigekawa and N. Yoshida: *Littlewood-Paley-Stein inequality for a symmetric diffusion*, J. Math. Soc. Japan, **44** (1992), 251–280.
- [17] E. M. Stein: Topics in harmonic analysis, related to Littlewood-Paley theory, Annals of Math. Studies, **63**, Princeton Univ. Press, 1974.
- [18] N. Yoshida: *Sobolev spaces on a Riemannian manifold and their equivalence*, J. Math. Kyoto Univ. **32** (1992), 621–654.
- [19] N. Yoshida: *The Littlewood-Paley-Stein inequality on an infinite dimensional manifold*, J. Funct. Anal. **122** (1994), 402–427.

Department of Mathematics  
Graduate School of Science  
Kyoto University  
Kyoto 606-8502  
Japan  
e-mail: [ichiro@kusm.kyoto-u.ac.jp](mailto:ichiro@kusm.kyoto-u.ac.jp)  
URL: <http://www.kusm.kyoto-u.ac.jp/~ichiro/>