



Title	Chern characters on compact Lie groups of low rank
Author(s)	Watanabe, Takashi
Citation	Osaka Journal of Mathematics. 1985, 22(3), p. 463-488
Version Type	VoR
URL	<a href="https://doi.org/10.18910/10102">https://doi.org/10.18910/10102</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## CHERN CHARACTERS ON COMPACT LIE GROUPS OF LOW RANK

Dedicated to Professor Minoru Nakaoka on his sixtieth birthday

TAKASHI WATANABE

(Received August 8, 1984)

### 0. Introduction

Let  $G$  be a compact, simply connected, simple Lie group of rank  $l$ .  $G$  has  $l$  irreducible representations  $\rho_1, \dots, \rho_l$ , whose highest weights are the fundamental weights  $\omega_1, \dots, \omega_l$  respectively (see [19]). Then the representation ring  $R(G)$  of  $G$  is a polynomial algebra  $Z[\rho_1, \dots, \rho_l]$ . By the theorem of Hodgkin [16], the  $Z/2$ -graded  $K$ -theory  $K^*(G)$  of  $G$  is an exterior algebra  $\Lambda_Z(\beta(\rho_1), \dots, \beta(\rho_l))$ , where  $\beta: R(G) \rightarrow K^*(G)$  is the map introduced in [16]. Therefore the Chern character  $ch: K^*(G) \rightarrow H^*(G; Q)$  is injective [5]. We may write

$$H^*(G; Q) = \Lambda_Q(x_{2m_1-1}, x_{2m_2-1}, \dots, x_{2m_l-1})$$

where  $2 = m_1 \leq m_2 \leq \dots \leq m_l$  and  $\deg x_{2m_j-1} = 2m_j - 1$ . If each  $x_{2m_j-1}$  is chosen to be integral and not divisible by any other integral classes, we can assign to a representation  $\lambda: G \rightarrow U(n)$  the rational numbers  $a(\lambda, 1), \dots, a(\lambda, l)$  by the equation

$$ch\beta(\lambda) = \sum_{j=1}^l a(\lambda, j) x_{2m_j-1}.$$

In view of [21] and [23], the  $a(\lambda, j)$  are closely related to the *Dynkin coefficients* of  $\lambda$  [14]. On the other hand, as is noted by Atiyah [4, Proposition 1], the determinant of the  $l \times l$  matrix  $(a(\rho_i, j))$  is equal to 1. We remark that for any system of generators  $\{\lambda_1, \dots, \lambda_l\}$  of the ring  $R(G)$ , the determinant of  $(a(\lambda_i, j))$  is also 1.

In this paper, with a suitable system of generators of  $R(G)$ , we shall describe the resulting matrix explicitly for the groups  $G$  with  $l \leq 4$  without using the above informations. Indeed, we deal with the following cases:

$$\begin{aligned} l = 2, \quad G &= \mathrm{SU}(3), & \mathrm{Sp}(2), & & G_2. \\ l = 3, \quad G &= \mathrm{SU}(4), \mathrm{Spin}(7), \mathrm{Sp}(3). \\ l = 4, \quad G &= \mathrm{SU}(5), \mathrm{Spin}(9), \mathrm{Sp}(4), \mathrm{Spin}(8), F_4. \end{aligned}$$

Results are stated in Theorems 2 ( $SU(l+1)$ ), 3 ( $Sp(l)$ ), 4 ( $Spin(7)$ ), 5 ( $Spin(8)$ ), 6 ( $Spin(9)$ ), 7 ( $G_2$ ) and 8 ( $F_4$ ).

The careful reader should notice that "up to sign" is implicitly added to some of the statements of this paper.

For later use we fix some notations. Let  $T$  be a maximal torus of  $G$ . The inclusion  $i: T \rightarrow G$  induces a map of classifying spaces  $\rho = Bi: BT \rightarrow BG$ . The action of the normalizer  $N_G(T)$  on  $T$  induces that of the Weyl group  $\Phi(G) = N_G(T)/T$  on  $BT$  and hence on  $H^*(BT; Z) = Z[\omega_1, \dots, \omega_l]$  (see [9]). Let  $H^*(BT; Z)^{\Phi(G)}$  denote the module of  $\Phi(G)$ -invariants. For a based space  $X$ , let  $\Omega X$  be its loop space, and let  $\sigma^*: H^i(X; Z) \rightarrow H^{i-1}(\Omega X; Z)$  be the cohomology suspension. For the rational cohomology, by [8] and [10] we have

$$\begin{aligned} \text{Im } \rho^* &= H^*(BT; Q)^{\Phi(G)} = Q[f_{2m_1}, \dots, f_{2m_l}] \\ &\cong \downarrow \\ H^*(BG; Q) &= Q[y_{2m_1}, \dots, y_{2m_l}] \\ &\downarrow \sigma^* \\ H^*(G; Q) &= \Lambda_Q(x_{2m_1-1}, \dots, x_{2m_l-1}) \\ &\downarrow \sigma^* \\ H^*(\Omega G; Q) &= Q[u_{2m_1-2}, \dots, u_{2m_l-2}] \end{aligned}$$

where all the generators, whose degrees are indicated by a subscript, are chosen to be integral and not divisible by any other integral classes.

The paper is organized as follows. The key point of our work is to characterize the generator  $x_{2m_j-1}$ . For this purpose we present two methods in Section 1: in the first method we characterize the generator  $y_{2m_j}$  and relate it to  $x_{2m_j-1}$ ; in the second method we characterize the generator  $u_{2m_j-2}$  and relate it to  $x_{2m_j-1}$ . Moreover in Section 1 we prove a lemma which is very useful if the  $\lambda$ -ring structure of  $R(G)$  is known. Subsequent sections are devoted to practical computations. In Section 2 we treat the most elementary cases, i.e.,  $G = SU(l+1)$ ,  $Sp(l)$  ( $l=2, 3, 4$ ) where  $H^*(G; Z)$  has no torsion. In Section 3 we consider the cases  $G = Spin(m)$  ( $m=7, 8, 9$ ) where  $H^*(G; Z)$  has only 2-torsion. In Section 4 we discuss the cases  $G = G_2$  and  $G = F_4$ .

I would like to thank my colleague H. Minami for showing me a computation of  $(a(\rho_i, j))$  for the case  $G = G_2$  and many helpful suggestions.

## 1. Methods

### Method I

For any group  $H$  let  $\alpha: R(H) \rightarrow K^*(BH)$  be the homomorphism of [5]. Let  $\sigma: K^i(X) \rightarrow K^{i-1}(\Omega X)$  be the suspension map. Then there is a commutative diagram

$$\begin{array}{ccccc}
 R(T) & \xrightarrow{\alpha} & K^*(BT) & \xrightarrow{ch} & H^*(BT; Q) \\
 i^* \uparrow & & \rho^* \uparrow & & \rho^* \uparrow \\
 R(G) & \xrightarrow{\alpha} & K^*(BG) & \xrightarrow{ch} & H^*(BG; Q) \\
 & \searrow \beta & \sigma \downarrow & & \sigma^* \downarrow \tau \\
 & & K^*(G) & \xrightarrow{ch} & H^*(G; Q)
 \end{array}
 \quad \begin{array}{c} \leftarrow \tau' \\ \leftarrow \end{array}$$

where  $\tau$  (resp.  $\tau'$ ) is the cohomology transgression in the Serre spectrum of the universal fibration  $G \rightarrow EG \rightarrow BG$  (resp. the fibration  $G \rightarrow G/T \rightarrow BT$ ). For  $j=1, \dots, l$  we may set (modulo decomposables)

$$\sigma^*(y_{2m_j}) = b(m_j)x_{2m_j-1} \quad \text{for some } b(m_j) \in \mathbb{Z}$$

and

$$\rho^*(y_{2m_j}) = c(m_j)f_{2m_j} \quad \text{for some } c(m_j) \in \mathbb{Z}.$$

Since  $\sigma^*$  and  $\tau$  are inverse to each other insofar as they are defined, it follows that

$$\begin{aligned}
 \tau'(x_{2m_j-1}) &= \frac{c(m_j)}{b(m_j)}f_{2m_j} + \text{decomposables} \\
 \text{in } H^*(BT; Q)^{\Phi(G)} &= \mathcal{Q}[f_{2m_1}, \dots, f_{2m_l}].
 \end{aligned}$$

Let  $\lambda: G \rightarrow U(n)$  be a representation with weights  $\mu_1, \dots, \mu_n$ . So

$$ch\alpha i^*(\lambda) = \sum_{i=1}^n \exp(\mu_i) = \sum_{m \geq 0} \sum_{i=1}^n \mu_i^m / m!$$

where  $\mu_i \in H^2(BT; \mathbb{Z})$  (see [9]). Set

$$(1.1) \quad ch\beta(\lambda) = \sum_{j=1}^l a(\lambda, j)x_{2m_j-1} \quad \text{where } a(\lambda, j) \in \mathcal{Q}.$$

Apply  $\tau'$  to this equation. Then the left hand side becomes

$$\begin{aligned}
 \tau' ch\beta(\lambda) &= \rho^* \tau ch\sigma\alpha(\lambda) \\
 &= \rho^* \tau \sigma^* ch\alpha(\lambda) \\
 &= \rho^* ch\alpha(\lambda) \\
 &= ch\alpha i^*(\lambda)
 \end{aligned}$$

and the right hand side becomes

$$\begin{aligned}
 \tau' \left( \sum_{j=1}^l a(\lambda, j)x_{2m_j-1} \right) &= \sum_{j=1}^l a(\lambda, j)\tau'(x_{2m_j-1}) \\
 &= \sum_{j=1}^l \frac{a(\lambda, j)c(m_j)}{b(m_j)}f_{2m_j} + \text{decomposables}.
 \end{aligned}$$

Hence

$$ch\alpha i^*(\lambda) = \sum_{j=1}^l \frac{a(\lambda, j)c(m_j)}{b(m_j)} f_{2m_j} + \text{decomposables}.$$

This argument shows that, in order to compute  $a(\lambda, j)$ , it suffices to settle  $f_{2m_j}$ , determine  $b(m_j)$ ,  $c(m_j)$  and find the coefficients of  $f_{2m_j}$  in the expression of  $ch\alpha i^*(\lambda)$  as a polynomial of the  $f_{2m_j}$ . We will use this method in all cases that concern us.

REMARK. In general we choose the  $f_{2m_j}$  as follows. Let  $\{f'_{2m_1}, \dots, f'_{2m_l}\}$  be a system of generators of the ring  $H^*(BT; Q)^{\Phi(G)}$ . First we take

$$f_{2m_1} = b_1 f'_{2m_1} \in H^{2m_1}(BT; Q)^{\Phi(G)}, \quad b_1 \in Q,$$

so that

(i)  $f_{2m_1}$  is integral;

(ii) for any  $b \in Q$  with  $|b| < |b_1|$ ,  $bf'_{2m_1}$  cannot be integral.

Assume inductively that we have chosen  $f_{2m_1}, \dots, f_{2m_{j-1}}$ . Then we take

$$f_{2m_j} = b_j f'_{2m_j} + \text{decomposables} \in H^{2m_j}(BT; Q)^{\Phi(G)}, \quad b_j \in Q,$$

so that

(i)  $f_{2m_j}$  is integral;

(ii) for any  $b \in Q$  with  $|b| < |b_j|$ ,  $bf'_{2m_j} + \text{decomposables} \in H^{2m_j}(BT; Q)^{\Phi(G)}$  cannot be integral.

Note that the choice of the  $f'_{2m_j}$  has no crucial influence on that of the  $f_{2m_j}$ . As will be seen in Sections 3 and 4, this settlement of the  $f_{2m_j}$  is not trivial but important.

## Method II

There is a commutative diagram

$$\begin{array}{ccccc} R(G) & \xrightarrow{\beta} & K^*(G) & \xrightarrow{ch} & H^*(G; Q) \\ & & \sigma \downarrow & & \sigma^* \downarrow \\ & & K^*(\Omega G) & \xrightarrow{ch} & H^*(\Omega G; Q) \end{array}$$

which is natural with respect to group homomorphisms. For  $j=1, \dots, l$  we may set

$$\sigma^*(x_{2m_j-1}) = d(m_j)u_{2m_j-2} \quad \text{for some } d(m_j) \in Z.$$

Applying  $\sigma^*$  to (1.1), we have

$$ch\sigma\beta(\lambda) = \sum_{j=1}^l a(\lambda, j)d(m_j)u_{2m_j-2}.$$

Let us now consider the case  $G = \text{SU}(n+1)$ ; then  $m_j = j+1$  for  $j=1, \dots, n$  and

$$PH^*(\Omega SU(n+1); Z) = Z\{u_{2i} | 1 \leq i \leq n\}$$

where  $P$  denotes the primitive module functor. Furthermore,  $d(j+1)=1$  for all  $j$  (e.g., see [28, Lemma 3]). Let  $\lambda_1: SU(n+1) \rightarrow U(n+1)$  be the natural inclusion, and consider the case  $\lambda=\lambda_1$ . Then it follows from (2.2) of the next section that

$$(1.2) \quad ch\sigma\beta(\lambda_1) = \sum_{i=1}^n \frac{(-1)^i}{i!} u_{2i}.$$

We return to the general case. Take the inclusion  $k: U(n) \rightarrow SU(n+1)$  such that  $SU(n+1)/U(n) = CP^n$  (see [12, §3]). In [28] it was shown that for the composite

$$\begin{aligned} PH^*(\Omega SU(n+1); Z) &\xrightarrow{(\Omega k)^*} PH^*(\Omega U(n); Z) \\ &\xrightarrow{(\Omega \lambda)^*} PH^*(\Omega G; Z) = Z\{u_{2m_1-2}, \dots, u_{2m_l-2}\}, \end{aligned}$$

the following statements are equivalent:

- (i)  $(\Omega \lambda)^*(\Omega k)^*(u_{2m_j-2}) = e(\lambda, j) u_{2m_j-2}$  for some  $e(\lambda, j) \in Z$ ;
- (ii) the element  $\theta_s(c_{m_j}(\lambda)) \in H^{2m_j-2}(G/C_s; Z)$  is exactly divisible by  $e(\lambda, j) \in Z$  (where  $H^*(G/C_s; Z)$  has no torsion; for notations and details see [28, §2]).

Applying  $(\Omega \lambda^*)(\Omega k)^*$  to (1.2), we have

$$ch\sigma\beta(\lambda) = \sum_{j=1}^l \frac{(-1)^{m_j-1} e(\lambda, j)}{(m_j-1)!} u_{2m_j-2}.$$

Hence

$$a(\lambda, j) d(m_j) = \frac{(-1)^{m_j-1} e(\lambda, j)}{(m_j-1)!}.$$

This argument shows that, in order to compute  $a(\lambda, j)$ , it suffices to determine  $d(m_j)$  and  $e(\lambda, j)$ . In particular, to find  $e(\lambda, j)$  one must examine the divisibility of  $\theta_s(c_{m_j}(\lambda))$  in  $H^{2m_j-2}(G/C_s; Z)$ .

Define a map  $\varphi: Z_+ \times Z_+ \times Z_+ \rightarrow Z$  by

$$\varphi(n, k, q) = \sum_{i=1}^k (-1)^{i-1} \binom{n}{k-i} i^{q-1}$$

where  $Z_+$  denotes the set of positive integers and we use the convention that  $\binom{x}{y} = 0$  if  $y < 0$  or  $x < y$ . Let  $\Lambda^k: R(G) \rightarrow R(G)$  be the  $k$ -th exterior power operation. Then we have

**Lemma 1.** *If  $\lambda$  is a representation of  $G$  of dimension  $n$ , then*

$$a(\Lambda^k \lambda, j) = \varphi(n, k, m_j) a(\lambda, j)$$

for  $j=1, \dots, l$ .

Proof. Let  $ch^q$  be the  $2q$ -th component of  $ch$ , i.e.,  $ch(x) = \sum_{i \geq 0} ch^i(x)$  with  $ch^q(x) \in H^{2q}(X; \mathcal{Q})$  for any  $x \in K^0(X)$ . Consider the element  $1_n \in R(U(n))$  which comes from the identity  $1_{U(n)}: U(n) \rightarrow U(n)$ . Then we assert that

$$(1.3) \quad \begin{aligned} ch^q \alpha(\Lambda^k 1_n) &= \varphi(n, k, q) ch^q \alpha(1_n) + \text{decomposables} \\ \text{in } H^*(BU(n); \mathcal{Q}) &= \mathcal{Q}[y_2, y_4, \dots, y_{2n}]. \end{aligned}$$

This assertion implies the result. For since  $\beta = \sigma \alpha$  and  $\sigma^*$  sends a decomposable element into zero, applying  $\sigma^*$  to (1.3) yields the desired result for the case  $G = U(n)$ . Then the general case follows from naturality.

To prove (1.3) we proceed by induction on  $k$ . The case  $k=1$  is clear. Suppose that it is true for  $k \leq m-1$ , and consider the case  $k=m$ . Let us recall the following relations:

$$\begin{aligned} \psi^k(x) + \sum_{i=1}^{k-1} (-1)^i \psi^{k-i}(x) \Lambda^i(x) + (-1)^k k \Lambda^k(x) &= 0; \\ ch^q(xy) &= \sum_{r=0}^q ch^r(x) ch^{q-r}(y); \\ ch^q \psi^k(x) &= k^q ch^q(x) \end{aligned}$$

where  $x, y \in K^0(X)$  [1]. Since  $\alpha$  is a  $\lambda$ -ring homomorphism, we have

$$\begin{aligned} ch^q \alpha(m \Lambda^m(1_n)) &= ch^q \alpha((-1)^{m-1} \psi^m(1_n) + \sum_{i=1}^{m-1} (-1)^{m-1-i} \psi^{m-i}(1_n) \Lambda^i(1_n)) \\ &= (-1)^{m-1} ch^q \alpha \psi^m(1_n) + \sum_{i=1}^{m-1} (-1)^{m-1-i} ch^q(\alpha \psi^{m-i}(1_n) \alpha \Lambda^i(1_n)) \\ &= (-1)^{m-1} ch^q \alpha \psi^m(1_n) + \sum_{i=1}^{m-1} (-1)^{m-1-i} \left[ \sum_{r=0}^q ch^r \alpha \psi^{m-i}(1_n) ch^{q-r} \alpha \Lambda^i(1_n) \right] \\ &= (-1)^{m-1} ch^q \alpha \psi^m(1_n) + \sum_{i=1}^{m-1} (-1)^{m-1-i} \left[ \binom{n}{i} ch^q \alpha \psi^{m-i}(1_n) + n ch^q \alpha \Lambda^i(1_n) \right] \\ &\quad \text{modulo decomposables} \\ &= (-1)^{m-1} ch^q \alpha \psi^m(1_n) + \sum_{i=1}^{m-1} (-1)^{m-1-i} \left[ \binom{n}{i} ch^q \psi^{m-i} \alpha(1_n) + n ch^q \alpha(\Lambda^i 1_n) \right] \\ &= (-1)^{m-1} m^q ch^q \alpha(1_n) + \sum_{i=1}^{m-1} (-1)^{m-1-i} \left[ \binom{n}{i} (m-i)^q ch^q \alpha(1_n) \right. \\ &\quad \left. + n \varphi(n, i, q) ch^q \alpha(1_n) \right] \\ &= \left[ \sum_{i=0}^{m-1} (-1)^{m-1-i} \binom{n}{i} (m-i)^q + \sum_{i=1}^{m-1} (-1)^{m-1-i} n \varphi(n, i, q) \right] ch^q \alpha(1_n) \end{aligned}$$

$$= [\sum_{j=1}^m (-1)^{j-1} \binom{n}{m-j} j^q + n \sum_{i=1}^{m-1} (-1)^{m-1-i} \varphi(n, i, q)] ch^q \alpha(1_n).$$

Thus it is sufficient to prove that

$$(1.4) \quad \varphi(n, m, q+1) + n \sum_{i=1}^{m-1} (-1)^{m-1-i} \varphi(n, i, q) = m \varphi(n, m, q).$$

From Pascal's triangle

$$\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$$

we deduce that

$$\sum_{i=0}^{k-1-j} (-1)^i \binom{n}{i} = (-1)^{k-1-j} \binom{n-1}{k-1-j}.$$

Using this, we have

$$\begin{aligned} \varphi(n-1, m-1, q) &= \sum_{j=1}^{m-1} (-1)^{j-1} \binom{n-1}{m-1-j} j^{q-1} \\ &= \sum_{j=1}^{m-1} [(-1)^m \sum_{i=0}^{m-1-j} (-1)^i \binom{n}{i}] j^{q-1} \\ &= \sum_{i=1}^{m-1} (-1)^{m-1-i} [\sum_{j=1}^i (-1)^{j-1} \binom{n}{i-j} j^{q-1}] \\ &= \sum_{i=1}^{m-1} (-1)^{m-1-i} \varphi(n, i, q). \end{aligned}$$

Therefore

$$\begin{aligned} &n \varphi(n-1, m-1, q) + \varphi(n, m, q+1) \\ &= n \sum_{j=1}^{m-1} (-1)^{j-1} \binom{n-1}{m-1-j} j^{q-1} + \sum_{j=1}^m (-1)^{j-1} \binom{n}{m-j} j^q \\ &= \sum_{j=1}^{m-1} (-1)^{j-1} n \binom{n-1}{m-1-j} j^{q-1} + \sum_{j=1}^m (-1)^{j-1} \binom{n}{m-j} j^q \\ &= \sum_{j=1}^{m-1} (-1)^{j-1} \binom{n}{m-j} (m-j) j^{q-1} + \sum_{j=1}^m (-1)^{j-1} \binom{n}{m-j} j^q \\ &= \sum_{j=1}^{m-1} (-1)^{j-1} \binom{n}{m-j} m j^{q-1} - \sum_{j=1}^{m-1} (-1)^{j-1} \binom{n}{m-j} j^q \\ &\quad + \sum_{j=1}^m (-1)^{j-1} \binom{n}{m-j} j^q \\ &= m \sum_{j=1}^{m-1} (-1)^{j-1} \binom{n}{m-j} j^{q-1} + (-1)^{m-1} \binom{n}{0} m^q \\ &= m \sum_{j=1}^m (-1)^{j-1} \binom{n}{m-j} j^{q-1} \\ &= m \varphi(n, m, q). \end{aligned}$$

This proves (1.4) and completes the proof.



## 2. The special unitary groups and the symplectic groups

Let us first consider the case of  $SU(l+1)$ . In this case,  $m_j = j+1$  for  $j=1, \dots, l$ . As is well known we can choose elements  $t_1, t_2, \dots, t_{l+1} \in H^2(BT; Z)$  so that

$$H^*(BT; Z) = Z[t_1, \dots, t_{l+1}]/(c_1)$$

and

$$H^*(BT; Z)^{\Phi(SU(l+1))} = Z[c_2, \dots, c_{l+1}]$$

where  $c_i = \sigma_i(t_1, \dots, t_{l+1})$  ( $\sigma_i(\cdot)$  denotes the  $i$ -th elementary symmetric function). It is evident that  $f_{2j+2} = c_{j+1}$  for  $j=1, \dots, l$ . Since  $H^*(SU(l+1); Z)$  has no torsion, the theorem of Borel [6] assures us that  $b(j+1) = c(j+1) = 1$  for all  $j$ . Thus we have  $\tau'(x_{2j+1}) = c_{j+1}$  for  $j=1, \dots, l$ .

Let us recall from [17] that

$$(2.1) \quad R(SU(l+1)) = Z[\lambda_1, \lambda_2, \dots, \lambda_l] \quad \text{where}$$

- (a)  $\dim \lambda_k = \binom{l+1}{k}$ ;
- (b) relations  $\Lambda^t \lambda_1 = \lambda_k$  hold;
- (c) the set of weights of  $\lambda_1$  is given by  $\{t_i \mid 1 \leq i \leq l+1\}$ .

Put

$$s_m = s_m(t_1, \dots, t_{l+1}) = \sum_{i=1}^{l+1} t_i^m.$$

From Newton's formula

$$s_m + \sum_{i=1}^{m-1} (-1)^i s_{m-i} c_i + (-1)^m m c_m = 0$$

(where  $c_m = 0$  if  $m > l+1$ ) it follows that

$$ch\alpha^*(\lambda_1) = l+1 + \sum_{m=1}^l \frac{(-1)^m}{m!} c_{m+1} + \text{decomposables.}$$

Therefore

$$(2.2) \quad ch\beta(\lambda_1) = \sum_{m=1}^l \frac{(-1)^m}{m!} x_{2m+1}$$

(cf. [20, Theorem 1]). By Lemma 1, if we evaluate  $\varphi(l+1, k, j+1)$ ,  $ch\beta(\lambda_k)$  can be calculated. Thus we have

**Theorem 2.** *The Chern characters on  $SU(l+1)$  for  $l=2,3,4$  are given by:*

$$\begin{aligned} l=2 \quad ch\beta(\lambda_1) &= -x_3 + (1/2!)x_5 \\ ch\beta(\lambda_2) &= -x_3 + (-1/2!)x_5 \end{aligned}$$

$$\begin{array}{ll}
l=3 & \begin{array}{l}
ch\beta(\lambda_1) = -x_3 + (1/2!)x_5 + (-1/3!)x_7 \\
ch\beta(\lambda_2) = -2x_3 + (4/3!)x_7 \\
ch\beta(\lambda_3) = -x_3 + (-1/2!)x_5 + (-1/3!)x_7
\end{array} & -1 \\
l=4 & \begin{array}{l}
ch\beta(\lambda_1) = -x_3 + (1/2!)x_5 + (-1/3!)x_7 + (1/4!)x_9 \\
ch\beta(\lambda_2) = -3x_3 + (1/2!)x_5 + (3/3!)x_7 + (-11/4!)x_9 \\
ch\beta(\lambda_3) = -3x_3 + (-1/2!)x_5 + (3/3!)x_7 + (11/4!)x_9 \\
ch\beta(\lambda_4) = -x_3 + (-1/2!)x_5 + (-1/3!)x_7 + (-1/4!)x_9
\end{array} & 1
\end{array}$$

where the number on the right hand side indicates the determinant of the corresponding matrix on the left hand side.

Let us consider the case of  $Sp(l)$ . In this case,  $m_j = 2j$  for  $j = 1, \dots, l$ . We can choose elements  $t_1, t_2, \dots, t_l \in H^2(BT; Z)$  so that

$$H^*(BT; Z) = Z[t_1, \dots, t_l]$$

and

$$H^*(BT; Z)^{\oplus(Sp(l))} = Z[q_1, \dots, q_l]$$

where  $q_i = \sigma_i(t_1^2, \dots, t_l^2)$ . It is evident that  $f_{4j} = q_j$  for  $j = 1, \dots, l$ . Since  $H^*(Sp(l); Z)$  has no torsion, it follows that  $b(2j) = c(2j) = 1$  for all  $j$ . Thus we have  $\tau'(x_{4j-1}) = q_j$  for  $j = 1, \dots, l$ .

Let us recall that

$$(2.3) \quad R(Sp(l)) = Z[\lambda_1, \lambda_2, \dots, \lambda_l] \quad \text{where}$$

- (a)  $\dim \lambda_k = \binom{2l}{k}$ ;
- (b) relations  $\Lambda^k \lambda_1 = \lambda_k$  hold;
- (c) the set of weights of  $\lambda_1$  is given by  $\{\pm t_i \mid 1 \leq i \leq l\}$ .

Put

$$s_{2m} = s_m(t_1^2, \dots, t_l^2) = \sum_{i=1}^l t_i^{2m}.$$

From Newton's formula

$$s_{2m} + \sum_{i=1}^{m-1} (-1)^i s_{2m-2i} q_i + (-1)^m m q_m = 0$$

it follows that

$$ch\alpha i^*(\lambda_1) = 2l + \sum_{m=1}^l \frac{(-1)^{m-1}}{(2m-1)!} q_m + \text{decomposables}.$$

Therefore

$$ch\beta(\lambda_1) = \sum_{m=1}^l \frac{(-1)^{m-1}}{(2m-1)!} x_{4m-1}$$

and by Lemma 1 we obtain

**Theorem 3.** *The Chern characters on  $Sp(l)$  for  $l=2, 3, 4$  are given by:*

$$\begin{array}{ll}
 l=2 & \begin{array}{l} ch\beta(\lambda_1) = x_3 + (-1/3!)x_7 \\ ch\beta(\lambda_2) = 2x_3 + (4/3!)x_7 \end{array} & 1 \\
 l=3 & \begin{array}{l} ch\beta(\lambda_1) = x_3 + (-1/3!)x_7 + (1/5!)x_{11} \\ ch\beta(\lambda_2) = 4x_3 + (2/3!)x_7 + (-26/5!)x_{11} \\ ch\beta(\lambda_3) = 6x_3 + (6/3!)x_7 + (66/5!)x_{11} \end{array} & 1 \\
 l=4 & \begin{array}{l} ch\beta(\lambda_1) = x_3 + (-1/3!)x_7 + (1/5!)x_{11} + (-1/7!)x_{15} \\ ch\beta(\lambda_2) = 6x_3 + (-24/5!)x_{11} + (120/7!)x_{15} \\ ch\beta(\lambda_3) = 15x_3 + (9/3!)x_7 + (15/5!)x_{11} + (-1191/7!)x_{15} \\ ch\beta(\lambda_4) = 20x_3 + (16/3!)x_7 + (80/5!)x_{11} + (2416/7!)x_{15} \end{array} & 1
 \end{array}$$

where the number on the right hand side indicates the determinant of the corresponding matrix on the left hand side.

### 3. The spinor groups

Let us first consider the case of  $Spin(7)$ . In this case,  $(m_1, m_2, m_3) = (2, 4, 6)$ . We can choose elements  $t_1, t_2, t_3, \gamma \in H^2(BT; Z)$  so that

$$H^*(BT; Z) = Z[t_1, t_2, t_3, \gamma]/(c_1 - 2\gamma)$$

and

$$H^*(BT; Q)^{\Phi(Spin(7))} = Q[p_1, p_2, p_3]$$

where  $c_i = \sigma_i(t_1, t_2, t_3)$  and  $p_i = \sigma_i(t_1^2, t_2^2, t_3^2)$ . In the light of the Remark in Section 1, using the formula

$$p_i = \sum_{j=0}^{2i} (-1)^{i+j} c_{2i-j} c_j,$$

we have

$$\begin{aligned}
 (3.1) \quad f_4 &= \frac{1}{2} p_1 = -c_2 + 2\gamma^2, \\
 f_8 &= \frac{1}{4} p_2 - \frac{1}{4} f_4^2 = -c_3\gamma + c_2\gamma^2 - \gamma^4, \\
 f_{12} &= p_3 = c_3^2.
 \end{aligned}$$

Let us determine  $b(2), b(4), b(6) \in Z$ . To do so we use the Serre spectral sequence  $\{E_r(Z)\}$  for the integral cohomology of the universal fibration

$$F = Spin(7) \rightarrow E = E Spin(7) \rightarrow B = B Spin(7).$$

Furthermore, to investigate it, we use the Serre spectral sequence  $\{E_r(Z/p)\}$  for the mod  $p$  cohomology of the same fibration, where  $p$  runs over all primes.

Recall that  $H^*(\text{Spin}(7); Z)$  has no  $p$ -torsion for  $p > 2$ . Let  $\Delta_{Z/2}(\ )$  denote a  $Z/2$ -algebra having a set in parentheses as a simple system of generators. Then it follows from [6] and [7] that

$$H^*(\text{Spin}(7); Z/p) = \begin{cases} \Delta_{Z/2}(x_3, x_5, x_6, x_7) & (p = 2) \\ \Delta_{Z/p}(x_3, x_7, x_{11}) & (p > 2) \end{cases}$$

and

$$H^*(B\text{Spin}(7); Z/p) = \begin{cases} Z/2[\bar{y}_4, \bar{y}_6, \bar{y}_7, \bar{y}_8] & (p = 2) \\ Z/p[\bar{y}_4, \bar{y}_8, \bar{y}_{12}] & (p > 2) \end{cases}$$

where  $x_i$  transgresses to  $\bar{y}_{i+1}$  for all  $i$  and  $\beta_2(x_5) = x_6$  ( $\beta_p$  denotes the mod  $p$  Bockstein homomorphism). For a based space  $X$ , let  $\pi_p: H^i(X; Z) \rightarrow H^i(X; Z/p)$  be the mod  $p$  reduction homomorphism. Then if  $i = 3$  or  $7$ ,  $\pi_p(x_i) = x_i$  and  $\pi_p(y_{i+1}) = \bar{y}_{i+1}$  for every prime  $p$ . Therefore we conclude that  $\tau(x_3) = y_4$  and  $\tau(x_7) = y_8$ . In other words,  $b(2) = b(4) = 1$ .

It remains to determine  $b(6)$ . Since

$$\pi_p(x_{11}) = \begin{cases} x_5 x_6 & (p = 2) \\ x_{11} & (p > 2) \end{cases} \quad \text{and} \quad \pi_p(y_{12}) = \begin{cases} \bar{y}_6^2 & (p = 2) \\ \bar{y}_{12} & (p > 2) \end{cases},$$

an analogous argument to the above yields that

$$(0) \quad \text{if } p > 2, \quad v_p(b(6)) = 0$$

where  $v_p(m)$  is the power of  $p$  in  $m$ . To get  $v_2(b(6))$  we consider  $\{E_r(Z/2)\}$ , which satisfies

$$E_2^{s,t}(Z/2) \cong H^s(B; Z/2) \otimes H^t(F; Z/2)$$

and  $E_\infty^{s,t}(Z/2) = 0$  unless  $(s, t) = (0, 0)$ . Then it is easy to see that

- (i)  $d_6(1 \otimes x_5 x_6) = \bar{y}_6 \otimes x_6$ .
- (ii)  $d_6(\bar{y}_6 \otimes x_5) = \bar{y}_6^2 \otimes 1$ .

Let

$$\beta_2^F: E_1^{s,t}(Z/2) \rightarrow E_1^{s,t+1}(Z/2)$$

be the map induced by  $\beta_2: H^t(F; Z/2) \rightarrow H^{t+1}(F; Z/2)$  through the isomorphism

$$E_1^{s,t}(Z/2) \cong C^s(B; H^t(F; Z/2)).$$

Then we have

$$(iii) \quad \beta_2^F(\bar{y}_6 \otimes x_5) = \bar{y}_6 \otimes x_6.$$

Denote again by  $\pi_p: \{E_r(Z)\} \rightarrow \{E_r(Z/p)\}$  the morphism of spectral sequences induced by  $\pi_p$ . By virtue of the isomorphism

$$E_2^{s,t}(Z) \cong H^s(B; H^t(F; Z)),$$

we find that there exist elements  $\{x_{11}\} \in E_2^{0,11}(Z)$ ,  $\{v_{12}\} \in E_2^{6,6}(Z)$  and  $\{y_{12}\} \in E_2^{12,0}(Z)$  which satisfy  $\pi_2(\{x_{11}\}) = 1 \otimes \bar{x}_5 \bar{x}_6$ ,  $\pi_2(\{v_{12}\}) = \bar{y}_6 \otimes \bar{x}_6$  and  $\pi_2(\{y_{12}\}) = \bar{y}_6^2 \otimes 1$  respectively. Then the conditions (0), (i), (ii), (iii) imply that in  $\{E_r(Z)\}$

$$(iv) \quad d_6(\{x_{11}\}) = \{v_{12}\}.$$

$$(v) \quad d_{12}(\{2x_{11}\}) = \{y_{12}\}.$$

In fact, (iv) is an immediate consequence of (i). In what follows we roughly state a proof of (v). Let us begin by recalling the construction of the Serre spectral sequence  $\{E_r(R)\}$  in cohomology with  $R$ -coefficients of a fibration  $F \rightarrow E \rightarrow B$ , where  $R = Z$  or  $Z/p$  (for details see [24]). There is a cochain complex  $\text{Hom}(C_*(E), R)$  which is filtered by its subcomplexes  $A^s(R) = \sum_t A^{s,t}(R)$  such that  $A^{s,t}(R) \subset A^{s-1,t+1}(R)$  and  $\delta(A^{s,t}(R)) \subset A^{s,t+1}(R)$  for all  $(s, t)$  (where  $\delta$  is the differential in  $\text{Hom}(C_*(E), R)$ ). This filtered cochain complex gives rise to  $\{E_r(R)\}$ , i.e.,

$$Z_r^{s,t}(R) = A^{s,t}(R) \cap \delta^{-1}(A^{s+r,t-r+1}(R)),$$

$$B_r^{s,t}(R) = A^{s,t}(R) \cap \delta A^{s-r,t+r-1}(R),$$

$$E_r^{s,t}(R) = Z_r^{s,t}(R) / (Z_{r-1}^{s+1,t-1}(R) + B_{r-1}^{s,t}(R)).$$

Note that there is an exact sequence

$$0 \rightarrow A^{s,t}(Z) \xrightarrow{\cdot p} A^{s,t}(Z) \xrightarrow{\pi_p} A^{s,t}(Z/p) \rightarrow 0$$

for all  $(s, t)$ . Since  $d_r: E_r^{s,t}(R) \rightarrow E_r^{s+r,t-r+1}(R)$  is induced by  $\delta$ , by (iv) we see that there exists a representative  $x \in A^{0,11}(Z)$  (resp.  $v \in A^{6,6}(Z)$ ) of  $\{x_{11}\}$  (resp.  $\{v_{12}\}$ ) such that

$$(3.2) \quad \delta(x) = v.$$

Let  $u \in A^{6,5}(Z/2)$  be a representative of  $\bar{y}_6 \otimes \bar{x}_5$ . Then by (iii) we observe that there exists  $u \in A^{6,5}(Z)$  such that  $\pi_2(u) = \bar{u}$  and

$$(3.3) \quad \delta(u) = 2v$$

(see [2, Chapter III, §2]). Similarly by (ii) there is a representative  $\bar{y} \in A^{12,0}(Z/2)$  of  $\bar{y}_6^2 \otimes 1$  such that  $\delta(\bar{u}) = \bar{y}$ . This implies that there exists a representative  $y \in A^{12,0}(Z)$  of  $\{y_{12}\}$  such that  $\pi_2(y) = \bar{y}$  and

$$(3.4) \quad \delta(y) = y.$$

By (3.2), (3.3) and (3.4), we have

$$\delta(2x) = 2v = \delta(u) = y$$

which gives (v). It is equivalent to  $b(6) = 2$ .

We discuss the problem of determining  $c(2)$ ,  $c(4)$ ,  $c(6) \in Z$  in a general form. Indeed, we claim that  $c(m_j)=1$  for  $j=1, \dots, l$  in all cases that concern us. To prove this we use the integral cohomology spectral sequence  $\{E_r\}$  of the fibration

$$G/T \rightarrow BT \xrightarrow{\rho} BG.$$

Then the homomorphism  $\rho^*: H^m(BG; Z) \rightarrow H^m(BT; Z)$  can be regarded as the composite

$$H^m(BG; Z) = E_2^{m,0} \rightarrow E_\infty^{m,0} = D^{m,0} \subset \dots \subset D^{0,m} = H^m(BT; Z)$$

where  $D^{i,m-i}/D^{i+1,m-i-1} = E_\infty^{i,m-i}$ . According to [6], the class  $\{y_{2m_j}\} \in E_2^{2m_j,0}$  survives to  $E_\infty$ . What we have to verify is to observe that no extension problems occur on the class  $\{y_{2m_j}\} \in E_\infty^{2m_j,0}$ . This is an essentially easy work, because all structures of  $H^*(G/T; Z)$ ,  $H^*(BT; Z)$  and  $H^*(BG; Z)$  were explicitly described (for  $H^*(BG; Z)$  see [7] and [25]; for  $H^*(G/T; Z)$  see [27] and also [26]). For example, consider the case  $G = \text{Spin}(7)$ . Then it is not hard to see that if  $m=4, 8$  or  $12$ ,  $E_\infty^{i,m-i}$  is trivial or torsion free for all  $i$ . This assures us that  $c(2)=c(4)=c(6)=1$ . In the future we omit such checks for the other cases, for our claim (except for the case  $G=F_4$ ) has been proved in a more general setting by [13] and [15].

Let us recall from [17] that

$$(3.5) \quad R(\text{Spin}(7)) = Z[\lambda'_1, \lambda'_2, \Delta_7] \quad \text{where}$$

- (a)  $\dim \lambda'_k = \binom{7}{k}$  and  $\dim \Delta_7 = 8$ ;
- (b) relations  $\Lambda^k \lambda'_1 = \lambda'_k$  and  $\Delta_7^2 = \lambda'_3 + \lambda'_2 + \lambda'_1 + 1$  hold;
- (c) the set of weights of  $\lambda'_1$  is given by  $\{\pm t_i, 0 \mid 1 \leq i \leq 3\}$ .

By the same calculation as in the case of  $Sp(l)$ , we have

$$\begin{aligned} ch^2 \alpha i^*(\lambda'_1) &= p_1, \\ ch^4 \alpha i^*(\lambda'_1) &= -\frac{1}{6} p_2 + \text{decomposables}, \\ ch^6 \alpha i^*(\lambda'_1) &= \frac{1}{120} p_3 + \text{decomposables}. \end{aligned}$$

On the other hand, from (3.1) and the results on  $b(m_j)$  and  $c(m_j)$  it follows that

$$\begin{aligned} \tau'(x_3) &= f_4 = \frac{1}{2} p_1, \\ \tau'(x_7) &= f_8 = \frac{1}{4} p_2 + \text{decomposables}, \end{aligned}$$

$$\tau'(x_{11}) = \frac{1}{2}f_{12} = \frac{1}{2}p_3.$$

Combining these, we have

$$ch\beta(\lambda'_1) = 2x_3 - \frac{2}{3}x_7 + \frac{1}{60}x_{11}.$$

Therefore by Lemma 1,

$$ch\beta(\lambda'_2) = 10x_3 + \frac{2}{3}x_7 - \frac{5}{12}x_{11}$$

and

$$ch\beta(\lambda'_3 + \lambda'_2 + \lambda'_1 + 1) = 32x_3 + \frac{16}{3}x_7 + \frac{4}{15}x_{11}.$$

On the other hand, by the formula (2) of [16, p. 8],

$$\beta(\Delta_7^2) = 8\beta(\Delta_7) + 8\beta(\Delta_7) = 16\beta(\Delta_7).$$

Thus from the relation  $\Delta_7^2 = \lambda'_3 + \lambda'_2 + \lambda'_1 + 1$  we deduce that

$$ch\beta(\Delta_7) = 2x_3 + \frac{1}{3}x_7 + \frac{1}{60}x_{11}.$$

**Theorem 4.** *The Chern characters on  $Spin(7)$  are given by:*

$$ch\beta(\lambda'_1) = 2x_3 + (-4/3!)x_7 + (2/5!)x_{11}$$

$$ch\beta(\lambda'_2) = 10x_3 + (4/3!)x_7 + (-50/5!)x_{11}$$

$$ch\beta(\Delta_7) = 2x_3 + (2/3!)x_7 + (2/5!)x_{11}$$

and the determinant of the corresponding matrix is 1.

Let us next consider the case of  $Spin(8)$ . In this case,  $(m_1, m_2, m_3, m_4) = (2, 4, 4, 6)$ . We can choose elements  $t_1, t_2, t_3, t_4, \gamma \in H^2(BT; Z)$  so that

$$H^*(BT; Z) = Z[t_1, \dots, t_4, \gamma]/(c_1 - 2\gamma)$$

and

$$H^*(BT; Q)^{\Phi(Spin(8))} = Q[p_1, c_4, p_2, p_3]$$

where  $c_i = \sigma_i(t_1, \dots, t_4)$  and  $p_i = \sigma_i(t_1^2, \dots, t_4^2)$ . By a similar calculation to the before, we have

$$f_4 = \frac{1}{2}p_1 = -c_2 + 2\gamma^2,$$

$$f'_8 = c_4,$$

$$f_8 = \frac{1}{4}p_2 - \frac{1}{2}f'_8 - \frac{1}{4}f_4^2 = -c_3\gamma + c_2\gamma^2 - \gamma^4,$$

$$f_{12} = p_3 = -2c_4c_2 + c_3^2.$$

Let us determine  $b(2)$ ,  $b(4)'$ ,  $b(4)$ ,  $b(6) \in Z$ . But, since  $H^*(\text{Spin}(8); Z)$  has no  $p$ -torsion for  $p > 2$  and

$$H^*(\text{Spin}(8); Z/2) = \Delta_{Z/2}(\mathfrak{x}_3, \mathfrak{x}_5, \mathfrak{x}_6, \mathfrak{x}'_7, \mathfrak{x}_7)$$

where all the  $\mathfrak{x}_i$  are universally transgressive and  $\beta_2(\mathfrak{x}_5) = \mathfrak{x}_6$  [7], the situation is quite similar to that for  $G = \text{Spin}(7)$ , and so we get a similar result, i.e.,  $b(2) = b(4)' = b(4) = 1$  and  $b(6) = 2$ . On the other hand, as mentioned earlier,  $c(2) = c(4)' = c(4) = c(6) = 1$ . Thus we have

$$\begin{aligned} (3.6) \quad \tau'(x_3) &= f_4 = \frac{1}{2}p_1, \\ \tau'(x'_7) &= f'_8 = c_4, \\ \tau'(x_7) &= f_8 = \frac{1}{4}p_2 - \frac{1}{2}c_4 + \text{decomposables}, \\ \tau'(x_{11}) &= \frac{1}{2}f_{12} = \frac{1}{2}p_3. \end{aligned}$$

Let us recall from [17] that

- (3.7)  $R(\text{Spin}(8)) = Z[\lambda_1, \lambda_2, \Delta_8^+, \Delta_8^-]$  where
- (a)  $\dim \lambda_k = \binom{8}{k}$  and  $\dim \Delta_8^+ = \dim \Delta_8^- = 8$ ;
  - (b) relations  $\Lambda^k \lambda_1 = \lambda_k$  and  $\Delta_8^+ \Delta_8^- = \lambda_3 + \lambda_1$  hold;
  - (c) the set of weights of  $\lambda_1$  is given by  $\{\pm t_i \mid 1 \leq i \leq 4\}$  and that of  $\Delta_8^+$  is given by  $\{\pm \gamma, \gamma - t_i - t_j \mid 1 \leq i < j \leq 4\}$ .

By direct calculations we have

$$\begin{aligned} (3.8) \quad ch^2 \alpha i^*(\lambda_1) &= p_1, \\ ch^4 \alpha i^*(\lambda_1) &= \frac{1}{12}(-2p_2 + p_1^2), \\ ch^6 \alpha i^*(\lambda_1) &= \frac{1}{360}(3p_3 - 3p_2p_1 + p_1^3) \end{aligned}$$

and

$$\begin{aligned} (3.9) \quad ch^2 \alpha i^*(\Delta_8^+) &= p_1, \\ ch^4 \alpha i^*(\Delta_8^+) &= \frac{1}{48}(4p_2 + 24c_4 + p_1^2). \end{aligned}$$

There are involutive automorphisms  $\kappa$  and  $\tilde{\kappa}$  of  $T$  and  $\text{Spin}(8)$  respectively, which make the diagram



$$\begin{array}{ccc}
 T & \xrightarrow{\kappa} & T \\
 i \downarrow & & \downarrow i \\
 \text{Spin}(8) & \xrightarrow{\tilde{\kappa}} & \text{Spin}(8)
 \end{array}$$

commute, such that the automorphism  $(B\kappa)^*$  of  $H^*(BT; Z)$  satisfies

$$(B\kappa)^*(t_i) = \begin{cases} t_i & (1 \leq i \leq 3) \\ -t_4 & (i = 4). \end{cases}$$

Therefore  $(B\kappa)^*(p_i) = p_i$ ,  $(B\kappa)^*(c_4) = -c_4$  and the automorphism  $\tilde{\kappa}^*$  of  $R(\text{Spin}(8))$  satisfies  $\tilde{\kappa}^*(\Delta_8^+) = \Delta_8^-$ . Applying  $(B\kappa)^*$  to (3.9), it follows that

$$\begin{aligned}
 (3.10) \quad ch^2 \alpha^*(\Delta_8^-) &= p_1, \\
 ch^4 \alpha^*(\Delta_8^-) &= \frac{1}{48}(4p_2 - 24c_4 + p_1^2).
 \end{aligned}$$

Combining (3.8), (3.9), (3.10) with (3.6), we have

$$\begin{aligned}
 ch\beta(\lambda_1) &= 2x_3 - \frac{1}{3}x'_7 - \frac{2}{3}x_7 + \frac{1}{60}x_{11}, \\
 ch\beta(\Delta_8^+) &= 2x_3 + \frac{2}{3}x'_7 + \frac{1}{3}x_7 + ax_{11}, \\
 ch\beta(\Delta_8^-) &= 2x_3 - \frac{1}{3}x'_7 + \frac{1}{3}x_7 + ax_{11}
 \end{aligned}$$

for some  $a \in Q$ . From Lemma 1 and the relation  $\Delta_8^+ \Delta_8^- = \lambda_3 + \lambda_1$  we deduce that  $a = 1/60$ .

**Theorem 5.** *The Chern characters on  $\text{Spin}(8)$  are given by:*

$$\begin{aligned}
 ch\beta(\lambda_1) &= 2x_3 + (-2/3!)x'_7 + (-4/3!)x_7 + (2/5!)x_{11} \\
 ch\beta(\lambda_2) &= 12x_3 & + (-48/5!)x_{11} \\
 ch\beta(\Delta_8^+) &= 2x_3 + (4/3!)x'_7 + (2/3!)x_7 + (2/5!)x_{11} \\
 ch\beta(\Delta_8^-) &= 2x_3 + (-2/3!)x'_7 + (2/3!)x_7 + (2/5!)x_{11}
 \end{aligned}$$

and the determinant of the corresponding matrix is  $-1$ .

REMARK. The equation  $ch\beta(\Delta_8^+ - \Delta_8^-) = x'_7$  confirms the fact that  $\text{Spin}(8)/\text{Spin}(7) = S^7$  (see [22, Proposition 6.2]).

Let us lastly consider the case of  $\text{Spin}(9)$ . In this case,  $(m_1, m_2, m_3, m_4) = (2, 4, 6, 8)$ . We can choose  $t_1, t_2, t_3, t_4, \gamma \in H^2(BT; Z)$  so that

$$H^*(BT; Z) = Z[t_1, \dots, t_4, \gamma]/(c_1 - 2\gamma)$$

and

$$H^*(BT; \mathbb{Q})^{\Phi(\text{Spin}(9))} = \mathbb{Q}[p_1, p_2, p_3, p_4]$$

where  $c_i = \sigma_i(t_1, \dots, t_4)$  and  $p_i = \sigma_i(t_1^2, \dots, t_4^2)$ . By a straightforward calculation we have

$$\begin{aligned} f_4 &= \frac{1}{2} p_1 = -c_2 + 2\gamma^2, \\ f_8 &= \frac{1}{2} p_2 - \frac{1}{2} f_4^2 = c_4 + 2(-c_3\gamma + c_2\gamma^2 - \gamma^4), \\ f_{12} &= p_3 = -2c_4c_2 + c_3^2, \\ f_{16} &= \frac{1}{4} p_4 - \frac{1}{4} f_8^2 = c_4c_3\gamma - c_4c_2\gamma^2 - c_3^2\gamma^2 + 2c_3c_2\gamma^3 \\ &\quad + c_4\gamma^4 - c_2^2\gamma^4 - 2c_3\gamma^5 + 2c_2\gamma^6 - \gamma^8. \end{aligned}$$

Since  $H^*(\text{Spin}(9); \mathbb{Z})$  has no  $p$ -torsion for  $p > 2$  and

$$H^*(\text{Spin}(9); \mathbb{Z}/2) = \Delta_{\mathbb{Z}/2}(\bar{x}_3, \bar{x}_5, \bar{x}_6, \bar{x}_7, \bar{x}_{15})$$

where all the  $\bar{x}_i$  are universally transgressive and  $\beta_2(\bar{x}_5) = \bar{x}_6$  [7], as in the case of  $\text{Spin}(7)$ , it follows that  $b(2) = b(4) = 1$ ,  $b(6) = 2$  and  $b(8) = 1$ . On the other hand,  $c(2) = c(4) = c(6) = c(8) = 1$ . Thus we have

$$\begin{aligned} (3.11) \quad \tau'(x_3) &= f_4 = \frac{1}{2} p_1, \\ \tau'(x_7) &= f_8 = \frac{1}{2} p_2 + \text{decomposables}, \\ \tau'(x_{11}) &= \frac{1}{2} f_{12} = \frac{1}{2} p_3, \\ \tau'(x_{15}) &= f_{16} = \frac{1}{4} p_4 + \text{decomposables}. \end{aligned}$$

REMARK. Let  $j: \text{Spin}(8) \rightarrow \text{Spin}(9)$  be the natural inclusion. Then by (3.6) and (3.11) we see that the homomorphism  $j^*: H^i(\text{Spin}(9); \mathbb{Z}) \rightarrow H^i(\text{Spin}(8); \mathbb{Z})$  satisfies

$$j^*(x_i) = \begin{cases} x_i & (i = 3, 11) \\ x'_7 + 2x_7 & (i = 7) \\ 0 & (i = 15). \end{cases}$$

Let us recall that

$$(3.12) \quad R(\text{Spin}(9)) = \mathbb{Z}[\lambda'_1, \lambda'_2, \lambda'_3, \Delta_9] \quad \text{where}$$

- (a)  $\dim \lambda'_k = \binom{9}{k}$  and  $\dim \Delta_9 = 16$ ;
- (b) relations  $\Lambda^k \lambda'_1 = \lambda'_k$  and  $\Delta_9^2 = \lambda'_4 + \lambda'_3 + \lambda'_2 + \lambda'_1 + 1$  hold;

(c) the set of weights of  $\lambda'_1$  is given by  $\{\pm t_i, 0 | 1 \leq i \leq 4\}$ .

The rest of the argument is parallel to that for  $G = \text{Spin}(7)$ . We only exhibit the result.

**Theorem 6.** The Chern characters on  $\text{Spin}(9)$  are given by:

$$\begin{aligned} ch\beta(\lambda'_1) &= 2x_3 + (-2/3!)x_7 + (2/5!)x_{11} + (-4/7!)x_{15} \\ ch\beta(\lambda'_2) &= 14x_3 + (-2/3!)x_7 + (-46/5!)x_{11} + (476/7!)x_{15} \\ ch\beta(\lambda'_3) &= 42x_3 + (18/3!)x_7 + (-18/5!)x_{11} + (-4284/7!)x_{15} \\ ch\beta(\Delta_9) &= 4x_3 + (2/3!)x_7 + (4/5!)x_{11} + (34/7!)x_{15} \end{aligned}$$

and the determinant of the corresponding matrix is 1.

#### 4. The exceptional Lie groups $G_2$ and $F_4$

Let us first consider the case of  $G_2$ . In this case,  $(m_1, m_2) = (2, 6)$ . We use the root system  $\{\alpha_1, \alpha_2\}$  of [11]. Let  $\omega_1, \omega_2$  be the fundamental weights. If we put

$$t_1 = \omega_1, t_2 = \omega_1 - \omega_2, t_3 = -2\omega_1 + \omega_2,$$

then

$$H^*(BT; Z) = Z[t_1, t_2, t_3]/(c_1)$$

where  $c_i = \sigma_i(t_1, t_2, t_3)$ , on which  $\Phi(G_2)$  acts as follows:

	$R_1$	$R_2$
$t_1$	$-t_2$	$t_1$
$t_2$	$-t_1$	$t_3$
$t_3$	$-t_3$	$t_2$

where  $R_j$  ( $j=1, 2$ ) is the reflection to the hyperplane  $\alpha_j=0$ , and  $\{R_1, R_2\}$  generates  $\Phi(G_2)$ . Therefore

$$H^*(BT; Q)^{\Phi(G_2)} = Q[p_1, p_3].$$

where  $p_i = \sigma_i(t_1^2, t_2^2, t_3^2)$ , and it follows that

$$f_4 = \frac{1}{2}p_1 = -c_2,$$

$$f_{12} = p_3 = c_3^2.$$

Since  $H^*(G_2; Z)$  has no  $p$ -torsion for  $p > 2$  and

$$H^*(G_2; Z/2) = \Delta_{Z/2}(\bar{x}_3, \bar{x}_5, \bar{x}_6)$$

where all the  $x_i$  are universally transgressive and  $\beta_2(x_5)=x_6$  [7], as in the case of  $\text{Spin}(7)$ , it follows that  $b(2)=1$  and  $b(6)=2$ . On the other hand,  $c(2)=c(6)=1$ . Thus we have

$$\begin{aligned}\tau'(x_3) &= f_4 = \frac{1}{2}p_1, \\ \tau'(x_{11}) &= \frac{1}{2}f_{12} = \frac{1}{2}p_3.\end{aligned}$$

Let us recall that

- (4.1)  $R(G_2) = Z[\rho_1, \Lambda^2\rho_1]$  where
- (a)  $\dim \Lambda^k \rho_1 = \binom{7}{k}$  (and  $\dim \rho_2 = 14$ );
  - ((b) a relation  $\Lambda^2 \rho_1 = \rho_1 + \rho_2$  holds;)
  - (c) the set of weights of  $\rho_1$  is given by  $\{\pm t_i (1 \leq i \leq 3), 0\}$ .

By a calculation we have

$$\begin{aligned}ch^2 \alpha i^*(\rho_1) &= p_1, \\ ch^6 \alpha i^*(\rho_1) &= \frac{1}{120}p_3 + \text{decomposables}.\end{aligned}$$

Therefore

$$ch\beta(\rho_1) = 2x_3 + \frac{1}{60}x_{11}$$

and by Lemma 1 we get

**Theorem 7.** *The Chern characters on  $G_2$  are given by :*

$$\begin{aligned}ch\beta(\rho_1) &= 2x_3 + (2/5!)x_{11} \\ ch\beta(\Lambda^2\rho_1) &= 10x_3 + (-50/5!)x_{11}\end{aligned}$$

and the determinant of the corresponding matrix is  $-1$ .

REMARK. Consider the following fibration

$$G_2 \xrightarrow{k} \text{Spin}(7) \rightarrow \text{Spin}(7)/G_2 = S^7.$$

Then it is easy to see that  $k^*: H^i(\text{Spin}(7); \mathbb{Z}) \rightarrow H^i(G_2; \mathbb{Z})$  satisfies

$$k^*(x_i) = \begin{cases} x_i & (i=3, 11) \\ 0 & (i=7) \end{cases}.$$

On the other hand,  $k^*: R(\text{Spin}(7)) \rightarrow R(G_2)$  satisfies

$$\begin{aligned}k^*(\lambda'_i) &= \Lambda^i \rho_1 \quad (i=1, 2) \\ k^*(\Delta_7) &= \rho_1 + 1\end{aligned}$$

(see [31]). Using these facts, we find that Theorem 7 follows from Theorem 4.

$H^*(\Omega G_2; Z)$  (for degrees  $\leq 10$ ) was calculated implicitly by Bott [10]. Using it and the cohomology spectral sequence of the path fibration  $\Omega G_2 \rightarrow PG_2 \rightarrow G_2$ , we can show that

$$d(2) = 1 \quad \text{and} \quad d(6) = 2$$

(see [12] and [28, p. 474]).

Let us now consider the case of  $F_4$ . In this case,  $(m_1, m_2, m_3, m_4) = (2, 6, 8, 12)$ . We can choose elements  $t_1, t_2, t_3, t_4, \gamma \in H^2(BT; Z)$  so that

$$H^*(BT; Z) = Z[t_1, \dots, t_4, \gamma] / (c_1 - 2\gamma)$$

and the action of  $\Phi(F_4)$  on it is as described in [9, §19] (see [18] and [29]). Let  $c_i = \sigma_i(t_1, \dots, t_4)$  and  $p_i = \sigma_i(t_1^2, \dots, t_4^2)$ . If we put

$$\begin{aligned} I_4 &= p_1, \\ I_{12} &= -6p_3 + p_2p_1, \\ I_{16} &= 12p_4 - 3p_3p_1 + p_2^2, \\ I_{24} &= -72p_4p_2 + 27p_4p_1^2 + 27p_3^2 - 9p_3p_2p_1 + 2p_2^3, \end{aligned}$$

then we have

$$H^*(BT; Q)^{\Phi(F_4)} = Q[I_4, I_{12}, I_{16}, I_{24}].$$

For a proof see [27, Lemma 5.1], however, its main part is accomplished by a pure calculation; see (4.7) and (4.8) below. By a troublesome calculation we obtain

$$\begin{aligned} f_4 &= \frac{1}{2} I_4 = -c_2 + 2\gamma^2, \\ f_{12} &= -\frac{1}{2} I_{12} \\ &= -4c_4c_2 + 3c_3^2 + c_2^3 - 4c_3c_2\gamma - 4c_4\gamma^2 - 2c_2^2\gamma^2 + 8c_3\gamma^3, \\ f_{16} &= \frac{1}{16} (I_{16} + 2f_{12}f_4 + f_4^4) \\ &= c_4^2 - c_4c_3\gamma + c_4c_2\gamma^2 + c_3^2\gamma^2 - 2c_3c_2\gamma^3 - c_4\gamma^4 + c_2^2\gamma^4 + 2c_3\gamma^5 - 2c_2\gamma^6 + \gamma^8, \\ f_{24} &= -\frac{1}{64} (I_{24} + 16f_{16}f_4^2 - 3f_{12}^2 + f_4^6) \\ &= 2c_4^3 - c_4^2c_2^2 - 3c_4^2c_3\gamma + c_4c_3c_2^2\gamma + 7c_4^2c_2\gamma^2 - 3c_4c_3^2\gamma^2 - c_4c_2^3\gamma^2 + 2c_4c_3c_2\gamma^3 \\ &\quad + 2c_3^3\gamma^3 + 2c_3c_2^3\gamma^3 - 7c_4^2\gamma^4 + 2c_4c_2^2\gamma^4 - 2c_3^2c_2\gamma^4 - c_2^4\gamma^4 - 2c_4c_3\gamma^5 - 4c_3c_2^2\gamma^5 \\ &\quad - 2c_4c_2\gamma^6 - c_2^3\gamma^6 + 4c_3^3\gamma^6 + 4c_3c_2\gamma^7 + c_4\gamma^8 - 7c_2^2\gamma^8 - 2c_3\gamma^9 + 6c_2\gamma^{10} - 2\gamma^{12}. \end{aligned}$$

Let us determine  $b(2)$ ,  $b(6)$ ,  $b(8)$ ,  $b(12) \in Z$ . Recall that  $H^*(F_4; Z)$  has no  $p$ -torsion for  $p > 3$ . Since

$$H^*(F_4; Z/2) = \Delta_{Z/2}(\bar{x}_3, \bar{x}_5, \bar{x}_6, \bar{x}_{15}, \bar{x}_{23})$$

where all the  $\bar{x}_i$  are universally transgressive and  $\beta_2(\bar{x}_5) = \bar{x}_6$  [7], it follows that  $\nu_2(b(2)) = 0$ ,  $\nu_2(b(6)) = 1$ ,  $\nu_2(b(8)) = 0$  and  $\nu_2(b(12)) = 0$ . Consider the case  $p = 3$ . Recall from [7] and [25] that

$$H^*(F_4; Z/3) = Z/3[\bar{x}_8]/(\bar{x}_8^3) \otimes \Lambda_{Z/3}(\bar{x}_3, \bar{x}_7, \bar{x}_{11}, \bar{x}_{15})$$

$$H^*(BF_4; Z/3) = Z/3[\bar{y}_{36}, \bar{y}_{48}] \otimes C,$$

$$C = Z/3[\bar{y}_4, \bar{y}_8] \otimes \{1, \bar{y}_{20}, \bar{y}_{20}^2\} + \Lambda_{Z/3}(\bar{y}_9) \otimes Z/3[\bar{y}_{26}] \otimes \{1, \bar{y}_{20}, \bar{y}_{21}, \bar{y}_{25}\}$$

where  $\tau(\bar{x}_i) = \bar{y}_{i+1}$  for  $i = 3, 7, 8$  and  $\beta_3(\bar{x}_7) = \bar{x}_8$ . Here we may suppose that

$$\begin{aligned} \pi_3(\bar{x}_3) &= \bar{x}_3, & \pi_3(\bar{y}_4) &= \bar{y}_4, \\ \pi_3(\bar{x}_{11}) &= \bar{x}_{11}, & \pi_3(\bar{y}_{12}) &= \bar{y}_4 \bar{y}_8, \\ \pi_3(\bar{x}_{15}) &= \bar{x}_{15}, & \pi_3(\bar{y}_{16}) &= \bar{y}_8^2, \\ \pi_3(\bar{x}_{23}) &= \bar{x}_7 \bar{x}_8^2, & \pi_3(\bar{y}_{24}) &= \bar{y}_8^3. \end{aligned}$$

In the mod 3 cohomology spectral sequence  $\{E_r(Z/3)\}$  of the universal fibration

$$F = F_4 \rightarrow E = EF_4 \rightarrow B = BF_4,$$

if

$$\beta_3^B: E_2^{s,t}(Z/3) \rightarrow E_2^{s+1,t}(Z/3)$$

is the map induced by  $\beta_3: H^s(B; Z/3) \rightarrow H^{s+1}(B; Z/3)$  through the isomorphism

$$E_2^{s,t}(Z/3) \cong H^s(B; H^t(F; Z/3)),$$

then we have

$$(4.2) \quad \begin{cases} d_9(1 \otimes \bar{x}_{11}) = \bar{y}_9 \otimes \bar{x}_3 \dots\dots\dots (*) \\ \beta_3^B(\bar{y}_8 \otimes \bar{x}_3) = \bar{y}_9 \otimes \bar{x}_3 \\ d_4(\bar{y}_8 \otimes \bar{x}_3) = \bar{y}_4 \bar{y}_8 \otimes 1 \end{cases}$$

$$(4.3) \quad \begin{cases} d_9(1 \otimes \bar{x}_{15}) = \bar{y}_9 \otimes \bar{x}_7 \dots\dots\dots (*) \\ \beta_3^B(\bar{y}_8 \otimes \bar{x}_7) = \bar{y}_9 \otimes \bar{x}_7 \\ d_8(\bar{y}_8 \otimes \bar{x}_7) = \bar{y}_8^2 \otimes 1 \end{cases}$$

$$(4.4) \quad \begin{cases} d_8(1 \otimes \bar{x}_7 \bar{x}_8^2) = \bar{y}_8 \otimes \bar{x}_8^2 \\ \beta_3^B(\bar{y}_8 \otimes \bar{x}_7 \bar{x}_8) = \bar{y}_8 \otimes \bar{x}_8^2 \\ d_8(\bar{y}_8 \otimes \bar{x}_7 \bar{x}_8) = \bar{y}_8^2 \otimes \bar{x}_8 \\ \beta_3^B(\bar{y}_8^2 \otimes \bar{x}_7) = \bar{y}_8^2 \otimes \bar{x}_8 \\ d_8(\bar{y}_8^2 \otimes \bar{x}_7) = \bar{y}_8^3 \otimes 1 \end{cases}$$

where the asterisks are due to [3]. Generally, with the obvious notation, since  $d_1: E_1^{s,t}(Z/3) \rightarrow E_1^{s+1,t}(Z/3)$  can be identified with the differential  $\delta_B: C^s(B; Z/3) \rightarrow C^{s+1}(B; Z/3)$ , if  $\beta_3^s(\{\bar{u}\}) = \{\emptyset\}$ , then there exist  $u, v \in A^{*,*}(Z)$  such that  $\pi_3(u) = \bar{u}$ ,  $\pi_3(v) = \emptyset$  and  $\delta(u) = 3v$ . In this way the same argument as in the case of  $\text{Spin}(7)$  is valid. Therefore the conditions (4.2), (4.3) and (4.4) imply that  $\nu_3(b(6)) = 1$ ,  $\nu_3(b(8)) = 1$  and  $\nu_3(b(12)) = 2$  respectively. Summarizing these, we have

$$b(2) = 1, \quad b(6) = 6, \quad b(8) = 3 \quad \text{and} \quad b(12) = 9.$$

On the other hand,  $c(2) = c(6) = c(8) = c(12) = 1$ . Thus we obtain

$$\begin{aligned} (4.5) \quad \tau'(x_3) &= f_4 = \frac{1}{2} I_4, \\ \tau'(x_{11}) &= \frac{1}{6} f_{12} = -\frac{1}{12} I_{12}, \\ \tau'(x_{15}) &= \frac{1}{3} f_{16} = \frac{1}{48} I_{16} + \text{decomposables}, \\ \tau'(x_{23}) &= \frac{1}{9} f_{24} = -\frac{1}{576} I_{24} + \text{decomposables}. \end{aligned}$$

Let us recall from [30] that

$$(4.6) \quad R(F_4) = Z[\rho_4, \Lambda^2 \rho_4, \Lambda^3 \rho_4, \rho_1] \quad \text{where}$$

- (a)  $\dim \Lambda^k \rho_4 = \binom{26}{k}$  and  $\dim \rho_1 = 52$ ;
- (b) the set of weights of  $\rho_4$  is given by

$$\{\pm t_i (1 \leq i \leq 4), \frac{1}{2} (\pm t_1 \pm t_2 \pm t_3 \pm t_4), 0, 0\}$$

and that of  $\rho_1$  is given by

$$\{\pm t_i \pm t_j (1 \leq i < j \leq 4), \pm t_i (1 \leq i \leq 4), \frac{1}{2} (\pm t_1 \pm t_2 \pm t_3 \pm t_4), 0, 0, 0, 0\}.$$

We have to calculate  $\text{chai}^*(\rho_4)$  and  $\text{chai}^*(\rho_1)$ . Consider the inclusion  $k: \text{Spin}(9) \rightarrow F_4$  such that  $F_4/\text{Spin}(9) = \Pi$ , the Cayley projective plane (see, e.g., [9, §19]). Then  $k^*: R(F_4) \rightarrow R(\text{Spin}(9))$  satisfies  $k^*(\rho_4) = \lambda'_1 + \Delta_9 + 1$  and  $k^*(\rho_1) = \lambda'_2 + \Delta_9$ ; see (4.6) (b). Let us calculate  $\text{chai}^*(\Delta_9)$ , where the set of weights of  $\Delta_9$  is  $\{1/2(\pm t_1 \pm t_2 \pm t_3 \pm t_4)\}$ . To do so we first calculate  $\text{chai}^*(\Delta_5)$ , where the set of weights of  $\Delta_5$  is  $\{1/2(\pm t_1 \pm t_2)\}$ ; using it, we calculate  $\text{chai}^*(\Delta_7)$ ; and using it, we calculate  $\text{chai}^*(\Delta_9)$ . Our final result is

$$ch^2\alpha i^*(\Delta_9) = 2p_1,$$

$$ch^6\alpha i^*(\Delta_9) = \frac{1}{2880}(48p_3 + 12p_2p_1 + p_1^3),$$

$$ch^8\alpha i^*(\Delta_9) = \frac{1}{645120}(1088p_4 + 256p_3p_1 + 16p_2^2 + 24p_2p_1^2 + p_1^4),$$

$$ch^{12}\alpha i^*(\Delta_9) = \frac{1}{122624409600}(31488p_4p_2 + 42432p_4p_1^2 + 3072p_3^2 + 4608p_3p_2p_1 \\ + 1920p_3p_1^3 + 64p_2^3 + 240p_2^2p_1^2 + 60p_2p_1^4 + p_1^6).$$

By a similar calculation to the before, we have

$$ch^2\alpha i^*(\lambda'_1) = p_1,$$

$$ch^6\alpha i^*(\lambda'_1) = \frac{1}{360}(3p_3 - 3p_2p_1 + p_1^3),$$

$$ch^8\alpha i^*(\lambda'_1) = \frac{1}{20160}(-4p_4 + 4p_3p_1 + 2p_2^2 - 4p_2p_1^2 + p_1^4),$$

$$ch^{12}\alpha i^*(\lambda'_1) = \frac{1}{239500800}(6p_4p_2 - 6p_4p_1^2 + 3p_3^2 - 12p_3p_2p_1 + 6p_3p_1^3 - 2p_2^3 \\ + 9p_2^2p_1^2 - 6p_2p_1^4 + p_1^6).$$

Thus we have

$$(4.7) \quad ch^2\alpha i^*(\rho_4) = 3p_1,$$

$$ch^6\alpha i^*(\rho_4) = \frac{1}{960}(24p_3 - 4p_2p_1 + 3p_1^3),$$

$$ch^8\alpha i^*(\rho_4) = \frac{1}{645120}(960p_4 + 384p_3p_1 + 80p_2^2 - 104p_2p_1^2 + 33p_1^4),$$

$$ch^{12}\alpha i^*(\rho_4) = \frac{1}{40874803200}(11520p_4p_2 + 13120p_4p_1^2 + 1536p_3^2 - 512p_3p_2p_1 \\ + 1664p_3p_1^3 - 320p_2^3 + 1616p_2^2p_1^2 - 1004p_2p_1^4 + 171p_1^6).$$

On the other hand,  $ch\alpha i^*(\rho_1 - \rho_4)$  was calculated in [27, §5] (with certain indeterminacy). Following it, we have

$$(4.8) \quad ch^2\alpha i^*(\rho_1 - \rho_4) = 6p_1,$$

$$ch^6\alpha i^*(\rho_1 - \rho_4) = \frac{1}{60}(-12p_3 + 2p_2p_1 - p_1^3),$$

$$ch^8\alpha i^*(\rho_1 - \rho_4) = \frac{1}{10080}(240p_4 - 156p_3p_1 + 20p_2^2 + 16p_2p_1^2 + 3p_1^4),$$



$$ch^{12}\alpha i^*(\rho_1 - \rho_4) = \frac{1}{39916800}(-720p_4p_2 + 1270p_4p_1^2 + 366p_3^2 - 122p_3p_2p_1 \\ - 346p_3p_1^3 + 20p_2^3 + 86p_2^2p_1^2 + 16p_2p_1^4 + p_1^6).$$

Thus we get

$$\begin{aligned} ch^2\alpha i^*(\rho_1) &= 9I_4, \\ ch^6\alpha i^*(\rho_1) &= \frac{7}{240}I_{12} + \text{decomposables}, \\ ch^8\alpha i^*(\rho_1) &= \frac{17}{8064}I_{16} + \text{decomposables}, \\ ch^{12}\alpha i^*(\rho_1) &= \frac{1}{4055040}I_{24} + \text{decomposables}. \end{aligned}$$

Combining these with (4.5), it follows that

$$\begin{aligned} ch\beta(\rho_4) &= 6x_3 + \frac{1}{20}x_{11} + \frac{1}{168}x_{15} + \frac{1}{443520}x_{23}, \\ ch\beta(\rho_1) &= 18x_3 - \frac{7}{20}x_{11} + \frac{17}{168}x_{15} - \frac{1}{7040}x_{23} \end{aligned}$$

and by Lemma 1 we obtain

**Theorem 8.** *The Chern characters on  $F_4$  are given by:*

$$\begin{aligned} ch\beta(\rho_4) &= 6x_3 + (6/5!)x_{11} + (30/7!)x_{15} + (90/11!)x_{23} \\ ch\beta(\Lambda^2\rho_4) &= 144x_3 + (-36/5!)x_{11} + (-3060/7!)x_{15} + (-181980/11!)x_{23} \\ ch\beta(\Lambda^3\rho_4) &= 1656x_3 + (-1584/5!)x_{11} + (-24480/7!)x_{15} + (11180160/11!)x_{23} \\ ch\beta(\rho_1) &= 18x_3 + (-42/5!)x_{11} + (510/7!)x_{15} + (-5670/11!)x_{23} \end{aligned}$$

and the determinant of the corresponding matrix is 1.

$H^*(\Omega F_4; Z)$  (for degrees  $\leq 22$ ) was calculated implicitly in [28]. Using it and the cohomology spectral sequence of the path fibration  $\Omega F_4 \rightarrow PF_4 \rightarrow F_4$ , we can show that

$$d(2) = 1, \quad d(6) = 2, \quad d(8) = 1 \quad \text{and} \quad d(12) = 3.$$

## References

- [1] J.F. Adams: *Vector fields on spheres*, Ann. of Math. **75** (1962), 603–632.
- [2] S. Araki: *Steenrod reduced powers in the spectral sequences associated with a fibering*, Mem. Fac. Sci. Kyusyu Univ. **11** (1957), 15–64.
- [3] S. Araki: *On the non-commutativity of Pontrjagin rings mod 3 of some compact*

- exceptional groups*, Nagoya Math. J. **26** (1960), 225–260.
- [4] M.F. Atiyah: *On the K-theory of compact Lie groups*, Topology **4** (1965), 95–99.
  - [5] M.F. Atiyah and F. Hirzebruch: *Vector bundles and homogeneous spaces*, Proc. Sympos. Pure Math. Vol. 3, 7–38, Amer. Math. Soc., 1961.
  - [6] A. Borel: *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math. **57** (1953), 115–207.
  - [7] A. Borel: *Sur l'homologie et la cohomologie des groupes de Lie compacts connexes*, Amer. J. Math. **76** (1954), 273–342.
  - [8] A. Borel: *Topology of Lie groups and characteristic classes*, Bull. Amer. Math. Soc. **61** (1955), 397–432.
  - [9] A. Borel and F. Hirzebruch: *Characteristic classes and homogeneous spaces*, I, Amer. J. Math. **80** (1958), 458–538.
  - [10] R. Bott: *The space of loops on a Lie group*, Michigan Math. J. **5** (1958), 35–61.
  - [11] N. Bourbaki: *Groupes et algèbres de Lie IV–VI*, Hermann, 1968.
  - [12] F. Clarke: *On the K-theory of the loop space of a Lie group*, Proc. Cambridge Philos. Soc. **76** (1974), 1–20.
  - [13] M. Curtis, A. Wiederhold and B. Williams: *Normalizers of maximal tori*, Localization in Group Theory and Homotopy Theory, Lecture Notes in Math. Vol. 418, 31–47, Springer, 1974.
  - [14] E.B. Dynkin: *Topological characteristics of homomorphisms of compact Lie groups*, Amer. Math. Soc. Transl. **12** (1959), 301–342.
  - [15] M. Feshbach: *The image of  $H^*(BG; \mathbb{Z})$  in  $H^*(BT; \mathbb{Z})$  for  $G$  a compact Lie group with maximal torus  $T$* , Topology **20** (1981), 93–95.
  - [16] L. Hodgkin: *On the K-theory of Lie groups*, Topology **6** (1967), 1–36.
  - [17] D. Husemoller: *Fibre bundles*, McGraw-Hill, 1966.
  - [18] K. Ishitoya and H. Toda: *On the cohomology of irreducible symmetric spaces of exceptional type*, J. Math. Kyoto Univ. **17** (1977), 225–243.
  - [19] N. Jacobson: *Lie algebras*, Interscience, 1962.
  - [20] O.V. Manturov: *Geometric models of vector bundles over compact homogeneous spaces*, Soviet Math. Dokl. **12** (1971), 1646–1650.
  - [21] O.V. Manturov: *Generators in the complex K-functor of compact homogeneous spaces*, Math. USSR Sb. **19** (1973), 47–84.
  - [22] H. Minami: *K-groups of symmetric spaces I*, Osaka J. Math. **12** (1975), 623–634.
  - [23] C.M. Naylor: *On the  $\beta$ -construction in K-theory*, Canad. J. Math. **24** (1972), 819–824.
  - [24] J.P. Serre: *Homologie singulière des espaces fibrés*, Ann. of Math. **54** (1951), 425–505.
  - [25] H. Toda: *Cohomology mod 3 of the classifying space  $BF_4$  of the exceptional group  $F_4$* , J. Math. Kyoto Univ. **13** (1972), 97–115.
  - [26] H. Toda: *On the cohomology ring of some homogeneous spaces*, J. Math. Kyoto Univ. **15** (1975), 185–199.
  - [27] H. Toda and T. Watanabe: *The integral cohomology rings of  $F_4/T$  and  $E_6/T$* , J. Math. Kyoto Univ. **14** (1974), 257–286.
  - [28] T. Watanabe: *The homology of the loop space of the exceptional group  $F_4$* , Osaka J. Math. **15** (1978), 463–474.
  - [29] T. Watanabe: *Cohomology operations in the loop space of the compact exceptional*

- group  $F_4$* , Osaka J. Math. **16** (1979), 471–478.
- [30] I. Yokota: *Exceptional Lie group  $F_4$  and its representation rings*, J. Fac. Sci. Shinshu Univ. **3** (1968), 35–60.
- [31] I. Yokota: *Groups and representations*, Shōkabō, (in Japanese), 1973.

Department of Mathematics  
Osaka City University  
Sugimoto, Sumiyoshi-ku  
Osaka 558, Japan