

Title	L^∞ boundedness of nonlinear eigenfunction under singular variation of domains
Author(s)	Ozawa, Shin
Citation	Osaka Journal of Mathematics. 1995, 32(2), p. 363–371
Version Type	VoR
URL	https://doi.org/10.18910/10245
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Ozawa, S. Osaka J. Math. 32 (1995), 363-371

L^{∞} BOUNDEDNESS OF NONLINEAR EIGENFUNCTION UNDER SINGULAR VARIATION OF DOMAINS

Shin OZAWA

(Received September 20, 1993)

1. Introduction.

Let M be a bounded domain in \mathbb{R}^3 with smooth boundary ∂M . Let \tilde{w} be a point in M. We remove ball B_{ε} of radius ε with the center \tilde{w} from M and we get $M_{\varepsilon} = M \setminus \overline{B}_{\varepsilon}$.

Fix $p \in (1,2)$. We fix k > 0. We put

(1.1)
$$\lambda(\varepsilon) = \inf_{u \in X} \left(\int_{M_{\varepsilon}} |\nabla u|^2 dx + k \int_{\partial B_{\varepsilon}} u^2 d\sigma_x \right),$$

where $X = \{u \in H^1(M_{\varepsilon}), u = 0 \text{ on } \partial M \text{ and } u \ge 0 \text{ in } M_{\varepsilon}, \|u\|_{p+1,\varepsilon} = 1\}$. Here $\|u\|_{L^q(M_{\varepsilon})} = \|u\|_{q,\varepsilon}$. We see that there exists at least one solution v_{ε} of the above problem which attains $(1.1)_{\varepsilon}$. We know that v_{ε} satisfies $-\Delta v_{\varepsilon}(x) = \lambda(\varepsilon)v_{\varepsilon}(x)^p$ in M_{ε} , $v_{\varepsilon}(x) = 0$ on ∂M and $kv_{\varepsilon}(x) + (\partial/\partial v_x)v_{\varepsilon}(x) = 0$ on ∂B_{ε} . Here $\partial/\partial v_x$ denotes the derivative along the exterior normal vector with respect to M_{ε} .

Let S_{ε} denote the set of positive function u_{ε} which attains the minimum of (1.1). Main result of this paper is the following

Theorem 1. Fix $p \in (1,2)$. Then, there exists a constant C independent of ε such that

$$\sup_{u_{\varepsilon}\in S_{\varepsilon}}\sup_{x\in M_{\varepsilon}}|v_{\varepsilon}(x)| < C.$$

Related topics are discussed in Lin [1], Ozawa [4].

2. Preliminary Lemmas.

We have the following Lemma 2.1.

Lemma 2.1. Consider the following equations.

(2.1)
$$\Delta v_{\varepsilon}(x) = 0 \qquad x \in M \setminus \bar{B}_{\varepsilon}$$

S. OZAWA

$$(2.2) v_{\varepsilon}(x) = 0 x \in \partial M$$

(2.3)
$$k v_{\varepsilon}(x) + (\partial v_{\varepsilon}/\partial v_{x}) = \alpha(\omega) \qquad x = w + \varepsilon \omega \in \partial B_{\varepsilon}.$$

Here $\omega \in S^2$. Then, the solution of (2.1), (2.2), (2.3) satisfies

$$|v_{\varepsilon}(x)| \le C\varepsilon^2 r^{-1} \|\alpha\|_{C^{(\sigma/2)}(S^2)}$$

for any $\sigma > 1$. Here C may depend on σ but independent of ε . Here $r = |x - \tilde{w}|$.

Proof. Let Δ_{S^2} denote the Laplace-Beltrami oporator on S^2 . It is well known that -n(n+1) is an eigenvalue of Δ_{S^2} whose multiplicity is (2n+1). We can write it explicitly by using the Legendre polynomial but we do not do it. We write complete orthonomal basis of $L^2(S^2)$ consisting of eigenfunction by $\{\varphi_j(\omega)\}_{j=1}^{\infty}$. If $\Delta \varphi_j(\omega) = -n(n+1)\varphi_j(\omega)$, then we write j as $j \in (n)$. Thus, $\#\{j; j \in (n)\} = 2n+1$.

First we want to construct a solution of (2.1), (2.3). We put

$$v^{*}(x) = \sum_{n=0}^{\infty} r^{-(n+1)} \sum_{j \in (n)} c_{j,n} \varphi_{j}(\omega).$$

Then,

$$\Delta v^*(x) = 0$$
 in $R^3 \setminus \overline{B}_{\varepsilon}$.

We expand α by $\varphi_i(\omega)$ as

$$\alpha(\omega) = \sum_{n=0}^{\infty} \sum_{j \in (n)} a_{j,n} \varphi_j(\omega).$$

By the equality (2.3), we have

$$(k c_{j,n} \varepsilon^{-(n+1)} + (n+1) c_{j,n} \varepsilon^{-(n+2)}) = a_{j,n}.$$

Then,

$$c_{j,n} = a_{j,n} \varepsilon^{n+1} (k + (n+1)\varepsilon^{-1})^{-1}.$$

Therefore,

$$|v^*(x)| \leq \varepsilon \sum_{n=0}^{\infty} \sum_{j \in (n)} a_{j,n} (\varepsilon/r)^{n+1} (\varepsilon k + (n+1))^{-1} \varphi_j(\omega).$$

We notice that we have

(2.5)
$$|\varphi_j(\omega)| \le C(n+1)^{1/2}$$

for $j \in (n)$ by the property of Legendre and associated Legendre polynomial. See Mizohata [2, p.312].

By the Schwarz inequality we have

$$|v_{\varepsilon}^{*}| \leq C\varepsilon^{2}r^{-1} \left(\sum_{n=0}^{\infty} (n+1)^{\sigma} \sum_{j \in (n)} a_{j,n}^{2}\right)^{1/2} \\ \times \left(\sum_{n=0}^{\infty} (n+1)^{-\sigma} (2n+1)n(n+1)^{-2}\right)^{1/2}.$$

If we take $\sigma > 1$, then

(2.6)
$$|v^*(x)| \le C\varepsilon^2 r^{-1} \left(\sum_{n=0}^{\infty} (n+1)^{\sigma} \sum_{j \in (n)} a_{j,n}^2 \right)^{1/2} \le C\varepsilon^2 r^{-1} \|\alpha\|_{H^{\sigma/2}(S^2)}.$$

Here we note that $\|\alpha\|_{H^2(S^2)}^2$ is equivalent to $\left(\sum_{n=0}^{\infty} (n+1)^4 \sum_{j \in (n)} a_{j,n}^2\right)$, since the eigenvalue of $-\Delta_{S^2}$ is n(n+1). We used representation of norm of fractional Sobolev space. Notice that $C^{\sigma'/2}(S^2) \subset H^{\sigma/2}(S^2)$ for $\sigma < \sigma'$.

Now $v^*(x)$ does not satisfy $v^*(x)=0$ on ∂M . By the same procedure as in the repeated construction of the function $v_{\varepsilon}^{(n)}$ in Proposition 1 of Ozawa [3] we proved Lemma 2.1.

The following Lemm is very useful.

Lemma 2.3. There exists an extension operator $E: H^1(M_k) \ni u \rightarrow Eu = \tilde{u} \in H^1(M)$ satisfying the followings;

(2.9)
$$\tilde{u}(x) = u(x) \qquad a.e. \ M_{\varepsilon}$$

holds for any $u \in H^1(M_{\epsilon})$,

$$\|\tilde{u}\|_{L^{s}(M)} \leq C \|u\|_{s,\varepsilon} \qquad (1 \leq s \leq \infty)$$

holds for any $u \in H^1(M_{\varepsilon}) \cap L^s(M_{\varepsilon})$,

(2.11)
$$\|\tilde{u}\|_{H^{1}(M)} \leq C \|u\|_{H^{1}(M_{\varepsilon})} + \varepsilon^{((s-2)3/(2s))-1} \|u\|_{s,\varepsilon}$$

for any $u \in H^1(M_{\epsilon}) \cap L^s(M_{\epsilon})$ with $2 \leq s < \infty$,

 $\|\tilde{u}\|_{H^{1}(M)} \leq C \|u\|_{H^{1}(M_{\varepsilon})} + C \varepsilon^{(1/2)} \|u\|_{\infty,\varepsilon}$

holds for any $u \in H^1(M_{\varepsilon}) \cap L^{\infty}(M_{\varepsilon})$.

Proof. Without loss of generality, we may assume that $\tilde{w} = 0$. We take an arbitrary $u \in H^1(M_s)$ and put

$$\widetilde{u}(x) = u(x) \qquad x \in M_{\varepsilon}$$
$$= u(\varepsilon^2 x |x|^{-2}) \eta_{\varepsilon}(x) \qquad x \in B_{\varepsilon},$$

where $\eta_{\varepsilon} \in C^{\infty}(\mathbb{R}^3)$ satisfies $0 \le \eta_{\varepsilon} \le 1$, $\eta_{\varepsilon} = 1$ on $\mathbb{R}^3 \setminus \overline{B}_{\varepsilon/2}$, $\eta_{\varepsilon} = 0$ on $B_{\varepsilon/4}$ and $|\nabla \eta_{\varepsilon}| \le 8\varepsilon^{-1}$. Notice that both $\eta_{\varepsilon}(\varepsilon^2 x |x|^{-2})$ and $(\nabla \eta_{\varepsilon})(\varepsilon^2 x |x|^{-2})$ vanish on $\mathbb{R}^3 \setminus B_{4\varepsilon}$. Then, by using the Kelvin transformation of co-ordinates, $y = \varepsilon^2 x |x|^{-2}$, we have

$$\int_{B_{\varepsilon}} |\tilde{u}(x)|^{s} dx \leq \int_{\mathbb{R}^{3} \setminus \overline{B}_{\varepsilon}} |u(y)|^{s} \eta_{\varepsilon} (\varepsilon^{2} y |y|^{-2})^{s} (\varepsilon |y|^{-1})^{6} dy$$
$$\leq \int_{M_{\varepsilon}} |u(y)|^{s} dy \qquad (1 \leq s < \infty),$$

and

$$\begin{split} \int_{B_{\varepsilon}} |\nabla u(x)|^2 dx &\leq C \int_{B_{\varepsilon}} |u(\varepsilon^2 x |x|^{-2})|^2 |(\nabla \eta_{\varepsilon})(x)|^2 dx \\ &+ C \int_{B_{\varepsilon}} (\varepsilon |x|^{-1})^4 |(\nabla u)(\varepsilon^2 x |x|^{-2})|^2 \eta_{\varepsilon}(x)^2 dx \\ &\leq C \varepsilon^4 \int_{M_{\varepsilon}} |u(y)|^2 |y|^{-6} dy + C \int_{M_{\varepsilon}} |\nabla u|^2 dy. \end{split}$$

By Hölder's inequality, we see that

$$\int_{M_{\varepsilon}} |u(y)|^{2} |y|^{-6} dy \leq \begin{cases} C \varepsilon^{-(1+(2/s))3} \|u\|_{L^{s}(M_{\varepsilon})}^{2} & (2 \leq s < \infty) \\ C \varepsilon^{-3} \|u\|_{L^{\infty}(M_{\varepsilon})}^{2} & (s = \infty). \end{cases}$$

Thus, we get Lemma 2.3.

3. The Green function.

Let G(x,y) be the Green function of the Laplacian in M under the Dirichlet condition on ∂M . We introduce the following kernel $p_{\varepsilon}(x,y)$.

$$p_{\varepsilon}(x,y) = G(x,y) + g(\varepsilon)G(x,\tilde{w})G(\tilde{w},y) + h(\varepsilon)\langle \nabla_{w}G(x,\tilde{w}), \nabla_{w}G(\tilde{w},y)\rangle,$$

where

$$\langle \nabla_{w} u(\tilde{w}), \nabla_{w} v(\tilde{w}) \rangle = \sum_{n=1}^{3} \frac{\partial u}{\partial w_{i}} \frac{\partial v}{\partial w_{i}}|_{w=\hat{w}},$$

$$g(\varepsilon) = -(\gamma + (4\pi\varepsilon)^{-1} + k^{-1}(4\pi)^{-1}\varepsilon^{-2})^{-1},$$

where

$$\gamma = \lim_{x \to \tilde{w}} (G(x, \tilde{w}) - (4\pi)^{-1} |x - \tilde{w}|^{-1})$$

and

$$h(\varepsilon) = k^{-1} / ((4\pi)^{-1} \varepsilon^{-2} + k^{-1} (2\pi)^{-1} \varepsilon^{-3}).$$

We put

$$G(x,y) - (4\pi)^{-1}|x-y|^{-1} = S(x,y).$$

Then, $S(x,y) \in C^{\infty}(M \times M)$. We have the following:

$$\begin{split} \langle \nabla_{w} G(x, \tilde{w}), \nabla_{w} G(\tilde{w}, y) \rangle \\ &= (4\pi)^{-1} \varepsilon^{-2} \frac{\partial}{\partial w_{1}} G(\tilde{w}, y) + \langle \nabla_{w} S(x, \tilde{w}), \nabla_{w} G(\tilde{w}, y) \rangle \\ \\ &\frac{\partial}{\partial x_{1}} \langle \nabla_{w} G(x, \tilde{w}), \nabla_{w} G(\tilde{w}, y) \rangle \\ &= -(2\pi)^{-1} \varepsilon^{-3} \frac{\partial}{\partial w_{1}} G(\tilde{w}, y) + \frac{\partial}{\partial x_{1}} \langle \nabla_{w} S(x, \tilde{w}), \nabla_{w} G(\tilde{w}, y) \rangle \end{split}$$

for $x = (\varepsilon, 0, 0)$. Then,

$$kp_{\varepsilon}(x,y) + \frac{\partial}{\partial v_{x}} p_{\varepsilon}(x,y)$$

$$= kG(x,y) + kg(\varepsilon)G(x,\tilde{w})G(\tilde{w},y) + kh(\varepsilon)\langle \nabla_{w}G(x,\tilde{w}), \nabla_{w}G(\tilde{w},y)\rangle$$

$$- \frac{\partial}{\partial x_{1}}G(x,y) - g(\varepsilon) \left(\frac{\partial}{\partial x_{1}}G(x,\tilde{w})\right)G(\tilde{w},y)$$

$$- h(\varepsilon)\frac{\partial}{\partial x_{1}}\langle \nabla_{w}G(x,\tilde{w}), \nabla_{w}G(\tilde{w},y)\rangle$$

$$= kG(x,y) - kG(\tilde{w},y) - g(\varepsilon)\frac{\partial}{\partial x_{1}}S(x,\tilde{w})G(\tilde{w},y)$$

S. OZAWA

$$-\frac{\partial}{\partial x_{1}}G(x,y) + \frac{\partial}{\partial w_{1}}G(\tilde{w},y) + kh(\varepsilon) \langle \nabla_{w}S(x,\tilde{w}), \nabla_{w}G(\tilde{w},y) \rangle$$
$$-h(\varepsilon)\frac{\partial}{\partial x_{1}} \langle \nabla_{w}S(x,\tilde{w}), \nabla_{w}G(\tilde{w},y) \rangle + O(\varepsilon)g(\varepsilon)G(\tilde{w},y)$$

for $x = (\varepsilon, 0, 0)$.

Let $G_{\varepsilon}(x,y)$ denote the Green function of Δ in M_{ε} satisfying $G_{\varepsilon}(x,y)=0$ on ∂M and $kG_{\varepsilon}(x,y) + \frac{\partial}{\partial v_{x}}G_{\varepsilon}(x,y)=0$ on ∂B_{ε} .

Let G_{ε} and \tilde{P}_{ε} be the operator defined by

$$G_{\varepsilon}f(x) = \int_{M_{\varepsilon}} G_{\varepsilon}(x, y) f(y) dy$$
$$P_{\varepsilon}f(x) = \int_{M_{\varepsilon}} p_{\varepsilon}(x, y) f(y) dy.$$

We put $Q_{\epsilon}f(x) = P_{\epsilon}f(x)$. Then, it satisfies

$$\Delta Q_{\varepsilon} f(x) = 0 \qquad x \in M_{\varepsilon}$$
$$Q_{\varepsilon} f(x) = 0 \qquad x \in \partial M$$
$$k Q_{\varepsilon} f(x) + \frac{\partial}{\partial v_{x}} Q_{\varepsilon} f(x) = k P_{\varepsilon} f(x) + \frac{\partial}{\partial v_{x}} P_{\varepsilon} f(x) \qquad x \in \partial B_{\varepsilon}$$

We know that

$$(3.1) k P_{\varepsilon}f(x) + \frac{\partial}{\partial v_{x}}P_{\varepsilon}f(x) \\ = k(Gf)(x) - (Gf)(\tilde{w}) - \frac{\partial}{\partial x_{1}}Gf(x) + \frac{\partial}{\partial w_{1}}Gf(\tilde{w}) \\ + O(1)g(\varepsilon)Gf(\tilde{w}) + O(h(\varepsilon))\frac{\partial}{\partial x_{1}}\langle \nabla_{w}S(x,\tilde{w}), \nabla_{w}Gf(\tilde{w}) \rangle \\ + O(h(\varepsilon))\langle \nabla_{w}S(x,\tilde{w}), \nabla_{w}Gf(\tilde{w}) \rangle.$$

4. Proof of Theorem 1.

We have the following decomposition of u_{e} .

(4.1)
$$u_{\varepsilon} = -\lambda(\varepsilon) Q_{\varepsilon} u_{\varepsilon}^{p} + \lambda(\varepsilon) P_{\varepsilon} u_{\varepsilon}^{p}.$$

Lemma 4.1. Fix $p \in (1,2)$. Then,

(4.2)
$$\sup_{x \in M_{\varepsilon}} |\boldsymbol{P}_{\varepsilon} \boldsymbol{u}_{\varepsilon}^{p}| \leq C(1 + \varepsilon \|\boldsymbol{u}_{\varepsilon}\|_{\infty, \varepsilon}^{\tau})$$

holds for any $\tau < 1$ which is close enough to 1.

Proof. Recall that

$$P_{\varepsilon}u_{\varepsilon}^{p}(x) = G\hat{u}_{\varepsilon}^{p}(x) + g(\varepsilon)G(x,\tilde{w})G\hat{u}_{\varepsilon}^{p}(\tilde{w}) + h(\varepsilon)\langle \nabla_{w}G(x,\tilde{w}), \nabla_{w}G\hat{u}_{\varepsilon}^{p}(\tilde{w})\rangle.$$

Here $\hat{u}_{\varepsilon}^{p}(x)$ is the extension of u_{ε}^{p} to M which is zero outside M_{ε} .

We have $|Gu_{\varepsilon}^{p}(\tilde{w})| \leq C ||u_{\varepsilon}^{p}||_{t,\varepsilon} \leq C' ||u_{\varepsilon}||_{tp,\varepsilon}^{p}$ for any t > (3/2). Notice that we can take t > (3/2) as close as 3/2 so that tp < p+1 for $p \in (1,2)$. Therefore, $|G\hat{u}_{\varepsilon}^{p}(w)|$ is bounded.

We have

$$\varepsilon |\nabla_{w} G u_{\varepsilon}^{p}(\tilde{w})| \leq C \varepsilon ||u_{\varepsilon}||_{\infty,\varepsilon}^{\tau} \left(\int_{M_{\varepsilon}} u_{\varepsilon}^{p-\tau}(y) |y-\tilde{w}|^{-2} dy \right)$$
$$\leq C' \varepsilon ||u_{\varepsilon}||_{\infty,\varepsilon}^{\tau} ||u_{\varepsilon}||_{p+1,\varepsilon}^{(p-\tau)/(p+1)} \left(\int_{M_{\varepsilon}} |y-w|^{-2r} dy \right)^{1/r}$$

where $r = (p+1)/(1+\tau)$. Thus, it does not exceed $\varepsilon ||u_{\varepsilon}||_{\infty,\varepsilon}^{\tau}$, since 2r < 3, if we take $p \in (1,2)$ and $\tau < 1$ as close as 1.

Lemma 4.2. Fix $p \in (1,2)$. Then,

$$(4.3) \qquad |\mathbf{Q}_{\varepsilon}u_{\varepsilon}^{p}| \le C\varepsilon^{1-p+(\mu/q)} \|u_{\varepsilon}\|_{\infty,\varepsilon}^{(\mu/q)}$$

holds for q > 6, $q > \mu$.

Proof. By Lemma 2.1 we see that

$$(4.4) \qquad |\mathbf{Q}_{\varepsilon}f(x)| \leq C\varepsilon^2 r^{-1} \|\mathbf{G}\hat{f}\|_{C^{1+(\sigma/2)}(M)}$$

for $\sigma > 1$ by using Lemma 2.1 and (3.1). Here \hat{f} is the extension of f to M which is zero outside M_{e} .

We have

$$\|Gf\|_{C^{1+(\sigma/2)}(M)} \le C \|f\|_{q,\varepsilon}$$

for $q > 3/(1 - (\sigma/2))$. We take $\sigma > 1$ as close as 1. Then, we have

 $(4.5) |Q_{\varepsilon}f(x)| \le C\varepsilon ||f||_{q,\varepsilon}$

for q > 6, $x \in M_{\varepsilon}$.

We have

$$\varepsilon \|u_{\varepsilon}^{p}\|_{q,\varepsilon} \leq \varepsilon \|u_{\varepsilon}\|_{\infty,\varepsilon}^{(\mu/q)} \left(\int_{M_{\varepsilon}} |u_{\varepsilon}|^{pq-\mu} dx \right)^{1/q}.$$

We take $\mu < q$ as close as q and q > 6 as close as 6. Then, $pq - \mu < 6$ for $p \in (1,2)$. On the other hand we have

$$\|u_{\varepsilon}\|_{pq-\mu,\varepsilon} \leq \|\tilde{u}_{\varepsilon}\|_{L^{pq-\mu}(M)} \leq C \|\tilde{u}_{\varepsilon}\|_{H^{1}(M)}$$

by the Sobolev embedding theorem using $pq - \mu < 6$. Here \tilde{u}_{ε} is an extension of u_{ε} in Lemma 2.3 which is different from \hat{u}_{ε} . It should be noted, since $\lambda(\varepsilon)$ is difined as an infimum of a functional so that it is easy to see that $\limsup_{\varepsilon \to 0} \lambda(\varepsilon) < \infty$. Therefore,

$$\|\tilde{u}_{\varepsilon}\|_{H^{1}(M)} \leq \|u_{\varepsilon}\|_{H^{1}(M_{\varepsilon})} + C\varepsilon^{-1}\|u_{\varepsilon}\|_{2,\varepsilon}$$

by taking s=2 in Lemma 2.3. Since u_{ε} is a minimizer of (1.1) and

 $\|u_{\varepsilon}\|_{2,\varepsilon} \leq C$

we see that

$$\|\tilde{u}_{\varepsilon}\|_{H^1(M)} \leq C \varepsilon^{-1}.$$

Therefore,

$$\|u_{\varepsilon}\|_{pq-\mu,\varepsilon} \leq C''\varepsilon^{-1}.$$

Therefore, we get the desired redult.

Proof of Theorem 1. We put $\xi = ||u_{\varepsilon}||_{\infty,\varepsilon}$. By Lemma 4.1 and 4.2 we get

$$\xi \leq C'(1 + \varepsilon \xi^{\tau} + \varepsilon^{1-p+(\mu/q)} \xi^{\mu/q}).$$

Since $p \in (1,2)$, we can take $\mu < q$ as close as q so that $1-p+(\mu/q)>0$. We take $\tau < 1$. Therefore, $\xi \le C''$. We get the desired result.

References

[1] S.S. Lin: Semilinear elliptic equations on singularly perturbed domains, Commun. in Partial Diff. Equations., 16, 617-645 (1991).

NONLINEAR EIGENFUNCTION

- [2] S. Mizohata: Theory of Partial Differential Equations (in Japanese), Iwanami, Tokyo 1965.
- [3] S. Ozawa: Spectra of domains with small spherical Neumann boundary, J. Fac. Sci. Univ. Tokyo Sec IV 30, 53-62 (1983).
- S. Ozawa: Singular variation of the ground state eigenvalue for a semilinear elliptic equation, Tôhoku Math. J. 45, 359-368 (1993).

Department of Mathematics Faculty of Science, Tokyo Institute of Technology Oh-okayama, Meguro-ku Tokyo 152, Japan