



Title	On non-commutative quadratic extensions of a commutative ring
Author(s)	Kanzaki, Teruo
Citation	Osaka Journal of Mathematics. 1973, 10(3), p. 597-605
Version Type	VoR
URL	<a href="https://doi.org/10.18910/10259">https://doi.org/10.18910/10259</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Kanzaki, T.  
Osaka J. Math.  
10 (1973), 597-605

## ON NON-COMMUTATIVE QUADRATIC EXTENSIONS OF A COMMUTATIVE RING

TERUO KANZAKI

(Received January 30, 1973)

Let  $A$  be a ring with the identity element and  $B$  an extension ring of  $A$  with the common identity element.  $B$  is called a quadratic extension of  $A$ , if the residue module  $B/A$  is an invertible  $A$ - $A$ -bimodule, i.e.  $B/A \otimes_A \text{Hom}_A(B/A, A) \approx \text{Hom}_A(B/A, A) \otimes_A B/A \approx A$ . In [4], [5] and [9], one has studied about commutative quadratic extensions. We like to extend these results to non-commutative quadratic extensions. But, in general, it is difficult. In this note, we shall study non-commutative quadratic extensions of a commutative ring. Let  $A$  be a commutative ring with the identity element. Let  $D$  be an  $A$ -algebra with the identity element such that  $D$  is a quadratic extension of a commutative subring  $B$  and  $B$  is a separable quadratic extension of  $A$ . In the section 1, we shall show that if  $A$  has no idempotents other than 0 and 1, then such an  $A$ -algebra  $D$  is either a commutative ring or a central  $A$ -algebra having  $B$  as a maximal commutative subring. We shall say that  $A$ -algebra  $D$  is a *quaternion  $A$ -algebra* with a maximal commutative and separable subalgebra  $B$ , if  $D$  is an  $A$ -algebra mentioned above and is a central separable  $A$ -algebra. In the section 2, we shall show that a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$  is characterized by the separable quadratic extension  $B$  of  $A$  and a non-degenerate hermitian  $B$ -module  $(V, \Phi)$  of rank one. Let  $(V, q)$  be a non-degenerate quadratic  $A$ -module such that  $V$  is a finitely generated projective  $A$ -module with a constant rank two. Then the Clifford algebra  $C(V, q) = C_0(V, q) \oplus C_1(V, q)$  is a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $C_0(V, q)$ . And, the quadratic  $A$ -module  $(V, q)$  is hyperbolic if and only if  $[C_0(V, q)] = 1$  in  $Q_S(A)$ , where  $Q_S(A)$  is the group of separable quadratic extensions of  $A$  (cf. [4], [5] and [9]).

1. Let  $A$  be a commutative ring with the identity element, and  $B$  a commutative and separable quadratic extension of  $A$ . Then  $B$  is characterized by an invertible  $A$ -module  $U$ , an  $A$ -linear map  $f: U \rightarrow A$  and a quadratic form  $q: U \rightarrow A$ , as  $B = A \oplus U$  and  $x^2 = f(x)x + q(x)$  for  $x \in U$  (cf. [4]). Let  $\tau$  be an  $A$ -algebra automorphism of  $B$  defined by  $\tau(a+x) = a + f(x) - x$  for  $a \in A$ ,  $x \in U$ . Then we have  $B^\tau = A$ . Because, if  $x$  is in  $U$  and  $\tau(x) = x$ , then  $f(x) = 0$ , and  $2x = 0$ . From the fact that a bilinear form  $D_{f, q}: U \times U \rightarrow A$ ;  $(x, y) \mapsto f(x)f(y) + 2B_q(x, y)$

is non-degenerate (Theorem 1 in [4]),  $x$  is 0, consequently we have  $B^\tau=A$ . Therefore,  $B$  is a Galois extension of  $A$  with the Galois group  $G(B/A)=\{I, \tau\}$ . If  $A$  has no idempotents other than 0 and 1, then  $G(B/A)=\{I, \tau\}$  is the group of all  $A$ -algebra automorphisms of  $B$ .

Let  $D$  be an  $A$ -algebra which is a quadratic extension of  $B$ . Then we have

(1.1) **Theorem.** *Let  $D$  and  $B$  be as above. If  $A$  has no idempotents other than 0 and 1, then  $D$  is either a commutative ring or a central  $A$ -algebra having the subalgebra  $B$  as a maximal commutative subring.*

Proof. Since the residue  $B$ - $B$ -bimodule  $D/B$  is invertible, there exists an  $A$ -algebra automorphism  $\sigma$  of  $B$  such that  $xb \equiv \sigma(b)x \pmod{B}$  for all  $x \in D$  and  $b \in B$ . Then  $\sigma$  is either  $I$  or  $\tau$ . If  $\sigma=I$ , then for each  $x$  in  $D$ ,  $d_x(b)=xb-bx$  is in  $B$  for all  $b \in B$ . The map  $d_x: B \rightarrow B$  becomes a derivation of  $B$  over  $A$ .  $B$  is separable over  $A$ , hence every derivation of  $B$  over  $A$  is 0, and so  $d_x=0$ . Therefore,  $D$  is a  $B$ -algebra. Since  $D$  is a quadratic extension of  $B$ ,  $D$  is a commutative ring. If  $\sigma=\tau$ , then for each  $x \in D$ ,  $d_x(b)=xb-\tau(b)x$  is in  $B$  for all  $b \in B$ , and the map  $d_x: B \rightarrow B$  is a  $(\tau, I)$ -derivation of  $B$  over  $A$ , i.e.  $d_x(b_1 b_2)=d_x(b_1)b_2+\tau(b_1)d_x(b_2)$  for  $b_1, b_2$  in  $B$ , (cf. p. 170 in [6]). Since  $D/B$  is a projective left  $B$ -module, the exact sequence  $0 \rightarrow B \rightarrow D \rightarrow D/B \rightarrow 0$  is split, i.e. there exists an invertible left  $B$ -submodule  $V$  of  $D$  such that  $D=B \oplus V$ . We consider the commutator ring  $V_D(B)=\{x \in D; xb=bx \text{ for all } b \in B\}$ , then  $V_D(B) \supset B$ . Now, we shall show  $V_D(B) \cap V=0$ . If  $x$  is in  $V_D(B) \cap V$ , we have  $d_x(b)=xb-\tau(b)x=bx-\tau(b)x \in B \cap V=0$ , and so  $\tau(b)x=bx$  for all  $b \in B$ . Since  $B \supset A$  is a Galois extension with the Galois group  $G(B/A)=\{I, \tau\}$ , there exist  $b_1, b_2, \dots, b_r$  and  $c_1, c_2, \dots, c_r$  in  $B$  such that  $\sum_i c_i b_i = 1$  and  $\sum_i c_i \tau(b_i) = 0$ . Then  $x=\sum_i c_i b_i x = \sum_i c_i \tau(b_i) x = 0$ . Consequently, we get  $V_D(B)=B$ , i.e.  $B$  is a maximal commutative subring of  $D$ . Finally, we shall show that the center of  $D$  is  $A$ . Let  $c$  be an element of the center.  $c$  is contained in  $B=V_D(B)$ . For any  $x \in V$ ,  $cx=xc=d_x(c)+\tau(c)x$  in  $B \oplus V=D$ . Therefore, we have  $cx=\tau(c)x$ . Since  $V$  is faithful over  $B$ ,  $c=\tau(c)$ , and  $c$  is contained in  $B^{G(B/A)}=A$ . Therefore,  $A$  is the center of  $D$ .

2. Let  $B$  be a commutative and separable quadratic extension of  $A$ , and  $D$  an  $A$ -algebra such that  $D$  is a quadratic extension of  $B$ . If  $D$  is central separable over  $A$ , then  $B$  is a maximal commutative subring of  $D$ . Because, when we regard  $D$  as  $D \otimes_A B$ -left module by  $d \otimes b \cdot x = dxb$  for  $d \otimes b \in D \otimes_A B$  and  $x \in D$ ,  $D$  is a finitely generated projective  $D \otimes_A B$ -module and  $\text{Hom}_{D \otimes B}(D, D) \approx V_D(B) \supset B$ . For every maximal ideal  $m$  of  $A$ ,  $\text{Hom}_{D \otimes B}(D, D) \otimes_A A_m \approx \text{Hom}_{D_m \otimes B_m}(D_m, D_m) \approx V_{D_m}(B_m) \supset B_m$ . But,  $A_m$  has no idempotents without 0 and 1,  $V_{D_m}(B_m)=B_m$ . Therefore,  $V_D(B)=B$ .  $B$  is a maximal commutative subring of  $D$ . We shall say that  $D$  is a *quaternion  $A$ -algebra* with a maximal commutative and separable subalgebra  $B$ , if  $D$  is an  $A$ -algebra defined above and is central separable over  $A$ . If  $A$  has no idempotents other than 0 and 1, and if  $D$  is non-commutative

and separable over  $A$ , then by (1.1),  $D$  is a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ .

(2.1) **Proposition.** *Let  $D$  be a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ . Then  $D$  is a generalized crossed product of  $B$  and  $G(B/A)$  (defined in [3]). Therefore, there exists an invertible  $B$ - $B$ -submodule of  $D$  such that  $D=B\oplus V$  and  $B=V\cdot V\approx V\otimes_B V$ .*

Proof.  $D$  is a central separable  $A$ -algebra and contains a maximal commutative subalgebra  $B$  which is a Galois extension of  $A$  with the Galois group  $G(B/A)=\{I, \tau\}$ . By Proposition 3 in [3],  $D$  is a generalized crossed product of  $B$  and  $G(B/A)$ , and so  $D$  is written as  $D=J_I\oplus J_\tau$ , where  $J_I=B$  and  $J_\tau=\{x\in D; \tau(b)x=xb \text{ for all } b\in B\}$  are invertible  $B$ - $B$ -bimodules. Furthermore, the map  $f_{\tau,\tau}: J_\tau\otimes_B J_\tau \rightarrow J_I; x\otimes y \mapsto xy$  is a  $B$ - $B$ -isomorphism. Put  $V=J_\tau$ ,  $V$  is the required  $B$ - $B$ -bimodlue.

**DEFINITION.** Let  $B\supset A$  be a commutative and separable quadratic extension which is a Galois extension with Galois group  $G(B/A)=\{I, \tau\}$ . For a left  $B$ -module  $M$  with an  $A$ -bilinear form  $\Phi: M\times M\rightarrow B$ , we shall call  $(M, \Phi)$  a *hermitian  $B$ -module* if it satisfies

- 1)  $\Phi(bx, y)=b\Phi(x, y)$ ,
- 2)  $\Phi(x, y)=\tau(\Phi(y, x))$  for every  $b\in B$  and  $x, y\in M$ .

We shall say that a hermitian  $B$ -module  $(M, \Phi)$  is non-degenerate, if the  $A$ -linear map  $M\rightarrow \text{Hom}_B(M, B); x\mapsto \Phi(-, x)$  is an isomorphism. Let  $(M_1, \Phi_1)$  and  $(M_2, \Phi_2)$  be hiermitian  $B$ -modules. The product  $(M_1, \Phi_1)\otimes(M_2, \Phi_2)$  is defined by  $(M_1\otimes_B M_2, \Phi_1\otimes\Phi_2)$  where  $\Phi_1\otimes\Phi_2: (M_1\otimes_B M_2)\times(M_1\otimes_B M_2)\rightarrow B; (x_1\otimes x_2, y_1\otimes y_2)\mapsto \Phi_1(x_1, y_1)\cdot\Phi_2(x_2, y_2)$ . We denote by  $(B, I)$  a hermitian  $B$ -module defined by  $I(b, b')=b\cdot\tau(b')$  for  $b, b'\in B$ .

If  $M_1$  and  $M_2$  are finitely generated projective  $B$ -modules, and if  $(M_1, \Phi_1)$  and  $(M_2, \Phi_2)$  are non-degenerate hermitian  $B$ -modules, then the product  $(M_1, \Phi_1)\otimes(M_2, \Phi_2)$  is also non-degenerate.

(2.2) **Theorem.** *Let  $D$  be a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ . Then there exists a non-degenerate hermitian  $B$ -module  $(V, \Phi)$  with an invertible  $B$ -bimodule  $V$  such that  $D=B\oplus V$ ,  $xb=\tau(b)x$  for  $b\in B$ ,  $x\in V$  and  $xy=\Phi(x, y)$  for  $x, y\in V$ . Conversely, if  $(V, \Phi)$  is any non-degenerate hermitian  $B$ -module with an invertible  $B$ -left module  $V$ , then an  $A$ -algebra  $D=B\oplus V$  which is defined by  $(b+x)\cdot(b'+x')=bb'+\Phi(x, x')+bx'+\tau(b')x$  for  $b, b'\in B$  and  $x, x'\in V$ , is a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ .*

Proof. Let  $D$  be a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ . By (2.1), there exsists an invertible  $B$ - $B$ -bimodule  $V$

such that  $D=B\oplus V$  and  $V \cdot V=B$ . We define an  $A$ -bilinear map  $\Phi: V \times V \rightarrow B$  by  $\Phi(x, y)=xy$  for  $x, y \in V$ . We shall show that  $(V, \Phi)$  is a non-degenerate hermitian  $B$ -module. Put  $\Psi(x, y)=\Phi(x, y)-\tau(\Phi(y, x))$  for  $x, y$  in  $V$ . For any maximal ideal  $m$  of  $A$ , the localization  $B_m$  is a semilocal ring, therefore  $V_m$  is a free  $B_m$ -module of rank 1. Let  $V_m=B_m v$ ,  $\Psi_m=\Psi \otimes I_m$ ,  $\Phi_m=\Phi \otimes I_m$  and  $\tau_m=\tau \otimes I_m$ . Then we have  $\Psi_m(bv, b'v)v=\Phi_m(bv, b'v)v-\tau_m(\Phi_m(b'v, bv))v=(vb'v)v-v(b'vbv)=b\tau_m(b')v^3-\tau_m(b')bv^3=0$  in  $D_m$ . Therefore, we have  $\Psi_m=0$  for any maximal ideal  $m$  of  $A$ , and so  $\Psi=0$ , i.e.  $\Phi(x, y)=\tau(\Phi(y, x))$  for every  $x, y$  in  $V$ . Since  $V \otimes_B V \rightarrow B$ ;  $x \otimes y \rightsquigarrow xy$  is  $B$ - $B$ -isomorphism from (2.1),  $(V, \Phi)$  is non-degenerate. Conversely, let  $(V, \Phi)$  be any non-degenerate hermitian  $B$ -module with an invertible left  $B$ -module  $V$ . We can make a  $B$ - $B$ -bimodule  $V$  by  $xb=\tau(b)x$  for  $b \in B$ ,  $x \in V$ . Since  $(V, \Phi)$  is non-degenerate, the map  $f_{\tau, \tau}: V \otimes_B V \rightarrow B$ ;  $x \otimes y \rightsquigarrow \Phi(x, y)$  is a  $B$ - $B$ -isomorphism as  $B$ - $B$ -bimodules. By [3], we can construct a generalized crossed product  $\Delta(f, B, \Psi, G)$  of  $B$  and  $G=G(B/A)=\{I, \tau\}$  provided  $\Psi; \Psi(I)=B$ ,  $\Psi(\tau)=V$ , and a factor set  $f=\{I=f_{I, I}, f_{\tau, I}, f_{I, \tau}, f_{\tau, \tau}\}$ , where  $f_{I, \tau}: B \otimes_B V \rightarrow V$ ;  $b \otimes x \rightsquigarrow bx$ ,  $f_{\tau, I}: V \otimes_B B \rightarrow V$ ;  $x \otimes b \rightsquigarrow xb$ . To show the commutativity of the diagrams of the factor set, we need only to show the following commutative diagram:

$$\begin{array}{ccc} V \otimes_B V \otimes_B V & \xrightarrow{I \otimes f_{\tau, \tau}} & V \otimes_B B \\ \downarrow f_{\tau, \tau} \otimes I & & \downarrow f_{\tau, I} \\ B \otimes_B V & \xrightarrow{f_{I, \tau}} & V \end{array}$$

we shall show it by taking the localization with respect to a maximal ideal  $m$  of  $A$ . Then we have  $f_{\tau, \tau} \circ (I \otimes f_{\tau, \tau}) (av \otimes bv \otimes cv)=f_{\tau, I} (av \otimes f_{\tau, \tau} (bv \otimes cv))=av \cdot \Phi(bv, cv)=a\tau(b)c\tau(\Phi(v, v))v=a\tau(b)c\Phi(v, v)v=\Phi(av, bv)cv=f_{\tau, \tau}(av \otimes bv)cv=f_{\tau, \tau} \circ (f_{\tau, \tau} \otimes I) (av \otimes bv \otimes cv)$  for all  $av, bv, cv$  in  $V_m=A_m v$ . Therefore, the diagram is commutative. Thus,  $D=B\oplus V=\Delta(f, B, \Psi, G)$  is an  $A$ -algebra defined the multiplication by  $(b+x) \cdot (b'+x')=bb'+\Phi(x, x')+bx'+\tau(b')x$  for  $b+x, b'+x'$  in  $B \oplus V=D$ . By Proposition 3 in [3],  $D$  is an Azumaya  $A$ -algebra, accordingly  $D=B\oplus V$  is a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ .

We shall call  $(V, \Phi)$  a non-degenerate hermitian  $B$ -module of rank 1 if  $(V, \Phi)$  is a non-degenerate hermitian  $B$ -module and  $V$  is an invertible left  $B$ -module. For a non-degenerate hermitian  $B$ -module of rank 1, we denote by  $D(B, V, \Phi)$  the quaternion  $A$ -algebra  $D$  with a maximal commutative and separable subalgebra  $B$  defined by  $(V, \Phi)$  in (2.2)

(2.3) **Corollary.** *Let  $(V, \Phi)$  and  $(V', \Phi')$  be non-degenerate hermitian  $B$ -modules of rank 1. Then  $(V, \Phi)$  and  $(V', \Phi')$  are isometric if and only if there exists an  $A$ -algebra isomorphism of  $D(B, V, \Phi)$  to  $D(B, V', \Phi')$  which is identity map on  $B$ .*

Let  $(P, q)$  be a quadratic  $A$ -module with a quadratic form  $q: P \rightarrow A$ . We shall call that  $(P, q)$  is a non-degenerate quadratic  $A$ -module of rank  $n$ , if  $P$  is a finitely generated projective  $A$ -module with constant rank  $n$ , i.e.  $[P_m : A_m] = n$  for every maximal ideal  $m$  of  $A$ , and  $q: P \rightarrow A$  is non-degenerate.

(2.4) **Proposition.** *Let  $(V, q)$  be a non-degenerate  $A$ -module of rank 2. Then the Clifford algebra  $C(V, q) = C_0(V, q) \oplus C_1(V, q)$  is a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $C_0(V, q)$ , where  $C_0(V, q)$  (resp.  $C_1(V, q)$ ) is the subalgebra of  $C(V, q)$  of homogeneous elements of degree 0 (resp. degree 1).)*

Proof.  $C(V, q)$  is an Azumaya  $A$ -algebra, and  $C_0(V, q)$  is a commutative and separable quadratic extension of  $A$  (Lemma 6 in [7]). Therefore,  $V \approx C_1(V, q)$  is a finitely generated projective  $C_0(V, q)$ -module. We shall show that  $C_1(V, q)$  is an invertible  $C_0(V, q)$ -module. It suffices to show that for the case where  $A$  is a local ring. Assume that  $A$  is a local ring. Then,  $V = Au \oplus Av$  is a free  $A$ -module of rank 2. Since  $(V, q)$  is non-degenerate, we may assume that  $q(u)$  is invertible in  $A$ . Then we have  $C_0(V, q) = A \oplus Auv$  and  $C_1(V, q) \approx V = Au \oplus Av = C_1(V, q)u$ . Since  $u$  is invertible in  $C(V, q)$ ,  $C_1(V, q)$  is a free  $C_0(V, q)$ -module of rank 1.

(2.5) **Lemma.** *Let  $\Lambda$  be a Galois extension of a ring  $\Gamma$  with a Galois group  $G$ , and  $P$  a  $\Lambda$ -module. Then we have  $\text{Hom}_\Gamma(P, \Gamma) = \text{Tr} \circ \text{Hom}_\Lambda(P, \Lambda)$ , where  $\text{Tr}(x) = \sum_{\sigma \in G} \sigma(x)$  for  $x \in \Lambda$ .*

Proof. Since  $\Lambda \supset \Gamma$  is a Galois extension with a Galois group  $G$ , there exist  $x_1, x_2, \dots, x_n$  and  $y_1, y_2, \dots, y_n$  in  $\Lambda$  such that  $\sum_i \sigma(x_i)y_i = \begin{cases} 1, & \sigma = \text{I} \\ 0, & \sigma \neq \text{I} \end{cases}$ . Then, for  $f$  in  $\text{Hom}_\Gamma(P, \Gamma)$ ,  $F(-) = \sum_i x_i f(y_i -)$  is contained in  $\text{Hom}_\Lambda(P, \Lambda)$ , and  $\text{Tr} \circ F(z) = \sum_i \text{Tr}(x_i f(y_i z)) = f(\sum_i \text{Tr}(x_i) y_i z) = f(z)$  for all  $z \in P$ . Therefore,  $f$  is in  $\text{Tr} \circ \text{Hom}_\Lambda(P, \Lambda)$ . The converse is clear.

(2.6) **Lemma.** *Let  $(P, \Phi)$  be a non-degenerate hermitian  $B$ -module. Then  $(P, \text{Tr} \circ \Phi)$  is a non-degenerate bilinear  $A$ -module.*

Proof.  $\text{Tr} \circ \Phi: P \times P \rightarrow A; (x, y) \mapsto \text{Tr}(\Phi(x, y)) = \Phi(x, y) + \tau(\Phi(x, y))$  is an  $A$ -bilinear form. We show that  $P \rightarrow \text{Hom}_A(P, A); x \mapsto \text{Tr}(\Phi(-, x))$  is an  $A$ -isomorphism. If  $x$  is in  $P$  such that  $\text{Tr}(\Phi(-, x)) = 0$ ,  $\Phi(P, x)$  is an ideal of  $B$  and  $\text{Tr}(\Phi(P, x)) = 0$ . Let  $b_1, b_2, \dots, b_n$  and  $b'_1, b'_2, \dots, b'_n$  be elements in  $B$  such that  $\sum_i b_i b'_i = 1$  and  $\sum_i \tau(b_i) b'_i = 0$ . Then, we have  $b = \sum_i \text{Tr}(bb_i) b'_i = 0$  for every  $b$  in  $\Phi(P, x)$ , hence  $\Phi(P, x) = 0$ . Therefore,  $x = 0$ . From Lemma (2.5),  $(P, \text{Tr} \circ \Phi)$  is non-degenerate.

(2.7) **Theorem.** *Let  $D = D(B, V, \Phi)$  be quaternion  $A$ -algebra with a maximal*

commutative and separable subalgebra  $B$ . Then there exists an involution  $\sigma: D \rightarrow D$  which is defined by  $\sigma(b+v) = \tau(b) - v$  for  $b \in B, v \in V$ . We put  $N(x) = x \cdot \sigma(x)$  and  $T(x) = x + \sigma(x)$  for  $x$  in  $D$ . Then  $N: D \rightarrow A$  is a quadratic form,  $(D, N)$  is a non-degenerate quadratic  $A$ -module of rank 4, and  $D = B \perp V$ .  $T: D \rightarrow A$  is an  $A$ -linear map and  $B_N(x, y) = T(x \cdot \sigma(y))$  for  $x, y \in D$ .

Proof. Let  $x = b + v$  and  $x' = b' + v'$  be elements in  $D = B \oplus V$ . Then we have  $\sigma(xx') = \sigma(bb' + \Phi(v, v') + bv' + \tau(b')v) = \tau(bb' + \Phi(v, v')) - (bv' + \tau(b')v) = \tau(b) \cdot \tau(b') + \Phi(v, v') - v'\tau(b) - \tau(b')v = \sigma(x') \cdot \sigma(x)$ , and  $\sigma^2(x) = x$ . Therefore,  $\sigma$  is an involution. Furthermore,  $N(b+v) = b\sigma(b) - \Phi(v, v)$  and  $T(b+v) = b + \tau(b)$  are contained in  $B^G = A$ , hence  $N: D \rightarrow A$  is a quadratic form, and the bilinear form is  $B_N(x, x') = N(x+x') - N(x) - N(x') = x\sigma(x') + x'\sigma(x) = T(x\sigma(x))$  for  $x, x' \in D$ . Therefore, we have  $D = B \perp V$ . To prove that  $(D, N)$  is non-degenerate, it suffices to show that  $(B, N|B)$  and  $(V, N|V)$  are non-degenerate. From Lemma (2.6),  $\text{Tr} \circ \Phi$  and  $\text{Tr} \circ I$  are non-degenerate, and  $B_N(b, b') = T(b\tau(b')) = \text{Tr}(b\tau(b')) = \text{Tr} \circ I(b, b')$  for  $b, b' \in B$  and  $B_N(v, v') = T(v(-v')) = \text{Tr}(-\Phi(v, v')) = -\text{Tr} \circ \Phi(v, v')$  for  $v, v' \in V$ , hence  $(B, N|B)$  and  $(V, N|V)$  are non-degenerate.

In Theorem (2.7), we put  $Q = -N|V$ . Then  $(V, Q)$  is a non-degenerate quadratic  $A$ -module of rank 2.

(2.8) **Theorem.** Let  $D(B, V, \Phi)$  be a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ , and  $N: D \rightarrow A$  and  $Q = -N|V$  as defined before. Then  $D(B, V, \Phi)$  is isomorphic to the Clifford algebra  $C(V, Q)$  of the quadratic module  $(V, Q)$  as  $A$ -algebras.

Proof. Since  $Q(x)$  is equal to  $x^2 = N(x)$  in  $D(B, V, \Phi)$  for every  $x \in V$ , the inclusion map  $V \rightarrow D(B, V, \Phi) = B \oplus V$  can be extended to an  $A$ -algebra homomorphism  $\rho: C(V, Q) \rightarrow D(B, V, \Phi)$ . From the fact that  $C(V, Q)$  and  $D(B, V, \Phi)$  are Azumaya algebras over  $A$  and are generated by  $V$ , we obtain that  $\rho$  is an  $A$ -isomorphism.

(2.9) **Lemma.** Let  $V$  be any invertible  $B$ -module. Then for any  $f$  in  $\text{Hom}_B(V, B)$  and  $x, y$  in  $V$ , we have  $f(x)y = f(y)x$ .

Proof. Put  $\Psi(x, y) = f(y)x - f(x)y$  for every  $x, y \in V$ , then  $\Psi: V \times V \rightarrow V$  is a  $B$ -bilinear form. By taking the localization of  $V$  with respect to a maximal ideal  $m$  of  $A$ , we get easily  $\Psi_m = 0$ . Therefore,  $\Psi = 0$ .

(2.10) **Proposition.** Let  $(V, \Phi)$  be a non-degenerate hermitian  $B$ -module of rank 1. Then, the quaternion  $A$ -algebra  $D(B, V \otimes_B V, \Phi \otimes \Phi)$  which is determined by  $(V, \Phi) \otimes (V, \Phi) = (V \otimes_B V, \Phi \otimes \Phi)$ , is  $A$ -algebra isomorphic to  $\text{Hom}_A(V, V)$ , and this isomorphism preserves the structure of  $B$ -modules.

**Proof.** We can define a map  $\theta: D(B, V \otimes_B V, \Phi \otimes \Phi) = B \oplus V \otimes_B V \rightarrow \text{Hom}_A(V, V)$  as follows:  $\theta(b)(x) = bx$  for  $b \in B$ ,  $x \in V$ , and  $\theta(u \otimes v)(x) = \Phi(u, x)v$  for  $u \otimes v \in V \otimes_B V$ ,  $x \in V$ . Then  $\theta$  is an  $A$ -algebra homomorphism. Because, for  $b \in B$ ,  $u \otimes v \in V \otimes_B V$  and  $x \in V$ , we have  $\theta(bu \otimes v)(x) = \Phi(bu, x)v = b\Phi(u, x)v = \theta(b) \circ \theta(u \otimes v)(x)$  and  $\theta(u \otimes vb)(x) = \Phi(u, x)vb = \tau(b)\Phi(u, x)v = \Phi(u, bx)v = \theta(u \otimes v) \circ \theta(b)(x)$ . And, for  $u \otimes v, u' \otimes v' \in V \otimes_B V$  and  $x \in V$ ,  $\theta(u \otimes v) \circ \theta(u' \otimes v')(x) = \theta(u \otimes v)(\Phi(u', x)v') = \Phi(u, \Phi(u', x)v')v = \Phi(u, v')\Phi(x, u')v$ . On the other hand,  $\Phi(-, v')$  and  $\Phi(-, u')$  are in  $\text{Hom}_B(V, B)$ , by Lemma (2.9) we get  $\Phi(x, u')\Phi(u, v')v = \Phi(x, u')\Phi(v, v')u = \Phi(u, u')\Phi(v, v')x = \theta(\Phi(u, u')\Phi(v, v'))(x) = \theta((u \otimes v)(u' \otimes v'))(x)$ . Thus,  $\theta$  is an  $A$ -algebra homomorphism. Now we check that  $\theta$  is an epimorphism. From Lemma (2.5), we have  $\text{Hom}_A(V, V) \approx \text{Hom}_A(V, A) \otimes_A V \approx \text{Tr} \circ \text{Hom}_B(V, B) \otimes_A V \approx (\text{Tr} \circ \Phi(-, V)) \otimes_A V$ . Therefore, any element  $f$  in  $\text{Hom}_A(V, V)$  is written as  $f = \sum_i \text{Tr} \circ \Phi(-, u_i)v_i = \sum_i (\Phi(-, u_i)v + \Phi(u_i, -)v_i)$  for some  $u_i, v_i \in V$ , and by Lemma (2.9),  $f(x) = \sum_i \Phi(x, u_i)v_i + \sum_i \Phi(u_i, x)v_i = \sum_i \Phi(v_i, u_i)x + \theta(\sum_i u_i \otimes v_i)(x)$  for  $x \in V$ . Thus, we get  $f = \theta(\sum_i \Phi(v_i, u_i) + \sum_i u_i \otimes v_i)$ . Since  $D(B, V \otimes_B V, \Phi \otimes \Phi)$  and  $\text{Hom}_A(V, V)$  are Azumaya  $A$ -algebras,  $\theta$  is an  $A$ -algebra isomorphism.

(2.11) **Corollary.**  $D(B, B, I) \approx \text{Hom}_A(B, B)$  as  $A$ -algebras.

(2.12) **Corollary.** For any non-degenerate hermitian  $B$ -modules of rank 1  $(V, \Phi)$  and  $(V, \Phi')$ ,  $(V \otimes_B V, \Phi \otimes \Phi)$  and  $(V \otimes_B V, \Phi' \otimes \Phi')$  are isometric.

(2.13) **Theorem.** Let  $D(B, V, \Phi)$  be a quaternion  $A$ -algebra with a maximal commutative and separable subalgebra  $B$ , and  $(V, Q)$  a non-degenerate quadratic  $A$ -module of rank 2 defined by  $D(B, V, \Phi)$  in (2.8). Then,  $(V, Q)$  is hyperbolic if and only if  $[B]=1$  in  $Q_S(A)$  (cf. [4]).

**Proof.** In (2.8), we obtained  $D(B, V, \Phi) = C(V, Q) = B \oplus V$ ,  $C_0(V, Q) = B$  and  $C_1(V, Q) = V$ . We assume that  $(V, Q)$  is hyperbolic. Then we may assume that  $V = P \oplus P^*$  for some invertible  $A$ -module  $P$  and  $P^* = \text{Hom}_A(P, A)$ , and  $Q(x+f) = f(x)$  for  $x \in P, f \in P^*$ . Since  $P \cdot P = P^* \cdot P^* = 0$  in  $C(V, Q)$ , we get  $C_0(V, Q) = A \oplus P \cdot P^* \approx A \oplus P \otimes_A P^*$ . For any  $\sum_i x_i f_i$  in  $P \cdot P^*$ , we have  $(\sum_i x_i f_i)^2 = \sum_{i,j} x_i f_i x_j f_j = \sum_{i,j} x_i (f_i(x_j) - x_j f_i) f_j = \sum_{i,j} f_i(x_j) x_i f_j = \sum_{i,j} f_i(x_i) x_j f_j = (\sum_i f_i(x_i)) (\sum_i x_i f_i)$  using Lemma (2.9). We consider an  $A$ -isomorphism  $\mu: P \cdot P^* (\approx P \otimes_B P^*) \rightarrow A$  defined by  $\mu(\sum_i x_i f_i) = \sum_i f_i(x_i)$  for  $\sum_i x_i f_i$  in  $P \cdot P^*$ . Then we have  $(\sum_i x_i f_i)^2 = \mu(\sum_i x_i f_i) \sum_i x_i f_i$  for every  $\sum_i x_i f_i$  in  $P \cdot P^*$ , hence  $B = C_0(V, Q) \approx (P \otimes_A P^*, \mu, 0) \approx (A, 1, 0)$  as  $A$ -algebras (cf. [4]). Accordingly,  $[B]=1$  in  $Q_S(A)$ . Conversely, we assume  $[B]=1$  in  $Q_S(A)$ . Then the quadratic extension  $B$  of  $A$  has idempotents  $e_1$  and  $e_2$  such that  $1 = e_1 + e_2$ ,  $e_1 e_2 = 0$  and  $B = Ae_1 \oplus Ae_2$ . Furthermore,  $A$ -module  $V$  is written as a direct sum of  $A$ -submodules  $e_1 V$  and  $e_2 V$ . Since the Galois group  $G = G(B/A) = \{I, \tau\}$  is permutations of  $\{e_1, e_2\}$ , we have  $Q(e_1 x) = \Phi(e_1 x, e_1 x) = e_1 \tau(e_1) \Phi(x, x) = e_1 e_2 \Phi(x, x) = 0$  for every

$x \in V$ . Therefore,  $e_1 V$  is totally isotropic. We have  $(e_1 V)^\perp = e_1 V$ , because, for  $e_1 y + e_2 z \in (e_1 V)^\perp$ ,  $0 = B_Q(e_1 y + e_2 z, e_1 x) = \Phi(e_1 y + e_2 z, e_1 x) + \Phi(e_1 x, e_1 y + e_2 z) = \Phi(e_2 z, e_1 x) + \Phi(e_1 x, e_2 z) = e_2 \Phi(z, x) + e_1 \Phi(x, z)$  in  $e_2 A \oplus e_1 A = B$ , hence  $e_1 \Phi(x, z) = \Phi(x, e_2 z) = 0$  for all  $z$  in  $V$ . Therefore, we get  $e_2 z = 0$ . Accordingly,  $(V, Q)$  is hyperbolic (cf. [2]).

(2.14) **Corollary.** *Let  $(P, q)$  be any non-degenerate quadratic  $A$ -module of rank 2. Then  $(P, q)$  is hyperbolic if and only if  $[C_0(P, q)] = 1$  in  $Q_S(A)$ .*

(2.15) **Corollary.** *If  $B$  is a quadratic extension of  $A$  such that  $[B] = 1$  in  $Q_S(A)$ , then every quaternion  $A$ -algebra  $D(B, V, \Phi)$  with a maximal commutative and separable subalgebra  $B$  is split, i.e.  $[D(B, V, \Phi)] = 1$  in the Brauer group  $B(A)$ .*

(2.16) **Corollary.** *If  $A$  is commutative ring such that  $Q_S(A) = 1$ , then every non-degenerate quadratic  $A$ -module of rank 2 is hyperbolic.*

(2.17) **EXAMPLE.** *If  $A$  is the integers  $\mathbb{Z}$  or the gaussian integers  $\mathbb{Z}[i]$ , then every non-degenerate quadratic  $A$ -module of rank 2 is hyperbolic (cf. [5], [7]).*

(2.18) **REMARK.** Let  $K$  be a field, and  $(V, q)$  and  $(V', q')$  non-degenerate quadratic  $K$ -modules of rank 2. Then,  $(V, q)$  and  $(V', q')$  are isometric if and only if  $[C(V, q)] = [C(V', q')]$  in the Brauer group  $B(K)$  and  $[C_0(V, q)] = [C_0(V', q')]$  in  $Q_S(A)$ .

**Proof.** For a field of characteristic  $\neq 2$ , this is obtained from Theorem 58:4 in [8], and for a field of characteristic 2, is obtained from Theorem 3 in [1].

OSAKA CITY UNIVERSITY

---

### References

- [1] C. Arf: *Untersuchungen über quadratische Formen in Körpern der Charakteristik 2*, J. Reine Angew. Math. **183** (1941), 148–167.
- [2] H. Bass: *Lectures on Topics in Algebraic K-theory*, Tata Institute of Fundamental Research, Bombay, 1967.
- [3] T. Kanzaki: *On generalized crossed product and Brauer group*, Osaka J. Math. **5** (1968), 175–188.
- [4] ———: *On the quadratic extensions and the extended Witt ring of a commutative ring*, Nagoya Math. J. **49** (1973), 127–141.
- [5] K. Kitamura: *On free quadratic extensions over a commutative ring*, Osaka J. Math. **10** (1973), 15–20.
- [6] N. Jacobson: *Structure of Rings*, Providence, R.I., 1964.
- [7] A. Micali and O.E. Villamayor: *Sur les algèbres de Clifford. II*, J. Reine Angew. Math. **242** (1970), 61–90.

- [8] O.T. O'Meara: *Introduction to Quadratic Forms*, Berlin-Göttingen-Heidelberg: Springer, 1963.
- [9] C. Small: *The group of quadratic extensions*, *J. Pure Appl. Algebra* **2** (1972), 83–105.

