



| | |
|--------------|--|
| Title | CONNECTIVITY OF JULIA SETS OF HÉNON MAPS NEAR THE BOUNDARY OF THE MANDELBROT SET |
| Author(s) | Yagi, Takumi |
| Citation | Osaka Journal of Mathematics. 2025, 62(4), p. 609-632 |
| Version Type | VoR |
| URL | https://doi.org/10.18910/102980 |
| rights | |
| Note | |

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

CONNECTIVITY OF JULIA SETS OF HÉNON MAPS NEAR THE BOUNDARY OF THE MANDELBROT SET

TAKUMI YAGI

(Received May 27, 2024)

Abstract

In this paper, we prove some new connectivity of the Julia sets J of the complex Hénon maps $H(x, y) = (x^2 + c + ay, ax)$ with sufficiently small $|a|$. We investigate the connectivity of J for the parameters near the boundary of the Mandelbrot set. We first give some conditions related to the connectivity of J for sufficiently small $|a|$, which are useful for considering the connectivity of J for the parameters near the boundary of the Mandelbrot set. We consider a perturbation $\{H_{a,\lambda_t}\}_{a \in \mathbb{D}_{\delta_0}, 0 \leq t < \delta_0}$ of dissipative semi-parabolic Hénon maps H_{a,λ_0} such that $\det DH_{a,\lambda_t} = -a^2$ and H_{a,λ_t} has a fixed point \mathbf{q}_{a,λ_t} for which $(DH_{a,\lambda_t})_{\mathbf{q}_{a,\lambda_t}}$ has an eigenvalue λ_t . Assume that $\lambda_t \rightarrow \lambda_0 = \exp(2\pi i m/l) \in \partial \mathbb{D}$ as $t \rightarrow 0$ and λ_t^l can be represented by $\exp(L_t + i\theta_t)$ with $L_t \neq 0$ for $0 < t < \delta_0$. We prove that if $\theta_t = O(L_t)$, then the Julia sets J_{a,λ_t} for $a \in \mathbb{D}_{\delta_0}, 0 < t < \delta_0$ are connected by using the conditions above.

1. Introduction

In this paper, we deal with the connectivity of the Julia sets of complex Hénon maps.

In one-dimensional (1-D) complex dynamics, we consider a complex polynomial $f_c(x) = x^2 + c, c \in \mathbb{C}$ and the *Julia set* J_{f_c} of f_c . The Julia set J_{f_c} of f_c is defined by the boundary of the *filled Julia set* $K_{f_c} := \{z \in \mathbb{C} : \{f_c^n(z)\}_{n \in \mathbb{N}} \text{ is bounded}\}$. Note that the notation f_c^n is the n -fold composition of f_c . The Mandelbrot set \mathcal{M} is defined by $\{c \in \mathbb{C} : J_{f_c} \text{ is connected}\}$. It is known that the Julia set J_{f_c} of a polynomial f_c is connected if and only if K_{f_c} contains the critical point 0 of f_c in \mathbb{C} (see [13]). By using the fact, it is easy to find the boundary of the connectedness locus for parameters $c \in \mathbb{C}$. For example, the Julia set $J_{f_{1/4}}$ is connected, and the parameter $1/4$ belongs to the boundary of the connectedness locus. Indeed, $J_{f_{1/4+\varepsilon}}$ is connected if $\varepsilon < 0$, and $J_{f_{1/4+\varepsilon}}$ is disconnected if $\varepsilon > 0$. The parameter $c = 1/4$ is called a *parabolic parameter* since $f_{1/4}$ has a parabolic fixed point $1/2$. Let us consider how the parameters c for which J_{f_c} is connected can approach parabolic parameters. Let us explain this by using perturbations of multipliers of parabolic fixed points. We say that a point $\alpha \in \mathbb{C}$ is a *parabolic fixed point* of f_c if $f_c(\alpha) = \alpha$ and $f'_c(\alpha)$ is a root of unity. Here we consider the case where f_{c_0} has a parabolic fixed point α_0 with multipliers $\lambda_0 := \exp(2\pi i m/l)$, where $l \in \mathbb{Z} \setminus \{0\}$, $m \in \mathbb{Z}$ and $(m, l) = 1$. Consider a one-parameter continuous family $\{\lambda_t\}_{t \in [0, \delta_0]}$, where $\delta_0 > 0$. Assume that $\lambda_t^l = \exp(L_t + i\theta_t)$ and $\mathbb{R} \ni \theta_t \rightarrow 0$ as $t \rightarrow 0$, where $L_t \in \mathbb{R} \setminus \{0\}$ and $\theta_t \in \mathbb{R}$ for $0 < t < \delta_0$. Let $\{f_{c_t}\}_{t \in [0, \delta_0]}$ satisfy that f_{c_t} has a fixed point α_t with multiplier λ_t . We say that λ_t^l converges to 1 *radially* if $\theta_t = O(L_t)$. If $\theta_t = O(L_t)$, then $\{f_{c_t}\}_{t \in [0, \delta_0]}$ has nice properties (such as continuity of $J_{f_{c_t}}$, continuity of the Hausdorff dimension of $J_{f_{c_t}}$) (see

[12]). In particular, we see that J_{c_t} is connected for each $t \in (0, \delta_0)$, taking a smaller $\delta_0 > 0$ if necessary.

Let us explain radial convergence by observing the main cardioid $\mathcal{M}_0 := \{c = \lambda/2 - \lambda^2/4 : |\lambda| \leq 1\}$ of the Mandelbrot set \mathcal{M} (Figure 1). For this purpose, we set $p_\lambda(x) := x^2 + \lambda/2 - \lambda^2/4$, which has a fixed point $\lambda/2$ with multiplier λ . The parameter $c = 1/4$ is a parabolic parameter. That is, the polynomial $p_1(x) = x^2 + 1/4$ has a parabolic fixed point $1/2$. Consider a family $\{p_{\lambda_t}\}$ with $\lambda_t \rightarrow 1$ as $t \rightarrow 0$. We set $c_t = \lambda_t/2 - \lambda_t^2/4$ and $\lambda_t = \exp(L_t + i\theta_t)$. If $\theta_t = 0$, then the parameters c_t approach $1/4$ in $\text{Int } \mathcal{M}_0 \cap \mathbb{R}$ (the first of Figure 1). If parameters c_t in the sector in the second of Figure 1 approach $1/4$, then λ_t satisfies $\theta_t = O(L_t)$. We remark that there is a family $\{\lambda_t\}$ such that the corresponding parameters $c_t \in \text{Int } \mathcal{M}_0$ approach $1/4$ as $t \rightarrow 0$. For example, if parameters c_t approach $1/4$ in the curve near $\partial \mathcal{M}_0$ in the third of Figure 1, then λ_t satisfies $\theta_t^2 = o(L_t)$ and $\theta_t \neq O(L_t)$. On the other hand, if parameters c_t approach $1/4$ in $\mathbb{R}_{>1/4}$, then $J_{f_{c_t}}$ is disconnected and $\theta_t \neq O(L_t)$ (see fourth of Figure 1).




Fig. 1. $\theta_t = 0$ (first), $\limsup_{t \rightarrow 0} |\theta_t/L_t| \leq 1/\sqrt{3}$ (second), $\theta_t^2 = o(L_t), \theta_t \neq O(L_t)$ (third) and disconnected (fourth)

In the case of two-dimensional (2-D) dynamics, for $(c, a) \in \mathbb{C}^2$, we consider the quadratic *Hénon map* of the form $H(x, y) = (x^2 + c + ay, ax)$. For a diffeomorphism $F(x, y) = (F_1(x, y), F_2(x, y))$ from an open set $U \subset \mathbb{C}^2$ to \mathbb{C}^2 , we set

$$(DF)_{(x_0, y_0)}(\zeta, \eta) := \begin{pmatrix} (F_1)_x(x_0, y_0) & (F_1)_y(x_0, y_0) \\ (F_2)_x(x_0, y_0) & (F_2)_y(x_0, y_0) \end{pmatrix} \begin{pmatrix} \zeta \\ \eta \end{pmatrix},$$

for $(x_0, y_0) \in U$ and $(\zeta, \eta) \in T_{(x_0, y_0)}U$. We have

$$(DH)_{(x, y)} = \begin{pmatrix} 2x & a \\ a & 0 \end{pmatrix},$$

for $(x, y) \in \mathbb{C}^2$. The map H has constant Jacobian $-a^2$, i.e., $\det(DH)_{(x, y)} = -a^2$ for all $(x, y) \in \mathbb{C}^2$. Unlike 1-D dynamics, we can consider the inverse H^{-1} of H if $a \neq 0$. Let K^\pm be the set of all points $(x, y) \in \mathbb{C}^2$ such that $\{H^{\pm n}(x, y)\}_{n \in \mathbb{N}}$ is bounded in \mathbb{C}^2 . We consider the *Julia sets* $J^\pm := \partial K^\pm$ of H . Furthermore we denote J by the intersection of J^+ and J^- . It is known that J^\pm are connected (see [2]). The *Hénon connectedness locus* is the set of parameters $(c, a) \in \mathbb{C} \times \mathbb{C} \setminus \{0\}$ for which the Julia set J is connected. Let us consider the condition that J is connected. Unlike one-dimensional dynamics, Hénon maps H do not have critical points for $a \neq 0$. Instead, it was suggested to consider critical points of the Green functions along the unstable manifolds of the saddle points to compute the connectivity of the Julia sets (see [4]). The Julia set J is connected if and only if the restriction of the Green function $G^+(x, y) := \lim_{n \rightarrow \infty} (1/2^n) \log^+ \|H^n(x, y)\|$ on the unstable manifold of some saddle point has no critical points in K^+ (see [4] and [10, Theorem 3.3]). However, it is not easy

to find the boundary of the Hénon connectedness locus. Our result (Theorem 1.2) describes the local geometry near semi-parabolic parameters (c, a) if $|a|$ is sufficiently small.

Let $\lambda \in \mathbb{C} \setminus \{0\}$. To consider the connectivity of J for the parameters near the boundary of the Mandelbrot set, we consider a Hénon family for which each element of the family has a fixed point such that one of the eigenvalues of DH at the fixed point is λ . Then, the set \mathcal{P}_λ of parameters $(c, a) \in \mathbb{C}^2$ for which the Hénon map $H(x, y) = (x^2 + c + ay, ax)$ has a fixed point \mathbf{q} such that λ is an eigenvalue of $(DH)_{\mathbf{q}}$ is the curve of equation

$$(1) \quad c = (1 - a^2) \left(\frac{\lambda}{2} - \frac{a^2}{2\lambda} \right) - \left(\frac{\lambda}{2} - \frac{a^2}{2\lambda} \right)^2.$$

We denote the right hand side of equation (1) by $c(a, \lambda)$. Moreover, we set $H_{a, \lambda}(x, y) = (x^2 + c(a, \lambda) + ax, ay)$ and $p_\lambda(x) = x^2 + c(0, \lambda)$. We denote the Julia sets of $H_{a, \lambda}$ by $J_{a, \lambda}^\pm, J_{a, \lambda}$ instead of J^\pm, J respectively. Based on the above notations, we now present the first main result of this paper.

Theorem 1.1. *Assume that a Hénon family $\{H_{a, \lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the vertical condition $(VC)_{\varepsilon, r}$ with respect to $\varepsilon > 0, r > 0$ (see Definition 3.1). Suppose that $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a, \lambda}^+ = \emptyset$ for each $a \in \mathbb{D}_{\delta_0}$. Then the Julia sets $J_{a, \lambda}$ of the Hénon maps $H_{a, \lambda}$ for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$ are connected if and only if the Julia set J_{p_λ} of the polynomial p_λ is connected.*

We regard $\mathbb{D}_\varepsilon \times \mathbb{D}_r$ as a neighborhood of the critical point $z = 0$ of p_λ in two dimensions. Note that most families $\{H_{a, \lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the $(VC)_{\varepsilon, r}$. Indeed, if $\lambda \neq 1$, then there is $\delta_0 > 0$ such that $\{H_{a, \lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the $(VC)_{\varepsilon, r}$ (see Lemma 3.2). The assumptions of Theorem 1.1 imply that the stable manifold of a saddle fixed point of $H_{a, \lambda}$ intersects transversely horizontal direction in $\mathbb{D}_r \times \mathbb{D}_r$. From this, we can construct a holomorphic motion of $J_{a, \lambda}^+ \cap (\mathbb{C} \times \{y\})$ over $a \in \mathbb{D}_{\delta_0}$ for each $y \in \mathbb{D}_r$ and can show that $J_{a, \lambda}^+ \cap (\mathbb{C} \times \{y\})$ is homeomorphic to the Julia set J_{p_λ} of p_λ .

It is known that for a hyperbolic polynomial $x^2 + c$, there is a positive constant $\delta(c) > 0$ such that a small perturbation $\{H(x, y) = (x^2 + c + ay, ax) : 0 < |a| < \delta(c)\}$ of $H(x, y) = (x^2 + c, 0)$ is hyperbolic (see [6] and [9]). In particular, the Julia sets J of the Hénon maps $H(x, y) = (x^2 + c + ay, ax)$ for $0 < |a| < \delta(c)$ are connected if and only if J_{x^2+c} is connected. However, the proofs in [6] and [9] do not give any uniform estimate on the constant $\delta(c)$ from below for c near the boundary of the Mandelbrot set. For example, it may be $\delta(c_n) \rightarrow 0$ as $\text{Int } \mathcal{M}_0 \ni c_n \rightarrow \lambda/2 - \lambda^2/4 \in \partial \mathcal{M}_0$, where $|\lambda| = 1$. Therefore, we cannot apply methods of [6] and [9] to compute the connectivity of J for the parameters near the boundary of the Mandelbrot set. In our result, we only need to check that the $(VC)_{\varepsilon, r}$ and the condition $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a, \lambda}^+ = \emptyset$ hold for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$. We can deduce the connectivity of J for the parameters near the boundary of the Mandelbrot set by using Theorem 1.1 (see the following Theorem 1.2 and Figure 2).

To present the second main result, we recall radial convergence. Let $\lambda_0 = \exp(2\pi i m/l)$, where $l \in \mathbb{Z} \setminus \{0\}$, $m \in \mathbb{Z}$ and $(m, l) = 1$. Consider a one-parameter continuous family $\{\lambda_t\}_{t \in [0, \delta_0]}$, where $\delta_0 > 0$. Assume that $\lambda_t^l = \exp(L_t + i\theta_t)$ and $\mathbb{R} \ni \theta_t \rightarrow 0$ as $t \rightarrow 0$, where $L_t \in \mathbb{R} \setminus \{0\}$ and $\theta_t \in \mathbb{R}$ for $0 < t < \delta_0$. We say that $RD_{\lambda_t, \delta_0} := \{H_{a, \lambda_t} : a \in \mathbb{D}_{\delta_0} \text{ and } 0 < t < \delta_0\}$ is a *radial* perturbation if $\theta_t = O(L_t)$. For each $0 < t < \delta_0$, we will show that the section $\{H_{a, \lambda_t}\}_{a \in \mathbb{D}_{\delta_0}}$ of RD_{λ_t, δ_0} satisfies the $(VC)_{\varepsilon, r}$ and that $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a, \lambda_t}^+ = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$. By applying Theorem 1.1 to the family RD_{λ_t, δ_0} , we can show the second main result:

Theorem 1.2. *There is $\delta_0 > 0$ such that each $H_{a,\lambda_t} \in RD_{\lambda_t, \delta_0}$ with $a \neq 0$ has connected Julia set J_{a,λ_t} .*

Note that H_{a,λ_0} does not belong to RD_{λ_t, δ_0} . A Hénon map H_{a,λ_0} has connected Julia set for $a \in \mathbb{D}_{\delta_0}$ (see [17]). Radu and Tanase showed that there is $\delta_0 > 0$ such that H_{a,λ_t} is hyperbolic for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$ if $\theta_t = 0$ for each $0 < t < \delta_0$ in [16]. In [16], by using hyperbolicity, the Julia sets are connected for the parameters if $\theta_t = 0$ for $0 < t < \delta_0$ (see the left of Figure 2). In our case, we consider a much wider range of eigenvalues than $\theta_t = 0$. In this case, we will show the Julia set J_{a,λ_t} of $H_{a,\lambda_t} \in RD_{\lambda_t, \delta_0}$ is connected without using hyperbolicity.




Fig. 2. $\theta_t = 0$ (left) and radial perturbations (right). The set \mathcal{P}_1 (resp. \mathcal{P}_{-1}) is a semi-parabolic parameter given by equation (1) with $\lambda = 1$ (resp. $\lambda = -1$).

The rest of this paper is organized as follows. In Section 2, we present fundamental facts for Hénon maps. In Section 3, we introduce the *vertical condition* $(VC)_{\varepsilon,r}$ with respect to ε, r , and the condition $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda}^+ = \emptyset$. By using these conditions, we construct a holomorphic motion of $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y\})$ over $a \in \mathbb{D}_{\delta_0}$ for each $y \in \mathbb{D}_r$. Using these, we show Theorem 1.1. In Section 4, we show Theorem 1.2 by using Theorem 1.1. In particular, we check the condition $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda}^+ = \emptyset$ holds for H_{a,λ_t} by using local coordinates near semi-parabolic fixed points.

2. Preliminary

In this section, we recall some basic results on the dynamics of Hénon maps. See [2], [8], [14] and [16] for more details.

DEFINITION 2.1. For $(c, a) \in \mathbb{C}^2$, let $H: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be the map of the form

$$H(x, y) = (p(x) + ay, ax), \text{ where } p(x) = x^2 + c.$$

We call $H: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ a *Hénon map*. If $a \neq 0$, the inverse is

$$H^{-1}(x, y) = \frac{1}{a}(y, x - p(y/a)).$$

REMARK 2.2. In [14], a holomorphic automorphism of \mathbb{C}^2 of the form

$$F: (x, y) \mapsto (y, y^2 + c - \delta x), \quad \delta, c \in \mathbb{C}, \quad \delta \neq 0,$$

is called a Hénon map. The form of the Hénon map H in Definition 2.1 differs from the form $(x, y) \mapsto (y, y^2 + c - \delta x)$ given in [14]; however, H is conjugate by a polynomial

automorphism to $(x, y) \mapsto (y, y^2 + c + a^2 x)$.

In [8], the dynamical space \mathbb{C}^2 is divided into the following three sets.

DEFINITION 2.3 ([8]). Let $r > 0$ be a large number. Consider the following three subsets of \mathbb{C}^2 ,

$$\mathbb{D}_r \times \mathbb{D}_r := \{(x, y) \in \mathbb{C}^2 : |x| < r, |y| < r\},$$

$$V^+ := \{(x, y) \in \mathbb{C}^2 : |x| \geq \max\{|y|, r\}\} \text{ and } V^- := \{(x, y) \in \mathbb{C}^2 : |y| \geq \max\{|x|, r\}\}.$$

Let H be a Hénon map with $a \neq 0$. We define the *escaping sets* U^\pm of H by

$$U^+ := \bigcup_{k \geq 0} H^{-k}(V^+), \text{ and } U^- := \bigcup_{k \geq 0} H^k(V^-).$$

We consider the Julia sets and the filled Julia sets of Hénon maps.

DEFINITION 2.4. For a Hénon map $H(x, y) = (p(x) + ay, ax)$ with $a \neq 0$, we define the *filled Julia sets* K^\pm of H as follows:

$$K^\pm := \{(x, y) \in \mathbb{C}^2 : \{H^{\pm n}(x, y)\}_{n \in \mathbb{N}} \text{ is bounded in } \mathbb{C}^2\}.$$

We define the *Julia sets* J^\pm and J of H as follows:

$$J^\pm := \partial K^\pm, \text{ and } J := J^+ \cap J^-.$$

REMARK 2.5. For $a = 0$, we can also define K^+ and J^+ . In this case, K^+ (resp. J^+) is the product set of the filled Julia set (resp. the Julia set) of p and \mathbb{C} .

Bedford and Smillie [2] showed that there is a positive constant $r > 0$ depending on H such that

$$(2) \quad H(V^+) \subset V^+, H^{-1}(V^-) \subset V^-, U^+ = \mathbb{C}^2 \setminus K^+, \text{ and } U^- = \mathbb{C}^2 \setminus K^-.$$

It is easy to see that for a polynomial $x^2 + c_0$, there is $r > 0$ and δ such that $H(x, y) = (x^2 + c + ay, ax)$ satisfies the condition (2) with respect to r for $(c, a) \in \mathbb{D}_\delta(c_0) \times \mathbb{D}_\delta \setminus \{0\}$.

In this paper, we consider the following three types of fixed points.

DEFINITION 2.6. Suppose that a Hénon map H has a fixed point \mathbf{q} . Let λ and ν be the eigenvalues of $(DH)_{\mathbf{q}}$. We say that the fixed point \mathbf{q} is

- (i) *attracting* if $|\lambda| < 1$ and $|\nu| < 1$,
- (ii) *semi-parabolic* if $|\nu| < 1$ and $\lambda = \exp(2\pi i p/l)$ for some $p/l \in \mathbb{Q}$,
- (iii) *a saddle* if $|\nu| < 1$ and $|\lambda| > 1$.

We write that $A_n = O(B_n)$ if there are a positive constant $K > 0$ and a positive integer $N \in \mathbb{N}$ such that $|A_n| \leq K|B_n|$ for $n \geq N$. We set $\text{Pr}_1: \mathbb{C}^2 \rightarrow \mathbb{C}$, $\text{Pr}_1(x, y) := x$ and $\text{Pr}_2: \mathbb{C}^2 \rightarrow \mathbb{C}$, $\text{Pr}_2(x, y) := y$. We recall *stable manifolds* (see [14], [18]).

DEFINITION 2.7 ([14, p.311]). Let H be a Hénon map and $r > 0$ satisfy the condition (2) with respect to H . For a saddle fixed point \mathbf{q} of H , the *stable manifold* $W^s(\mathbf{q})$ of \mathbf{q} is defined

as

$$W^s(\mathbf{q}) := \{z \in \mathbb{C}^2 : \lim_{n \rightarrow \infty} \|H^n(z) - \mathbf{q}\| = 0\},$$

where $\|\cdot\|$ is the Euclidean metric of \mathbb{C}^2 .

Let $\lambda \in \mathbb{C} \setminus \{0\}$. To consider the connectivity of J for the parameters near the boundary of the Mandelbrot set, we consider a Hénon family for which each element of the family has a fixed point such that one of the eigenvalues of DH at the fixed point is λ . A Hénon map $H(x, y) = (x^2 + c + ay, ax)$ has a fixed point \mathbf{q} such that $\lambda \neq 0$ is an eigenvalue of $(DH)_{\mathbf{q}}$ if and only if

$$(3) \quad c = (1 - a^2) \left(\frac{\lambda}{2} - \frac{a^2}{2\lambda} \right) - \left(\frac{\lambda}{2} - \frac{a^2}{2\lambda} \right)^2.$$

Let \mathcal{P}_λ be the set of parameters $(c, a) \in \mathbb{C}^2$ satisfying (3). We denote the right hand side of equation (3) by $c(a, \lambda)$. Moreover, we set $H_{a, \lambda}(x, y) = (x^2 + c(a, \lambda) + ay, ax)$ and $p_\lambda(x) = x^2 + c(0, \lambda)$. We denote the filled Julia sets and the Julia sets of $H_{a, \lambda}$ by $K_{a, \lambda}^\pm, J_{a, \lambda}^\pm, J_{a, \lambda}$ instead of K^\pm, J^\pm, J respectively. We see that $H_{a, \lambda}$ has a fixed point

$$(4) \quad \mathbf{q}_{a, \lambda} := \left(\frac{\lambda}{2} - \frac{a^2}{2\lambda}, a \left(\frac{\lambda}{2} - \frac{a^2}{2\lambda} \right) \right),$$

with eigenvalues λ and $\nu := -a^2/\lambda$. We give the parametrization of $W^{ss}(\mathbf{q}_{a, \lambda})$.

Lemma 2.8 ([7], [15, the proofs of Propositions 3.16, 3.17], [17, Proposition 5.2]). *Let $v = (-a/\lambda, 1)$ be an eigenvector for ν . Assume that λ satisfies $|-a^2/\lambda| < 1$ and $|\lambda| > |-a^2/\lambda|$. Then there exists the injective holomorphic map*

$$(5) \quad \Phi_{a, \lambda}: \mathbb{C} \rightarrow \mathbb{C}^2, \Phi_{a, \lambda}(z) = \lim_{k \rightarrow \infty} H_{a, \lambda}^{-k}(\mathbf{q}_{a, \lambda} + \nu^k z v),$$

such that $\Phi_{a, \lambda}(vz) = H_{a, \lambda}(\Phi_{a, \lambda}(z))$ for $z \in \mathbb{C}$ and $a \neq 0$.

Let $\Phi_{0, \lambda}(z) := (\Pr_1 \mathbf{q}_{0, \lambda}, z)$. Then, $\Phi_{a, \lambda}$ is analytic with respect to a and $\sup_{z \in K} \|\Phi_{a, \lambda}(z) - \Phi_{0, \lambda}(z)\| = O(a)$ for each compact subset K of \mathbb{C} .

REMARK 2.9. Fix $\lambda_0 \neq 0$. Consider a family $\{H_{a, \lambda}\}_{(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)}$. Since the fixed point $\mathbf{q}_{a, \lambda}$ of $H_{a, \lambda}$ depends holomorphically on a, λ , we see that $\Phi_{a, \lambda}(z)$ is holomorphic with respect to $(a, \lambda, z) \in \mathbb{D}_{\delta_0} \setminus \{0\} \times \mathbb{D}_{\delta_0}(\lambda_0) \times \mathbb{C}$ (see the proof of Theorem 6.43 in [14]). Since $\Phi_{a, \lambda}$ is holomorphic with respect to each variable separately when the other variables are fixed, $\Phi_{a, \lambda}(z)$ is holomorphic with $(a, \lambda, z) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0) \times \mathbb{C}$, taking a smaller $\delta_0 > 0$ if necessary.

DEFINITION 2.10. The curve $W^{ss}(\mathbf{q}) := \Phi(\mathbb{C})$ is called the *strong stable manifold* of a fixed point \mathbf{q} for $a \neq 0$, where Φ is given by (5), and the definition of (5) is valid if at least one eigenvalue of the fixed point has absolute value less than 1 (see the proof of Theorem 6.43 in [14]).

For a fixed point \mathbf{q} of H , the *local strong stable manifold* $W_{loc}^{ss}(\mathbf{q})$ of \mathbf{q} is defined by the component of $W^{ss}(\mathbf{q}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ that contains \mathbf{q} , with the topology induced by $\Phi: \mathbb{C} \rightarrow W^{ss}(\mathbf{q})$.

REMARK 2.11. When $a = 0$, we set $W^{ss}(\mathbf{q}) = \bigcup_{j \geq 0} p^{-j}(\{\text{Pr}_1 \mathbf{q}\}) \times \mathbb{C}$ and $W_{\text{loc}}^{ss}(\mathbf{q}) = \{\text{Pr}_1 \mathbf{q}\} \times \mathbb{D}_r$.

REMARK 2.12. If \mathbf{q} is a saddle, then $W^s(\mathbf{q}) = W^{ss}(\mathbf{q})$. In this case, the local stable manifold $W_{\text{loc}}^s(\mathbf{q})$ of \mathbf{q} is defined by $W_{\text{loc}}^{ss}(\mathbf{q})$.

By using Lemma 2.8, we have the following.

Lemma 2.13. *If $\lambda_0 \in \mathbb{C} \setminus \{0, 1\}$, then there is a positive constant $\delta_0 > 0$ such that $H_{a,\lambda}$ has a saddle fixed point $\mathbf{s}_{a,\lambda}$ depending holomorphically on $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$. Let $\tilde{\Phi}_{a,\lambda}(z) := (\text{Pr}_1 \mathbf{s}_{a,\lambda}, z)$, and $\tilde{\Phi}_{a,\lambda}(z) := \lim_{k \rightarrow \infty} H_{a,\lambda}^{-k}(\mathbf{s}_{a,\lambda} + \tilde{v}^k z \tilde{v})$, where \tilde{v} is the eigenvalue of $(DH_{a,\lambda})_{\mathbf{s}_{a,\lambda}}$ with $|\tilde{v}| < 1$ and \tilde{v} is the eigenvector of \tilde{v} of the form $(\cdot, 1)$ for $(a, \lambda) \in (\mathbb{D}_{\delta_0} \setminus \{0\}) \times \mathbb{D}_{\delta_0}(\lambda_0)$ (see (5)). Then, $\tilde{\Phi}_{a,\lambda}(z)$ is holomorphic with respect to $(a, \lambda, z) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0) \times \mathbb{C}$.*

Proof. If $|\lambda_0| > 1$, then $\mathbf{q}_{a,\lambda}$ is saddle for $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$, taking a smaller δ_0 if necessary. By taking $\mathbf{s}_{a,\lambda}$ as $\mathbf{q}_{a,\lambda}$ and applying Lemma 2.8, we obtain the statement of Lemma 2.13.

We may assume that $\lambda_0 \neq 1, \lambda_0 \neq 0$ and $|\lambda_0| \leq 1$. Then $c(0, \lambda_0) \neq 1/4$ and $H_{0,\lambda_0}(x, y)$ has a saddle fixed point. By the implicit function theorem, there are a positive constant δ_0 and a saddle fixed point $\mathbf{s}_{a,\lambda}$ of $H_{a,\lambda}$ depending holomorphically on $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$.

We show that $\tilde{\Phi}_{a,\lambda}(z)$ is holomorphic with respect to $(a, \lambda, z) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0) \times \mathbb{C}$ by using Lemma 2.8. It follows from $\mathbf{s}_{0,\lambda_0} \neq \mathbf{q}_{0,\lambda_0}$ that $\mathbf{s}_{a,\lambda} \neq \mathbf{q}_{a,\lambda}$ for $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$, taking δ_0 small enough. By $\mathbf{s}_{a,\lambda} \neq \mathbf{q}_{a,\lambda}$ and the fixed point equation $(x^2 + c(a, \lambda) + ay, ax) = (x, y)$, we have

$$\mathbf{s}_{a,\lambda} = (1 - a^2 - \text{Pr}_1 \mathbf{q}_{a,\lambda}, a(1 - a^2 - \text{Pr}_1 \mathbf{q}_{a,\lambda})).$$

Consider the characteristic equation $\det((DH_{a,\lambda})_{\mathbf{s}_{a,\lambda}} - \tilde{\lambda}I) = 0$, which is equivalent to $-(2\text{Pr}_1 \mathbf{s}_{a,\lambda} - \tilde{\lambda})\tilde{\lambda} - a^2 = 0$. We set $f(a, \lambda, \tilde{\lambda}) := -(2\text{Pr}_1 \mathbf{s}_{a,\lambda} - \tilde{\lambda})\tilde{\lambda} - a^2$. We see that $f(0, \lambda_0, \tilde{\zeta}) = 0$, where $\tilde{\zeta} = 0, 2 - \lambda_0$ (see (4)). By $\partial_{\tilde{\lambda}} f(0, \lambda_0, 2 - \lambda_0) = 2 - \lambda_0 \neq 0$ and the Implicit Function Theorem, there exists a holomorphic map $\tilde{\lambda}(a, \lambda)$ with $|\tilde{\lambda}(a, \lambda)| > 1$ for $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$ such that $f(a, \lambda, \tilde{\lambda}(a, \lambda)) = 0$ and $\tilde{\lambda}(0, \lambda_0) = 2 - \lambda_0$, taking a smaller δ_0 if necessary. We see that $c(a, \lambda) = c(a, \tilde{\lambda}(a, \lambda))$ for each $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$ since $H_{a,\lambda}$ has the fixed point $\mathbf{s}_{a,\lambda}$ with one eigenvalue $\tilde{\lambda}(a, \lambda)$ (see (3)). In particular, $H_{a,\lambda} = H_{a,\tilde{\lambda}(a,\lambda)}$ for each $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$. By the identity theorem, we have $\mathbf{s}_{a,\lambda} = \mathbf{q}_{a,\tilde{\lambda}(a,\lambda)}$ for $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$. Thus $\tilde{\Phi}_{a,\lambda} = \Phi_{a,\tilde{\lambda}(a,\lambda)}$ for $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$, where $\Phi_{a,\tilde{\lambda}(a,\lambda)}$ is the map in Lemma 2.8. By Remark 2.9 and the fact that $\tilde{\lambda}(a, \lambda)$ depends holomorphically on $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$, it follows that $\tilde{\Phi}_{a,\lambda}(z)$ is holomorphic with respect to $(a, \lambda, z) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0) \times \mathbb{C}$. \square

3. Vertical Condition

For the rest of the section, we assume that the Jacobians of Hénon maps $H_{a,\lambda}$ are less than 1 in the absolute value, and $\lambda \neq 0$. In this section, we show the first main result (Theorems 3.10 and 3.11). We construct a holomorphic motion of $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y\})$ over $a \in \mathbb{D}_{\delta_0}$ for each $y \in \mathbb{D}_r$ to obtain the first main result. In order to construct it, we consider the *vertical condition* and the condition $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda}^+ = \emptyset$. We consider the vertical cone

field $\{C_{(x,y)}^v\}_{(x,y) \in \mathbb{D}_r \times \mathbb{D}_r}$ given by

$$(6) \quad C_{(x,y)}^v := \{(\zeta, \eta) \in T_{(x,y)}\mathbb{C}^2 : |\eta| > |\zeta|\},$$

for $(x, y) \in \mathbb{D}_r \times \mathbb{D}_r$. We first introduce the *vertical condition* as follows.

DEFINITION 3.1. Let $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ be a Hénon family. Fix $\varepsilon > 0$, and fix $r > 0$ such that $H_{a,\lambda}$ satisfies the condition (2) with respect to r for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$. Moreover, we assume that $\delta_0 < \min\{1/2, \varepsilon\}$. We say that $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the *vertical condition* $(VC)_{\varepsilon,r}$ with respect to ε, r if the following three conditions hold:

- (i) $H_{a,\lambda}^{-1}((\mathbb{D}_r \times \mathbb{D}_r) \setminus (\mathbb{D}_r \times \mathbb{D}_{r/2})) \subset V^-$ and $|\text{Pr}_2 H_{a,\lambda}^{-1}(x, y)| > 2|y|$ for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$ and $(x, y) \in V^- \cup ((\mathbb{D}_r \times \mathbb{D}_r) \setminus (\mathbb{D}_r \times \mathbb{D}_{r/2}))$.
- (ii) $(DH_{a,\lambda}^{-1})_{(x,y)}(C_{(x,y)}^v) \subset C_{H_{a,\lambda}^{-1}(x,y)}^v$ and $|\text{Pr}_2 (DH_{a,\lambda}^{-1})_{(x,y)}(\zeta, \eta)| > 2|\eta|$ for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$, $(x, y) \in H_{a,\lambda}((\mathbb{D}_r \times \mathbb{D}_r) \setminus (\mathbb{D}_r \times \mathbb{D}_{r/2})) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ and $(\zeta, \eta) \in C_{(x,y)}^v$.
- (iii) There are a saddle fixed point $\mathbf{s}_{a,\lambda}$ of $H_{a,\lambda}$ depending holomorphically on $a \in \mathbb{D}_{\delta_0}$ and a holomorphic map $f_a: \mathbb{D}_r \rightarrow \mathbb{C}$ depending holomorphically on $a \in \mathbb{D}_{\delta_0}$ such that $W_{loc}^s(\mathbf{s}_{a,\lambda}) = \{(f_a(y), y) : y \in \mathbb{D}_r\}$ and $T_{(x,y)} W_{loc}^s(\mathbf{s}_{a,\lambda}) \subset C_{(x,y)}^v$ for $(x, y) \in W_{loc}^s(\mathbf{s}_{a,\lambda})$.

We see that most families $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfy the condition $(VC)_{\varepsilon,r}$ by the following lemma.

Lemma 3.2. Fix $\lambda \neq 1$. Then there is a positive constant $\delta_0 > 0$ such that $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the $(VC)_{\varepsilon,r}$.

Proof. Fix $\varepsilon > 0$. We can take $r > 0$ such that each $H_{a,\lambda}$ satisfies the condition (2) with respect to r for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$ by taking a smaller δ_0 if necessary. Assume that $\delta_0 < \min\{1/2, \varepsilon\}$.

We show that the condition (i) in Definition 3.1 holds. For $(x, y) \in V^- \cup ((\mathbb{D}_r \times \mathbb{D}_r) \setminus (\mathbb{D}_r \times \mathbb{D}_{r/2}))$, we set $(x_1, y_1) := H_{a,\lambda}^{-1}(x, y) = (y/a, (x - y^2/a^2 - c(a, \lambda))/a)$. We show $|y_1| > 2|y|$. Since $|x| \leq \max\{|y|, r\}$, we have

$$|ay_1| = |x - y^2/a^2 - c(a, \lambda)| \geq |y/a|^2 - |c(a, \lambda)| - \max\{|y|, r\}.$$

To obtain $|y_1| > 2|y|$, it suffices to show that $|y/a|^2 - 2|a||y| - |c(a, \lambda)| - \max\{|y|, r\} > 0$ which is equivalent to $|y|^2 - 2|a|^3|y| - |a|^2|c(a, \lambda)| - |a|^2 \max\{|y|, r\} > 0$. Note that $|y| \geq r/2$ by $(x, y) \in V^- \cup ((\mathbb{D}_r \times \mathbb{D}_r) \setminus (\mathbb{D}_r \times \mathbb{D}_{r/2}))$. If $a = 0$, we have $|y|^2 \geq r^2/4 > 0$. Thus, we can take $\delta_0 > 0$ such that $|y_1| > 2|y|$ for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$. Similarly, we have $(x_1, y_1) \in V^-$ if $(x, y) \in (\mathbb{D}_r \times \mathbb{D}_r) \setminus (\mathbb{D}_r \times \mathbb{D}_{r/2})$ for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$, taking a smaller δ_0 if necessary. Thus, $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0} \setminus \{0\}}$ satisfies the condition (i) in Definition 3.1.

We next show that the condition (ii) in Definition 3.1 holds. Fix $(x, y) \in H_{a,\lambda}((\mathbb{D}_r \times \mathbb{D}_r) \setminus (\mathbb{D}_r \times \mathbb{D}_{r/2})) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ and $(\zeta, \eta) \in C_{(x,y)}^v$. We set $(x_1, y_1) = H_{a,\lambda}^{-1}(x, y)$ and $(\zeta_1, \eta_1) = (DH_{a,\lambda}^{-1})_{(x,y)}(\zeta, \eta)$. By $(x_1, y_1) = H_{a,\lambda}^{-1}(x, y) = (y/a, (x - y^2/a^2 - c(a, \lambda))/a)$, we have $\zeta_1 = \eta/a$ and $\eta_1 = (\zeta - 2y\eta/a^2)/a$. By $x_1 = y/a$, we have $\eta_1 = (\zeta - 2x_1\eta/a)/a$. Thus we have

$$|\eta_1| > \frac{1}{|a|} \left(\frac{|2x_1\eta|}{|a|} - |\zeta_1| \right) \geq \frac{1}{|a|} \left(\frac{2\varepsilon}{|a|} - 1 \right) |\eta| > \frac{1}{\delta_0} \left(\frac{2\varepsilon}{\delta_0} - 1 \right) |\eta| > 2|\eta|,$$

by $\delta_0 < \min\{1/2, \varepsilon\}$. In particular, we have $|\eta_1| > 1/|a|(2\varepsilon - |a|)|\zeta_1| > (2\varepsilon/|\delta_0| - 1)|\zeta_1| > |\zeta_1|$

by $\eta = a\zeta_1$ and $\delta_0 < \min\{1/2, \varepsilon\}$. Thus the condition (ii) holds for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$.

Finally, we show that the condition (iii) in Definition 3.1 holds. There exists a saddle fixed point $\mathbf{s}_{a,\lambda}$ of $H_{a,\lambda}$ depending holomorphically on $a \in \mathbb{D}_{\delta_0}$ by Lemma 2.13. By Lemmas 2.8 and 2.13, $W_{\text{loc}}^s(\mathbf{s}_{a,\lambda}) = \{(f_a(y), g_a(y)) : y \in g_a^{-1}(\mathbb{D}_r)\}$ for some holomorphic maps f_a, g_a depending holomorphically on $a \in \mathbb{D}_{\delta_0}$. Here, we remark that by Rouché's theorem and $g_0(y) = y$, there exists g_a^{-1} in \mathbb{D}_r for $a \in \mathbb{D}_{\delta_0}$, taking a smaller δ_0 if necessary. Thus we have $W_{\text{loc}}^s(\mathbf{s}_{a,\lambda}) = \{(f_a(g_a^{-1}(y)), y) : y \in \mathbb{D}_r\}$. In the case of $a = 0$, we have $W_{\text{loc}}^s(\mathbf{s}_{0,\lambda}) = \{\text{Pr}_1 \mathbf{s}_{0,\lambda}\} \times \mathbb{D}_r$ and $T_{(x,y)} W_{\text{loc}}^s(\mathbf{s}_{0,\lambda}) = \{(\zeta, \eta) : \zeta = 0\} \subset C_{(x,y)}^v$. Since $W_{\text{loc}}^s(\mathbf{s}_{a,\lambda})$ depends on a holomorphically, by taking δ_0 sufficiently small, we have $|f_a(g_a^{-1}(y))'| < 1$ for $y \in \mathbb{D}_r$. \square

REMARK 3.3. The proof of Lemma 3.2 is still valid for perturbations $\{H_{a,\lambda}\}_{(a,\lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)}$, where $\lambda_0 \neq 1$. That is, there is $\delta_0 > 0$ such that the family $\{H_{a,\lambda}\}_{(a,\lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)}$ satisfies (i), (ii), and (iii) in $(\text{VC})_{\varepsilon,r}$.

REMARK 3.4. By the proof of Lemma 3.2, (i) and (ii) in Definition 3.1 hold for $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0} \setminus \{0\}}$ without the condition $\lambda \neq 1$ in Lemma 3.2. Moreover, for each $\varepsilon > 0$, there is $\delta_0 > 0$ such that (ii) in Definition 3.1 holds for $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$.

Consider a Hénon family $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfying the $(\text{VC})_{\varepsilon,r}$, and

$$(7) \quad (\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda}^+ = \emptyset,$$

for $a \in \mathbb{D}_{\delta_0}$ (see Remark 2.5 for the case $a = 0$). Let $\mathbf{s}_{a,\lambda}$ be a saddle fixed point of $H_{a,\lambda}$ depending holomorphically on $a \in \mathbb{D}_{\delta_0}$. Let $\mathbf{v}_{a,\lambda} := (\bigcup_{j \in \mathbb{Z}_{\geq 0}} H_{a,\lambda}^{-j}(W_{\text{loc}}^s(\mathbf{s}_{a,\lambda}))) \cap (\mathbb{D}_r \times \mathbb{D}_r)$. We say that v is a *vertical component* of $\mathbf{v}_{a,\lambda}$ if v is a connected component of $H_{a,\lambda}^{-m}(W_{\text{loc}}^s(\mathbf{s}_{a,\lambda})) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ for some $m \in \mathbb{Z}_{\geq 0}$. Under the assumption (7), we have

$$(8) \quad \mathbf{v}_{a,\lambda} \subset (\mathbb{D}_r \setminus \mathbb{D}_\varepsilon) \times \mathbb{D}_r,$$

for $a \in \mathbb{D}_{\delta_0}$ by $W^s(\mathbf{s}_{a,\lambda}) \subset J_{a,\lambda}^+$ and (2). To construct a holomorphic motion of $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y\})$ over $a \in \mathbb{D}_{\delta_0}$ for each $y \in \mathbb{D}_r$, we prove the following two lemmas.

Lemma 3.5. Suppose that $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the $(\text{VC})_{\varepsilon,r}$ and the condition (7) holds for $a \in \mathbb{D}_{\delta_0}$. Let v_a be a vertical component of $\mathbf{v}_{a,\lambda}$ represented by $\{(f_a(y), y) : y \in \mathbb{D}_r\}$ for some holomorphic map f_a depending holomorphically on $a \in \mathbb{D}_{\delta_0}$. Then for each $a \in \mathbb{D}_{\delta_0}$, $H_{a,\lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ is the union of two distinct vertical components $v_{a,1}$ and $v_{a,2}$. Moreover, $v_{a,j}$ can be represented by $\{(f_{a,j}(y), y) : y \in \mathbb{D}_r\}$ for some holomorphic map $f_{a,j} : \mathbb{D}_r \rightarrow \mathbb{C}$ depending holomorphically on $a \in \mathbb{D}_{\delta_0}$ for $j = 1, 2$.

Proof. Let v_a be a vertical component of $\mathbf{v}_{a,\lambda}$ represented by $\{(f_a(y), y) : y \in \mathbb{D}_r\}$ for some holomorphic map f_a depending holomorphically on $a \in \mathbb{D}_{\delta_0}$.

We first show that the set $H_{a,\lambda}^{-1}(v_a) \cap (\mathbb{C} \times \{w\})$ consists of exactly two points for $w \in \mathbb{D}_r$ and $a \in \mathbb{D}_{\delta_0}$. For $a = 0$, f_0 is constant and $H_{0,\lambda}(x, y) = (x^2 + c(0, \lambda), 0)$. Clearly, $H_{0,\lambda}^{-1}(v_0) \cap (\mathbb{C} \times \{w\})$ consists of exactly two points for $w \in \mathbb{D}_r$ since the critical value $c(0, \lambda)$ of $x^2 + c(0, \lambda)$ does not belong to $f_0(\mathbb{D}_r)$ (see (7)). For $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$, recall that $H_{a,\lambda}^{-1}(x, y) = (y/a, (x - y^2/a^2 - c(a, \lambda))/a)$. Consider the equation

$$(9) \quad f_a(y) - y^2/a^2 - c(a, \lambda) = aw,$$

for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$. We set $g_a(y) := f_a(y) - y^2/a^2 - c(a, \lambda) - aw$ and $h_a(y) := -f_a(y) + c(a, \lambda) + aw$ for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$. By $v_a \subset \mathbf{v}_{a, \lambda} \subset \mathbb{D}_r \times \mathbb{D}_r$, we have $|f_a(y)| \leq r$ for $y \in \mathbb{D}_r$. If $|a| < 1$, then for $y \in \partial \mathbb{D}_{\sqrt{(4r+4|c(a, \lambda))|a|^2}}$, we have

$$\begin{aligned} |g_a(y)| &\geq |y^2/a^2| - |c(a, \lambda)| - |aw| - |f_a(y)| \geq (3 - |a|)r + 3|c(a, \lambda)| > r + |c(a, \lambda)| + |a|r \\ &\geq |f_a(y)| + |c(a, \lambda)| + |aw| \geq |h_a(y)|. \end{aligned}$$

By Rouché's theorem, g_a and $g_a + h_a$ have the same number of zeros inside $\mathbb{D}_{\sqrt{(4r+4|c(a, \lambda))|a|^2}}$. Since $g_a(y) + h_a(y) = -y^2/a^2$, the map g_a has two zeros in $\mathbb{D}_{\sqrt{(4r+4|c(a, \lambda))|a|^2}}$. Moreover, the map g_a has two distinct zeros by using the condition (ii) and (iii) in Definition 3.1. Indeed, let $g_a(y_0) = 0$. Then we have $(y_0/a, w) \in H_{a, \lambda}^{-1}(v_a) \cap (\mathbb{C} \times \{w\})$. By (8), we have $|y_0/a| > \varepsilon$. Moreover, we have $|f'_a(y_0)| < 1$ by the condition (ii), (iii) in Definition 3.1, and (6). Hence we have

$$(10) \quad |g'_a(y_0)| \geq |2y_0/a^2| - |f'_a(y_0)| > 2\varepsilon/|a| - 1 > 2\varepsilon/\delta_0 - 1 > 0,$$

by $\delta_0 < \min\{\varepsilon, 1/2\}$ (see Definition 3.1). Thus, g_a has two distinct zeros in $\mathbb{D}_{\sqrt{(4r+4|c(a, \lambda))|a|^2}}$. On the other hand, by $\delta_0 < \min\{\varepsilon, 1/2\}$, if $y \in \mathbb{D}_r$ satisfies $|y| \geq \sqrt{(4r+4|c(a, \lambda))|a|^2}$, then

$$|\text{Pr}_2 H_{a, \lambda}^{-1}(f_a(y), y)| = \frac{|f_a(y) - y^2/a^2 - c(a, \lambda)|}{|a|} \geq \frac{|y^2/a^2| - |f_a(y)| - |c(a, \lambda)|}{|a|} \geq \frac{3r + 3|c(a, \lambda)|}{|a|} > r.$$

Hence, there are exactly two distinct solutions of (9) with respect to $y \in \mathbb{D}_r$, which belong to $\mathbb{D}_{\sqrt{(4r+4|c(a, \lambda))|a|^2}}$. This implies that the set $H_{a, \lambda}^{-1}(v_a) \cap (\mathbb{C} \times \{w\})$ consists of exactly two points for $w \in \mathbb{D}_r$. Note that $H_{a, \lambda}^{-1}(v_a) \cap ((\mathbb{C} \setminus \mathbb{D}_r) \times \{w\}) = \emptyset$ by (2), otherwise $W^s(\mathbf{s}_{a, \lambda}) \cap V^+ \neq \emptyset$. Thus, $H_{a, \lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ is the union of two vertical components of $\mathbf{v}_{a, \lambda}$.

Recall that f_0 is constant. Assume $f_0 \equiv A$ for some $A \in \mathbb{C}$. Let $z^2 + c(0, \lambda) - A = (z - A_1)(z - A_2)$. Then we have $H_{0, \lambda}^{-1}(v_0) \cap (\mathbb{D}_r \times \mathbb{D}_r) = (\{A_1\} \cup \{A_2\}) \times \mathbb{D}_r$. There are positive constants $\delta_1, \varepsilon_1 > 0$ such that $\delta_1 < \delta_0$, $\mathbb{D}_{\varepsilon_1}(A_1) \cap \mathbb{D}_{\varepsilon_1}(A_2) = \emptyset$, $H_{a, \lambda}^{-1}(v_a) \cap (\mathbb{D}_{\varepsilon_1}(A_j) \times \mathbb{D}_r) \neq \emptyset$ for $j = 1, 2$ and $a \in \mathbb{D}_{\delta_1}$, and $H_{a, \lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r) \subset (\mathbb{D}_{\varepsilon_1}(A_1) \cup \mathbb{D}_{\varepsilon_1}(A_2)) \times \mathbb{D}_r$ for $a \in \mathbb{D}_{\delta_1}$. For $a \in \mathbb{D}_{\delta_1}$ and $j \in \{1, 2\}$, we let $v_{a, j}$ be the component of $H_{a, \lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ which is contained in $\mathbb{D}_{\varepsilon_1}(A_j) \times \mathbb{D}_r$.

We show that $v_{a, j} \cap (\mathbb{C} \times \{w\})$ moves holomorphically over \mathbb{D}_{δ_1} for each fixed $w \in \mathbb{D}_r$ by using the implicit function theorem. We set $F(a, y) := f_a(y) - y^2/a^2 - c(a, \lambda) - aw$ for $a \in \mathbb{D}_{\delta_1} \setminus \{0\}$. Since $T_{(f_a(y), y)}v \subset C_{(f_a(y), y)}^v$ for $y \in \mathbb{D}_r$, we have $|\partial_y f_a(y)| < 1$ for $y \in \mathbb{D}_r$. Fix arbitrary points $a \in \mathbb{D}_{\delta_1} \setminus \{0\}$ and $\tilde{z}_j = \tilde{z}_j(a)$ with $F(a, \tilde{z}_j(a)) = 0$ and $\tilde{z}_j(a)/a \in \mathbb{D}_{\varepsilon_1}(A_j)$. Then, we have $(\tilde{z}_j/a, (f_a(\tilde{z}_j) - \tilde{z}_j^2/a^2 - c(a, \lambda))/a) = (\tilde{z}_j/a, w) \in v_{a, j} \cap (\mathbb{C} \times \{w\})$. By (8), we have $|\tilde{z}_j/a| > \varepsilon$. Since $F(a, \tilde{z}_j) = g_a(\tilde{z}_j)$, we have $|\partial_y F(a, \tilde{z}_j)| > 0$ (see (10)). By the implicit function theorem, $\{\tilde{z}_j/a, w\} = v_{a, j} \cap (\mathbb{C} \times \{w\})$ moves holomorphically over $\mathbb{D}_{\delta_1} \setminus \{0\}$. Moreover, $\tilde{z}_j(a)^2/a^2 = f_a(\tilde{z}_j(a)) - c(a, \lambda) - aw$ since $F(a, \tilde{z}_j(a)) = 0$. Note that $\tilde{z}_j(a) \in \mathbb{D}_{r/2}$ by $r > |w| = |\text{Pr}_2 H_{a, \lambda}^{-1}(f_a(\tilde{z}_j), \tilde{z}_j)|$ and (i) in Definition 3.1. Since $f_a(z)$ is holomorphic with respect to $a \in \mathbb{D}_{\delta_0}$ and $z \in \mathbb{D}_r$, $f_a(z) \rightarrow f_0(z)$ uniformly on $\overline{\mathbb{D}_{r/2}}$ as $a \rightarrow 0$. This implies that $\tilde{z}_j(a)/a \rightarrow A_j$ as $a \rightarrow 0$ since $f_a(z) \rightarrow f_0(z) \equiv A$ as $a \rightarrow 0$. Thus $\{\tilde{z}_j/a\} = v_{a, j} \cap (\mathbb{C} \times \{w\})$ moves holomorphically over \mathbb{D}_{δ_1} .

For each $a \in \mathbb{D}_{\delta_0} \setminus \mathbb{D}_{\delta_1}$ and each $j \in \{1, 2\}$, $v_{a, j} \cap (\mathbb{C} \times \{w\})$ can be analytically continued along a path connecting a and a point in \mathbb{D}_{δ_1} . By the monodromy theorem, for each $a \in \mathbb{D}_{\delta_0}$

and each $j \in \{1, 2\}$, there is a component $v_{a,j}$ of $H_{a,\lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ such that $v_{a,j} \cap (\mathbb{C} \times \{w\})$ moves holomorphically over \mathbb{D}_{δ_0} for each fixed $w \in \mathbb{D}_r$. We show that $v_{a,1} \neq v_{a,2}$ for each $a \in \mathbb{D}_{\delta_0} \setminus \mathbb{D}_{\delta_1}$. Assume that $v_{a,1} = v_{a,2}$ for some $a \in \mathbb{D}_{\delta_0} \setminus \mathbb{D}_{\delta_1}$. There exists a sequence $\{a_n\}_{n \in \mathbb{N}}$ in $\{a \in \mathbb{D}_{\delta_0} : v_{a,1} = v_{a,2}\}$ such that $a_n \rightarrow a_0$ as $n \rightarrow \infty$ and

$$(11) \quad |a_0| = \inf\{|a| \in \mathbb{D}_{\delta_0} : v_{a,1} = v_{a,2}\}.$$

Clearly, we have $|a_0| \geq \delta_1$ by the argument above. Moreover, we have $v_{a_0,1} = v_{a_0,2}$. Otherwise, $v_{a,1} \neq v_{a,2}$ for all a in a small neighborhood of a_0 , which implies that $a_n \notin \{a \in \mathbb{D}_{\delta_0} : v_{a,1} = v_{a,2}\}$ for sufficiently large n . Consider the vertical component v_{a_0} of $\mathbf{v}_{a_0,\lambda}$. Let $H_{a_0,\lambda}^{-1}(v_{a_0}) \cap (\mathbb{D}_r \times \mathbb{D}_r) = \tilde{v}_{a_0,1} \cup \tilde{v}_{a_0,2}$ for some vertical components $\tilde{v}_{a_0,1}, \tilde{v}_{a_0,2}$ of $\mathbf{v}_{a,\lambda}$. There are open neighborhoods U_1, U_2 of $\tilde{v}_{a_0,1}, \tilde{v}_{a_0,2}$ respectively such that $U_1 \cap U_2 = \emptyset$. We may assume that $v_{a_0,1} (= v_{a_0,2}) = \tilde{v}_{a_0,1}$. We take a positive constant $\delta_2 > 0$ such that $H_{a,\lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r) \cap U_j \neq \emptyset$ for $j = 1, 2$ and $a \in \mathbb{D}_{\delta_2}(a_0)$, and $H_{a,\lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r) \subset U_1 \cup U_2$ for $a \in \mathbb{D}_{\delta_2}(a_0)$. Recall that $F(a, y) = f_a(y) - y^2/a^2 - c(a, \lambda) - aw$ for $w \in \mathbb{D}_r$. We see that $|\partial_y F(a_0, \tilde{z}_1)| > 0$, where \tilde{z}_1 satisfies that $F(a_0, \tilde{z}_1) = 0$ and $(\tilde{z}_1/a_0, w) \in U_1$ (see (10)). By the implicit function theorem, $v_{a,1} \cap (\mathbb{C} \times \{w\}) \subset U_1$ moves holomorphically over $\mathbb{D}_{\delta_2}(a_0)$ for each fixed $w \in \mathbb{D}_r$. We can take $\tilde{a} \in \mathbb{D}_{\delta_2}(a_0)$ with $|\tilde{a}| < |a_0|$ such that $v_{\tilde{a},1} = v_{\tilde{a},2}$. This contradicts (11). Thus, $H_{a,\lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r) = v_{a,1} \cup v_{a,2}$ and $v_{a,j} \cap (\mathbb{C} \times \{w\})$ moves holomorphically over \mathbb{D}_{δ_0} for each fixed $w \in \mathbb{D}_r$ and $j \in \{1, 2\}$.

Finally, we show that there are holomorphic maps $f_{a,1}, f_{a,2}$ such that $v_{a,j} = \{(f_{a,j}(y), y) : y \in \mathbb{D}_r\}$ for $j = 1, 2$. Since $\text{Pr}_2: v_{a,j} \rightarrow \mathbb{D}_r$ is a bijective holomorphic map for $a \in \mathbb{D}_{\delta_0}$ and $j \in \{1, 2\}$, there are holomorphic maps $f_{a,1}, f_{a,2}$ such that $v_{a,j} = \{(f_{a,j}(y), y) : y \in \mathbb{D}_r\}$ for $a \in \mathbb{D}_{\delta_0}$ and $j = 1, 2$. \square

Let $W_{a,\lambda}$ be the vertical component of $\mathbf{v}_{a,\lambda}$ such that $H_{a,\lambda}(W_{a,\lambda}) \subset W_{\text{loc}}^s(\mathbf{s}_{a,\lambda})$ and $W_{a,\lambda} \cap W_{\text{loc}}^s(\mathbf{s}_{a,\lambda}) = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$.

Lemma 3.6. *Suppose that $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the $(\text{VC})_{\varepsilon,r}$ and the condition (7) holds for $a \in \mathbb{D}_{\delta_0}$. Then the two sets $H_{a,\lambda}^{-n}(W_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ and $H_{a,\lambda}^{-m}(W_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ do not intersect for $n \neq m \in \mathbb{Z}_{\geq 0}$ and $a \in \mathbb{D}_{\delta_0}$.*

Proof. Fix $a \in \mathbb{D}_{\delta_0}$. Assume that there are components v_n and v_m of $H_{a,\lambda}^{-n}(W_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ and $H_{a,\lambda}^{-m}(W_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ respectively such that $v_n \cap v_m \neq \emptyset$. We may assume that $n > m$. We see that $H_{a,\lambda}^m(v_n)$ is a subset of some component of $H_{a,\lambda}^{-n+m}(W_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$. On the other hand, $H_{a,\lambda}^m(v_m) \subset W_{a,\lambda}$. Thus, an intersection point of v_n and v_m is mapped under $H_{a,\lambda}^m$ into $W_{a,\lambda} \cap H_{a,\lambda}^{-n+m}(W_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$. Hence, we have $H_{a,\lambda}^{n-m}(W_{a,\lambda}) \cap W_{a,\lambda} \neq \emptyset$. This contradicts $H_{a,\lambda}(W_{a,\lambda}) \subset W_{\text{loc}}^s(\mathbf{s}_{a,\lambda})$, $H_{a,\lambda}(W_{\text{loc}}^s(\mathbf{s}_{a,\lambda})) \subset W_{\text{loc}}^s(\mathbf{s}_{a,\lambda})$ and $W_{\text{loc}}^s(\mathbf{s}_{a,\lambda}) \cap W_{a,\lambda} = \emptyset$. Thus we have proved Lemma 3.6. \square

Let us consider the section $S_{a,\lambda,y} := W^s(\mathbf{s}_{a,\lambda}) \cap (\mathbb{C} \times \{y\})$ for $y \in \mathbb{D}_r$. Recall that $H_{0,\lambda}(x, y) = (p_\lambda(x), 0)$. We denote the Julia set of p_λ by J_{p_λ} . We now construct holomorphic motions.

Lemma 3.7. *Suppose that $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the $(\text{VC})_{\varepsilon,r}$ and the condition (7) holds for each $a \in \mathbb{D}_{\delta_0}$. Then there exists a holomorphic motion $h_{\lambda,y}: \mathbb{D}_{\delta_0} \times S_{0,\lambda,y} \rightarrow \mathbb{C} \times \{y\}$ such that $h_{\lambda,y}(a, (x, y)) \in S_{a,\lambda,y}$ for $y \in \mathbb{D}_r$, $a \in \mathbb{D}_{\delta_0}$ and $(x, y) \in S_{0,\lambda,y}$. In particular, $h_{\lambda,y}$ can be*

extended to a holomorphic motion $\tilde{h}_{\lambda,y}: \mathbb{D}_{\delta_0} \times (\mathbb{C} \times \{y\}) \rightarrow \mathbb{C} \times \{y\}$ and $\tilde{h}_{\lambda,y}(\{a\} \times (J_{p_\lambda} \times \{y\})) = \overline{S_{a,\lambda,y}}$.

Proof. By Lemma 3.5, $H_{a,\lambda}^{-j}(W_{a,\lambda}) \cap (\mathbb{C} \times \{y\})$ has exactly 2^j elements for each $j \in \mathbb{Z}_{\geq 0}$. By Lemma 3.6, we have

$$S_{a,\lambda,y} = (W_{\text{loc}}^s(\mathbf{s}_{a,\lambda}) \cap (\mathbb{C} \times \{y\})) \sqcup \bigsqcup_{j \in \mathbb{Z}_{\geq 0}} (H_{a,\lambda}^{-j}(W_{a,\lambda}) \cap (\mathbb{C} \times \{y\})).$$

By (iii) in Definition 3.1, we have $W_{\text{loc}}^s(\mathbf{s}_{a,\lambda}) = \{(f_a(y), y) : y \in \mathbb{D}_r\}$ for some holomorphic map $f_a: \mathbb{D}_r \rightarrow \mathbb{C}$ depending holomorphically on $a \in \mathbb{D}_{\delta_0}$.

Let v_a be a vertical component of $\mathbf{v}_{a,\lambda}$ such that $v_a \cap (\mathbb{C} \times \{w\})$ moves holomorphically over \mathbb{D}_{δ_0} for each $w \in \mathbb{D}_r$. By Lemma 3.5, there are two vertical components $v_{a,1}$ and $v_{a,2}$ of $\mathbf{v}_{a,\lambda}$ such that $H_{a,\lambda}^{-1}(v_a) \cap (\mathbb{D}_r \times \mathbb{D}_r) = v_{a,1} \cup v_{a,2}$, and $v_{a,j} \cap (\mathbb{C} \times \{w\})$ moves holomorphically over \mathbb{D}_{δ_0} for each fixed $w \in \mathbb{D}_r$ and $j \in \{1, 2\}$. Thus, we can construct $h_{\lambda,y}: \mathbb{D}_{\delta_0} \times S_{0,\lambda,y} \rightarrow \mathbb{C} \times \{y\}$ such that $h_{\lambda,y}(a, (x, y)) \in S_{a,\lambda,y}$ for $y \in \mathbb{D}_r$, $a \in \mathbb{D}_{\delta_0}$ and $(x, y) \in S_{0,\lambda,y}$. By Lemma 3.6, for each fixed $a \in \mathbb{D}_{\delta_0}$ and $y \in \mathbb{D}_r$, $h_{\lambda,y}(a, (x, y))$ is injective with respect to x with $(x, y) \in S_{0,\lambda,y}$. The map $h_{\lambda,y}$ can be extended to a holomorphic motion $\tilde{h}_{\lambda,y}: \mathbb{D}_{\delta_0} \times \mathbb{C} \times \{y\} \rightarrow \mathbb{C} \times \{y\}$ and $\tilde{h}_{\lambda,y}(\{a\} \times (J_{p_\lambda} \times \{y\})) = \overline{S_{a,\lambda,y}}$ (see [11] and [19]). \square

Corollary 3.8. Suppose that $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the $(\text{VC})_{\varepsilon,r}$ and the condition (7) holds for $a \in \mathbb{D}_{\delta_0}$. Then $J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r) = \bigcup_{y \in \mathbb{D}_r} \overline{S_{a,\lambda,y}}$. In particular, $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y\})$ is path connected for $y \in \mathbb{D}_r$ if J_{p_λ} is connected.

Proof. We first show that

$$(12) \quad \overline{W^s(\mathbf{s}_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)} \cap (\mathbb{D}_r \times \mathbb{D}_r) = \overline{W^s(\mathbf{s}_{a,\lambda})} \cap (\mathbb{D}_r \times \mathbb{D}_r),$$

as follows. Let $(z_1, w_1) \in \overline{W^s(\mathbf{s}_{a,\lambda})} \cap (\mathbb{D}_r \times \mathbb{D}_r)$. Since $(z_1, w_1) \in \mathbb{D}_r \times \mathbb{D}_r$, for each $n \in \mathbb{N}$, there exists $(z_{1,n}, w_{1,n}) \in W^s(\mathbf{s}_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ such that $(z_{1,n}, w_{1,n}) \rightarrow (z_1, w_1)$ as $n \rightarrow \infty$. Thus $(z_1, w_1) \in \overline{W^s(\mathbf{s}_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r)}$. Hence $(z_1, w_1) \in \overline{W^s(\mathbf{s}_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r) \cap (\mathbb{D}_r \times \mathbb{D}_r)}$. The opposite inclusion is obvious. Thus we have shown (12).

We next show that

$$(13) \quad \overline{\bigcup_{y \in \mathbb{D}_r} (S_{a,\lambda,y})} \cap (\mathbb{D}_r \times \mathbb{D}_r) = \bigcup_{y \in \mathbb{D}_r} \overline{S_{a,\lambda,y}},$$

as follows. Let $(z_2, w_2) \in \overline{\bigcup_{y \in \mathbb{D}_r} (S_{a,\lambda,y})} \cap (\mathbb{D}_r \times \mathbb{D}_r)$. By $(z_2, w_2) \in \mathbb{D}_r \times \mathbb{D}_r$, we can take $(z_{2,n}, w_{2,n}) \in (\bigcup_{y \in \mathbb{D}_r} S_{a,\lambda,y}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ such that $(z_{2,n}, w_{2,n}) \rightarrow (z_2, w_2)$ as $n \rightarrow \infty$. Let v_n be the vertical component $\mathbf{v}_{a,\lambda}$ which contains $(z_{2,n}, w_{2,n})$. Let $x_n \in \mathbb{D}_r$ such that $(x_n, w_2) \in v_n$. To show $(z_2, w_2) \in \overline{\bigcup_{y \in \mathbb{D}_r} S_{a,\lambda,y}}$, we show that $(x_n, w_2) \rightarrow (z_2, w_2)$ as $n \rightarrow \infty$. Since $\|(x_n, w_2) - (z_2, w_2)\| \leq \|(x_n, w_2) - (z_{2,n}, w_{2,n})\| + \|(z_{2,n}, w_{2,n}) - (z_2, w_2)\|$, it suffices to show that $\|(x_n, w_2) - (z_{2,n}, w_{2,n})\| \rightarrow 0$ as $n \rightarrow \infty$. Since $\text{Pr}_2: v_n \rightarrow \mathbb{D}_r$ is a homeomorphism, we can take a curve $\gamma: [0, 1] \rightarrow v_n$, represented by $\gamma(s) = (f(s), g(s))$, between (x_n, w_2) and $(z_{2,n}, w_{2,n})$ such that $\text{length}_E(\text{Pr}_2 \gamma) = |w_{2,n} - w_2|$, where $\text{length}_E(\text{Pr}_2 \gamma)$ is the length of the curve $\text{Pr}_2 \gamma$ with respect to the Euclidean metric. Note that $\text{Pr}_2 \gamma$ is the segment between $w_{2,n}$ and w_2 . By $(f'(s), g'(s)) \in C_{\gamma(s)}^v$ (see (ii) in Definition 3.1), we have

$$\|(x_n, w_2) - (z_{2,n}, w_{2,n})\| \leq \text{length}_E \gamma \leq \sqrt{2} \int_0^1 \max\{|f'(s)|, |g'(s)|\} ds = \sqrt{2} \int_0^1 |g'(s)| ds.$$

Since $\int_0^1 |g'(s)|ds = |w_{2,n} - w_2| \rightarrow 0$ as $n \rightarrow \infty$, we have that $\|(x_n, w_2) - (z_{2,n}, w_{2,n})\| \rightarrow 0$ as $n \rightarrow \infty$. Hence we have $(z_2, w_2) \in \bigcup_{y \in \mathbb{D}_r} \overline{S_{a,\lambda,y}}$. Therefore $\bigcup_{y \in \mathbb{D}_r} (S_{a,\lambda,y}) \cap (\mathbb{D}_r \times \mathbb{D}_r) \subset \bigcup_{y \in \mathbb{D}_r} \overline{S_{a,\lambda,y}}$. The opposite inclusion is obvious. Thus we have shown (13).

By $J_{a,\lambda}^+ = \overline{W^s(\mathbf{s}_{a,\lambda})}$ (see [3]), $W^s(\mathbf{s}_{a,\lambda}) \cap (\mathbb{D}_r \times \mathbb{D}_r) = \bigcup_{y \in \mathbb{D}_r} S_{a,\lambda,y}$, (12) and (13), we have

$$J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r) = \overline{W^s(\mathbf{s}_{a,\lambda})} \cap (\mathbb{D}_r \times \mathbb{D}_r) = \overline{\bigcup_{y \in \mathbb{D}_r} (S_{a,\lambda,y})} \cap (\mathbb{D}_r \times \mathbb{D}_r) = \bigcup_{y \in \mathbb{D}_r} \overline{S_{a,\lambda,y}}.$$

By the above, we have $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y\}) = \overline{S_{a,\lambda,y}}$ for $y \in \mathbb{D}_r$. Note that the condition $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda}^+ = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$ implies that $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap (J_{p_\lambda} \times \mathbb{C}) = \emptyset$ since $J_{0,\lambda}^+ = J_{p_\lambda} \times \mathbb{C}$. In particular, the critical point 0 of p_λ belongs to $\mathbb{C} \setminus J_{p_\lambda}$. Since J_{p_λ} is connected, p_λ has an attracting or parabolic periodic point. Hence J_{p_λ} is the image of unit circle under a continuous map (see [5]). In particular, J_{p_λ} is path connected. Since $\overline{S_{a,\lambda,y}}$ is homeomorphic to J_{p_λ} (see Lemma 3.7), the section $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y\})$ is path connected for each $y \in \mathbb{D}_r$. \square

The following lemma is useful for checking whether $J_{a,\lambda}$ is disconnected.

Lemma 3.9. *Suppose that $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the condition $(VC)_{\varepsilon,r}$ and the condition (7) holds for $a \in \mathbb{D}_{\delta_0}$. Then each vertical component v of $\mathbf{v}_{a,\lambda}$ contains a point of $J_{a,\lambda}$ for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$.*

Proof. Let $v_0 := v$ be a vertical component of $\mathbf{v}_{a,\lambda}$. Inductively, let v_n be a component of $H_{a,\lambda}^{-1}(v_{n-1}) \cap (\mathbb{D}_r \times \mathbb{D}_r)$ for each $n \in \mathbb{N}$. Then we have the nested compact sequence $\{H_{a,\lambda}^n(\overline{v_n})\}_{n \geq 0}$ with $H_{a,\lambda}^n(\overline{v_n}) \subset H_{a,\lambda}^{n-1}(\overline{v_{n-1}})$ for $n \in \mathbb{N}$. For any $a \in \mathbb{D}_1$, we have $J_{a,\lambda}^- = K_{a,\lambda}^-$ (see Lemma 5.5 in [2]). A point of $\bigcap_{n \in \mathbb{N}} H_{a,\lambda}^{n-1}(\overline{v_{n-1}})$ belongs to $J_{a,\lambda}$ since its backward orbit is bounded and \overline{v} is a subset of $J_{a,\lambda}^+$. \square

We now prove the first main result of this paper, divided into Theorem 3.10 and Theorem 3.11. Theorem 3.10 relates to the connected case, and Theorem 3.11 to the disconnected case.

Theorem 3.10. *Let $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ be a Hénon family satisfying the $(VC)_{\varepsilon,r}$. Assume that the condition (7) holds for $a \in \mathbb{D}_{\delta_0}$. If the Julia set J_{p_λ} of the polynomial p_λ is connected, then the Julia set $J_{a,\lambda}$ of the Hénon map $H_{a,\lambda}$ is connected for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$.*

Proof. Assume that J_{p_λ} is connected and $\lambda \neq 1$. We first show that $J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r)$ is connected. We take any distinct points (x_1, y_1) and (x_2, y_2) in $J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r)$. We construct a path between these points as follows. Let v be a vertical component of $\mathbf{v}_{a,\lambda}$, and (z_j, y_j) be the intersection of $v \cap (\mathbb{C} \times \{y_j\})$ for $j = 1, 2$. Since $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y_j\})$ is homeomorphic to J_{p_λ} for $j = 1, 2$, and J_{p_λ} is a path connected, there exists a path between (z_j, y_j) and (x_j, y_j) in $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y_j\})$ for each $j = 1, 2$. We can take a path between (z_1, y_1) and (z_2, y_2) in v since v is path connected. Thus, $J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r)$ is path connected, which implies that it is connected.

We now show that $J_{a,\lambda}$ is connected. We see that $J_{a,\lambda} = \bigcap_{k \geq 0} \overline{(H_{a,\lambda}^k(J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r)))}$. Moreover, we have $\overline{H_{a,\lambda}^{k+1}(J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r))} \subset \overline{H_{a,\lambda}^k(J_{a,\lambda}^+ \cap (\mathbb{D}_r \times \mathbb{D}_r))}$. Hence $J_{a,\lambda}$ is a nested intersection of connected compact subsets. Thus $J_{a,\lambda}$ is connected. \square

We next show the following theorem.

Theorem 3.11. *Let $\{H_{a,\lambda}\}_{a \in \mathbb{D}_{\delta_0}}$ be a Hénon family satisfying the condition $(VC)_{\varepsilon,r}$. Assume that the condition (7) holds for $a \in \mathbb{D}_{\delta_0}$. If the Julia set J_{p_λ} of the polynomial p_λ is disconnected, then the Julia set $J_{a,\lambda}$ of the Hénon map $H_{a,\lambda}$ is disconnected for $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$.*

Proof. Assume that J_{p_λ} is disconnected. That is, it is a Cantor set. We show that $J_{a,\lambda}$ is disconnected. We can take a Jordan curve $\gamma_j \subset \mathbb{C}$ for each $j = 1, 2$ with $\gamma_1 \cap \gamma_2 = \emptyset$ such that $J_{p_\lambda} \subset U_1 \cup U_2$ and $J_{p_\lambda} \cap U_j \neq \emptyset$ for $j = 1, 2$, where U_j is the bounded domain with the boundary γ_j for $j = 1, 2$. Let $\tilde{h}_{\lambda,y} : \mathbb{D}_{\delta_0} \times (\mathbb{C} \times \{y\}) \rightarrow \mathbb{C} \times \{y\}$ be the holomorphic motion given in Lemma 3.7.

Fix $a \in \mathbb{D}_{\delta_0} \setminus \{0\}$. Let $\gamma_{a,y,j} := \text{Pr}_1 \tilde{h}_{\lambda,y}(\{a\} \times (\gamma_j \times \{y\})) \subset \mathbb{C}$ and $U_{a,y,j} = \text{Pr}_1 \tilde{h}_{\lambda,y}(\{a\} \times (U_j \times \{y\})) \subset \mathbb{C}$ for $j = 1, 2$ and $y \in \mathbb{D}_r$. Since $(\gamma_{a,y,j} \times \{y\}) \cap J_{a,\lambda}^+ = \emptyset$ for $j = 1, 2$ and $y \in \mathbb{D}_r$, there exists $\varepsilon_1(a, y)$ with $0 < \varepsilon_1(a, y) < r - |y|$ such that

$$(14) \quad \left(\bigcup_{y \in \mathbb{D}_r} (\gamma_{a,y,j} \times \overline{\mathbb{D}_{\varepsilon_1(a,y)}(y)}) \right) \cap J_{a,\lambda}^+ = \emptyset,$$

for each $j = 1, 2$ and each $y \in \mathbb{D}_r$. We now show that

$$(15) \quad J_{a,\lambda}^+ \cap (\mathbb{C} \times \overline{\mathbb{D}_{\varepsilon_1(a,y)}(y)}) \subset \left(\bigcup_{j=1}^2 U_{a,y,j} \right) \times \overline{\mathbb{D}_{\varepsilon_1(a,y)}(y)},$$

for each $y \in \mathbb{D}_r$. Fix $y_0 \in \mathbb{D}_r$. Let $(z, w) \in (\mathbb{C} \setminus \bigcup_{j=1}^2 U_{a,y_0,j}) \times \overline{\mathbb{D}_{\varepsilon_1(a,y_0)}(y_0)}$. In order to show (15), it suffices to show that $(z, w) \notin J_{a,\lambda}^+ \cap (\mathbb{C} \times \overline{\mathbb{D}_{\varepsilon_1(a,y_0)}(y_0)})$. If $(z, w) \in \bigcup_{j=1}^2 \gamma_{a,y_0,j} \times \overline{\mathbb{D}_{\varepsilon_1(a,y_0)}(y_0)}$, then $(z, w) \notin J_{a,\lambda}^+$ by (14). Thus we may assume that $(z, w) \in (\mathbb{C} \setminus \bigcup_{j=1}^2 U_{a,y_0,j}) \times \overline{\mathbb{D}_{\varepsilon_1(a,y_0)}(y_0)}$. By assuming that $(z, w) \in J_{a,\lambda}^+ \cap (\mathbb{C} \times \overline{\mathbb{D}_{\varepsilon_1(a,y_0)}(y_0)})$, we derive a contradiction as follows. Note that $w \in \mathbb{D}_r$ by $\varepsilon_1(a, y_0) < r - |y_0|$. By $\overline{W^s(\mathbf{s}_{a,\lambda})} = J_{a,\lambda}^+$, there exists a vertical component v_0 of $\mathbf{v}_{a,\lambda}$ and a point $\hat{z} \in \mathbb{C}$ such that $\{(\hat{z}, w)\} = v_0 \cap (\mathbb{C} \times \{w\}) \subset (\mathbb{C} \setminus \bigcup_{j=1}^2 \overline{U_{a,y_0,j}}) \times \{w\}$. Let $x_0 \in \mathbb{C}$ be the point such that $\{(x_0, y_0)\} = v_0 \cap (\mathbb{C} \times \{y_0\})$. By $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y_0\}) = \overline{S_{a,\lambda,y_0}}$ (see Corollary 3.8), we have $J_{a,\lambda}^+ \cap (\mathbb{C} \times \{y_0\}) \subset (\bigcup_{j=1}^2 U_{a,y_0,j}) \times \{y_0\}$. Hence, we have $\{(x_0, y_0)\} = v_0 \cap (\mathbb{C} \times \{y_0\}) \subset (\bigcup_{j=1}^2 U_{a,y_0,j}) \times \{y_0\}$. Since $\text{Pr}_2 : v_0 \rightarrow \mathbb{D}_r$ is a homeomorphism, we can take a path $\gamma_0 \subset v_0 \cap (\mathbb{C} \times \overline{\mathbb{D}_{\varepsilon_1(a,y_0)}(y_0)})$ between (x_0, y_0) and (\hat{z}, w) . Then we have $(\text{Pr}_1 \gamma_0) \cap (\bigcup_{j=1}^2 \gamma_{a,y_0,j}) \neq \emptyset$ which implies that $v_0 \cap ((\bigcup_{j=1}^2 \gamma_{a,y_0,j}) \times \overline{\mathbb{D}_{\varepsilon_1(a,y_0)}(y_0)}) \neq \emptyset$. By $v_0 \subset J_{a,\lambda}^+$ and (14), we have a contradiction. Thus we have (15) for each $y \in \mathbb{D}_r$.




Fig.3. The case $(x_0, y_0) \in U_{a,y_0,1} \times \{y_0\}$

Let $\mathbf{U}_1 = \bigcup_{y \in \mathbb{D}_r} (U_{a,y,1} \times \overline{\mathbb{D}_{\varepsilon_1(a,y)/2}(y)})$. Since $\overline{\mathbb{D}_{r/2}}$ is compact, we can take a positive integer

$N \in \mathbb{N}$ and points $y_1, y_2, \dots, y_N \in \mathbb{D}_r$ such that $\mathbf{U}_{1,N} := \bigcup_{j=1}^N (U_{a,y_j,1} \times \mathbb{D}_{\varepsilon_1(a,y_j)/2}(y_j))$ satisfies $\overline{\mathbb{D}_{r/2}} \subset \text{Pr}_2 \mathbf{U}_{1,N}$. Let $\mathbf{v}_{a,\lambda,j}$ be the union of all vertical components of $\mathbf{v}_{a,\lambda}$ which intersect $U_{a,0,j} \times \{0\}$ for $j = 1, 2$. We now show that

$$(16) \quad \mathbf{v}_{a,\lambda,1} \cap (\overline{\mathbb{D}_r} \times \mathbb{D}_{r/2}) \subset \mathbf{U}_{1,N}.$$

Take a vertical component $v_1 \subset \mathbf{v}_{a,\lambda,1}$ of $\mathbf{v}_{a,\lambda}$ and a point $w_1 \in \mathbb{D}_{r/2}$. To show (16), it suffices to show that $v_1 \cap (\mathbb{D}_r \times \{w_1\}) \subset \mathbf{U}_{1,N}$. Let $\tilde{h}_{a,\lambda,y} : \mathbb{C} \times \{y\} \rightarrow \mathbb{C} \times \{y\}$, $\tilde{h}_{a,\lambda,y}(x, y) := \tilde{h}_{a,\lambda,y}(a, (x, y))$ for $y \in \mathbb{D}_r$ and $x \in \mathbb{C}$. By the construction of $\tilde{h}_{a,\lambda,y}$ (see the proof of Lemma 3.7), $\text{Pr}_1 \bigcup_{y \in \mathbb{D}_r} \tilde{h}_{a,\lambda,y}^{-1}(v_1 \cap (\mathbb{C} \times \{y\}))$ consists of a single point of J_{p_λ} , say b_1 . We have $b_1 \in U_1$ by $v_1 \cap (U_{a,0,1} \times \{0\}) \neq \emptyset$. Since $w_1 \in \mathbb{D}_{r/2}$, we can take a positive integer k_1 with $1 \leq k_1 \leq N$ such that $w_1 \in \mathbb{D}_{\varepsilon_1(a,y_{k_1})/2}(y_{k_1})$. Let $x_1 \in \mathbb{C}$ be the point such that $\{(x_1, y_{k_1})\} = v_1 \cap (\mathbb{C} \times \{y_{k_1}\})$. By $\text{Pr}_1 \tilde{h}_{a,\lambda,y_{k_1}}^{-1}(U_{a,y_{k_1},1} \times \{y_{k_1}\}) = U_1 \ni b_1$, we have $(x_1, y_{k_1}) \in U_{a,y_{k_1},1} \times \mathbb{D}_{\varepsilon_1(a,y_{k_1})/2}(y_{k_1})$. This implies that $v_1 \cap (U_{a,y_{k_1},1} \times \{y_{k_1}\}) \neq \emptyset$. By (14) and (15), we have $\{(z_1, w_1)\} := v_1 \cap (\mathbb{D}_r \times \{w_1\}) \subset U_{a,y_{k_1},1} \times \{w_1\}$. Otherwise, for any path in $v_1 \cap (\mathbb{C} \times \mathbb{D}_{\varepsilon_1(a,y_{k_1})/2}(y_{k_1}))$ between (z_1, w_1) and (x_1, y_{k_1}) , the path necessarily intersects $\gamma_{a,y_{k_1},1} \times \mathbb{D}_{\varepsilon_1(a,y_{k_1})/2}(y_{k_1})$. This leads to a contradiction (see the proof of (15)). Thus we have $v_1 \cap (\mathbb{D}_r \times \{w_1\}) \subset U_{a,y_{k_1},1} \times \{w_1\}$, which implies that $v_1 \cap (\mathbb{D}_r \times \{w_1\}) \subset U_{a,y_{k_1},1} \times \mathbb{D}_{\varepsilon_1(a,y_{k_1})/2}(y_{k_1}) \subset \mathbf{U}_{1,N}$. Thus we have (16).

By using (16), we show that

$$(17) \quad \overline{\mathbf{v}_{a,\lambda,1}} \cap (\overline{\mathbb{D}_r} \times \mathbb{D}_{r/2}) \subset \mathbf{U}_{1,N}.$$

We have $\overline{\mathbf{v}_{a,\lambda,1}} \cap (\mathbb{D}_r \times \mathbb{D}_{r/2}) = \bigcup_{y \in \mathbb{D}_{r/2}} \overline{\mathbf{v}_{a,\lambda,1} \cap (\mathbb{D}_r \times \{y\})}$ by the same argument as in the proof of (13). Thus, to show that (17), it suffices to show that $\overline{\mathbf{v}_{a,\lambda,1} \cap (\mathbb{D}_r \times \{y\})} \subset \mathbf{U}_{1,N}$ for $y \in \mathbb{D}_{r/2}$. Fix $\hat{w}_1 \in \mathbb{D}_{r/2}$. By using the argument in the proof of (16), we can take $\hat{k}_1 \in \mathbb{N}$ with $1 \leq \hat{k}_1 \leq N$ such that $\hat{w}_1 \in \mathbb{D}_{\varepsilon_1(a,y_{\hat{k}_1})/2}(y_{\hat{k}_1})$ and $\mathbf{v}_{a,\lambda,1} \cap (\mathbb{D}_r \times \{\hat{w}_1\}) \subset U_{a,y_{\hat{k}_1},1} \times \{\hat{w}_1\}$. By (14), $\overline{\mathbf{v}_{a,\lambda,1} \cap (\mathbb{D}_r \times \{\hat{w}_1\})} \subset U_{a,y_{\hat{k}_1},1} \times \{\hat{w}_1\}$. In particular, we have $\overline{\mathbf{v}_{a,\lambda,1} \cap (\mathbb{D}_r \times \{\hat{w}_1\})} \subset U_{a,y_{\hat{k}_1},1} \times \mathbb{D}_{\varepsilon_1(a,y_{\hat{k}_1})/2}(y_{\hat{k}_1}) \subset \mathbf{U}_{1,N}$. Thus we have (17).

We next show that

$$(18) \quad \mathbf{v}_{a,\lambda,2} \cap (\overline{\mathbb{D}_r} \times \mathbb{D}_{r/2}) \cap \overline{\mathbf{U}_{1,N}} = \emptyset.$$

Note that $\overline{\mathbf{U}_{1,N}} = \bigcup_{j=1}^N (\overline{U_{a,y_j,1}} \times \overline{\mathbb{D}_{\varepsilon_1(a,y_j)/2}(y_j)})$. Assume that there is $(z_2, w_2) \in \mathbf{v}_{a,\lambda,2} \cap (\overline{\mathbb{D}_r} \times \mathbb{D}_{r/2}) \cap \overline{\mathbf{U}_{1,N}}$. Let $v_2 \subset \mathbf{v}_{a,\lambda,2}$ be the vertical component of $\mathbf{v}_{a,\lambda}$ which contains (z_2, w_2) . By (14), we may assume that $(z_2, w_2) \in U_{a,y_{k_2},1} \times \overline{\mathbb{D}_{\varepsilon_1(a,y_{k_2})/2}(y_{k_2})}$ for some k_2 with $1 \leq k_2 \leq N$. Let $x_2 \in \mathbb{C}$ be the point such that $\{(x_2, y_{k_2})\} = v_2 \cap (\mathbb{C} \times \{y_{k_2}\})$. By (14) and $z_2 \in U_{a,y_{k_2},1}$, we have $\{(x_2, y_{k_2})\} = v_2 \cap (\mathbb{C} \times \{y_{k_2}\}) \subset U_{a,y_{k_2},1} \times \{y_{k_2}\}$. Otherwise, for any path in $v_2 \cap (\mathbb{C} \times \mathbb{D}_{\varepsilon_1(a,y_{k_2})/2}(y_{k_2}))$ between (x_2, y_{k_2}) and (z_2, w_2) , the path necessarily intersects $\gamma_{a,y_{k_2},1} \times \mathbb{D}_{\varepsilon_1(a,y_{k_2})/2}(y_{k_2})$. This leads to a contradiction (see the proof of (15)). Thus we have $v_2 \cap (\mathbb{C} \times \{y_{k_2}\}) \subset U_{a,y_{k_2},1} \times \{y_{k_2}\}$, which implies that $\text{Pr}_1 \tilde{h}_{a,\lambda,y_{k_2}}^{-1}(v_2 \cap (\mathbb{C} \times \{y_{k_2}\})) \subset U_1$. On the other hand, $v_2 \cap (U_{a,0,2} \times \{0\}) \neq \emptyset$ since $v_2 \subset \mathbf{v}_{a,\lambda,2}$ is a vertical component of $\mathbf{v}_{a,\lambda}$. Thus $\text{Pr}_1 \tilde{h}_{a,\lambda,0}^{-1}(v_2 \cap (\mathbb{C} \times \{0\})) \subset U_2$. Since $U_1 \cap U_2 = \emptyset$ and $\text{Pr}_1 \bigcup_{y \in \mathbb{D}_r} \tilde{h}_{a,\lambda,y}^{-1}(v_2 \cap (\mathbb{C} \times \{y\}))$ consists of a single point, we have a contradiction. This contradiction implies that (18) holds.

By using (18), we show that

$$(19) \quad \overline{\mathbf{v}_{a,\lambda,2}} \cap (\overline{\mathbb{D}_r} \times \mathbb{D}_{r/2}) \cap \overline{\mathbf{U}_{1,N}} = \emptyset.$$

By the same argument as in the proof of (17), it suffices to show that $\overline{\mathbf{v}_{a,\lambda,2} \cap (\mathbb{D}_r \times \{y\})} \cap$

$\overline{\mathbf{U}_{1,N}} = \emptyset$ for $y \in \mathbb{D}_{r/2}$. Fix $\hat{w}_2 \in \mathbb{D}_{r/2}$. By (18), we have $\mathbf{v}_{a,\lambda,2} \cap (\mathbb{D}_r \times \{\hat{w}_2\}) \cap \overline{\mathbf{U}_{1,N}} = \emptyset$. Note that $\overline{\mathbf{U}_{1,N} \cap (\mathbb{D}_r \times \{\hat{w}_2\})}$ can be represented by a finite union of the sets $\overline{\mathbf{U}_{a,y_j,1}} \times \{\hat{w}_2\}$. Thus, by (14), we have $\overline{\mathbf{v}_{a,\lambda,2} \cap (\mathbb{D}_r \times \{\hat{w}_2\})} \cap \overline{\mathbf{U}_{1,N} \cap (\mathbb{D}_r \times \{\hat{w}_2\})} = \emptyset$. Hence we have (19) by $\overline{\mathbf{v}_{a,\lambda,2} \cap (\mathbb{D}_r \times \{\hat{w}_2\})} \cap \overline{\mathbf{U}_{1,N} \cap (\mathbb{D}_r \times \{\hat{w}_2\})} = \overline{\mathbf{v}_{a,\lambda,2} \cap (\mathbb{D}_r \times \{\hat{w}_2\})} \cap \overline{\mathbf{U}_{1,N}}$.

Finally, we show that $J_{a,\lambda}$ is disconnected. We set $\mathbf{U}_2 := (\mathbb{C} \times \mathbb{D}_{r/2}) \setminus \overline{\mathbf{U}_{1,N}}$. Then $J_{a,\lambda} \subset \mathbf{U}_{1,N} \cup \mathbf{U}_2$. Indeed, we have $J_{a,\lambda} \subset \mathbb{D}_r \times \mathbb{D}_{r/2}$ by (2) and (i) in Definition 3.1. Thus $J_{a,\lambda} \subset (\overline{\mathbf{v}_{a,\lambda,1}} \cup \overline{\mathbf{v}_{a,\lambda,2}}) \cap (\mathbb{D}_r \times \mathbb{D}_{r/2})$. Therefore, by (17) and (19), $J_{a,\lambda} \subset \mathbf{U}_{1,N} \cup \mathbf{U}_2$. Clearly, we have $J_{a,\lambda} \cap \mathbf{U}_{1,N} \neq \emptyset$ by (16) and Lemma 3.9. Similarly, we have $J_{a,\lambda} \cap \mathbf{U}_2 \neq \emptyset$ by (18) and Lemma 3.9. Thus $J_{a,\lambda}$ is disconnected. \square

4. Application for radial perturbations of semi-parabolic Hénon maps

In this section, we apply Theorem 3.10 to perturbations of semi-parabolic Hénon maps. To consider the connectivity of J for the parameters near the boundary of the Mandelbrot set, we consider perturbations of semi-parabolic Hénon maps by using a perturbation of one eigenvalue of semi-parabolic fixed points. Let $\lambda_0 = \exp(2\pi i m/l)$, where $l \in \mathbb{Z} \setminus \{0\}$, $m \in \mathbb{Z}$ and $(m, l) = 1$. Let $\{\lambda_t\}_{t \in [0, \delta_0]}$ be a one-parameter continuous family of complex numbers, where $\delta_0 > 0$. Assume that $\lambda_t^l = \exp(L_t + i\theta_t)$ and $\mathbb{R} \ni \theta_t \rightarrow 0$ as $t \rightarrow 0$, where $L_t \in \mathbb{R} \setminus \{0\}$ and $\theta_t \in \mathbb{R}$ for $0 < t < \delta_0$.

DEFINITION 4.1 (RADIAL PERTURBATIONS). We say that a family $RD_{\lambda_t, \delta_0} := \{H_{a, \lambda_t}\}_{a \in \mathbb{D}_{\delta_0}, 0 < t < \delta_0}$ is a *radial perturbation* of the semi-parabolic Hénon family $\{H_{a, \lambda_0}\}_{a \in \mathbb{D}_{\delta_0}}$ if $\theta_t = O(L_t)$.

In order to apply Theorem 3.10 to RD_{λ_t, δ_0} , we first check that the section $\{H_{a, \lambda_t}\}_{a \in \mathbb{D}_{\delta_0}}$ of RD_{λ_t, δ_0} satisfies the condition $(VC)_{\varepsilon, r}$ for each t with $0 < t < \delta_0$.

Lemma 4.2. *There is $\delta_0 > 0$ such that the section $\{H_{a, \lambda_t}\}_{a \in \mathbb{D}_{\delta_0}}$ of RD_{λ_t, δ_0} satisfies the condition $(VC)_{\varepsilon, r}$ for each t with $0 < t < \delta_0$.*

Proof. If $\lambda_0 \neq 1$, then the section $\{H_{a, \lambda_t}\}_{a \in \mathbb{D}_{\delta_0}}$ of RD_{λ_t, δ_0} satisfies the $(VC)_{\varepsilon, r}$ for $0 < t < \delta_0$ by Lemma 3.2, taking a smaller $\delta_0 > 0$ if necessary (see Remark 3.3).

Assume that $\lambda_0 = 1$. We may assume that each $H_{a, \lambda_t} \in RD_{\lambda_t, \delta_0}$ with $a \neq 0$ satisfies the condition (2) with respect to r by taking a smaller δ_0 and a larger r if necessary. Fix $\varepsilon > 0$. We can show that $\{H_{a, \lambda_t}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies (i), (ii) in Definition 3.1 for $0 < t < \delta_0$ in the same way as in the proof of Lemma 3.2, taking a smaller $\delta_0 > 0$ if necessary (see Remark 3.4). Thus it suffices to show that $\{H_{a, \lambda_t}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies (iii) in Definition 3.1 for $0 < t < \delta_0$, taking a smaller $\delta_0 > 0$ if necessary.

To show that (iii) in Definition 3.1 holds, we first show that H_{a, λ_t} has a saddle fixed point for $0 < t < \delta_0$. In order to show this, we show that H_{a, λ_t} has two distinct fixed points. Consider the equation $(x^2 + c(a, \lambda_t) + ay, ax) = (x, y)$. By $y = ax$, we have

$$(20) \quad x^2 + (a^2 - 1)x + c(a, \lambda_t) = 0.$$

Assume that $(x - \alpha)(x - \beta) = x^2 + (a^2 - 1)x + c(a, \lambda_t)$. If $\alpha = \beta$, then $\alpha = 1/2 - a^2/2$. In this case, H_{a, λ_t} has only one fixed point $(1/2 - a^2/2, a/2 - a^3/2)$. Since H_{a, λ_t} has a fixed point $\mathbf{q}_{a, \lambda_t} = (\lambda_t/2 - a^2/(2\lambda_t), a(\lambda_t/2 - a^2/(2\lambda_t)))$ (see (4)), we have $\lambda_t = 1$ or $\lambda_t = -a^2$. Since $\lambda_0 = 1$, we have $\lambda_t = 1$ for $a \in \mathbb{D}_{\delta_0}$, by taking $\delta_0 > 0$ so that $0 < \delta_0 < 1/2$ and

$|\lambda_t| > 1/2$ for $0 < t < \delta_0$. This contradicts $\lambda_t = \exp(L_t + i\theta_t)$ with $L_t \neq 0$ for $0 < t < \delta_0$. This contradiction shows that H_{a,λ_t} has two distinct fixed points. The other fixed point of $H_{a,\lambda}$ is $\mathbf{u}_{a,\lambda} := (1 - a^2 - \text{Pr}_1 \mathbf{q}_{a,\lambda}, a(1 - a^2 - \text{Pr}_1 \mathbf{q}_{a,\lambda}))$ by (20). These fixed points $\mathbf{q}_{a,\lambda}, \mathbf{u}_{a,\lambda}$ depend holomorphically for (a, λ) in a small neighborhood of $(0, 1)$. At least one of the fixed points $\mathbf{q}_{a,\lambda_t}, \mathbf{u}_{a,\lambda_t}$ is a saddle by $\mathbf{q}_{a,\lambda_t} \neq \mathbf{u}_{a,\lambda_t}$ (see [14, Theorem 7.1.16, p.234]). Hence, there is a fixed point $\mathbf{s}_{a,\lambda}$ of $H_{a,\lambda}$ depending holomorphically for (a, λ) in a small neighborhood of $(0, \lambda_0)$, and \mathbf{s}_{a,λ_t} is a saddle fixed point of $H_{a,\lambda_t} \in RD_{\lambda_t, \delta_0}$, by taking a smaller δ_0 if necessary.

Finally, we show that (iii) in Definition 3.1 holds. Let $\tilde{\Phi}_{0,\lambda}(z) := (\text{Pr}_1 \mathbf{s}_{0,\lambda}, z)$ and $\tilde{\Phi}_{a,\lambda}$ be the parametrization of $W^{ss}(\mathbf{s}_{a,\lambda})$ for $(a, \lambda) \in \mathbb{D}_{\delta_0} \setminus \{0\} \times \mathbb{D}_{\delta_0}(\lambda_0)$ given by (5). In the same way as in the proof of Lemma 2.13, $\tilde{\Phi}_{a,\lambda}(z)$ is holomorphic with $(a, \lambda, z) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0) \times \mathbb{C}$. We have $W_{\text{loc}}^{ss}(\mathbf{s}_{a,\lambda}) = \{(\tilde{\Phi}_{a,\lambda,1}(\tilde{\Phi}_{a,\lambda,2}^{-1}(y)), y) : y \in \mathbb{D}_r\}$ for $(a, \lambda) \in \mathbb{D}_{\delta_0} \times \mathbb{D}_{\delta_0}(\lambda_0)$, taking a smaller δ_0 if necessary, where $\tilde{\Phi}_{a,\lambda} = (\tilde{\Phi}_{a,\lambda,1}, \tilde{\Phi}_{a,\lambda,2})$ (see the proof of Lemma 3.2). We have $\mathbf{s}_{a,\lambda_0} = \mathbf{q}_{a,\lambda_0}$ since $\mathbf{u}_{a,\lambda_t}, \mathbf{q}_{a,\lambda_t} \rightarrow \mathbf{q}_{a,\lambda_0}$ as $t \rightarrow 0$. Since $W_{\text{loc}}^{ss}(\mathbf{q}_{0,\lambda_0}) = \{\text{Pr}_1 \mathbf{q}_{0,\lambda_0}\} \times \mathbb{D}_r$, we see that $T_{(x,y)} W_{\text{loc}}^s(\mathbf{s}_{a,\lambda_t}) \subset C_{(x,y)}^v$ for $(x, y) \in W_{\text{loc}}^s(\mathbf{s}_{a,\lambda_t})$, $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$, taking a smaller $\delta_0 > 0$ if necessary. Thus, $H_{a,\lambda_t} \in RD_{\lambda_t, \delta_0}$ satisfies (iii) in Definition 3.1 for $0 < t < \delta_0$. \square

We next prepare local coordinates near semi-parabolic fixed points to check that the condition (7) holds for RD_{λ_t, δ_0} .

Lemma 4.3 ([16, Theorem 3.5 and its proof]). *Let $r > 3$ be a fixed constant, $\lambda_0 := \exp(2\pi im/l)$ and $\lambda_t := (1 + t)\lambda_0$ where $t \in \mathbb{R}$. Then, there exist $\delta > 0, \delta' > 0$ such that for $|a| < \delta$ and $|t| < \delta'$ there exists a coordinate transformation $\phi_{a,t}: B = \mathbb{D}_{\rho'}(\text{Pr}_1 \mathbf{q}_{0,\lambda_t}) \times \mathbb{D}_r \rightarrow \mathbb{D}_{\rho} \times \mathbb{D}_{r+O(|a|)}$ such that $\phi_{a,t}(\mathbf{q}_{a,\lambda_t}) = (0, 0)$, $W_{\text{loc}}^{ss}(\mathbf{q}_{a,\lambda_t}) \subset \mathbb{D}_{\rho'}(\text{Pr}_1 \mathbf{q}_{0,\lambda_t}) \times \mathbb{D}_r$, $\phi_{a,t}(W_{\text{loc}}^{ss}(\mathbf{q}_{a,\lambda_t})) \subset \{0\} \times \mathbb{C}$, the image of any horizontal curve $\mathbb{D}_{\rho'}(\text{Pr}_1 \mathbf{q}_{0,\lambda_t}) \times \{y_1\}$ under $\phi_{a,t}$ is a subset of $\mathbb{C} \times \{y_2\}$ for some $y_2 \in \mathbb{D}_{r+O(|a|)}$, and $\tilde{H}_{a,\lambda_t} = \phi_{a,t} \circ H_{a,\lambda_t} \circ \phi_{a,t}^{-1}, \tilde{H}_{a,\lambda_t}(x, y) = (X_1, Y_1)$ has the form*

$$(21) \quad (X_1, Y_1) = (\lambda_t(x + x^{l+1} + C_{a,t}x^{2l+1} + b_{a,t,2l+2}(y)x^{2l+2} + \dots), v_{a,t}y + xh_{a,t}(x, y)),$$

where $C_{a,t}$ is a constant depending only on a and t , $xh_{a,t}(x, y) = O(a)$ and $v_{a,t}$ is the other eigenvalue of $(DH_{a,\lambda_t})_{\mathbf{q}_{a,\lambda_t}}$. Moreover, the transformation $\phi_{a,t}$ is analytic for a and t ,

$$\lim_{a \rightarrow 0} \phi_{a,t}(x, y) = (\phi_t(x), y),$$

uniformly for t . The map $\phi_t: \mathbb{D}_{\rho'}(\text{Pr}_1 \mathbf{q}_{0,\lambda_t}) \rightarrow \mathbb{D}_{\rho}$ is the transformation of the polynomial p_{λ_t} and

$$\phi_t \circ p_{\lambda_t} \circ \phi_t^{-1}(x) = \lambda_t(x + x^{l+1} + C_{0,t}x^{2l+1} + O(x^{2l+2})).$$

For $r > 3$, $\lambda_0 = \exp(2\pi im/l)$, if $(c, a) \in \mathcal{P}_{\lambda_0}$ with sufficiently small $|a|$, then the sets U^{\pm} given in Definition 2.3 satisfy the equations (2). The condition $\lambda_t = (1 + t)\lambda_0$ in Lemma 4.3 corresponds to $\theta_t = 0$ for $0 < t < \delta_0$ in Definition 4.1. To see that $H_{a,\lambda_t} \in RD_{\lambda_t, \delta_0}$ has the form (21) (Lemma 4.4), we sketch the proof of Lemma 4.3.

Sketch of the proof of Lemma 4.3. The proof of Lemma 4.3 breaks into four steps.

Step 1. Let $\Phi_{a,\lambda_t} = (\Phi_{a,\lambda_t,1}, \Phi_{a,\lambda_t,2})$ be given in Lemma 2.8. Then $\Phi_{a,\lambda_t}(y) = \Phi_{0,\lambda_t}(y) + O(a)$ by Lemma 2.8. For sufficiently small $|a|$ and $|t|$, we may assume that there exists $\Phi_{a,\lambda_t,2}^{-1}$ in \mathbb{D}_r by Rouché's theorem and $\Phi_{a,\lambda_t,2}(z) = z$. For $(x, y) \in \mathbb{C} \times \mathbb{D}_r$, consider the transformation (22)

$$(X, Y) = (x - \Phi_{a,\lambda_t,1}(\Phi_{a,\lambda_t,2}^{-1}(y)), \Phi_{a,\lambda_t,2}^{-1}(y)) \text{ with inverse } (x, y) = (X + \Phi_{a,\lambda_t,1}(Y), \Phi_{a,\lambda_t,2}(Y)),$$

which maps $W_{\text{loc}}^{ss}(\mathbf{q}_{a,\lambda_t})$ into $\{0\} \times \mathbb{C}$. By using the transformation (22), H_{a,λ_t} and H_{0,λ_t} have the forms

$$(b_{a,t,1}(Y)X + b_{a,t,2}(Y)X^2 + \cdots, v_{a,t}Y + Xh_{a,t}(X, Y)) \text{ and } (\lambda_t X + X^2, 0),$$

respectively. Note that $b_{a,t,1}(0) = \lambda_t$ and $b_{a,t,1}(Y) = \lambda_t + O(a)$ since $\Phi_{a,t}(y) = \Phi_{0,t}(y) + O(a)$.

Step 2. Suppose that $H_{a,\lambda_t}(x, y)$ has the form

$$(23) \quad (b_{a,t,1}(y)x + b_{a,t,2}(y)x^2 + \cdots, v_{a,t}y + xh_{a,t}(x, y)).$$

Let us reduce the function $b_{a,t,1}(y)$ to $b_{a,t,1}(0) = \lambda_t$ (see Proposition 3.2 in [17]). Since $\frac{b_{a,t,1}(v_{a,t}^n y)}{\lambda_t} = 1 + O(v_{a,t}^n y)$, the product

$$u_{a,t}(y) = \prod_{n \geq 0} \left(\frac{b_{a,t,1}(v_{a,t}^n y)}{\lambda_t} \right),$$

converges for $y \in \mathbb{D}_r$. By using $(X, Y) = (u_{a,t}(y)x, y)$ with inverse $(x, y) = (X/u_{a,t}(Y), Y)$, (23) has the form

$$\begin{aligned} & u_{a,t}(v_{a,t}Y + Xh_{a,t}(X/u_{a,t}(Y), Y)/u_{a,t}(Y)) \times (b_{a,t,1}(Y)X/u_{a,t}(Y) + b_{a,t,2}(Y)(X/u_{a,t}(Y))^2 + \cdots) \\ &= \frac{u_{a,t}(v_{a,t}Y)b_{a,t,1}(Y)}{u_{a,t}(Y)}X + O(X^2) = \lambda_t X + O(X^2), \end{aligned}$$

in the first coordinate.

Step 3. We may assume that H_{a,λ_t} has the form

$$(\lambda_t x + b_{a,t,2}(y)x^2 + b_{a,t,3}(y)x^3 + \cdots, v_{a,t}y + xh_{a,t}(x, y)).$$

We next reduce the function $b_{a,t,k}(y)$ to constants by induction on $2 \leq k \leq 2l + 1$. Consider

$$(24) \quad (\lambda_t x + b_{a,t,2}x^2 + b_{a,t,3}x^3 + \cdots + b_{a,t,k-1}x^{k-1} + b_{a,t,k}(y)x^k + \cdots, v_{a,t}y + xh_{a,t}(x, y)),$$

where $b_{a,t,j}$ is constant for $j = 1, 2, \dots, k-1$. We set

$$v_{a,t}(y) = \sum_{n=0}^{\infty} (b_{a,t,k}(v_{a,t}^n y) - b_{a,t,k}(0))\lambda_t^{n(k-1)-1}.$$

This series converges since $|v_{a,t}\lambda_t^{k-1}| < |v_{a,t}\lambda_t^{2l}| < 1$ for sufficiently small t and $|a|$. By using local coordinate $(X, Y) = (x + v_{a,t}(y)x^k, y)$ with inverse $(x, y) = (X - v_{a,t}(Y)X^k + \cdots, Y)$, (24) has the form

$$\begin{aligned} & \lambda_t X + \cdots + b_{a,t,k-1}X^{k-1} + (b_{a,t,k}(Y) + \lambda_t^k v_{a,t}(v_{a,t}Y) - \lambda_t v_{a,t}(Y))X^k + O(X^{k+1}) \\ &= \lambda_t X + \cdots + b_{a,t,k-1}X^{k-1} + b_{a,t,k}(0)X^k + O(X^{k+1}), \end{aligned}$$

in the first coordinate.

Step 4. We may assume that H_{a,λ_t} has the form

$$\lambda_t(x + b_{a,t,2}x^2 + \cdots + b_{a,t,2l+1}x^{2l+1} + b_{a,t,2l+2}(y)x^{2l+2} + \cdots),$$

in the first coordinate. We reduce $b_{a,t,k}$ to 0 for each k with $2 \leq k \leq 2l + 1$ and $k-1 \notin l\mathbb{N}$ by induction. We first assume that H_{a,λ_t} has the form

$$(25) \quad \lambda_t(x + b_{a,t,k}x^k + \cdots + b_{a,t,l+1}x^{l+1} + \cdots),$$

in the first coordinate. By the local coordinate

$$(26) \quad (X, Y) = (x - \lambda_t b_{a,t,k}x^k / (\lambda_t^k - \lambda_t), y) \text{ with inverse } (x, y) = (X + \lambda_t b_{a,t,k}X^k / (\lambda_t^k - \lambda_t) + \cdots, Y),$$

(25) has the form

$$\lambda_t X + (\lambda_t b_{a,t,k} - (\lambda_t^k - \lambda_t)\lambda_t b_{a,t,k} / (\lambda_t^k - \lambda_t))X^k + \cdots = \lambda_t X + O(X^{k+1}).$$

By induction and a linear transformation, we may assume that the first coordinate of H_{a,λ_t} has the form

$$\lambda_t(x + x^{l+1} + b_{a,t,l+2}x^{l+2} + \cdots + b_{a,t,2l+1}x^{2l+1} + b_{a,t,2l+2}(y)x^{2l+2} + \cdots).$$

We next assume that H_{a,λ_t} has the form

$$(27) \quad \lambda_t(x + x^{l+1} + b_{a,t,k}x^k + \cdots + b_{a,t,2l+1}x^{2l+1} + \cdots),$$

in the first coordinate. By induction, we reduce $b_{a,t,k}$ to 0 for each k with $l+2 \leq k \leq 2l$. By (26), (27) has the form

$$\lambda_t(X + X^{l+1}) + (\lambda_t b_{a,t,k} - (\lambda_t^k - \lambda_t)\lambda_t b_{a,t,k} / (\lambda_t^k - \lambda_t))X^k + \cdots = \lambda_t(X + X^{l+1}) + O(X^{k+1}),$$

in the first coordinate. Therefore, we may assume that H_{a,λ_t} has the form

$$\lambda_t(x + x^{l+1} + C_{a,t}x^{2l+1} + b_{a,t,2l+2}(y)x^{2l+2} + O(x^{2l+3})),$$

in the first coordinate. Hence we have Theorem 4.3. Note that, by repeating Step 3 and Step 4, H_{a,λ_t} has the form

$$(28) \quad (\lambda_t(x + x^{l+1} + C_{a,t}x^{2l+1} + b_{a,t,2l+2}(y)x^{2l+2} + \cdots), v_{a,t}y + xh_{a,t}(x, y)).$$

We obtain the following lemma by the same computation as in the proof of Theorem 3.5 in [16].

Lemma 4.4. *There is $\delta_0 > 0$ such that by a coordinate transformation $\phi_{a,t}$, $\phi_{a,t} \circ H_{a,\lambda_t} \circ \phi_{a,t}^{-1}$ has the form (21) for each $H_{a,\lambda_t} \in RD_{\lambda_t, \delta_0}$.*

We now prove the second main result of this paper.

Theorem 4.5. *There is $\delta_0 > 0$ such that each $H_{a,\lambda_t} \in RD_{\lambda_t, \delta_0}$ with $a \neq 0$ has connected Julia set J_{a,λ_t} .*

Proof. It suffices to show the statement of Theorem 4.5 for $H_{a,\lambda_t^{-1}}$ instead of H_{a,λ_t} . Moreover, by Theorem 3.10 and Lemma 4.2, it suffices to show that there are $\varepsilon > 0$ and $\delta_0 > 0$ such that $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda_t^{-1}}^+ = \emptyset$ for $H_{a,\lambda_t^{-1}} \in RD_{\lambda_t^{-1}, \delta_0}$, and $J_{p_{\lambda_t^{-1}}}$ is connected for $0 < t < \delta_0$.

We first consider the case $\lambda_0 = 1$. By a transformation $\phi_{a,t}$ (see Lemma 4.4), $\phi_{a,t} \circ H_{a,\lambda_t^{-1}} \circ \phi_{a,t}^{-1}$ is of the form

$$(29) \quad (x_1, y_1) = (\lambda_t^{-1}(x + x^2 + C_{a,t}x^3 + O_y(x^4)), v_{a,t}y + xh_{a,t}(x, y)),$$

in $\mathbb{D}_\rho \times \mathbb{D}_{r+O(|a|)}$, where the notation $O_y(x^\alpha)$ represents a holomorphic map of (x, y) which is bounded by $K|x|^\alpha$ for some K . By the transformation $\psi_t(x, y) = (-1/(\lambda_t x), y)$, the map $\psi_t \circ \phi_{a,t} \circ H_{a,\lambda_t^{-1}} \circ \phi_{a,t}^{-1} \circ \psi_t^{-1}(X, Y) = (X_1, Y_1)$ is of the form

(30)

$$(\lambda_t X + 1 + g_{a,t}(X, Y), \nu_{a,t} Y + f_{a,t}(X, Y)) = \left(\lambda_t X + 1 + D_{a,t}/X + O_Y(1/|X|^2), \nu_{a,t} Y + O_Y(1/|X|) \right).$$

We take a constant $M > 0$ such that

$$(31) \quad |g_{a,t}(X, Y)| \leq \frac{M}{|X|} \text{ and } |f_{a,t}(X, Y)| \leq \frac{M}{|X|},$$

for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$. Recall that $\phi_{a,t}$ is a transformation from $\mathbb{D}_{\rho'}(\Pr_1 \mathbf{q}_{0,\lambda_t}) \times \mathbb{D}_r$ to $\mathbb{D}_\rho \times \mathbb{D}_{r+O(|a|)}$. Since $\Pr_2 \phi_{a,t}(x, y) = y + O(a)$ (see the sketch of the proof of Lemma 4.3), we may assume that the forms (30) are defined in $\{|X| > 1/\rho\} \times \mathbb{D}_r$ and the inequalities (31) hold for $(X, Y) \in \{|X| > 1/\rho\} \times \mathbb{D}_r$. Note that $\psi_t \circ \phi_{a,t}(\mathbf{q}_{a,\lambda_t^{-1}}) = (\infty, 0)$. We set $(X_0, Y_0) := \psi_0 \circ \phi_{0,0} \circ H_{0,\lambda_0^{-1}}^N(0, 0)$ for some large $N \in \mathbb{N}$. We have $\operatorname{Re} X_0 > 1/\rho$, by taking a larger N if necessary, since the forward orbit of critical point 0 under $p_{\lambda_0^{-1}}$ converges to its parabolic fixed point (see [1], p.120). Let $\gamma > 0$ be a number such that $\mathbb{D}_\gamma(X_0) \times \mathbb{D}_r \subset \{X \in \mathbb{C} : \operatorname{Re} X > 1/\rho\} \times \mathbb{D}_r$. We set $E := \mathbb{D}_\gamma(X_0) \times \mathbb{D}_r$. We consider the affine transformations

$$Q_t(z, w) = \left(\frac{z - b_t}{X_0 - b_t}, w \right) \text{ and } Q_t^{-1}(z, w) = ((X_0 - b_t)z + b_t, w), \text{ where } b_t := \frac{1}{1 - \lambda_t}.$$

We set

$$F_{a,t}(z, w) = Q_t \circ \psi_t \circ \phi_{a,t} \circ H_{a,\lambda_t^{-1}} \circ \phi_{a,t}^{-1} \circ \psi_t^{-1} \circ Q_t^{-1}(z, w).$$

Then, we have

$$\Pr_1 F_{a,t}(z, w) = \Pr_1 Q_t \circ \psi_t \circ \phi_{a,t} \circ H_{a,\lambda_t^{-1}} \circ \phi_{a,t}^{-1} \circ \psi_t^{-1} \circ Q_t^{-1}(z, w) = \lambda_t z + \frac{g_{a,t}((X_0 - b_t)z + b_t, w)}{X_0 - b_t}.$$

We set

$$G_{a,t}(z, w) := \frac{g_{a,t}((X_0 - b_t)z + b_t, w)}{X_0 - b_t}.$$

Further we set $E' (= E'_t) := Q_t(E)$ and $U' (= U'_t) := Q_t(U)$, where $U = \{|z| < 1/\rho\} \times \mathbb{D}_r$. Then we have

$$E' = \left\{ z \in \mathbb{C} : |z - 1| < \frac{\gamma}{|b_t - X_0|} \right\} \times \mathbb{D}_r \text{ and } U' = \left\{ z \in \mathbb{C} : \left| z - \frac{b_t}{b_t - X_0} \right| < \frac{1}{\rho|b_t - X_0|} \right\} \times \mathbb{D}_r.$$

Clearly, if ρ and δ_0 are sufficiently small, then for each $a \in \mathbb{D}_{\delta_0}$ and each $t \in (0, \delta_0)$, we have

$$(32) \quad \Pr_2 F_{a,t}((\mathbb{C} \times \mathbb{D}_r) \setminus U') \subset \mathbb{D}_r.$$

Indeed, $|\Pr_2 \psi_t \circ \phi_{a,t} \circ H_{a,\lambda_t^{-1}} \circ \phi_{a,t}^{-1} \circ \psi_t^{-1}(X, Y)| \leq |\nu_{a,t} Y| + M/|X| < |\nu_{a,t}|r + M\rho$ for $(X, Y) \in \{|X| > 1/\rho\} \times \mathbb{D}_r$ (see (30) and (31)). Recall that $\nu_{a,t} = -a^2/\lambda_t^{-1}$ and $|\lambda_0| = 1$. By taking ρ and δ_0 with sufficiently small, we have $|\nu_{a,t}|r + M\rho < r$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$. Since $\Pr_2 Q_t(z, w) = w$, we have (32).

By $X_0 = \Pr_1 \psi_0 \circ \phi_{0,0} \circ H_{0,\lambda_0^{-1}}^N(0, 0)$, $E = \mathbb{D}_\gamma(X_0) \times \mathbb{D}_r$ and (32), we have $Q_t \circ \psi_t \circ \phi_{a,t} \circ H_{a,\lambda_t^{-1}}^N(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \subset E'$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$, by taking $\delta_0 > 0$ and $\varepsilon > 0$ sufficiently small if necessary. Thus, to show that there are $\varepsilon > 0$ and $\delta_0 > 0$ such that $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda_t^{-1}}^+ = \emptyset$ for $H_{a,\lambda_t^{-1}} \in RD_{\lambda_t^{-1}, \delta_0}$, it suffices to show that there is $\delta_0 > 0$ such that

$$(33) \quad \bigcup_{k \in \mathbb{N}} F_{a,t}^k(E') \cap U' = \bigcup_{k \in \mathbb{N}} F_{a,t}^k(E'_t) \cap U'_t = \emptyset,$$

for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$, taking smaller $\gamma > 0$, $\rho > 0$ and a larger $N \in \mathbb{N}$ if necessary. Indeed, if (33) holds for $F_{a,t}$, then $\bigcup_{n \geq 0} H_{a,\lambda_t^{-1}}^n(\mathbb{D}_\varepsilon \times \mathbb{D}_r)$ is a bounded set of \mathbb{C}^2 , which implies that $\mathbb{D}_\varepsilon \times \mathbb{D}_r \subset \text{Int } K_{a,\lambda_t^{-1}}$. To obtain (33), we show the following Claims 1,2,3.

Claim 1. For sufficiently small δ_0 , the following hold. If $L_t > 0$ for $0 < t < \delta_0$, then $U' \subset \mathbb{D} \times \mathbb{D}_r$ for $0 < t < \delta_0$. If $L_t < 0$ for $0 < t < \delta_0$, then $U' \cap (\mathbb{D} \times \mathbb{D}_r) = \emptyset$ for $0 < t < \delta_0$.




Fig. 4. $L_t > 0$ and $\theta_t > 0$ (left), $L_t < 0$ and $\theta_t > 0$ (right)

We prove Claim 1. Assume that $L_t > 0$ for each $0 < t < \delta_0$. We show that $\frac{|b_t|}{|b_t - X_0|} + \frac{1}{\rho|b_t - X_0|} < 1$ for sufficiently small t to obtain the conclusion. The inequality is equivalent to

$$(34) \quad |1 - X_0(1 - \lambda_t)| - |1 - \lambda_t|/\rho - 1 > 0.$$

Let $x_1 := \text{Re}(\lambda_t - 1)$ and $y_1 := \text{Im}(\lambda_t - 1)$. Then, $x_1 = L_t + O(L_t^2)$ and $y_1 = \theta_t + O(\theta_t L_t)$. First, we have

$$\begin{aligned} |1 + X_0(\lambda_t - 1)| &= \sqrt{(1 + \text{Re}(X_0)x_1 - \text{Im}(X_0)y_1)^2 + (\text{Re}(X_0)y_1 + \text{Im}(X_0)x_1)^2} \\ &= \sqrt{1 + 2\text{Re}(X_0)x_1 - 2\text{Im}(X_0)y_1 + O((x_1 + y_1)^2)} \\ &= \sqrt{1 + 2\text{Re}(X_0)x_1 - 2\text{Im}(X_0)y_1 + O(L_t^2)} \\ &= 1 + \text{Re}(X_0)x_1 - \text{Im}(X_0)y_1 + O(L_t^2). \end{aligned}$$

We now show the inequality (34). By the above computation, we have

$$|1 - X_0(1 - \lambda_t)| - |1 - \lambda_t|/\rho - 1 = \text{Re}(X_0)x_1 - \text{Im}(X_0)y_1 + O(L_t^2) - |1 - \lambda_t|/\rho.$$

Recall that $X_0 = \text{Pr}_1 \psi_0 \circ \phi_{0,0} \circ H_{0,\lambda_0^{-1}}^N(0,0)$. Since $\text{Re } X_0 > 0$, $0 < x_1 \asymp L_t$, $y_1 = O(\theta_t)$ and $|1 - \lambda_t| = O(L_t)$, we have the assertion, taking a large $N \in \mathbb{N}$ so that $|\text{Im } X_0|/\text{Re } X_0$ and $1/(\rho \text{Re } X_0)$ are sufficiently small (see [1], p.120), where $x_1 \asymp L_t$ means that there is $K > 0$ such that $L_t/K < x_1 < KL_t$ for sufficiently small t .

Assume that $L_t < 0$ for $0 < t < \delta_0$. We claim that $\frac{|b_t|}{|b_t - X_0|} - \frac{1}{\rho|b_t - X_0|} > 1$ for sufficiently small t . The inequality is equivalent to $1 - |1 - \lambda_t|/\rho - |1 - X_0(1 - \lambda_t)| > 0$. We have

$$1 - |1 - \lambda_t|/\rho - |1 - X_0(1 - \lambda_t)| = -|1 - \lambda_t|/\rho - \text{Re}(X_0)x_1 + \text{Im}(X_0)y_1 + O(L_t^2).$$

By $\operatorname{Re} X_0 > 0$ and $x_1 < 0$, we have the assertion, taking a larger $N \in \mathbb{N}$ so that $|\operatorname{Im} X_0|/\operatorname{Re} X_0$ and $1/(\rho \operatorname{Re} X_0)$ are sufficiently small if necessary (see [1], p.120). Thus, we have proved Claim 1.

Claim 2. For sufficiently small δ_0 , the following hold. If $L_t > 0$ for $0 < t < \delta_0$, then $|\Pr_1 F_{a,t}(z, w)| > e^{L_t/2}|z| > |z|$ for $a \in \mathbb{D}_{\delta_0}$, $0 < t < \delta_0$ and $(z, w) \in (\{|z| > 1/2\} \times \mathbb{D}_r) \setminus U'$. If $L_t < 0$ for $0 < t < \delta_0$, then $|\Pr_1 F_{a,t}(z, w)| < e^{L_t/2}|z| < |z|$ for $a \in \mathbb{D}_{\delta_0}$, $0 < t < \delta_0$ and $(z, w) \in (\{|z| > 1/2\} \times \mathbb{D}_r) \setminus U'$.

We prove Claim 2. Assume that $(z, w) \in (\{|z| > 1/2\} \times \mathbb{D}_r) \setminus U'$. By using the inequality (31), we have

$$\frac{|G_{a,t}(z, w)|}{|\lambda_t z| - e^{L_t/2}|z|} < \frac{2|G_{a,t}(z, w)|}{|e^{L_t} - e^{L_t/2}|} \leq \frac{2M\rho}{|e^{L_t} - e^{L_t/2}| |X_0 - b_t|} = \frac{2M\rho |1 - \lambda_t|}{e^{L_t/2} |e^{L_t/2} - 1| |X_0(1 - \lambda_t) - 1|}.$$

Since $|1 - \lambda_t| = O(L_t)$, there exists a positive constant δ_0 such that the ratio is less than $1/2$ if $0 < t < \delta_0$, taking $\rho > 0$ sufficiently small if necessary. We note that the constant δ_0 does not depend on z . By the inequality

$$(35) \quad |\lambda_t z| - e^{L_t/2}|z| - |G_{a,t}(z)| \leq |\Pr_1 F_{a,t}(z, w)| - e^{L_t/2}|z| \leq |\lambda_t z| - e^{L_t/2}|z| + |G_{a,t}(z, w)|,$$

the statement of Claim 2 holds.

Claim 3. For sufficiently small δ_0 , the following hold. If $L_t > 0$ for $0 < t < \delta_0$, then $F_{a,t}(E') \cap (\mathbb{D} \times \mathbb{D}_r) = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$. If $L_t < 0$ for $0 < t < \delta_0$, then $F_{a,t}(E') \subset \mathbb{D} \times \mathbb{D}_r$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$.

We prove Claim 3. Let $(z, w) \in E'$. Then, we have $z = 1 + z_0$, where $|z_0| < \gamma/|b_t - X_0|$. Since

$$|\Pr_1 F_{a,t}(z, w)| = |\lambda_t(1 + z_0) + G_{a,t}(1 + z_0, w)| = |\lambda_t + \lambda_t z_0 + G_{a,t}(1 + z_0, w)|,$$

we have

$$e^{L_t} - |\lambda_t z_0 + G_{a,t}(1 + z_0, w)| \leq |\Pr_1 F_{a,t}(z, w)| \leq e^{L_t} + |\lambda_t z_0 + G_{a,t}(1 + z_0, w)|.$$

Since $|z_0| < \gamma/|b_t - X_0| = O(L_t)$ and $|G_{a,t}(1 + z_0, w)| < M\rho/|X_0 - b_t| = O(L_t)$, we have the assertion, taking smaller γ, ρ if necessary. Hence, we have proved Claim 3.

We now show that there is $\delta_0 > 0$ such that $F_{a,t}^k(E') \cap U' = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$, $0 < t < \delta_0$ and each $k \in \mathbb{N}$ by using Claims 1, 2, and 3. First, assume that $L_t > 0$ for $0 < t < \delta_0$. By Claim 1, $U' \subset \mathbb{D} \times \mathbb{D}_r$. By Claim 3, $F_{a,t}(E') \cap (\mathbb{D} \times \mathbb{D}_r) = \emptyset$, and so $F_{a,t}(E') \cap U' = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$. By $U' \subset \mathbb{D} \times \mathbb{D}_r$ and Claim 2, we have $|\Pr_1 F_{a,t}(z, w)| > |z| > 1$ for $(z, w) \in (\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r$, and so $F_{a,t}((\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r) \subset (\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r$. By using $F_{a,t}(E') \subset (\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r$, for $k \geq 2$, we have

$$F_{a,t}^k(E') = F_{a,t}^{k-1}(F_{a,t}(E')) \subset F_{a,t}^{k-1}((\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r) \subset (\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r.$$

Hence, we have $F_{a,t}^k(E') \cap U' = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$, $0 < t < \delta_0$ each $k \in \mathbb{N}$ by $U' \subset \mathbb{D} \times \mathbb{D}_r$.

Assume that $L_t < 0$ for $0 < t < \delta_0$. Similarly, we have $F_{a,t}(E') \cap U' = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$. We note that $U' \subset (\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r$ by Claim 1. Therefore, $F_{a,t}$ is defined in $\mathbb{D} \times \mathbb{D}_r$. It is easy to see that

$$(36) \quad F_{a,t}(\overline{\mathbb{D}_{1/2}} \times \mathbb{D}_r) \subset \mathbb{D}_{2/3} \times \mathbb{D}_r,$$

for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$, taking a smaller δ_0 if necessary. Indeed, if $(z, w) \in \overline{\mathbb{D}_{1/2}} \times \mathbb{D}_r$, then $|\text{Pr}_1 F_{a,t}(z, w)| \leq |\lambda_t z| + |G_{a,t}(z, w)| \leq |\lambda_t|/2 + M\rho/(|X_0 - b_t|) < 2/3$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$, taking a smaller δ_0 if necessary. Thus we have (36). By Claim 2, $U' \subset (\mathbb{C} \setminus \mathbb{D}) \times \mathbb{D}_r$ and (36), we have $|\text{Pr}_1 F_{a,t}(z, w)| < |z| < 1$ or $|\text{Pr}_1 F_{a,t}(z, w)| < 2/3$ for $(z, w) \in \mathbb{D} \times \mathbb{D}_r$, and so $F_{a,t}(\mathbb{D} \times \mathbb{D}_r) \subset \mathbb{D} \times \mathbb{D}_r$. For $k \geq 2$, we have

$$F_{a,t}^k(E') = F_{a,t}^{k-1}(F_{a,t}(E')) \subset F_{a,t}^{k-1}(\mathbb{D} \times \mathbb{D}_r) \subset \mathbb{D} \times \mathbb{D}_r.$$

Hence, we have $F_{a,t}^k(E') \cap U' = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$, $0 < t < \delta_0$ and each $k \in \mathbb{N}$ by $U' \cap (\mathbb{D} \times \mathbb{D}_r) = \emptyset$. Hence we obtain (33). This implies that there are $\delta_0 > 0$ and $\varepsilon > 0$ such that $(\mathbb{D}_\varepsilon \times \mathbb{D}_r) \cap J_{a,\lambda_t^{-1}} = \emptyset$ for $a \in \mathbb{D}_{\delta_0}$ and $0 < t < \delta_0$. For the constant $\varepsilon > 0$, $\{H_{a,\lambda_t}\}_{a \in \mathbb{D}_{\delta_0}}$ satisfies the condition $(\text{VC})_{\varepsilon,r}$ for $0 < t < \delta_0$ by Lemma 4.2, taking a smaller δ_0 if necessary (see Remark 3.4).

In order to apply Theorem 3.10 to $\{H_{a,\lambda_t^{-1}}\}_{a \in \mathbb{D}_{\delta_0}}$ for $0 < t < \delta_0$, we show that $J_{p_{\lambda_t^{-1}}} \subset \mathbb{C}$ is connected for $0 < t < \delta_0$. It suffices to show that $p_{\lambda_t^{-1}}$ has an attracting fixed point. Since $p_{\lambda_t^{-1}}$ has a fixed point with multiplier $\lambda_t^{-1} = \exp(-L_t - i\theta_t)$, if $L_t > 0$ for $0 < t < \delta_0$, then $p_{\lambda_t^{-1}}$ has an attracting fixed point. If $L_t < 0$ for $0 < t < \delta_0$, then $F_{0,t}(\overline{\mathbb{D} \times \mathbb{D}_r}) \subset \mathbb{D} \times \mathbb{D}_r$ by Claim 2 and (36). Thus $p_{\lambda_t^{-1}}$ has an attracting fixed point.

In the case of $l \geq 2$, by the transformation $X = -1/(lx^l)$, $Y = y$ and a linear transformation, the form of (28) with λ_t replaced by λ_t^{-1} is conjugate to

$$(X_1, Y_1) = \lambda_t^l X + 1 + D_{a,t}/X + O_Y(1/X^2), v_n Y + O_Y(1/|X|^{1/l}).$$

Similar to the case of $l = 1$, we can show the statement in Theorem 4.5 in the case of $l \geq 2$. \square

ACKNOWLEDGEMENTS. The author is grateful to Prof. Tomoki Kawahira, who offered continuing support and constant encouragement. Further, the author thanks Prof. Shunsuke Morosawa and Prof. Hiroki Sumi for helping the author. The author is partially supported by JSPS Grant-in-Aid for JSPS Fellows Grant Number 23KJ1179.

References

- [1] A. Beardon: Iteration of Rational Functions, Complex analytic dynamical systems, Grad. Texts Math. **132**, Springer-Verlag, New York, 1991.
- [2] E. Bedford and J. Smillie: *Polynomial diffeomorphisms of \mathbb{C}^2 : currents, equilibrium measure and hyperbolicity*, Invent. Math. **103** (1991), 69–99.
- [3] E. Bedford and J. Smillie: *Polynomial diffeomorphisms of \mathbb{C}^2 . II: Stable manifolds and recurrence*, J. Amer. Math. Soc. **4** (1991), 657–679.
- [4] E. Bedford and J. Smillie: *Polynomial diffeomorphisms of \mathbb{C}^2 : VI. Connectivity of J* , Ann. of Math. **148** (1998), 695–735.
- [5] A. Douady and J.H. Hubbard: Étude dynamique des polynômes complexes. Partie I (French), Publications Mathématiques d’Orsay, vol. 84, Université de Paris-Sud, Département de Mathématiques, Orsay, 1984.

- [6] J.-E. Fornæss and N. Sibony: *Complex Hénon mappings in \mathbb{C}^2 and Fatou-Bieberbach domains*, Duke Math. J. **65** (1992), 345–380.
- [7] J.H. Hubbard: *Parametrizing unstable and very unstable manifolds*, Mosc. Math. J. **5** (2005), 105–124.
- [8] J.H. Hubbard and R.W. Oberste-Vorth: *Hénon mappings in the complex domain I: The global topology of dynamical space*, Inst. Hautes Études Sci. Publ. Math. No. **79** (1994), 5–46.
- [9] J.H. Hubbard and R.W. Oberste-Vorth: *Hénon mappings in the complex domain II: Projective and inductive limits of polynomials*; in *Real and Complex Dynamical Systems* (Hillerød, 1993), Kluwer Academic Publishers, Dordrecht, 1995, 89–132.
- [10] Y. Ishii: *Dynamics of polynomial diffeomorphisms of \mathbb{C}^2 : Combinatorial and topological aspects*, Arnold Math. J. **3** (2017), 119–173.
- [11] R. Mañé, P. Sad and D. Sullivan: *On the dynamics of rational maps*, Ann. Sci., École. Norm. Sup. **16** (1983), 193–217.
- [12] C. McMullen: *Hausdorff dimension and conformal dynamics. II. Geometrically finite rational maps*, Comment. Math. Helv. **75** (2000), 535–593.
- [13] J. Milnor: *Dynamics in one complex variable*, 3rd ed., Ann. of Math. Stud., vol. 160, Princeton University Press, Princeton, NJ, 2006.
- [14] S. Morosawa, Y. Nishimura, M. Taniguchi and T. Ueda: *Holomorphic dynamics*, Translated from the 1995 Japanese original and revised by the authors, Cambridge Studies in Advanced Mathematics, 66, Cambridge University Press, Cambridge, 2000.
- [15] R. Radu: *Topological Models for Hyperbolic and Semi-parabolic Complex Hénon Maps*, Ph.D. Thesis, Cornell University, 2013.
- [16] R. Radu and R. Tanase: *Semi-parabolic tools for hyperbolic Hénon maps and continuity of Julia sets in \mathbb{C}^2* , Trans. Amer. Math. Soc. **370** (2018), 3949–3996.
- [17] R. Radu and R. Tanase: *A structure theorem for semi-parabolic Hénon maps*, Adv. Math. **350** (2019), 1000–1058.
- [18] M. Shub: *Global Stability of Dynamical Systems*, Springer-Verlag, New York, 1987.
- [19] Z. Slodkowski: *Holomorphic motions and polynomial hulls*, Proc. Amer. Math. Soc. **111** (1991), 347–355.

Course of Mathematical Science, Department of Human Coexistence
 Graduate School of Human and Environmental Studies
 Kyoto University
 Yoshida Nihonmatsu-cho, Sakyo-ku
 Kyoto, 606–8501
 Japan
 e-mail: t.yagi125a@gmail.com