



Title	On the intermediate cohomology group of a holomorphic line bundle over a complex torus
Author(s)	Matsushima, Yozô
Citation	Osaka Journal of Mathematics. 1979, 16(3), p. 617-631
Version Type	VoR
URL	https://doi.org/10.18910/10327
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

ON THE INTERMEDIATE COHOMOLOGY GROUP OF A HOLOMORPHIC LINE BUNDLE OVER A COMPLEX TORUS

Yozo MATSUSHIMA

(Received August 7, 1978)

Let $E=V/L$ be a complex torus, where V is an n -dimensional complex vector space and L a lattice of V . Let H be a Hermitian form on V and A the imaginary part of H . Then A is an \mathbf{R} -bilinear alternating form on V . We say that H is a *Riemann form of signature* (s, r) for the torus E if

- (a) H is non-degenerate and of signature (s, r) ;
- (b) A is integral valued on the lattice L .

To a Riemann form H we associate a factor

$$(1) \quad J_{H,\psi}(g, z) = \psi(g) \varepsilon \left[\frac{1}{2i} H(z, g) + \frac{1}{4i} H(g, g) \right]$$

with $g \in L$, $z \in V$, where $\varepsilon[\cdot] = \exp 2\pi i \cdot$ and ψ is a map from L to $\mathbf{C}_1^* = \{z \in \mathbf{C} \mid |z|=1\}$ satisfying $\psi(g+h) = \psi(g)\psi(h) \varepsilon \left[\frac{1}{2} A(g, h) \right]$; the function ψ being called a semi-character of L for A .

The factor $J_{H,\psi}: L \times V \rightarrow \mathbf{C}^*$ satisfies the equation

$$J_{H,\psi}(g+f, z) = J_{H,\psi}(g, h+z) J_{H,\psi}(h, z),$$

where $g, h \in L$, $z \in V$.

Given the factor $J_{H,\psi}$ we define an action of the lattice group L on $V \times \mathbf{C}$ by the rule

$$(z, \xi) \cdot g = (z+g, J_{H,\psi}(g, z)\xi),$$

where $z \in V$, $\xi \in \mathbf{C}$ and $g \in L$. The action of L on $V \times \mathbf{C}$ is free and the quotient of $V \times \mathbf{C}$ by L has a natural structure of a holomorphic line bundle over the complex torus $E=V/L$. We shall denote this line bundle by $F(H, \psi)$.

The following vanishing theorem for the cohomology of $F(H, \psi)$ is well-known [2, 4]: If H is a Riemann form of signature (s, r) , then we have

$$H^q(E, F(H, \psi)) = 0$$

for $q \neq r$ (for a proof see Appendix 2 of this paper).

In particular if $r=0$, namely if H is positive definite, $H^q(E, F(H, \psi))=0$ except for $q=0$ and $H^0(E, F(H, \psi))$ is identified with the space of all holomorphic theta functions on V for the factor $J_{H, \psi}$. Replacing H by a suitable positive interger multiple of H , if necessary, these theta functions define a holomorphic imbedding of E into a complex projective space. A complex torus which admits a positive definite Riemann form is called an abelian variety. There exist complex tori which are not abelian varieties but which admit Riemann forms of signature (s, r) with $s>0$ and $r>0$ (see Appendix 1). For such a complex torus E , there exists no non-trivial theta function. However, there exists the non-trivial intermediate cohomology group $H^r(E, F(H, \psi))$ with $0<r<n$.

The purpose of this paper is to give an interpretation of the intermediate cohomology group $H^r(E, F(H, \psi))$. Namely we associate to a Riemann form H of signature (s, r) a family of polarized abelian varieties (E_b, H_b) parametrized by elements b of the Hermitian symmetric space $\mathcal{B}=U(H)/K$, where $U(H)$ is the unitary group of the Hermitian form H and K is a maximal compact subgroup of $U(H)$. Here E_b is of the form $E_b=V_b/L$, where V_b is an n -dimensional complex vector space with the same underlying real vector space as V and with a complex structure J_b distinct from that of V and parametrized by $b \in \mathcal{B}$ and H_b is a positive definite Riemann form for E_b whose imaginary part is equal to A . We then assign a family of line bundles $F(H_b, \psi)$ over E_b for each b . Finally we shall show that there exists a canonically defined isomorphism from $H^r(E, F(H, \psi))$ to $H^r(E_b, F(H_b, \psi))$ for each b . We also see that there exists a family Φ_b of differentiable imbedding of E into a complex projective space which is partially holomorphic and partially antiholomorphic.

It should be mentioned that C.L. Siegel [3] has associated to an indefinite quadratic form a family of theta series parametrized by a symmetric space. It is possible to interpret Siegel's family of theta series as a subfamily of the family $\{H^0(E_b, F(H_b, \psi))\}$ of theta functions attached to a certain complex torus E and a Riemann form H related with the given indefinite quadratic form.

1. A Riemann form of signature (s, r) and a family of polarized abelian varieties

Let $E=V/L$ be a complex torus, where V is an n -dimensional complex vector space and L a lattice of V . We shall denote by W the underlying $2n$ -dimensional real vector space of V and by J the complex structure of W defining the complex vector space V .

Let H be a non-degenerate Hermitian form on V of signature (s, r) , where $s+r=n$. We denote by A the imaginary part of H . Then we have

$$H(u, v) = A(Ju, v) + iA(u, v), \quad u, v \in W.$$

We assume that the alternating \mathbf{R} -bilinear form A to be integral valued on $L \times L$ and we call H a Riemann form of signature (s, r) for the complex tours E .

We shall denote by $U(H)$ the unitary group of the Hermitian form H . A basis $B = \{v_1, \dots, v_n\}$ of V is said to be a *privileged basis* for H if the matrix of H with respect to the basis B is of the form

$$1_{s,r} = \begin{pmatrix} 1_s & 0 \\ 0 & -1_r \end{pmatrix}$$

where 1_s and 1_r denote the unit matrix of size s and r respectively.

The group $U(H)$ acts simply transitively on the set of all privileged bases for H . We denote by $V_1(B)$ and $V_2(B)$ the subspaces of V spanned by $\{v_1, \dots, v_s\}$ and $\{v_{s+1}, \dots, v_n\}$ respectively. Then we have

$$(2) \quad W = V_1(B) \oplus V_2(B).$$

We say that two privileged bases B and B' are equivalent, $B \sim B'$, if $V_i(B) = V_i(B')$ for $i=1, 2$. We shall denote by \mathcal{B} the set of equivalence classes of privileged bases for H . Then the group $U(H)$ acts transitively on \mathcal{B} and \mathcal{B} is identified with the Hermitian symmetric space $U(H)/K$, where K is a maximal compact subgroup of $U(H)$.

Let $b \in \mathcal{B}$ and let B be a privileged basis representing b . We define a linear transformation J_b of W by requiring

$$J_b = J \quad \text{on} \quad V_1(B)$$

and

$$J_b = -J \quad \text{on} \quad V_2(B).$$

We have $J_b^2 = -1$ and hence J_b defines a complex structure on W . We shall denote by V_b the complex vector space defined by W and J_b .

Define the symbol $\varepsilon_k (k=1, 2, \dots, n)$ by

$$\varepsilon_k = \begin{cases} 1, & k \in [1, s], \\ -1, & k \in [s+1, n]. \end{cases}$$

If $B = \{v_1, \dots, v_n\}$ is a privileged basis representing b , then we have

$$J_b v_k = \varepsilon_k J v_k$$

We also have $H(v_k, v_j) = \varepsilon_k \cdot \delta_{kj}$ and since $H(v_k, v_j) = A(Jv_k, v_j) + iA(v_k, v_j)$, we get $A(v_k, v_j) = 0$ and $A(Jv_k, v_j) = \varepsilon_k \cdot \delta_{kj}$. It follows from these that the decomposition (2) is orthogonal for A and also for H . We have also $A(J_b u, J_b v) = A(u, v)$ for $u, v \in W$. For let $u = u_1 + u_2$, $v = v_1 + v_2$ with $u_1, v_1 \in V_1(B)$ and $u_2, v_2 \in V_2(B)$. Then $J_b u = Ju_1 - Ju_2$ and $J_b v = Jv_1 - Jv_2$ and $Ju_1, Jv_1 \in V_1(B)$ and $Ju_2, Jv_2 \in V_2(B)$. Hence $A(J_b u, J_b v) = A(Ju_1, Jv_1) + A(Ju_2, Jv_2) = A(u_1, v_1) + A(u_2, v_2)$. We can then

define a Hermitian form H_b on the complex vector space V_b by

$$H_b(u, v) = A(J_b u, v) + iA(u, v).$$

Then the imaginary part of H_b is A and we have $H_b(v_k, v_j) = A(J_b v_k, v_j) + iA(v_k, v_j) = \varepsilon_k A(J v_k, v_j) = \varepsilon_k^2 \delta_{kj} = \delta_{kj}$. This means that B is an orthonormal basis of V_b for the Hermitian form H_b and in particular H_b is positive definite and the decomposition (2) of W is also orthogonal for H_b .

Let now

$$E_b = V_b/L.$$

Then H_b is a positive definite Riemann form for E_b and hence E_b is an abelian variety. Thus we have associated to a complex torus E and a Riemann form H of signature (s, r) a family of polarized abelian varieties (E_b, H_b) parametrized by $b \in \mathcal{B} = U(H)/K$.

We need the following lemma in the next section.

Lemma. *We have*

$$H(u, v) = H_b(u, v) \quad \text{for } u \in V_1(B)$$

and

$$H(u, v) = -H_b(v, u) \quad \text{for } u \in V_2(B).$$

For we have $H_b(u, v) = A(J_b u, v) + iA(u, v)$ and $J_b u = Ju$ or $J_b u = -Ju$ according as $u \in V_1(B)$ or $u \in V_2(B)$.

2. The cohomology group $H^q(E, F(H, \psi))$

We associate to the Riemann form H of signature (s, r) for E the factor $J_{H, \psi}$ defined by (1) and the line bundle $F(H, \psi)$ over E . For the cohomology groups of $F(H, \psi)$ we have the following theorem.

Theorem 1. (i) *We have $H^q(E, F(H, \psi)) = 0$ for $q \neq r$.*

(ii) *Let (z_1, \dots, z_s) be the coordinates of the complex vector space V determined by a privileged basis B of V for the Hermitian form H . Then $H^r(E, F(H, \psi))$ is identified with the complex vector space of all C^∞ functions f on V satisfying the following conditions:*

1) *f is a differentiable theta function for the factor $J_{H, \psi}$; namely we have*

$$f(z+g) = J_{H, \psi}(z, g) \cdot f(z), \quad z \in V, \quad g \in L.$$

2) $\frac{\partial f}{\partial \bar{z}_k} = 0$ for $k \in [1, s]$

and

$$\frac{\partial f}{\partial z_{s+j}} + \pi \bar{z}_{s+j} \cdot f = 0 \quad \text{for } j \in [1, r].$$

The assertion (i) in Theorem 1 is a well-known vanishing theorem due to Mumford. We shall give a proof of Theorem 1 in the Appendix 2 based on the harmonic theory.

We denote by $H(B)$ the space of C^∞ functions f on W satisfying the above conditions (1) and (2) to make explicit its dependence of the condition (2) on the choice of the privileged basis B . We show that if B and B' are equivalent, then we have $H(B)=H(B')$. In fact, let (z'_1, \dots, z'_n) be the coordinates of V determined by B' . Then we have

$$z'_i = \sum_{j=1}^s a_{ij} z_j \quad (i=1, \dots, s)$$

and

$$z'_{s+i} = \sum_{j=1}^r b_{ij} z_{s+j} \quad (i=1, \dots, r)$$

where the matrices (a_{ij}) and (b_{ij}) are both unitary. We get

$$(*) \quad \frac{\partial f}{\partial \bar{z}_k} = \sum_{i=1}^s a_{ik} \frac{\partial f}{\partial \bar{z}'_i} \quad (k=1, \dots, s)$$

and

$$\frac{\partial f}{\partial z_{s+k}} + \pi \bar{z}_{s+k} f = \sum_{i=1}^r b_{ik} \frac{\partial f}{\partial z'_{s+i}} + \pi \left(\sum_{i=1}^r \bar{b}'_{ki} \bar{z}'_{s+i} \right) f,$$

where (b'_{ki}) is the inverse matrix of (b_{ki}) . Since (b_{ki}) is unitary, we have $(b'_{ki}) = {}^t(\bar{b}_{ki})$ and hence $\bar{b}'_{ki} = b_{ik}$. Hence we get

$$(**) \quad \frac{\partial f}{\partial z_{s+k}} + \pi \bar{z}_{s+k} f = \sum_{i=1}^r b_{ik} \left(\frac{\partial f}{\partial z'_{s+i}} + \pi \bar{z}'_{s+i} f \right)$$

From (*) and (**) we get $H(B)=H(B')$. Hence we can denote the space of C^∞ functions f satisfying (1) and (2) by $H(b)$, $b \in \mathcal{B}$.

Consider now the family of polarized abelian varieties (E_b, H_b) ($b \in \mathcal{B}$) defined in §1. We have the factor $J_{H_b, \psi}: L \times V_b \rightarrow \mathbf{C}^*$ defined by

$$J_{H_b, \psi}(g, u) = \psi(g) \varepsilon \left[\frac{1}{2i} H_b(u, g) + \frac{1}{4i} H_b(g, g) \right]$$

where $g \in L$ and $u \in V_b$; this is because the imaginary part of H_b is equal to A for any b . Let $F(H_b, \psi)$ be the line bundle over E_b associated with the factor $J_{H_b, \psi}$. Since H_b is positive definite, we have $H^q(E_b, F(H_b, \psi)) = 0$ for $q \neq 0$ and $H^0(E_b, F(H_b, \psi))$ is identified with the complex vector space of all holomorphic theta functions on V_b for the factor $J_{H_b, \psi}$.

Let $p_i: W \rightarrow V_i(B)$ ($i=1, 2$) be the projection of W onto $V_i(B)$ with respect

to the decomposition (2) of W and let

$$\phi_b(u) = \exp [-\pi H_b(p_2(u), p_2(u))].$$

We have

$$\phi_b(u+g) = \phi_b(u) \exp L(u, g), \quad u \in W, \quad g \in L,$$

where

$$L(u, g) = -\pi[H_b(p_2(u), p_2(g)) + H_b(p_2(g), p_2(u)) + H_b(p_2(g), p_2(g))].$$

Let θ be a holomorphic theta function on V_b for the factor $J_{H_b, \psi}$ and let

$$f = \phi_b \cdot \theta$$

We show that the function f satisfies the conditions (1) and (2) in Theorem 1,

i.e. $f \in H(b)$. We have

$$f(u+g) = f(u)\psi(g) \exp [L(u, g) + \pi H_b(u, g) + \frac{\pi}{2} H_b(g, g)].$$

Since the decomposition (2) is orthogonal for H_b we get

$$\begin{aligned} \pi H_b(u, g) + \frac{\pi}{2} H_b(g, g) &= \pi H_b(p_1(u), p_1(g)) + \pi H_b(p_2(u), p_2(g)) + \frac{\pi}{2} H_b(p_1(g), p_1(g)) \\ &+ \frac{\pi}{2} H_b(p_2(g), p_2(g)) \quad \text{and hence} \quad L(u, g) + \pi H_b(u, g) + \frac{\pi}{2} H_b(g, g) \\ &= \pi [H_b(p_1(u), p_1(g)) - H_b(p_2(g), p_2(u))] + \frac{\pi}{2} [H_b(p_1(g), p_1(g)) - H_b(p_2(g), p_2(g))]. \end{aligned}$$

From Lemma at the end of §1 and from the orthogonality of the decomposition (2) for H we see that the left hand side of the above equality is equal to $\pi H(u, g) + \frac{\pi}{2} H(g, g)$. Hence we get $f(u+g) = f(u) \cdot \psi(g) \exp [\pi H(u, g) + \frac{\pi}{2} H(g, g)] = f(u) J_{H, \psi}(g, u)$ which shows that f is a differentiable theta function for the factor $J_{H, \psi}$.

Now let B be any privileged basis representing b and let (z_1, \dots, z_n) be the coordinates of V determined by B . Then B is also an orthonormal basis of V_b for the Hermitian form H_b and let (w_1, \dots, w_n) be the coordinates of V_b determined by B . Then as functions on W we have

$$\begin{aligned} z_i &= w_i \quad \text{for } i \in [1, s], \\ \bar{z}_{s+i} &= w_{s+i} \quad \text{for } i \in [1, r]. \end{aligned}$$

Since θ is a holomorphic function on V_b we have

$$\frac{\partial \theta}{\partial \bar{w}_k} = 0, \quad \text{for } k \in [1, n]$$

and hence

$$\frac{\partial \theta}{\partial \bar{z}_i} = 0, \quad i \in [1, s]; \quad \frac{\partial \theta}{\partial z_{s+i}} = 0, \quad i \in [1, r],$$

If $u = \sum_{k=1}^n z_k v_k$, then $p_2(u) = \sum_{i=1}^r z_{s+i} v_{s+i}$ and hence $H_b(p_2(u), p_2(u)) = \sum_{i=1}^r |z_{s+i}|^2$ and so

$$\phi_b = \exp \left[-\pi \sum_{i=1}^r |z_{s+i}|^2 \right].$$

We see easily that we have $\frac{\partial f}{\partial \bar{z}_i} = 0$ for $i \in [1, s]$ and $\frac{\partial f}{\partial \bar{z}_{s+i}} + \pi \bar{z}_{s+i} f = 0$ and hence f belongs to $H(b)$. Analogously we can see that if f is a function belonging to $H(b)$, then the function θ defined by $\theta(u) = f(u) \cdot \phi_b(u)^{-1}$ is a holomorphic theta function on V_b for the factor $J_{H_b, \psi}$ and moreover the map $f \rightarrow \theta$ defines a bijection of $H(b)$ onto the space $H^0(E_b, F(H_b, \psi))$ of holomorphic theta functions on V_b for the factor $J_{H_b, \psi}$. Since $H(b)$ is canonically isomorphic to $H'(E, F(H, \psi))$ by Theorem 1, we obtain the following theorem.

Theorem 2. *Let H be a Riemann form of signature (s, r) for a complex torus E and let $F(H, \psi)$ be the holomorphic line bundle over E associated with the factor $J_{H, \psi}$ defined by (1). Let (E_b, H_b) and $(F(H_b, \psi))$ be the family of polarized abelian varieties and the family of line bundles over each E_b parametrized by $b \in \mathfrak{B}$. Then there exists a canonical isomorphism of $H'(E, F(H, \psi))$ onto $H^0(E_b, F(H_b, \psi))$.*

In particular, we have

$$\dim H'(E, F(H, \psi)) = \dim H^0(E_b, F(H_b, \psi))$$

and since the imaginary part of H_b is equal to the imaginary part A of H , we have $\dim H^0(E_b, F(H_b, \psi)) = e_1, \dots, e_n$, where e_1, \dots, e_n are the elementary divisors of the integral valued alternating form A on $L \times L$. Thus we get also

$$\dim H'(E, F(H, \psi)) = e_1 \cdots e_n.$$

Let $N+1 = \dim H'(E, F(H, \psi))$ and let (f_0, f_1, \dots, f_N) be a basis of the complex vector space $H(b)$ which is canonically isomorphic to $H'(E, F(H, \psi))$. The map $u \rightarrow [f_0(u) : \dots : f_N(u)]$ defines a differentiable map $\tilde{\Phi}$ from W into the complex projective space P^N . Since each f_i is a differentiable theta function on W for the factor $J_{H, \psi}$, the map $\tilde{\Phi}$ defines a map Φ from $E = V/L$ into P^N .

Let $\theta_i = \phi_b \cdot f_i$ for $i \in [0, N]$. Then we have:

$$[\theta_0(u) : \theta_1(u) : \dots : \theta_N(u)] = [f_0(u) : f_1(u) : \dots : f_N(u)].$$

It follows from this that Φ defines a holomorphic map from $E_b = V_b/L$ to P^N . We

may assume without loss of generality that Φ is a holomorphic imbedding (this can be achieved by replacing H by $3H$ and ψ by ψ^3). Then Φ defines a differentiable imbedding of E into P^N . Let (z_1, \dots, z_n) be the coordinates on V corresponding to a privileged basis of V for H . Then these coordinates define local coordinates of the complex torus E at each point of E . Since Φ is holomorphic as a map from E_b into P^N , we see that Φ is holomorphic in z_1, \dots, z_s and anti-holomorphic in z_{s+1}, \dots, z_n . Thus we get the following theorem.

Theorem 3. *Let H be a Riemann form of signature (s, r) for a complex torus E . Then the cohomology group $H^*(E, F(3H, \psi^3))$ of the holomorphic line bundle $F(3H, \psi^3)$ defines a differentiable imbedding of E into the complex projective space P^N with $N+1 = \dim H^*(E, F(3H, \psi^3))$ which is holomorphic in z_1, \dots, z_s and anti-holomorphic in z_{s+1}, \dots, z_n , where (z_1, \dots, z_n) are the coordinates of the complex vector space V determined by a privileged basis for H .*

Appendix 1. We give here an example of a complex 2-torus which is not an abelian variety and which admits a Riemann form of signature $(1, 1)$. Let

$$\omega_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \omega_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \omega_3 = \begin{pmatrix} i\sqrt{2} \\ i\sqrt{3} \end{pmatrix}, \quad \omega_4 = \begin{pmatrix} i\sqrt{3} \\ -i\sqrt{5} \end{pmatrix}.$$

These vectors are linearly independent over \mathbf{R} and they generate a lattice L of \mathbf{C}^2 .

The matrix J_0 of the complex structure of \mathbf{C}^2 with respect to the basis $\{\omega_1, \omega_2, \omega_3, \omega_4\}$ of \mathbf{C}^2 over \mathbf{R} is of the form

$$J_0 = \begin{pmatrix} 0 & J_1 \\ J_2 & 0 \end{pmatrix}$$

where

$$J_1 = \begin{pmatrix} -\sqrt{2} & -\sqrt{3} \\ -\sqrt{3} & \sqrt{5} \end{pmatrix}$$

and

$$J_2 = \frac{1}{d} \begin{pmatrix} -\sqrt{5} & -\sqrt{3} \\ -\sqrt{3} & \sqrt{2} \end{pmatrix}, \quad d = -\sqrt{10} - 3.$$

Let A be an alternating \mathbf{R} -bilinear form on $\mathbf{C}^2 \times \mathbf{C}^2$ which is integral valued on $L \times L$ and let A_0 be the matrix of A with respect to the basis $\{\omega_i\}$ and write

$$A_0 = \begin{pmatrix} P_1 & P_2 \\ -{}^t P_2 & P_3 \end{pmatrix} \quad P_1 = \begin{pmatrix} 0 & p \\ -p & 0 \end{pmatrix} \quad P_3 = \begin{pmatrix} 0 & q \\ -q & 0 \end{pmatrix},$$

where p and q are integers and P_2 is an integral 2×2 matrix.

The alternating form A is the imaginary part of a Riemann form if and only if the \mathbf{R} -bilinear form S on $\mathbf{C}^2 \times \mathbf{C}^2$ defined by $S(u, v) = A(iu, v)$ is symmetric and non-degenerate. Let S_0 be the matrix of S with respect to $\{\omega_i\}$. Then we have $S_0 = {}^t J_0 \cdot A_0$. We see easily that the condition that S_0 is symmetric is equivalent to the set of the following three conditions: (a) $P_1 J_1 = -{}^t J_2 P_3$; (b) $P_2 J_2$ is symmetric; (c) ${}^t P_2 J_1$ is symmetric. The conditions (b) and (c) are both equivalent to the single condition that P_2 is to be of the form

$$P_2 = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \quad a \in \mathbf{Z}.$$

The condition (a) is equivalent to the condition $pd = q$, where $d = -\sqrt{10} - 3$ and p and q are integers and so (a) is equivalent to the condition $p = q = 0$.

Thus we reached the conclusion that S_0 is symmetric if and only if A_0 is the form

$$(*) \quad A_0 = \begin{pmatrix} 0 & a1_2 \\ -a1_2 & 0 \end{pmatrix}$$

where $a \neq 0$ is an integer and 1_2 denote the 2×2 unit matrix. Then S_0 takes the form

$$S_0 = \begin{pmatrix} -aJ_2 & 0 \\ 0 & aJ_1 \end{pmatrix}$$

and S_0 is a non-singular matrix. However S_0 is not definite because the symmetric matrix J_1 is not definite. Thus A can be the imaginary part of a Riemann form if and only if A_0 is of the form (*) and when this is the case, the corresponding Riemann form is not definite but of signature (1,1). Hence $E = \mathbf{C}^2/L$ provides an example of a complex torus which is not an abelian variety and which admits a Riemann form of signature (1,1).

Appendix 2. Since the second assertion in Theorem 1 is an essential part of this article we give a proof of Theorem 1 in this appendix.

Let H be a Riemann form of signature (s, r) for a complex torus $E = V/L$ and $J_{H, \psi}$ the factor defined by (1) and $F(H, \psi)$ the holomorphic line bundle associated with $J_{H, \psi}$. Let D^q be the vector space of all $F(H, \psi)$ -valued differential forms of type $(0, q)$ on E . Then the cohomology group $H(E, F(H, \psi))$ of E with coefficient in the sheaf of germs of holomorphic sections of $F(H, \psi)$ is isomorphic to the cohomology group of the complex $D = \sum_{q=0}^n D^q$, where the coboundary operator is given by d'' (or $\bar{\partial}$). On the other hand, there exists a canonical identification of an $F(H, \psi)$ -valued $(0, q)$ -form α on E with a $(0, q)$ -form φ on V (of class \mathbf{C}^∞) satisfying the condition that

$$(*) \quad T_g^* \varphi = J_{H, \psi}(g, \cdot) \varphi$$

for $g \in L$, where T_g denotes the translation of V by g . Then $d''\varphi$ also satisfies the same condition $(*)$ and $d''\alpha$ is identified with $d''\varphi$. Denote by A^q the vector space of all $(0, q)$ -form on V (of class C^∞) satisfying the condition $(*)$. Then the cohomology group $H(E, F(H, \psi))$ is isomorphic to the cohomology group of the complex $A = \sum_{q=0}^n A^q$, where the coboundary operator is given by d'' . Notice that A^0 is the vector space of all differentiable theta functions on V . Let (z_1, \dots, z_n) be coordinates on V . Then a $(0, q)$ -form is expressed uniquely in the form

$$\varphi = \frac{1}{q!} \sum_J \varphi_J d\bar{z}_J,$$

where $J = (j_1, \dots, j_q)$ is a multi-index and each φ_J is alternating in the indices and $d\bar{z}_J = d\bar{z}_{j_1} \wedge \dots \wedge d\bar{z}_{j_q}$. Since $d\bar{z}_J$ is invariant by translation, φ satisfies the condition $(*)$ if and only if each component φ_J belongs to A^0 .

Lemma 1. *Let $f, g \in A^0$ and let $\langle f, g \rangle$ be defined by*

$$\langle f, g \rangle(u) = f(u)g(u) \exp [-\pi H(u, u)]$$

for $u \in V$. Then the function $\langle f, g \rangle$ is invariant by the translation T_g for any $g \in L$.

We can verify the lemma by a straightforward computation.

We may consider $\langle f, g \rangle$ as function on the torus $E = V/L$.

Corollary of Lemma 1. *If $f, g \in A^0$, then*

$$|f(u)| |g(u)| < C \exp \pi H(u, u)$$

for any $u \in V$, where C is a positive constant.

Let us choose a *positive definite* Hermitian form G and let

$$G = \sum_{i,j} g_{ij} z_i \bar{z}_j.$$

Let

$$dV = \left(\frac{i}{2}\right)^n \det(g_{ij}) dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_n$$

the volume element on V determined by G . The volume element dV is invariant by translation and so it defines a translation invariant volume element dv on E such that $\pi^* dv = dV$, where $\pi: V \rightarrow E$ is the canonical projection. We define the inner product (f, g) , where $f, g \in A^0$, by

$$(f, g) = \int_P \langle f, g \rangle dV,$$

where P is a fundamental parallelepiped for the lattice L , or equivalently by

$$(f, g) = \int_B \langle f, g \rangle dv$$

regarding $\langle f, g \rangle$ as function on E .

Let us write

$$H = \sum_{i,j} h_{ij} z_i \bar{z}_j$$

and introduce covariant derivations D'_i, D''_i ($i=1, \dots, n$) by the formula

$$\begin{aligned} (D'_i f)(z) &= \frac{\partial f}{\partial z_i}(z) - \pi \left(\sum_k h_{ik} \bar{z}_k \right) f(z), \\ (D''_i f)(z) &= \frac{\partial f}{\partial \bar{z}_i}(z). \end{aligned}$$

We can show without difficulty that if $f \in A^0$, then we have $D'_i f, D''_i f \in A^0$ for $i=1, \dots, n$. We have also the following formulas:

$$\begin{aligned} \langle D'_i f, g \rangle + \langle f, D''_i g \rangle &= \frac{\partial}{\partial z_i} \langle f, g \rangle, \\ \langle D''_i f, g \rangle + \langle f, D'_i g \rangle &= \frac{\partial}{\partial \bar{z}_i} \langle f, g \rangle. \end{aligned}$$

Integralating both sides of the equalities and using the Green's theorem we obtain

$$\begin{aligned} (D'_i f, g) + (f, D''_i g) &= 0, \\ (D''_i f, g) + (f, D'_i g) &= 0, \end{aligned}$$

where $f, g \in A^0$ and $i \in [1, n]$.

Denote by g^{ij} the (i, j) -entry of the inverse matrix of the Hermitian matrix (g_{ij}) and let

$$g^{IJ} = g^{i_1 j_1} \dots g^{i_q j_q}$$

where $I = (i_1, \dots, i_q)$ and $J = (j_1, \dots, j_q)$.

For $\varphi, \psi \in A^q$, we define the function $\langle \varphi, \psi \rangle$ by

$$\langle \varphi, \psi \rangle = \frac{1}{q!} \sum_{I, J} g^{IJ} \langle \varphi_I, \psi_J \rangle$$

Then $\langle \varphi, \psi \rangle$ is invariant by the translation $T_g(g \in L)$ and we define the inner

product (φ, ψ) by

$$(\varphi, \psi) = \int_P \langle \varphi, \psi \rangle dV = \int_{\#} \langle \varphi, \psi \rangle dv.$$

There exists the adjoint operator \mathfrak{d} for the operator $d'': A^q \rightarrow A^{q+1}$ so that we have

$$(d''\varphi, \psi) = (\varphi, \mathfrak{d}\psi)$$

for $\varphi \in A^q$ and $\psi \in A^{q+1}$

We define the Laplacian \square by

$$\square = d''\mathfrak{d} + \mathfrak{d}d''$$

Then \square is an operator from A^q to A^q for all q and a $(0, q)$ -form $\varphi \in A^q$ is said to be harmonic if $\square\varphi = 0$. Each element of the cohomology group $H^q(A)$ of the complex A is represented by a unique harmonic form. In this sense we can say that each element of the cohomology group $H^q(E, F(H, \psi))$ is represented by a unique harmonic form. *Thus we may identify $H^q(E, F(H, \psi))$ with the vector space of all harmonic forms in A^q .* We now introduce the following notation. For $I = (i_1, \dots, i_{q+1})$, I_u will denote the multi-index $(i_1, \dots, \hat{i}_u, \dots, i_{q+1})$, where the index i_u under \wedge is omitted. We also introduce the operator D'^i by

$$D'^i = \sum_j g^{ij} D'_j$$

We can prove the following three formulas.

A) Let $\varphi \in A^q$. Then the components $(d''\varphi)_I$, $I = (i_1, \dots, i_{q+1})$ of $d''\varphi \in A^{q+1}$ is given by the formula

$$(d''\varphi)_I = \sum_{u=1}^{q+1} (-1)^{u+1} D'_{i'_u} \varphi_{I_u}$$

B) Let $\psi \in A^{q+1}$. Then the component $(\mathfrak{d}\psi)_J$, $J = (j_1, \dots, j_q)$, of $\mathfrak{d}\psi$ is given by the formula

$$(\mathfrak{d}\psi)_J = - \sum_{j=1}^n D'^j \psi_{jJ},$$

where $jJ = (j, j_1, \dots, j_q)$.

C) Let $\varphi \in A^q$. Then the component $(\square\varphi)_I$ of $\square\varphi \in A^q$ is given by the formula

$$(\square\varphi)_I = - \left(\sum_{i=1}^n D'^i D'_{i'} \right) \cdot \varphi_I + \pi \sum_{u=1}^q (-1)^{u+1} \sum_{i=1}^n \left(\sum_{k=1}^n g^{ik} h_{ki_u} \right) \varphi_{iI_u}$$

where $iI_u = (i, i_1, \dots, \hat{i}_u, \dots, i_q)$.

We omit the proof of these formulas. Similar formulas had been proved in [1] in a somewhat different context, but the proof can be carried out quite

similarly.

Up to this point the choices of the coordinates (z_1, \dots, z_n) and the positive definite Hermitian form $G = \sum_{i,j} g_{ij} z_i \bar{z}_j$ are arbitrary. From now on we choose privileged coordinates (z_1, \dots, z_n) for the Hermitian form H so that we have

$$H = \sum_{i=1}^s |z_i|^2 - \sum_{i=1}^r |z_{s+i}|^2$$

and hence we have $h_{ij}=0$ for $i \neq j$ and $H_{ii} = \varepsilon_i$ (the symbol ε_i being defined in §1). We choose G such that

$$G = \frac{1}{a} (|z_1|^2 + \dots + |z_s|^2) + |z_{s+1}|^2 + \dots + |z_n|^2,$$

where $a > 0$ (cf. [4]). Then we have $g^{ij}=0$ for $i \neq j$ and

$$g^{ii} = \begin{cases} a & \text{for } i \in [1, s] \\ 1 & \text{for } i \in [s+1, n]. \end{cases}$$

Then we have

$$\sum_k g^{ik} h_{kj} = \begin{cases} 0, & i \neq j \\ a, & i=j \text{ and } i \in [1, s] \\ -1, & i=j \text{ and } i \in [s+1, n]. \end{cases}$$

Let

$$\alpha_i = \begin{cases} a & \text{for } i \in [1, s] \\ -1 & \text{for } i \in [s+1, n]. \end{cases}$$

Then we have $\sum_{i=1}^n (\sum_{k=1}^n g^{ik} h_{ki_u}) \varphi_{iI_u} = (-1)^{u+1} \alpha_{i_u} \varphi_I$ and we get

$$(\square \varphi)_I = -(\sum_i D'^i D'_i) \varphi_I + \pi (\sum_{u=1}^q \alpha_{i_u}) \varphi_I,$$

where

$$D'^i = g^{ii} D'_i \quad (\text{not summed}).$$

From this we obtain

$$((\square \varphi)_I, \varphi_I) = \sum_{i=1}^n g^{ii} (D'_i{}' \varphi_I, D'_i{}' \varphi_I) + \pi (\sum_{u=1}^q \alpha_{i_u}) (\varphi_I, \varphi_I).$$

Since the first term of the right hand side is non-negative, we get

$$((\square \varphi)_I, \varphi_I) \geq \pi \cdot \alpha(I) \cdot (\varphi_I, \varphi_I).$$

where we put

$$\alpha(I) = \sum_{u=1}^q \alpha_{i_u}.$$

Let us denote by N (resp. M) the number of indices i_u such that $i_u \leq s$ (resp. $i_u > s$). Then by the definition of α_k , we have

$$\alpha(I) = a \cdot N - M.$$

For the multi-index I we may assume that these q indices are distinct, otherwise we get $\varphi_I = 0$. Suppose $q > r$. Then at least one of the indices i_u must be less than or equal to s and hence $N \geq 1$. Then we get

$$\alpha(I) \geq a - r$$

Choose a such that $a > r$. Then we have $\alpha(I) > 0$ for $q > r$.

Suppose the $\square \varphi = 0$, where $\varphi \in A^q$ with $q > r$. Then

$$0 = ((\square \varphi)_I, \varphi_I) \geq \pi \alpha(I) \cdot (\varphi_I, \varphi_I)$$

with $\alpha(I) > 0$. Hence we must have $(\varphi_I, \varphi_I) = 0$ and hence $\varphi_I = 0$ for any I and this means $\varphi = 0$. This shows that $H^q(A) = 0$ and hence $H^q(E, F(H, \psi)) = 0$ for $q > r$.

On the other hand, by the Serre duality, we have

$$H^q(E, F(H, \psi)) \cong H^{n-q}(E, K \otimes F^*),$$

where K is the canonical line bundle of E and F^* is the dual of $F(H, \psi)$. It is easy to see that $F^* \cong F(-H, \varphi^{-1})$ and $-H$ is of signature (r, s) . Moreover since E is a complex torus, K is a trivial bundle and so we get

$$H^q(E, F(H, \psi)) \cong H^{n-q}(E, F(-H, \varphi^{-1})).$$

Since $-H$ is of signature (r, s) , we get from what we have already proved that $H^{n-q}(E, F(-H, \varphi^{-1})) = 0$ whenever $n - q > s$ or whenever $n - s = r > q$. Thus we get $H^q(E, F(H, \psi)) = 0$ for $q < r$ and these prove the first assertion in Theorem 1.

Consider now that case $q = r$, $\varphi \in A^r$ and $\square \varphi = 0$. Even in this case we get $\alpha(I) \geq a - r > 0$ except in the case where all of the r indices in I are greater than s , namely except in the case where I is a permutation of $(s+1, \dots, n)$. Then we get $\varphi_I = 0$ for each I which is not a permutation of $(s+1, \dots, n)$ and φ is of the form

$$(**) \quad \varphi = f d\bar{z}_{s+1} \wedge \dots \wedge d\bar{z}_n,$$

where $f = f_{s+1, \dots, n}$.

Conversely let φ be a $(0, r)$ -form on V of the type $(**)$ belonging to A^r . Then $f \in A^0$ and we have

$$d''\varphi = 0 \Leftrightarrow \frac{\partial f}{\partial \bar{z}_i} = 0 \quad \text{for } i \in [1, s]$$

and

$$(\mathfrak{d}\varphi)_I = -\sum_{i=1}^n g^{ii} D'_i \varphi_{iI}$$

where $I=(i_1, \dots, i_{r-1})$. If $I \neq (s+1, \dots, \hat{u}, \dots, n)$ for some u , (i, I) cannot be a permutation of $(s+1, \dots, n)$ and $\varphi_{iI}=0$ and hence $(\mathfrak{d}\varphi)_I=0$. If $I=(s+1, \dots, \hat{u}, \dots, n)$ for some u , then

$$(\mathfrak{d}\varphi)_I = \pm g^{uu} D'_u f$$

Therefore we have

$$\mathfrak{d}\varphi = 0 \Leftrightarrow D'_u f = 0 \quad \text{for } u = s+1, \dots, n.$$

It follows from our definition of the operator D'_u and from the fact $h_{ij}=\delta_{ij} \cdot \varepsilon_j$, we see that

$$D'_u f = \frac{\partial f}{\partial \bar{z}_u} + \pi \bar{z}_u f.$$

We have thus proved that the space of harmonic $(0, r)$ -form φ in A' consists of all the $(0, r)$ -form φ on V of the form

$$\varphi = f d\bar{z}_{s+1} \wedge \dots \wedge d\bar{z}_n,$$

where

- 1) f is a differentiable theta function for the factor $J_{H,\psi}$,
- 2) $\frac{\partial f}{\partial \bar{z}_i} = 0$ for $i \in [1, s]$

and

$$\frac{\partial f}{\partial \bar{z}_i} + \pi \bar{z}_i f = 0 \quad \text{for } i \in [s+1, n].$$

Then we can identify the cohomology group $H^r(A)$ with the vector space of functions f satisfying the conditions 1) and 2) and this proves the second assertion in Theorem 1.

UNIVERSITY OF NOTRE DAME

Bibliography

- [1] Y. Matsushima and S. Murakami: *On vector bundle valued harmonic forms and automorphic forms on symmetric Riemannian manifolds*, Ann. of Math. **78** (1963), 365-416.
- [2] D. Mumford: *Abelian varieties*, Tata Inst. Studies in Math., Oxford Univ. Press, 1970.
- [3] C.L. Siegel: *Indefinite quadratische Formen und Funktionentheorie*. I, Math. Ann. **124** (1951), 17-54.
- [4] H. Umemura: *Some results in the theory of vector bundles*, Nagoya Math. J. **52** (1973), 97-128.

