The University of Osaka
Institutional Knowledge Archive

Title	Label-Free Evaluation of Maturation and Hepatotoxicity of Human iPSC-Derived Hepatocytes Using Hyperspectral Raman Imaging
Author(s)	Li, Menglu; Ueyama-Toba, Yukiko; Lindley, Matthew et al.
Citation	Analytical Chemistry. 2023, 95(24), p. 9252-9262
Version Type	AM
URL	https://hdl.handle.net/11094/103302
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Supporting Information

Label-free Evaluation of Maturation and Hepatotoxicity of Human iPSC-derived Hepatocytes using Hyperspectral Raman Imaging

Menglu Li[1,2†*], Yukiko Ueyama-Toba[3,4,5†], Matthew Lindley[1], Gunganist Kongklad[1,2,6], Yasunori Nawa[1,2], Yasuaki Kumamoto[1,4], Seiichi Ishida[2,7], Yasunari Kanda[2,8], Satoshi Fujita[1,2], Hiroyuki Mizuguchi[3,4,5], Katsumasa Fujita[1,2,4*]

Affiliations:

[1] Laboratory of Nanophotonics, Department of Applied Physics, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

[2] AIST-Osaka University Advanced Photonics and Biosensing Open Innovation Laboratory, National Institute of Advanced Industrial Science and Technology (AIST), Suita, Osaka 565-0871, Japan.

[3] Laboratory of Biochemistry and Molecular Biology, Graduate School of Pharmaceutical Sciences, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

[4] Institute for Open and Transdisciplinary Research Initiatives, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

[5] Laboratory of Functional Organoid for Drug Discovery, Center for Drug Discovery Resources Research, National Institutes of Biomedical Innovation, Health and Nutrition, 7-6-8 Saito-Asagi, Ibaraki, Osaka 567-0085, Japan

[6] Department of Physics, Faculty of Science, Mahidol University, Bangkok 10400, Thailand

[7] Division of Applied Life Science, Graduate School of Engineering, Sojo University, 4-22-1, Ikeda, Nishi-ku, Kumamoto 860-0082, Japan.

[8] Division of Pharmacology, National Institute of Health Sciences, Kawasaki, Kanagawa 210-9501, Japan.

*Correspondence to: Menglu Li, Katsumasa Fujita

† Equal contributions

EMAIL: menglu.li@ap.eng.osaka-u.ac.jp; fujita@ap.eng.osaka-u.ac.jp

27 **Table S1 The primary antibodies used in this study**

Antigen	Type	Company	Catalog number	Dilution
NANOG	Mouse	Santa Cruz Biotechnology	sc-293121	1:300
CXCR4	Mouse	Santa Cruz Biotechnology	sc-53534	1:300
HNF4α	Rabbit	Cell Signaling Technology	CST3113	1:300
CYP3A4	Mouse	Sigma-Aldrich	SAB5300118	1:1000
ALB	Goat	Bethyl laboratories	A80-129A	1:500
CK-19	Rabbit	Abcam	ab52625	1:500
Cyt *c*	Mouse	Santa Cruz Biotechnology	Sc-13561	1:500

28

29

30 **Table S2 The secondary antibodies used in this study**

Antigen	Type	Fluorophore	Company	Catalog number	Dilution
Mouse IgG	Goat	Alexa Fluor 488	Invitrogen	A11001	1:500
Mouse IgG	Goat	Alexa Fluor 594	Invitrogen	A11005	1:500
Rabbit IgG	Goat	Alexa Fluor 488	Invitrogen	A11008	1:500
Rabbit IgG	Goat	Alexa Fluor 594	Invitrogen	A11012	1:500
Goat IgG	Donkey	Alexa Fluor 594	Invitrogen	A11058	1:500

31

32 **Table S3 The primers used for real-time RT-PCR in this study**

Gene Symbol	Primers (forward/reverse; 5' to 3')
SOX2	GGCAGCTACAGCATGATGATGCAGGAGC/CTGGTCATGGAGTTGTACTGCAGG
SOX17	GTGGACCGCACGGAATTTG/GAGGCCCATCTCAGGCTTG
HNF4α	CGTCATCGTTGCCAACACAAT/GGGCCACTCACACATCTGTC
CYP3A4	AAGTCGCCTCGAAGATACACA/AAGGAGAGAACACTGCTCGTG
ALB	GCACAGAATCCTTGGTGAACAG/GCACAGAATCCTTGGTGAACAG
GAPDH	GGTGGTCTCCTCTGACTTCAACA/GTGGTCGTTGAGGGCAATG

33

34

35



Figure S1 The gene expression levels of *SOX2*, *SOX17*, *HNF4α*, *CYP3A4*, and *ALB*. (a) hiPSCs and their derivatives cultured in plastic plates. (b) hiPSCs and their derivatives cultured in quartz dishes.



Figure S2 Time-course change of lipid-rich colony from D20 to D25 following immunofluorescence staining of CYP3A4. This result indicates the potential of using lipid as an early marker of CYP-positive HLCs. Red arrows indicate the same position at different time points. Scale bars, 50 μm.



Figure S3 Screening hepatotoxicity in iCell hepatocytes and HepaRG cells using Raman imaging. 0-10 mM APAP was applied to iCell hepatocytes (a) and HepaRG cells (b) culture. Reconstructed Raman images indicating cellular components at 642, 675, 1636, 940, 1440, and 2930 cm^{-1} are shown. Scale bars, 50 µm.