

Title	On a generalization of the theorem of Reisz- Frostman-Nevanlinna
Author(s)	Ikegami, Teruo
Citation	Osaka Journal of Mathematics. 1972, 9(3), p. 519–533
Version Type	VoR
URL	https://doi.org/10.18910/10339
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Ikegami, T. Osaka J. Math. 9 (1972), 519-533

# ON A GENERALIZATION OF THE THEOREM OF RIESZ-FROSTMAN-NEVANLINNA

To Professor Yukinari Tôki on the occasion of his 60th birthday

TERUO IKEGAMI

(Received December 17, 1971)

## Introduction

The original form of the theorem of Riesz-Frostman-Nevanlinna is stated as follows: let f(z) be regular and bounded in the unit disc. If  $\lim_{r \to 1} f(re^{i\theta})$  is equal to zero on a subset of positive measure of |z| = 1, then  $f(z) \equiv 0$ . R. Nevanlinna<sup>1)</sup> and O. Frostman<sup>2)</sup> extended independently this theorem to the case of meromorphic functions of bounded type. However, if we consider arbitrary regular functions this theorem does not hold in general as the example of Lusin-Priwalow shows<sup>3)</sup>.

Meanwhile, it has been made known by the recent studies of Constantinescu-Cornea<sup>4</sup><sup>3</sup> that the boundary behavior of analytic maps of Riemann surfaces depends deeply on the harmonic character of maps. In [7], they developed this idea to maps of harmonic spaces satisfying the Brelot's axioms.

In this paper, we shall generalize the theorem of Riesz-Frostman-Nevanlinna for maps of a Green space into a harmonic space. Generalizations are done in some points. One of them is the use of cluster sets along Green lines issuing from a fixed point and of a Green measure instead of radial limits and the Lebesgue measure, respectively. However, an essential point is the validity of the theorem for Fatou maps which include all Lindelöfian maps<sup>5)</sup> of hyperbolic Riemann surfaces<sup>6)</sup>.

In §1, we state the theorem and list up all notations which will be used in the sequel. §2 is devoted to auxiliary lemmas. They are needful to the proof of the theorem. The proof of the theorem is carried out in §3 divided into three

6) Cf. [5], p. 113 and [4], p. 72.

<sup>1)</sup> Cf. [13], p. 205.

<sup>2)</sup> Cf. [8], p. 96.

<sup>3)</sup> Cf. [11] and [14], p. 222.

<sup>4)</sup> Cf. [4] and [5].

<sup>5)</sup> Cf. [9].

cases. In the last section, as an application, we shall mention a result which is an improvement of my former one [10].

## 1. Preliminaries and the theorem

Let  $\Omega$  be a Green space in the sense of Brelot-Choquet<sup>7</sup>). We consider the Green lines issuing from a fixed point  $y_0$ . They are the maximal orthogonal trajectories of

$$\sum^{\lambda} = \{y \in \Omega; G_{y_0}(y) = \lambda\},\$$

where  $G_{y_0}$  is a Green function of  $\Omega$  with a pole at  $y_0$  and  $0 < \lambda < G_{y_0}(y_0)$ . We put

$$D^{\lambda} = \{ y \in \Omega; G_{y_0}(y) > \lambda \} \quad \text{for} \quad 0 < \lambda < G_{y_0}(y_0) \,.$$

On the set  $\mathcal{L}$  of all Green lines, we can define a topology homeomorphic to the unit sphere and a Radon measure g called the *Green measure*. A Green line l is called *regular*, if inf  $\{G_{y_0}(y); y \in l\} = 0$ . The set of all regular Green lines will be denoted by  $\mathcal{L}'$ .

Let X be a harmonic space in the sense of Brelot<sup>8</sup>, i.e., X is locally compact, connected and on which it is given a sheaf of continuous functions, called harmonic functions, satisfying the axioms 1, 2 and 3 in [2]. A Green space is a harmonic space in an obvious way. We denote by  $\mathcal{P}$  (resp.  $\mathcal{H}$ ) the class of harmonic spaces on which there exists a positive potential (resp. a positive harmonic function). A continuous map  $\varphi$  of a harmonic space X into a second harmonic space X' is called a *harmonic map*, if for any open set  $U' \subset X'$  and any harmonic function u' on U',  $u' \circ \varphi$  is harmonic on  $\varphi^{-1}(U')$ .

Let U be an open subset of X,  $U \in \mathcal{P}$  and f be a real function defined on U. We denote by  $\overline{w}_{T}^{U}$  the set of hyperharmonic functions s on U such that

a) s possesses a non-positive subharmonic minorant

and

b) s dominates f outside a compact subset of U.

We denote by  $\bar{h}_{I}^{U} = \inf \{s; s \in \overline{w}_{I}^{U} \}$ . Also we define

$$\underline{w}_{f}^{U} = \{-s; s \in \overline{w}_{(-f)}^{U}\}$$

and

$$\underline{h}_{J}^{U} = \sup \{s; s \in \underline{w}_{J}^{U}\}.$$

If  $\bar{h}_{r}^{U} = \underline{h}_{r}^{U}$  and is finite, f is called *harmonizable on U*. A finite continuous function f on X is called a *Wiener function*, if there exists an open set  $U \in \mathcal{P}$  with

<sup>7)</sup> Cf. [3]. For the following facts we refer to [3].

<sup>8)</sup> Cf. [2]. In [2], it is assumed that X is not compact. In this paper we do not require this.

compact complement such that f is harmonizable on U. A harmonic map  $\varphi$  of X into X' is called a *Fatou map* if for any bounded Wiener function f' on  $X', f' \circ \varphi$  is a Wiener function on X. All harmonic maps into  $X' \in \mathcal{P}$  are Fatou maps. When  $X \in \mathcal{P}$  and  $X' \in \mathcal{H} - \mathcal{P}$ , in order that a harmonic map  $\varphi$  of X into X' be a Fatou map, it is necessary and sufficient that there exists a closed non-polar set  $F' \subset X'$  such that  $\hat{R}_1^{\varphi - 1(F')}$  is a potential<sup>9</sup>.

A compactification  $X^*$  of X is a compact space containing X as a dense open subset. A subset A of  $X^*$  is called *polar*, if for any domain  $U \in \mathcal{P}$  of X there exists a positive superharmonic function s on U such that

$$\lim_{x\to z} s(x) = +\infty \quad \text{for any} \quad z \in A \cap \overline{U}.$$

We shall list up the notations which will be used in the sequel.

- $\Omega$ : a Green space.
- $\{\Omega_n\}$ : an exhaustion of  $\Omega$ , i.e.,  $\Omega_n$  is a relatively compact domain satisfying  $\overline{\Omega}_n \subset \Omega_{n+1}$  and  $\bigcup_{n=1}^{\infty} \Omega_n = \Omega$ .
- X: a harmonic space with countable basis, on which there exists a positive superharmonic function, i.e.,  $X \in \mathcal{P} \cup \mathcal{H}$ .
- $X^*$ : a compactification of X.

 $\{X_n\}$ : an exhaustion of X.

 $\Lambda$ : a set of regular Green lines.

 $\{\lambda_n\}$ : a decreasing sequence of positive numbers tending to zero.

 $\varphi$ : a harmonic map of  $\Omega$  into X.

 $\phi(\ell) = \bigcap_{n=1}^{\infty} \overline{\{\varphi((\lambda_m, \ell)); m \ge n\}},$  where the closure is taken in the topology of  $X^*$ .

A: a polar set in  $X^*$ .

In this paper, we shall prove the following theorem, which is a generalization of the theorem of Riesz-Frostman-Nevanlinna.

**Theorem.** Let X be a harmonic space. We assume the existence of

(\*) a countable basis of open sets for X

and

(\*\*) a superharmonic function with positive infimum on X. If X is compact, we assume further the existence of

(\*\*\*) a non-polar subset E of X each point of which is polar.

Let  $\varphi$  be a non-constant Fatou map of a Green space  $\Omega$  into X and X\* be an arbitrary compactification of X.

9) Cf. [7], p. 52, th. 6.1.

T. İkegami

If there exist a set  $\Lambda$  of regular Green lines issuing from  $y_0 \in \Omega$ , a decreasing sequence  $\{\lambda_n\}$  of positive numbers tending to zero and a polar set A of X\* such that

$$\phi(\ell) = \bigcap_{n=1}^{\infty} \overline{\{\varphi((\lambda_m, \ell)); m \ge n\}} \subset A \quad \text{for any} \quad \ell \in \Lambda ,$$

then the outer Green measure of  $\Lambda$  is zero, i.e.,  $g^*(\Lambda)=0$ , where  $(\lambda_m, \ell)$  denotes the point of  $\ell$  on which the value of a Green function  $G_{y_0}$  is  $\lambda_m$ , and the closure is taken in  $X^*$ .

#### 2. Lemmas

**2.1.** To prove the theorem stated above, we require some lemmas, which will be given in this section. Throughout this section we shall suppose

$$g^*(\Lambda) > 0$$
.

**Lemma 1.** Assume that X is non-compact and

$$\hat{\phi}(l) \subset A$$
 for any  $l \in \Lambda$ .

Then, there exists a sequence  $\{D_n\}$  of relatively compact domains in X such that

(2.1) 
$$g^*(\{\ell \in \Lambda; \phi(\ell) \cap \overline{D}_n = \phi\}) > 1/2 g^*(\Lambda) \quad (n = 1, 2, \cdots)$$

and

(2.2) 
$$(\varphi(\overline{\Omega}_n) \cup \bigcup_{k=1}^{n-1} \overline{D}_k \cup \overline{X}_n) \cap \overline{D}_n = \phi \qquad (n = 1, 2, \cdots).$$

Proof. Suppose defined  $D_1, D_2, \dots, D_{p-1}$ , relatively compact domains and let (2.1) and (2.2) hold for  $n=1, 2, \dots, p-1$ . Since X is not compact,

$$X - (\varphi(\overline{\Omega}_p) \cup \bigcup_{k=1}^{p-1} \overline{D}_k \cup \overline{X}_p)$$

is an open non-empty set, so that it is non-polar. Since  $A \cap X$  is polar, there exists x such that

(2.3) 
$$x \in X - (\varphi(\overline{\Omega}_p) \cup \bigcup_{k=1}^{p-1} \overline{D}_k \cup \overline{X}_p \cup A).$$

From the second axiom of countability for  $X^{10}$ , we have a sequence  $\{E_n\}$  of relatively compact neighbourhoods of x such that

(2.4) 
$$\begin{cases} \overline{E}_{m+1} \subset E_m \quad (m = 1, 2, \cdots), \\ \overline{E}_1 \cap [\varphi(\overline{\Omega}_p) \cup \bigcup_{k=1}^{p-1} \overline{D}_k \cup \overline{X}_p] = \phi, \\ \bigcap_{m=1}^{\infty} \overline{E}_m = \{x\}. \end{cases}$$

10) It is true that the same conclusion is derived from the first axiom of countability for X. However, in the harmonic space X the two countability axioms are equivalent. (Cf. [6]).

Put

$$L_{m} = \{ \ell \in \Lambda; \, \hat{\varphi}(\ell) \cap \bar{E}_{m} = \phi \}$$

for  $m = 1, 2, \dots$ . It is easy to see

$$(2.5) L_1 \subset L_2 \subset \cdots$$

and

(2.6) 
$$\bigcup_{m=1}^{\infty} L_m = \Lambda.$$

In fact, if there exists  $l \in \Lambda$  such that  $l \in L_m$  for all m, then

$$\phi(\ell) \cap \overline{E}_m \neq \phi \qquad (m = 1, 2, \cdots).$$

Since  $X^*$  is compact this means

$$\phi(l) \cap \bigcap_{m=1}^{\infty} \overline{E}_m \neq \phi$$
,

so that by (2.4)

$$x \in \phi(l) \subset A$$
 ,

which contradicts (2.3).

From (2.5), (2.6) and the regularity of the outer Green measure<sup>11)</sup> we have

$$\lim_{m\to\infty}g^*(L_m)=g^*(\Lambda)\,.$$

Thus, we have an  $m_0$  such that  $g^*(L_{m_0}) > (1/2)g^*(\Lambda)$ .  $E_{m_0} = D_p$  is the desired one, q.e.d..

**Lemma 2.** Suppose that X is non-compact and  $X \in \mathcal{H} - \mathcal{P}$ , and  $\varphi$  is a Fatou map of  $\Omega$  into X. Let  $\{D_n\}$  be a sequence of relatively compact domains satisfying (2.2). Then, there exists a closed subset F of X such that  $\hat{R}_1^{\varphi^{-1}(F)^{12}}$  is a potential,  $F_n = F \cap D_n$  is non-polar and compact, and

(2.7) 
$$\lim_{n\to\infty}\hat{R}_1^{\varphi-1(F_n)}=0 \quad on \quad \Omega.$$

Proof. Let  $f_n$  be a non-negative continuous function on X whose support is in  $D_n$  and whose maximum is 1. Since  $\{D_n\}$  are mutually disjoint and do not cluster at any point of  $X, f = \sum_{n=1}^{\infty} f_n$  is a non-negative bounded continuous function on X, so that f is a Wiener function on  $X^{13}$ . From the definition of a Fatou map,  $\overline{f} = f \circ \varphi$  is a Wiener function on  $\Omega$ . In virtue of a theorem of Constantinescu-Cornea<sup>14</sup>)  $\hat{R}_1^{N_{\alpha}}$  is a potential except for countable values of  $\alpha$ ,

<sup>11)</sup> Cf. [15], p. 51.

<sup>12)</sup> Cf. [2], p. 80, def. 9.

<sup>13)</sup> Cf. [7], p. 16.

T. Ikegami

where

$$N_{\alpha} = \{y \in \Omega; \overline{f}(y) = \alpha\}$$

We may take  $\alpha$  so that  $0 < \alpha < 1$  and  $\hat{R}_{1}^{N_{\alpha}}$  is a potential. Let us write

$$F = \{x \in X; f(x) = \alpha\}.$$

 $\hat{R}_1^{\varphi_{-1(F)}}$  is a potential and  $F_n = F \cap D_n$  is compact and non-polar, since

$$F \cap D_n = \{x \in X; f_n(x) = \alpha\}$$

and  $X - (F \cap D_n)$  is not connected.

For any  $y \in \Omega$  there exists  $n_0$  such that  $y \in \Omega_n$  for any  $n \ge n_0$ .  $\varphi^{-1}(F_n) \cap \Omega_n = \phi$  implies that  $\hat{R}_1^{\varphi^{-1}(F_n)}$  is bounded and harmonic in  $\Omega_n$ . Hence,

$$\hat{R}_{1}^{\varphi_{-1(F_{n})}}(y) = H_{\hat{R}_{1}^{\varphi_{-1}(F_{n})}}^{\mathcal{Q}_{n}}(y) \leq H_{\hat{R}_{1}^{\varphi_{-1}(F)}}^{\mathcal{Q}_{n}}(y) .$$

Since  $\hat{R}_1^{\varphi_{-1}(F)}$  is a potential, the last term tends to zero as  $n \to \infty$ , q.e.d..

2.2.

Let us take  $y_1 \in \Omega$  such that  $\varphi(y_1) \notin A \cup F$  and we shall fix it. This is possible for  $\varphi^{-1}(X \cap A)$  is polar. By the Harnack's inequality we can find K > 1satisfying

(2.8) 
$$K u(y_0) \ge u(y_1) \ge 1/K u(y_0)$$

for all non-negative harmonic functions u on  $\Omega$ .

Let  $\delta$  be a positive number less than 1/4. By Lemma 2, we have  $n_0$  such that

(2.9) 
$$\hat{R}_{1}^{\varphi_{-1}(F_{n_{0}})}(y_{1}) < \delta/(8K)g^{*}(\Lambda).$$

Since  $F_{n_0}$  is non-polar, each component of  $X - F_{n_0} \in \mathcal{P}$ . There exists a positive superharmonic function v on  $X - F_{n_0}$  such that

(2.10) 
$$\begin{cases} \lim_{x \to x'} v(x) = +\infty & \text{for any} \quad x' \in A \cap \overline{(X - F_{n_0})}, \\ v[\varphi(y_1)] < 1/(4K)g^*(\Lambda). \end{cases}$$

**Lemma 3.** Let E be a closed subset of  $\Omega$  such that  $\hat{R}_1^E$  is a potential. Then, for a fixed  $y_1 \in \Omega$  we have

(2.11)  $\hat{R}_{1}^{E}(y_{1}) = \inf \{ \hat{R}_{1}^{\omega \cap \Omega}(y_{1}); \omega \text{ is an open subset of } \Omega_{W}^{*} \text{ containing } \bar{E} \},$ where  $\Omega_{W}^{*}$  is a Wiener compactification<sup>15</sup> of  $\Omega$  and the closure is taken in  $\Omega_{W}^{*}$ .

<sup>14)</sup> Cf. [7], p. 14, th. 2.6.

<sup>15)</sup> Cf. [5], p. 98 and [7], p. 43.

Proof. Denoting by  $\alpha_1$  the right-hand side of (2.11), we have clearly

$$lpha_{\scriptscriptstyle 0}=\hat{R}_{\scriptscriptstyle 1}^{\scriptscriptstyle E}\!(y_{\scriptscriptstyle 1})\!\leq\!lpha_{\scriptscriptstyle 1}$$
 .

Suppose  $\alpha_0 < \alpha_1$ . In the first place, we shall show that there exists an open subset  $G_1$  of  $\Omega_W^*$  such that

$$\overline{E} \cap \Delta_{W}^{15} \subset G_{1}$$

and

$$R_1^{G_1 \cap \Omega}(y_1) < (\alpha_1 - \alpha_0)/4$$
.

In fact, since  $\hat{R}_1^E$  is a potential  $\overline{E} \cap \Delta_W \cap \Gamma_W = \phi^{16}$ , we have an open subset G of  $\Omega_W^*$  such that

$$\overline{E} \cap \Delta_{W} \subset G$$
 and  $\overline{G} \cap \Gamma_{W}^{(15)} = \phi$ .

 $\hat{R}_1^{G\cap\Omega}$  is a potential. Hence we have an  $\Omega_n$  such that

$$(\alpha_1 - \alpha_0)/4 > H^{\mathcal{Q}_{n-1}}_{\hat{\mathcal{R}}_1^{\mathcal{G}} \cap \Omega}(y_1) \ge H^{\mathcal{Q}_{n-1}}_{\hat{\mathcal{R}}_1^{\mathcal{G}} - \overline{\Omega}_n) \cap \Omega}(y_1) = \hat{\mathcal{R}}_1^{\mathcal{G} - \overline{\Omega}_n) \cap \Omega}(y_1) + \hat{\mathcal{R}}_1^{\mathcal{G}} - \overline{\Omega}_n \cap \Omega}(y_1) = \hat{\mathcal{R}}_1^{\mathcal{G}} - \overline{\Omega}_n \cap \Omega}(y_1) + \hat{\mathcal{R}}_1^{\mathcal{G}} - \overline{\Omega}_n \cap$$

We have  $G_1 = G - \overline{\Omega}_n$ .

On the other hand, since  $R_1^E(y_1)$  defines a capacity in the sense of Choquet<sup>17</sup>, there exists an open subset  $\omega_0$  of  $\Omega$  such that

$$E \subset \omega_0$$
 and  $R_1^{\omega_0}(y_1) \leq (\alpha_0 + \alpha_1)/2$ .

 $\omega = \omega_0 \cup G_1$  is open in  $\Omega_W^*$ ,  $\overline{E} \subset \omega$  and

$$\begin{array}{c} R_1^{\omega \cap \Omega}(y_1) \leq R_1^{\omega_0}(y_1) + R_1^{G_1 \cap \Omega}(y_1) \leq (\alpha_0 + \alpha_1)/2 + (\alpha_1 - \alpha_0)/4 \\ < \alpha_1, \end{array}$$

which contradicts the definition of  $\alpha_1$ . Hence  $\alpha_0 = \alpha_1$ , q.e.d..

**Lemma 4.** Let E be a closed subset of  $\Omega$ . If  $\hat{R}_1^E$  is a potential and  $\hat{R}_1^E(y_1) < \alpha$ , then there exists a closed non-polar subset Q of  $\Omega$  such that  $E \subset Q$ ,  $\hat{R}_1^Q$  is a continuous potential and

(2.12) 
$$R_1^Q(y_1) < \alpha$$
.

Proof. The proof is obvious if E is empty. We assume E is not empty. By Lemma 3, we have an open subset  $\omega$  of  $\Omega_W^*$  such that

$$\overline{E} \subset \omega$$
 and  $\widehat{R}_1^{\omega \cap \Omega}(y_1) < \alpha$ .

We have also an open subset  $\omega_1$  of  $\Omega^*_W$  such that

<sup>16)</sup> Cf. [7], p. 45, th. 5.6. .

<sup>17)</sup> Cf. [2], p. 122. .

$$\overline{E} \cap \Delta_W \subset \omega_1$$
,  $\omega_1 \subset \omega$  and  $\overline{\omega}_1 \cap \Gamma_W = \phi$ .

 $E-\omega_1$  is compact. Denoting by  $\omega_2$  a relatively compact neighbourhood of  $E-\omega_1$  whose closure is contained in  $\omega$  and putting

$$G = (\omega_1 \cap \Omega) \cup \omega_2$$

we have

$$\hat{R}_{1}^{G_{\Omega_{\Omega_{1}}}} \leq \hat{R}_{1}^{\omega_{1}\Omega_{\Omega_{1}}} + \hat{R}_{1}^{\omega_{2}}.$$

Since  $\overline{G} \cap \Delta_{W} \subset (\overline{\omega}_{1} \cap \Delta_{W}) \cup (\overline{\omega}_{2} \cap \Delta_{W}) = \overline{\omega}_{1} \cap \Delta_{W} \subset \Delta_{W} - \Gamma_{W}$ ,  $\hat{R}_{1}^{G \cap \Omega}$  is a potential and  $\hat{R}_{1}^{G \cap \Omega}(y_{1}) \leq \hat{R}_{1}^{\omega \cap \Omega}(y_{1}) < \alpha$ .

Next, for each point y of  $\overline{\Omega}_2 \cap E$  we assign a regular neighbourhood<sup>18</sup>) of y contained in  $G \cap \Omega_3$ . A finite number of them, say  $V_i$   $(1 \le i \le m_1)$ , covers  $\overline{\Omega}_2 \cap E$ . In general, for each point y of  $(\overline{\Omega}_{n+1} - \Omega_{n-1}) \cap E$  we assign a regular neighbourhood of y contained in  $G \cap (\Omega_{n+2} - \overline{\Omega}_{n-2})$  and cover  $(\overline{\Omega}_{n+1} - \Omega_{n-1}) \cap E$  by a finite number of them, say  $V_i(m_{n-1} + 1 \le i \le m_n)$ . Put  $Q = \bigcup_{i=1}^{\infty} V_i$ . Q is closed since  $\{V_i\}$  is locally finite. It is clear  $E \subset Q \subset G$ , therefore  $\hat{R}_1^Q$  is a potential and  $\hat{R}_1^Q(y_1) < \alpha$ . Since Q is not thin at every boundary point y of  $\Omega - Q$ , y is regular for  $\Omega - Q$  with respect to the Dirichlet problem. Thus  $\hat{R}_1^Q$  is continuous, q.e.d..

## 2.3.

By Lemma 4 we may construct a continuous potential  $p = \hat{R}_1^Q$  for  $E = \varphi^{-1}(F_{n_0})$ and  $\alpha = \delta/(8K)g^*(\Lambda)$  (see (2.9)). Put

$$V_{0} = \{ y \in \Omega; p(y) > 1 - \delta \},$$
  
$$V_{1} = \{ y \in \Omega; p(y) > 1 - 2\delta \}$$

and

$$p_1 = \min(v \circ \varphi, 1) \qquad (\text{see } (2.10))$$

 $\partial V_0 = \{x \in \Omega; p(y) = 1 - \delta\}$  and each point of  $\partial V_0$  is regular for  $\Omega - \overline{V}_0$  with respect to the Dirichlet problem.  $p_1$  is superharmonic on each component of  $\Omega - \varphi^{-1}(F_{n_0})$  and  $0 < p_1 \le 1$ .

Lemma 5.

$$s = \begin{cases} (1-\delta)/\delta & on \quad \overline{V}_{o} \\ (1-\delta)/\delta \cdot \hat{R}_{1}^{V_{0}} + (\overline{R}_{p_{1}}^{\Omega-\overline{V}_{1}})_{\Omega-\overline{V}_{0}}^{19)} & on \quad \Omega-\overline{V}_{o} \end{cases}$$

<sup>18)</sup> We use this terminology in the following sense: a neighbourhood V is regular if it is compact and its local image is a sphere. If V is regular, then both V and  $\mathcal{Q}-V$  are not thin at each point of  $\partial V$ .

<sup>19)</sup> Cf. [2], p. 82, def. 10.

is a superharmonic function on  $\Omega$ . Especially, on  $\Omega - \overline{V}_1$  we have

 $s \geq \min(v \circ \varphi, 1)$ .

Proof.  $(1-\delta)/\delta - p/\delta$  is a positive superharmonic function on  $\Omega - \bar{V}_0$  and  $\geq 1$  on  $\Omega - V_1$ . Therefore we have

$$(1-\delta)/\delta - p/\delta \ge (R_{p_1}^{\Omega-\overline{\nu}_1})_{\Omega-\overline{\nu}_0}$$
 on  $\Omega - \overline{V}_0$ .

/ .

On  $\Omega - \overline{V}_0$ , we have further

$$\begin{split} &(1-\delta)/\delta - (1-\delta)/\delta \cdot \hat{R}_{1}^{\overline{p}_{0}} - (\hat{R}_{p_{1}}^{\Omega-\overline{p_{1}}})_{\Omega-\overline{p}_{0}} \\ \geq &(1-\delta)/\delta - (1-\delta)/\delta \cdot \hat{R}_{1}^{\overline{p}_{0}} - [(1-\delta)/\delta - p/\delta] \\ \geq &(1-\delta)/\delta - (1-\delta)/\delta \cdot p/(1-\delta) - [(1-\delta)/\delta - p/\delta] = 0 , \end{split}$$

since  $p \ge 1 - \delta$  on  $\bar{V}_0$ . This means

$$(2.13) \qquad (1-\delta)/\delta \geq (1-\delta)/\delta \hat{R}_1^{\overline{Y}_0} + (\widehat{R}_{p_1}^{\Omega-\overline{Y}_1})_{\Omega-\overline{Y}_0} \qquad on \quad \Omega-\overline{V}_0$$

From the regularity of each point  $y \in \partial V_0$  for  $\Omega - \overline{V}_0$  we have

$$\lim_{z
ightarrow y}\left[(1\!-\!\delta)\!/\delta\;\hat{R}_1^{\overline{V}_0}\!(z)\!+\!(\widehat{R}_{p_1}^{\Omega-\overline{V}_1})_{\Omega-\overline{V}_0}\!(z)
ight]=(1\!-\!\delta)\!/\delta$$
 ,

so that s is continuous on  $\partial V_0$ . Combining this with (2.13), s is superharmonic. On  $\Omega - \overline{V}_1$ , we have

$$s \ge (\widehat{R_{p_1}^{\Omega-\overline{\nu}_1}})_{\Omega-\overline{\nu}_0} = p_1 = \min(v \circ \varphi, 1), \quad \text{q.e.d.}$$

2.4.

We shall put

$$A_n = \{ \ell \in \mathcal{L}'; (\lambda_n, \ell) \in \Omega - \bar{V}_1 \}$$

and

 $\widehat{A}_n = \{ (\lambda_n, l); l \in A_n \}.$ 

 $\widehat{A}_n$  is the set of points on which a Green line of  $A_n$  intersects  $\sum^{\lambda_n}$ . Correspondingly, we put

$$B_n = \{ l \in \mathcal{L}'; (\lambda_n, l) \in \overline{V}_1 \}$$

and

$$\widetilde{B}_n = \{ (\lambda_n, l); l \in B_n \} .$$

It is known that  $A_n$  (resp.  $B_n$ ) differs from an analytic set only in a set of dg-measure zero. They are dg-measurable. The difference between  $\tilde{B}_n$  and  $\bar{V}_1 \cap \sum^{\lambda_n}$  is within a set of  $d \omega_{\nu_0}^{\lambda_n}$ -measure<sup>20)</sup> zero.

20)  $\omega_{y_0}^{\lambda_n}$  denotes a harmonic measure on  $\sum_{n}^{\lambda_n}$  with respect to  $D^{\lambda_n}$  and  $y_0$ .

Lemma 6.

$$\lim_{n \to \infty} g(B_n) = 0.$$
Proof.  

$$g(B_n) = \int_{B_n} dg = \int_{\widetilde{B}_n} d\omega_{v_0}^{\lambda_n} = \int_{\Sigma^{\lambda_n} \cap \overline{\nu}_1} d\omega_{v_0}^{\lambda_n}$$

$$\leq \int_{\Sigma^{\lambda_n} \cap \overline{\nu}_1} p/(1-2\delta) d\omega_{v_0}^{\lambda_n}$$

$$\leq \int_{\Sigma^{\lambda_n}} p/(1-2\delta) d\omega_{v_0}^{\lambda_n}$$

$$\leq 1/(1-2\delta) H_p^{D_{\lambda_n}}(y_0).$$

Since p is a potential, we have  $\lim_{n\to\infty} H_p^{D\lambda_n}(y_0) = 0$ , q.e.d..

Put

$$U = \{x \in X - F_{n_0}; v(x) > 1\}$$

and

$$\Lambda' = \{\ell \in \Lambda; \, \phi(\ell) \subset A - D_{n_0}\}$$
 ,

where  $D_{n_0}$  is the domain defined in Lemma 1.  $A - \overline{D}_{n_0}$  is a polar subset of  $X^*$  and by Lemma 1 we have

(2.14) 
$$g^*(\Lambda') > 1/2g^*(\Lambda)$$
.

Lemma 7. If we put

$$C_n = \{ \ell \in \Lambda'; \varphi((\lambda_m, \ell)) \in U \quad \text{for any} \quad m \ge n \},$$

then we have

(2.15) 
$$C_1 \subset C_2 \subset \cdots$$
 and  $\bigcup_{n=1}^{\infty} C_n = \Lambda'$ 

and

(2. 16) 
$$\lim_{n \to \infty} g^*(A_n \cap C_n) = g^*(\Lambda').$$

Proof. First, we shall prove (2.15). The first part is obvious from the definition of  $C_n$ . Suppose we have  $l \in \Lambda'$  such that  $l \notin C_n$  for any *n*. Then, there should exist numbers  $\{\nu_n\}$  satisfying  $\nu_n \ge n$  and  $\varphi((\lambda_{\nu_n}, l)) \notin U$ . From

$$\lim_{x \to x'} v(x) = +\infty \quad \text{for any} \quad x' \in \hat{\varphi}(l)$$

we have an open neighbourhood W of  $\phi(l)$  in  $X^*$  such that

$$v > 2$$
 on  $W \cap X$ .

Then, we have an *n* such that  $W \supset \overline{\{\varphi((\lambda_m, l)); m \ge n\}}$ , so that

$$v[\varphi((\lambda_m, \ell))] \ge 2$$
 for any  $m \ge n$ .

On the other hand,  $v[\varphi((\lambda_m, l))] \leq 1$  for infinitely many m, which leads to a contradiction.

To prove (2. 16) we shall remark

(2.17) 
$$\lim_{n \to \infty} g^*(C_n) = g^*(\Lambda') \, .$$

This is an immediate consequence of (2.15) and the regularity of  $g^*$ . Since  $A_n$  is dg-measurable we have

$$g^{*}(C_{n}) = g^{*}(C_{n} \cap A_{n}) + g^{*}(C_{n} - A_{n}).$$

 $\mathcal{L}-A_n$  and  $B_n$  differ in a set of dg-measure zero each other, so that we have

(2.18) 
$$g^{*}(C_{n}) = g^{*}(C_{n} \cap A_{n}) + g^{*}(C_{n} \cap B_{n})$$
$$\leq g^{*}(C_{n} \cap A_{n}) + g^{*}(B_{n}).$$

Letting  $n \rightarrow \infty$  in (2.18) and in view of Lemma 6 and (2.17) we have

$$g^*(\Lambda') \leq \lim_{n \to \infty} g^*(C_n \cap A_n) \leq \overline{\lim_{n \to \infty}} g^*(C_n \cap A_n)$$
$$\leq \lim_{n \to \infty} g^*(C_n) = g^*(\Lambda'), \text{ q.e.d.}.$$

#### 3. The proof of the theorem

**3.1.** In this section, we shall give the proof of the theorem stated in §1. We consider three cases: (1)  $X \in \mathcal{H} - \mathcal{P}$  and non-compact, (2)  $X \in \mathcal{H} - \mathcal{P}$  and compact, and (3)  $X \in \mathcal{P}$ .

The proof of the case (1). Suppose, on the contrary,  $g^*(\Lambda) > 0$ . Denoting by s the function defined in Lemma 5,

$$s(y) \ge H_s^{D^{\lambda}}(y)$$
 for all  $\lambda$ .

Therefore

$$s(y) \geq \lim_{n \to \infty} H_s^{D^{\lambda_n}}(y) = u(y),$$

where u(y) is the best harmonic minorant<sup>21)</sup> of s in  $\Omega$ . It is clear that u is nonnegative. We assert u is positive and  $u(y_0) \ge g^*(\Lambda')$  (for the definition of  $\Lambda'$ , see 2.4, § 2). In fact,

$$u(y_{0}) = \lim_{n \to \infty} H^{D^{\lambda_{n}}}_{s}(y_{0}) = \lim_{n \to \infty} \int_{\Sigma^{\lambda_{n}}} sd\,\omega_{y_{0}}^{\lambda_{n}}$$
$$\geq \lim_{n \to \infty} \int_{\Sigma^{\lambda_{n}} \cap (\Omega - \overline{\psi}_{1})} \min(v \circ \varphi, 1) d\,\omega_{y_{0}}^{\lambda_{n}} \qquad (by \text{ Lemma 5})$$

21) Cf. [1], p. 434.

$$= \lim_{n \to \infty} \int_{\widetilde{A}_n} \min(v \circ \varphi, 1) d \omega_{y_0}^{\lambda_n}$$
  
$$\geq \lim_{n \to \infty} \overline{\int}_{\widetilde{A}_n \cap C_n} \min(v \circ \varphi, 1) d \omega_{y_0}^{\lambda_n}$$

where  $\widetilde{A_n \cap C_n} = \{(\lambda_n, l); l \in A_n \cap C_n\}$ . Since  $(\lambda_n, l) \in \widetilde{A_n \cap C_n}$  implies  $\varphi((\lambda_n, l)) \in U$ , so that  $v[\varphi((\lambda_n, l))] > 1$ , the last term is equal to

$$\underbrace{\lim_{n\to\infty}}_{n\to\infty}\overline{\int}_{A_n\cup C_n} d\omega_{y_0}^{\lambda_n} \\
= \underbrace{\lim_{n\to\infty}}_{n\to\infty}\overline{\int}_{A_n\cap C_n} dg = \underbrace{\lim_{n\to\infty}}_{n\to\infty} g^*(A_n\cap C_n) = g^*(\Lambda') \quad \text{(by Lemma 7)}.$$

Thus, we have  $u(y_0) \ge g^*(\Lambda')$ . From (2.8) and (2.14)

(3.1) 
$$s(y_1) \ge u(y_1) \ge 1/K u(y_0)$$
  
 $\ge 1/K g^*(\Lambda') \ge 1/(2K)g^*(\Lambda).$ 

On the other hand, in virtue of (2.12)

$$p(y_1) = \hat{R}_1^Q(y_1) < \delta/(8K)g^*(\Lambda)$$
$$\leq \delta/(8K) < 1-2\delta$$

which means  $y_1 \in \Omega - \overline{V}_1$ . Therefore

$$s(y_{1}) = (1-\delta)/\delta \cdot \hat{R}_{1}^{\overline{V}_{0}}(y_{1}) + (\hat{R}_{p_{1}}^{\Omega-\overline{v}_{1}})_{\Omega-\overline{V}_{0}}(y_{1})$$

$$\leq (1-\delta)/\delta \cdot p(y_{1})/(1-\delta) + p_{1}(y_{1})$$

$$= 1/\delta p(y_{1}) + \min(v[\varphi(y_{1})], 1)$$

$$\leq 1/\delta \cdot \delta/(8K)g^{*}(\Lambda) + 1/(4K)g^{*}(\Lambda) \quad (by (2.10))$$

$$\leq 3/(8K)g^{*}(\Lambda) .$$

This contradicts (3.1). Hence we conclude  $g^*(\Lambda) = 0$ .

## **3.2.** The proof of the case (2)

Next, we proceed to the case (2). From our assumption (\*\*\*), we have a non-polar set E each point of which is polar.

$$E - (A \cup \{\varphi(y_0)\}) \neq \phi,$$

for if  $E \subset A \cup \{\varphi(y_0)\}$ , then  $A \cup \{\varphi(y_0)\}$  is non-polar. Since A is polar, this implies  $\varphi(y_0) \in E$ , so that  $E \subset A$ , which is absurd. Let us take  $x_0 \in E - (A \cup \{\varphi(y_0)\})$ .  $\varphi^{-1}(\{x_0\})$  is a polar subset of  $\Omega$ . Let us write

$$\Omega_{0} = \Omega - \varphi^{-1}(\{x_{0}\}), X_{0} = X - \{x_{0}\}$$

and let  $\varphi_0$  denote the restriction of  $\varphi$  on  $\Omega_0$ .  $\varphi_0$  is a Fatou map of  $\Omega_0$  into  $X_0$ . In fact, since  $\varphi$  is a Fatou map, we have a closed non-polar subset F of X such that  $\hat{R}_1^{\varphi^{-1}(F)}$  is a potential.  $F_0 = F \cap X_0$  is closed and non-polar in  $X_0$ . Our assertion is derived at once from the facts  $\hat{R}_1^{\varphi^{-1}(F_0)}$  is a potential and  $X_0 \in \mathcal{H} - \mathcal{P}$ . The Green function of  $\Omega_0$  is the restriction on  $\Omega_0$  of the Green function of  $\Omega$ . Denoting by  $\Lambda_0$  the set of Green lines issuing from  $y_0$  and passing no points of  $\varphi^{-1}(\{x_0\})$ , the condition  $\hat{\varphi}(\ell) \subset A$  is reduced to

$$\phi_0(\ell) = \bigcap_{n=1}^{\infty} \overline{\{\varphi_0((\lambda_m, \ell)); m \ge n\}} \subset A$$

for any  $l \in \Lambda_0$  Thus, we can reduce the case (2) to the previous one, since a set of Green lines passing the points of  $\varphi^{-1}(\{x_0\})$  is of dg-measure zero.

## **3.3.** The proof of the case (3).

It remains to be proved the case (3), i.e.,  $X \in \mathcal{P}$ . In this case, the situation is rather simple and we can prove without resorting many lemmas.

There exists a positive superharmonic function v defined on the whole X such that

$$\lim_{x\to x'} v(x) = +\infty \quad \text{for any} \quad x' \in A.$$

 $s = v \circ \varphi$  defines a positive superharmonic function on  $\Omega$  and as before

$$u=\lim_{n\to\infty}H^{D^{\lambda_n}}_s$$

is non-negative and harmonic on  $\Omega$ .

Assuming, as in 3.1,  $g^*(\Lambda) > 0$ , let us take M > 0 such that

$$(3.2) M/2 g^*(\Lambda) > u(y_0).$$

We define

$$U = \{x \in X; v(x) > M\}$$

and

$$C_n = \{ \ell \in \Lambda; \varphi((\lambda_m, \ell)) \in U \quad \text{for} \quad m \ge n \} \quad (n = 1, 2, \cdots) .$$

Quite in the same way, we can prove

$$\lim_{n\to\infty} g^*(C_n) = g^*(\Lambda) .$$
$$u(y_0) = \lim_{n\to\infty} H_s^{D^{\lambda_n}}(y_0) = \lim_{n\to\infty} \overline{\int}_{\Sigma^n} sd\,\omega_{y_0}^{\lambda_n}$$
$$\geq \lim_{n\to\infty} \overline{\int}_{\widetilde{C}_n} sd\,\omega_{y_0}^{\lambda_n} \geq \lim_{n\to\infty} \overline{\int}_{\widetilde{C}_n} Md\,\omega_{y_0}^{\lambda_n} .$$

since  $\varphi((\lambda_n, l)) \in U$ , so that  $s = v \circ \varphi > M$  on  $C_n$ . The last term is equal to

$$\lim_{n\to\infty}\int_{C_n} Mdg = M\lim_{n\to\infty} g^*(C_n) = Mg^*(\Lambda).$$

Combining this with (3.2)

$$M/2g^*(\Lambda) \ge u(y_0) \ge Mg^*(\Lambda)$$
,

which is a contradiction. Thus, the proof is completed.

### 4. Consequences

In this section, we shall consider the case where  $\Omega$  is a hyperbolic Riemann surface and  $\varphi$  is an analytic map of  $\Omega$  into a Riemann surface X. A Fatou map in our definition is the same as in  $[5]^{22}$ . We have then

**Corollary 1.** Let R be a hyperbolic Riemann surface and  $\varphi$  be a Fatou map of R into a Riemann surface R'. If

$$\bigcap_{m=1}^{\infty} \overline{\{\varphi((\lambda_m, \ell)); m \ge n\}}$$

is polar in  $R'^*$  for every  $l \in \Lambda$ , where  $R'^*$  is an arbitrary compactification of R',  $\{\lambda_n\}$  is a decreasing sequence of positive numbers tending to zero and the outer Green measure of  $\Lambda$  is posive, then  $\varphi$  is a constant map.

Since an AD function on R (a holomorphic function with finite Dirichlet integral) is a Dirichlet map of R into a Riemann sphere<sup>23)</sup> and a Dirichlet map is a Fatou map, this is an extension of a result of M. Nakai<sup>24)</sup>.

A meromorphic function defined on |z| < 1 is a Lindelöfian map of  $R = \{|z| < 1\}$  into a Riemann sphere if and only if it is of bounded type. The theorem of Riesz-Frostman-Nevanlinna for these functions is classical. Our theorem is a generalization even in the classical case, since we have known an example of a Fatou map which is not a Lindelöfian map<sup>25)</sup>.

In [10], we have investigated the boundary behavior of harmonic functions on a Green space along Green lines. As an application, we have given there a theorem of Riesz type for holomorphic functions f in the Smirnov class (i.e.,  $\log^+ |f|$  has a quasi-bounded harmonic majorant) on a hyperbolic Riemann surface. For functions in the class AL (i.e.,  $\log^+ |f|$  has a harmonic majorant, or equivalently, f is lindelöfian) we have proved under some assumption. Now, we can remove the restriction:

**Corollary 2.** Let  $\Omega$  be a hyperbolic Riemann surface. Let  $f \in AL(\Omega)$ , that

- 24) Cf. [12], p. 19 and [16], p. 206.
- 25) Cf. [4], p. 72.

<sup>22)</sup> Cf. [5], p. 110 and [7], p. 52.

<sup>23)</sup> Cf. [5], p. 115, Folgesatz 10.3.

is, f is holomorphic on  $\Omega$  and  $\log^+ |f|$  has a harmonic majorant. If there exists a sequence  $\{\lambda_n\}$  of positive numbers tending to zero such that

$$\lim_{n\to\infty}f((\lambda_n, \ell))=0 \quad \text{for all} \quad \ell\in\alpha\,,$$

where  $\alpha$  is a set of Green lines of positive outer Green measure, then

 $f \equiv 0$ .

**OSAKA CITY UNIVERSITY** 

#### References

- [1] M. Brelot: Majorant harmoniques et principe du maximum, Archiv Math. 5 (1954), 429–440.
- [2] M. Brelot: Lectures on Potential Theory, Tata Inst, Bombay, 1960.
- [3] M. Brelot-G. Choquet: Espaces et lignes de Green, Ann. Inst. Fourier 3 (1951), 199-263.
- [4] C. Constantinescu-A. Cornea: Über das Verhalten der analytischer Abbildungen Riemannscher Flächen auf dem idealen Rand von Martin, Nagoya Math. J. 17 (1960), 1–87.
- [5] C. Constantinescu-A. Cornea: Ideale Ränder Riemannscher Flächen, Ergebnisse der Math. ihere Grenzgebiete, Neue Folge Bd. 32, Springer-Verlag, 1963.
- [6] C. Constantinescu-A. Cornea: On the axiomatic of harmonic functions I, Ann. Inst. Fourier 13 (1963), 373–388.
- [7] C. Constantinescu-A. Cornea: Compactifications of harmonic spaces, Nagoya Math. J. 25 (1965), 1-57.
- [8] O. Frostman: Potentiel d'equilibre et capacité des ensembles avec quelque applications à la théorie des fonctions, Medd. Lunds Univ. Mat. Sem. 3 (1953), 1–118.
- [9] M. Heins: Lindelöfian maps, Ann. of Math. 62 (1955), 418-446.
- [10] T. Ikegami: Dirichlet problem on Green lines related to the compactifications of Green spaces, Osaka J. Math. 8 (1971), 347-375.
- [11] N. Lusin-I. Priwalow: Sur l'unicité et la multiplicité des fonctions analytiques, Ann. Sci. École Norm. Sup. 42 (1925), 143-191.
- [12] M. Nakai: Behavior of Green lines at Royden's boundary of Riemann surfaces, Nagoya Math. J. 24 (1964), 1–27.
- [13] R. Nevanlinna: Analytic Functions, Grundlehren der Math. Wiss., Bd. 162, Springer, 1970.
- [14] I. Priwalow: Randeigenschaften Analytischer Funktionen, Deutscher Verlag der Wiss. Berlin, 1956.
- [15] S. Saks: Theory of the Integral, 1933.
- [16] L. Sario-M. Nakai: Classification Theory of Riemann Surfaces, Grundlehren der Math. Wiss., Bd. 164, Springer, 1970.