



Title	General constructions of normal numbers of Korobov type
Author(s)	Kano, Hiroyuki
Citation	Osaka Journal of Mathematics. 1993, 30(4), p. 909-919
Version Type	VoR
URL	https://doi.org/10.18910/10356
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

GENERAL CONSTRUCTIONS OF NORMAL NUMBERS OF KOROBV TYPE

HIROYUKI KANO

(Received July 20, 1992)

§1. Introduction

Let $b \geq 2$ be an integer given arbitrarily, and let $x = [x].a_1a_2\cdots = [x] + a_1b^{-1} + a_2b^{-2} + \cdots$, $a_i \in \{0, 1, \dots, b-1\}$ be the b -adic expansion of a real number x , where $[x]$ is the integral part of x . For any block $d_1\cdots d_s$ of (b -adic) digits $d_1, \dots, d_s \in \{0, 1, \dots, b-1\}$ of length S , $A(x, b, d_1\cdots d_s; N)$ denotes the number of $n (1 \leq n \leq N)$ such that $a_{n+i-1} = d_i (i=1, \dots, s)$. Then x is said to be normal to base b if $\lim_{N \rightarrow \infty} A(x, b, d_1\cdots d_s; N)/N = b^{-s}$ for any s and any block $d_1\cdots d_s$.

Normality can be defined also in terms of uniform distribution. For a set $E \subset [0, 1)$ with N elements we define the discrepancy $D(E)$ as

$$D(E) = \sup_{0 \leq u < v \leq 1} |\#(E \cap [u, v)) - N(v-u)|.$$

Further we write

$$D(N) = D(\{\{x\}, \{xb\}, \dots, \{xb^{N-1}\}\})$$

and

$$D(N, H) = D(\{\{xb^N\}, \{xb^{N+1}\}, \dots, \{xb^{N+H-1}\}\}),$$

where $\{x\} = x - [x]$. Then we note that

$$|A(x, b, d_1\cdots d_s; N) - Nb^{-s}| \leq D(N),$$

and that x is normal if and only if $D(N) = o(N)$ (cf. [6]). $\tau_b(a)$ denotes the order of an integer b mod a for any integers a and b with $(a, b) = 1$.

It is known that almost all real numbers are normal, however only few methods have been known to generate normal numbers. Among them, we mention arithmetic constructions of Stoneham and of Korobov. Historical surveys for another type of constructions of normal numbers can be found in [6], [8], and [9]. Stoneham ([10] Theorem 1) found the following normal numbers: Let a, b be relatively prime integers greater than 1, and let $\{Z_n\}_{n \geq 1}$ and $\{\alpha_n\}_{n \geq 1}$ be sequences of positive integers with $Z_n < a^n$, $(Z_n, a) = 1$, and $\lim_{n \rightarrow \infty} \alpha_n = \infty$. Assume

that $\alpha_0=Z_0=0$, and put $S(n, a)=\sum_{i=1}^n \alpha_i \tau_i(a^i)$ with $S(0, a)=0$. Then the number

$$\sum_{n=1}^{\infty} \frac{Z_n - aZ_{n-1}}{a^n b^{s(n-1, a)}} \tag{1}$$

is normal to base b . In particular

$$(a-1) \sum_{n=1}^{\infty} \frac{1}{a^n b^{((n-1)a^n - na^{n-1} + 1)/(a-1)}}$$

is normal to base b . A.N. Korobov ([3] Theorem 1) gave, independently of Stoneham, similar constructions: Let a, b be relatively prime integers greater than 1, and let $\{\lambda_n\}_{n \geq 1}$ and $\{\mu_n\}_{n \geq 1}$ be strictly increasing sequences of positive integers with $\mu_n \geq a^{\lambda_n}$. Then the number

$$\sum_{n=1}^{\infty} \frac{1}{a^{\lambda_n} b^{\mu_n}}$$

is normal to base b . Recently Wagner ([12] Theorem) constructed rings of normal numbers for the first time: Let a be an odd prime, b be an integer with $b \geq 2$ and $a \nmid b$, and let $\{\lambda_n\}_{n \geq 1}$ and $\{\mu_n\}_{n \geq 1}$ be strictly increasing sequences of positive integers with $\lim_{n \rightarrow \infty} \lambda_n / (n \mu_{n-1}) = \lim_{n \rightarrow \infty} (\log \mu_n) / \lambda_n = \infty$. Then any nonzero element of the ring generated by

$$\prod_{n=1}^{\infty} \left(1 + \frac{\epsilon_n}{a^{\lambda_n} b^{\mu_n}} \right) (\epsilon_n = \pm 1)$$

is normal to base b and nonnormal to base ab . The conditions in Wagner's theorem can be weakened, namely we need to assume only that a, b are relatively prime integers greater than 1 and $\lim_{n \rightarrow \infty} \lambda_n / \mu_{n-1} = \lim_{n \rightarrow \infty} (\log \mu_n) / \lambda_n = \infty$ (cf. [1] Theorem). More recently the author jointly with Shiokawa [2] gave rings of normal numbers of another type. To prove the normality of nonzero elements of the rings therein, we need a criterion ([2] Theorem 3) of normal numbers for numbers of the form given below, which can be rewritten as in the following: Let a, b be relatively prime integers greater than 1, and let $\{\lambda_n\}_{n \geq 1}$, $\{\mu_n\}_{n \geq 1}$, and $\{A_n\}_{n \geq 1}$ be sequences of integers such that $\lambda_{n+1} > \lambda_n$, $\mu_{n+1} > \mu_n$, $a^{\lambda_n} \leq \mu_n$ and

$$0 < |A_n| < a^{\lambda_n - \lambda_{n-1}} \tag{2}$$

for sufficiently large n . Then the number

$$x = \sum_{n=1}^{\infty} \frac{A_n}{a^{\lambda_n} b^{\mu_n}} \tag{3}$$

is normal to base b and nonnormal to base ab . In this paper, we give a wide

class of normal numbers which contains all the normal numbers of Stoneham, Korobov, Wagner, and ours mentioned above. Moreover, we shall discuss the discrepancy estimates, transcendency, irrationality measures, and non-Liouville property of these numbers. As for the proof of the normality, Stoneham's approach is considerably involved. Korobov's proof is quite different from that of Stoneham. Our method is an improvement of that of Korobov, by using Erdős-Turán inequality, as developed in [2].

§2. A class of normal numbers

Theorem 1. *Let a, b be relatively prime integers greater than 1, and let $\{\lambda_n\}_{n \geq 1}$, $\{\mu_n\}_{n \geq 1}$, and $\{A_n\}_{n \geq 1}$ be sequences of integers such that $\lambda_{n+1} > \lambda_n$, $\mu_{n+1} > \mu_n$, $a^{\lambda_n} \ll \mu_n$, and*

$$|A_n| \ll a^{\lambda_n}, a^{\lambda_n - \lambda_{n-1}} \not\asymp A_n \tag{4}$$

for sufficiently large n . Then the number defined by (3) is normal to base b and nonnormal to base ab .

REMARK 1. Putting $A_n=1$, we have Korobov's normal numbers. If $\lambda_n=n$, $\mu_n=\sum_{i=1}^{n-1} \alpha_i \tau_b(a^i)$, and $A_n=Z_n - aZ_{n-1}$, we get Stoneham's Theorem mentioned above, even without the condition $\lim_{n \rightarrow \infty} \alpha_n = \infty$ (using Lemma 1 below).

We remark that if $Z_n=1$ (n : odd), $=a^n - 1$ (n : even), $A_n=Z_n - aZ_{n-1}$ does not satisfy (2), but satisfies (4). Any nonzero element in the rings constructed by Wagner [12] and also by the author and Shiokawa [2] can be written in the form (3) with $\{A_n\}$ satisfying the conditions in Theorem 1 (cf. [1], [2]).

To prove Theorem 1 we shall need the following lemmas.

Lemma 1. (cf. [4] Lemma 1 (Remark)). *For any relatively prime integers a and b greater than one, there exists a positive integer n_0 and a rational number C such that*

$$\tau_b(a^n) = Ca^n$$

for all integers n with $n \geq n_0$.

Lemma 2. (cf. [4] Theorem 2). *Let a, b , and n_0 be as above. For any integers $n \geq n_0$ and c with $a^{n-n_0} \not\asymp c$, we have*

$$\sum_{j=1}^{\tau_b(c^n)} e^{2\pi i (cb^j/a^n)} = 0.$$

Lemma 3. (cf. [5] Lemma 2). *With the same conditions as in Lemma 2, we have for any positive integer $N \leq \tau_b(a^n)$,*

$$\left| \sum_{j=1}^N e^{2\pi i(cb^j/a^n)} \right| \ll na^{n/2}.$$

Lemma 4. (cf. [6] Theorem 2.5 (Erdős-Turán inequality)). *Let $E = \{x_1, \dots, x_N\} \in [0, 1)$. Then for any positive integer M ,*

$$D(E) \ll \frac{N}{M} + \sum_{\nu=1}^M \frac{1}{\nu} \left| \sum_{j=1}^N e^{2\pi i\nu x_j} \right|.$$

Proof of Theorem 1. We shall prove the normality. We may assume that

$$\begin{aligned} \lambda_{n-1} &> \max(\lambda_1, \dots, \lambda_{n-2}) > 2n_0, \\ \mu_{n-1} &> \max(\mu_1, \dots, \mu_{n-2}), \text{ and } a^{\lambda_n - \lambda_{n-1}} \not\in A_n \end{aligned}$$

for all $n \geq n_1$ for some n_1 . Let $n \geq n_1$. We put

$$x_n = \sum_{i=1}^n A_i a^{-\lambda_i} b^{-\mu_i} = B_n a^{-\lambda_n} b^{-\mu_n},$$

where

$$B_n = \sum_{i=1}^{n-1} A_i a^{\lambda_n - \lambda_i} b^{\mu_n - \mu_i} A_n,$$

and $\tau_m = \tau_b(a^{\lambda_m})$. For any integer $N > \mu_{n_1+1}$ we define n, h_m , and r_m by

$$\begin{aligned} \mu_n &< N \leq \mu_{n+1}, \\ \mu_{m+1} - \mu_m &= h_m \tau_m + r_m (n_1 \leq m \leq n-1, 1 \leq r_m \leq \tau_m), \end{aligned}$$

and

$$N - \mu_n = h_n \tau_n + r_n (1 \leq r_n \leq \tau_n),$$

so that we have

$$h_m < (\mu_{m+1} - \mu_m) / \tau_m (m < n), \quad h_n < (N - \mu_n) / \tau_n. \tag{5}$$

Then

$$\begin{aligned} D(N) &\ll \sum_{m=n_1}^{n-1} D(\mu_m, \mu_{m+1} - \mu_m) + D(\mu_n, N - \mu_n) + 1 \\ &\ll \sum_{m=n_1}^n \left\{ \sum_{h=0}^{h_m-1} D(\mu_m + h\tau_m, \tau_m) + D(\mu_m + h_m\tau_m, r_m) \right\} + 1. \end{aligned}$$

Here we write $e(u) = e^{2\pi iu}$ and

$$E_{\nu m h}(u, v) = \sum_{j=1}^u e(\nu v b^{\mu_m + h\tau_m + j}).$$

By Lemma 4 with $M = M_m = 2^{\lceil \lambda_{m-1} / 2 \rceil - n_0}$, we have

$$\begin{aligned} D(N) &\ll \sum_{m=n_1}^n \left[\sum_{h=0}^{h_m-1} \left\{ \tau_m / M_m + \sum_{\nu=1}^{M_m} \nu^{-1} |E_{\nu m h}(\tau_m, x)| \right\} + r_m / M_m \right. \\ &\quad \left. + \sum_{\nu=1}^{M_m} \nu^{-1} |E_{\nu m h_m}(r_m, x)| \right] + 1 \end{aligned}$$

$$\begin{aligned} &\leq \sum_{m=n_1}^n (h_m \tau_m + r_m) / M_m + \sum_{m=n_1}^n \sum_{\nu=1}^{M_m} \nu^{-1} \sum_{h=0}^{h_m-1} |E_{\nu m h}(\tau_m, x)| \\ &\quad + \sum_{m=n_1}^n \sum_{\nu=1}^{M_m} \nu^{-1} |E_{\nu m h_m}(r_m, x)| + 1 \\ &\leq \Sigma_1 + \Sigma_2 + \Sigma_3 + \Sigma_4 + \Sigma_5 + 1, \end{aligned} \tag{6}$$

where

$$\begin{aligned} \Sigma_1 &= \sum_{m=n_1}^{n-1} (h_m \tau_m + r_m) / M_m, \quad \Sigma_2 = (N - \mu_n) / M_n, \\ \Sigma_3 &= \sum_{m=n_1}^n \sum_{\nu=1}^{M_m} \nu^{-1} \sum_{h=0}^{h_m-1} |E_{\nu m h}(\tau_m, x_m)|, \\ \Sigma_4 &= \sum_{m=n_1}^n \sum_{\nu=1}^{M_m} \nu^{-1} |E_{\nu m h_m}(r_m, x_m)|, \\ \Sigma_5 &= \sum_{m=n_1}^n \sum_{\nu=1}^{M_m} \nu^{-1} \left\{ \sum_{h=0}^{h_m-1} |E_{\nu m h}(\tau_m, x) - E_{\nu m h}(\tau_m, x_m)| \right. \\ &\quad \left. + |E_{\nu m h_m}(r_m, x) - E_{\nu m h_m}(r_m, x_m)| \right\}. \end{aligned}$$

It is easily seen that

$$\Sigma_1 \ll \sum_{m=n_1}^{n-1} (\mu_{m+1} - \mu_m) 2^{-\lambda_{m-1}/2}, \tag{7}$$

$$\Sigma_2 \ll (N - \mu_n) 2^{-\lambda_{n-1}/2}. \tag{8}$$

We shall estimate Σ_3 using Lemma 2. Since $a^{\lambda_m - \lambda_{m-1}} \not\mid A_m$ and $a^{\lambda_m - \lambda_{m-1}} \mid A_i a^{\lambda_m - \lambda_i} b^{\mu_m - \mu_i} (i < m)$, we have $a^{\lambda_m - \lambda_{m-1}} \not\mid B_m$. For a prime p and an integer n , we denote by $v_p(m)$ the integer h for which $p^h \mid n$ and $p^{h+1} \nmid n$. Then there is a prime $p \mid a$ such that $\{p^{v_p(a)}\}^{\lambda_m - \lambda_{m-1}} \not\mid B_m$, so that

$$v_p(B_m) < v_p(a) (\lambda_m - \lambda_{m-1}).$$

Furthermor we have

$$v_p(\nu) \leq \log_2 M_m < \lambda_{m-1} - n_0.$$

Hence, sihce $(a, b) = 1$, we have

$$v_p(\nu B_m b^{h \tau_m + j}) < v_p(a) (\lambda_m - n_0).$$

Namely $p^{v_p(a) (\lambda_m - n_0)} \not\mid \nu B_m b^{h \tau_m + j}$, and so

$$a^{\lambda_m - n_0} \not\mid \nu B_m b^{h \tau_m + j}. \tag{9}$$

Therefore we get

$$E_{\nu m h}(\tau_m, x_m) = 0$$

by Lemma 2, which leads to

$$\Sigma_3 = 0. \tag{10}$$

Similarly we have by Lemma 3 with (9)

$$E_{\nu m h_m}(r_m, x_m) \ll \lambda_m a^{\lambda_m / 2},$$

and hence

$$\Sigma_4 \ll \sum_{m=n_1}^n \lambda_m a^{\lambda_m / 2} \log M_m \ll \lambda_n^2 a^{\lambda_n / 2}. \tag{11}$$

Finally we shall estimate Σ_5 . Since $|e(x) - e(x_m)| \ll |x - x_m| \ll b^{-\mu_{m+1}}$, we have

$$\begin{aligned} |E_{\nu m h}(\tau_m, x) - E_{\nu m h}(\tau_m, x_m)| &\leq \sum_{j=1}^{\tau_m} \nu |x - x_m| b^{\mu_m + h\tau_m + j} \\ &\ll \nu b^{\mu_m + h\tau_m - \mu_{m+1}} \sum_{j=1}^{\tau_m} b^{-j} \ll \nu \end{aligned}$$

and similarly

$$|E_{\nu m h_m}(r_m, x) - E_{\nu m h_m}(r_m, x_m)| \ll \nu.$$

Hence we obtain by Lemma 1 with (5) that

$$\begin{aligned} \Sigma_5 &\ll \sum_{m=n_1}^n M_m h_m + \sum_{m=n_1}^n M_m \\ &\ll \sum_{m=n_1}^{n-1} (\mu_{m+1} - \mu_m) 2^{-(\lambda_m - \lambda_{m-1} / 2)} + (N - \mu_n) 2^{-(\lambda_n - \lambda_{n-1} / 2)} + 2^{\lambda_{n-1} / 2}. \end{aligned} \tag{12}$$

Therefore it follows from (6), (7), (8), (10), (11), and (12) that

$$D(N) \ll \sum_{m=n_1}^{n-1} (\mu_{m+1} - \mu_m) 2^{-\lambda_m - 1/2} + (N - \mu_n) 2^{-\lambda_n - 1/2} + \lambda_n^2 a^{\lambda_n / 2}. \tag{13}$$

Here the first term is $o(\sum_{m=n_1}^{n-1} (\mu_{m+1} - \mu_m)) = o(\mu_n) = o(N)$, and the second term is also $o(N)$. Furthermore, since $a^{\lambda_n} \ll \mu_n$, the third term is $o(N)$. Therefore we obtain $D(N) = o(N)$, and the normality to base b is proved. Nonnormality to base ab can be proved similarly as in the proof of Theorem 2 in [2].

REMARK 2. We can estimate as in [2] the discrepancy of normal numbers in Theorem 1. Let x be given in Theorem 1. Then for any positive integer N with $\mu_n < N \leq \mu_{n+1}$ and any block $d_1 \cdots d_s \in \{0, 1, \dots, b-1\}^s$ we have by (13)

$$\begin{aligned} &\left| A(x, b, d_1 \cdots d_s; N) - \frac{N}{b^s} \right| (\ll D(N)) \\ &\ll \sum_{m=2}^{n-1} \frac{\mu_{m+1} - \mu_m}{(\sqrt{2})^{\lambda_{m-1}}} + \frac{N - \mu_n}{(\sqrt{2})^{\lambda_{n-1}}} + \lambda_n^2 a^{\lambda_n / 2}. \end{aligned}$$

On the other hand, if s is sufficiently large, we have for any $\varepsilon < 0$

$$(D(N) \gg) \left| A(x, b, d_1 \cdots d_s; N) - \frac{N}{b^s} \right| \gg \frac{\mu_{n+1} - (1 + \varepsilon)\mu_n}{a^{\lambda_n}}$$

for infinitely many n . This can be proved similarly as in [2] Theorem 1.

EXAMPLE 1. As these estimates are implicit, we give here an example: Under the same conditions as in Theorem 1, assume further that $\lim_{n \rightarrow \infty} \lambda_{n+1}/\lambda_n = \alpha$ and $\mu_n = f(\lambda_n)\beta^{\lambda_n}$ where $f(x)$ is any polynomial with $f(m) \in N$ ($m \in N$). Then for any $\varepsilon > 0$

$$D(N) \ll N^{1 - (\log \beta^2)/2\alpha^2 + \varepsilon}$$

and, if s is sufficiently large,

$$|A(x, b, d_1 \dots d_s; N) - Nb^{-s}| \gg N^{1 - (\log \beta \alpha)/\alpha^2 - \varepsilon}$$

for infinitely many n .

§3. Irrationality measure and transcendency

Stoneham ([10] Theorem 2) proved that the normal number (1) is non-Liouville and transcendental, if there exist positive constants δ and β such that $\delta < \alpha_n \tau_b(a^n) / S(n-1, a) < \beta$. And it is remarked in [2] that

$$\sum_{n=1}^{\infty} \frac{1}{a^n b^{c^n}}$$

($a, b, c \in N; a, b \geq 2; (a, b) = 1; c \geq a$) has the same properties. In this section we shall give a class of non-Liouville normal numbers containing these examples.

Theorem 2. *Make the same assumptions as in Theorem 1. Put $c = \overline{\lim}_{n \rightarrow \infty} \mu_{n+1} / \mu_n$ and $d = \underline{\lim}_{n \rightarrow \infty} \mu_{n+1} / \mu_n$. Assume that $c < \infty$ and $d > 1$, then x is non-Liouville and transcendental. More precisely for any $\varepsilon > 0$, we have*

$$\left| x - \frac{P}{Q} \right| \gg \frac{1}{Q^{\max(c, 1 + c/(d-1)) + \varepsilon}} \tag{14}$$

for all integers P, Q (≥ 1), and

$$\left| x - \frac{P}{Q} \right| \ll \frac{1}{Q^{c - \varepsilon}} \tag{15}$$

for infinitely many integers P, Q (≥ 1) with $(P, Q) = 1$. On the other hand if $c = \infty$, x is a Liouville number.

REMARK 3. Putting $\mu_n = S(n-1, a) = \sum_{i=1}^{n-1} a_i \tau_b(a^i)$ in Theorem 2, we have the Stoneham's result ([10] Theorem 2) mentioned above.

EXAMPLE 2. Let a, b, c be integers greater than 1 with $(a, b) = 1$ and $c \geq a$.

Then, for any polynomial $f(x)$ with $f(m) \in N$ ($m \in N$), the number

$$x = \sum_{n=1}^{\infty} \frac{1}{a^n b^{f(n)c^n}}$$

is normal to base b , nonnormal to base ab , non-Liouville, and transcendental. Furthermore we have for any $\varepsilon > 0$,

$$|x - P/Q| \gg Q^{-\max(c,3) - \varepsilon}$$

for all integers $P, Q (\geq 1)$, and

$$|x - P/Q| \ll Q^{-c + \varepsilon}$$

for infinitely many integers $P, Q (\geq 1)$ with $(P, Q) = 1$. In this sense Theorem 2, is the best possible.

To prove Theorem 2 we need the following lemmas.

Lemma 5 ([11] Lemma 5). *Let θ be real. Suppose that there exist sequences of integers $\{p_n\}$ and $\{q_n\}$ with $\lim_{n \rightarrow \infty} q_n = \infty$, $q_n \leq \kappa_1 q_{n-1}^\gamma$ for some constant $\kappa_1 > 0$ and $\gamma > 1$ such that for any $\delta > 0$*

$$\left| \theta - \frac{p_n}{q_n} \right| < \frac{\kappa_2}{q_n^{\xi - \delta}}$$

for some constants $\kappa_2 > 0$ and $\xi > 1$. Then for any $\varepsilon > 0$

$$\left| \theta - \frac{x}{y} \right| > \frac{\kappa_3}{y^{1 + \gamma / (\xi - 1) + \varepsilon}}$$

holds for all integers $x, y (\geq 1)$ with $x/y \notin \{p_n/q_n\}$ ($n \in N$) and for some constant $\kappa_3 > 0$ independent of n .

For completeness we give here the proof of this lemma.

Proof. Let y be large, and let $n = n(y)$ the least positive integer such that

$$|q_n \theta - p_n| \leq 1/(2y)$$

for all $m \leq n$. Let x/y ($x, y \in N$) be a rational number with $x/y \notin \{p_n/q_n\}$ ($n \in N$). Then

$$\begin{aligned} |\theta - x/y| &\geq |p_n/q_n - x/y| - |\theta - p_n/q_n| \\ &\geq 1/(yq_n) - 1/(2yq_n) = 1/(2yq_n). \end{aligned}$$

By the minimality of n , we get

$$1/(2y) < |q_{n-1} \theta - p_{n-1}| < \kappa_2 / q_{n-1}^{\xi - \delta},$$

which implies $2\kappa_2 y > q_n^{\xi-1-\delta}$, so that

$$q_{n-1} < (2\kappa_2 y)^{1/(\xi-1-\delta)},$$

for any $\delta > 0$. Hence

$$q_n \leq \kappa_1 q_{n-1}^\gamma < \kappa_4 y^{\gamma/(\xi-1)+\varepsilon}$$

for any $\varepsilon > 0$ and a constant $\kappa_4 > 0$. Therefore, we obtain

$$|\theta - x/y| \geq 1/(2yq_n) > \kappa_3/y^{1+\gamma/(\xi-1)+\varepsilon}$$

for any $\varepsilon > 0$ and a constant $\kappa_2 > 0$.

Lemma 6 (cf. [7] §5 p. 427). *Let θ be any real number. If there exist some constant $\kappa > 1$ and infinite sequence $\{p_n/q_n\}$ ($n \in \mathbb{N}$, $p_n \in \mathbb{Z}$, $q_n \in \mathbb{N}$) such that $q_n = q'_n q''_n$ where each of q'_n and q''_n is a power of an integer independent of n , $q_n < q_{n+1}$,*

$$0 < |\theta - p_n/q_n| < q_n^{-\kappa},$$

$$\overline{\lim}_{n \rightarrow \infty} \log q_{n+1}/\log q_n < \infty, \text{ and } \lim_{n \rightarrow \infty} \log q'_n/\log q_n = 0,$$

then θ is transcendental.

Proof of Theorem 2. We write

$$x_n = \sum_{i=1}^n A_i a^{-\lambda_i} b^{-\mu_i} = B_n a^{-\lambda_n} b^{-\mu_n} = p_n/q_n,$$

where $p_n = B_n$, $q_n = a^{\lambda_n} b^{\mu_n}$, and set $q'_n = a^{\lambda_n}$, $q''_n = b^{\mu_n}$. Then for sufficiently large n we have

$$q_n < q_{n+1}, \quad q_{n+1} \leq a^{\lambda_{n+1}} b^{(c+\varepsilon/2)\mu_n} \leq q_n^{c+\varepsilon}, \tag{16}$$

and

$$|x - p_n/q_n| = \sum_{i=n+1}^{\infty} A_i a^{-\lambda_i} b^{-\mu_i} \ll b^{-\mu_{n+1}}. \tag{17}$$

Here for any $\varepsilon > 0$ it follows that

$$\log b^{\mu_{n+1}}/\log q_n \geq \mu_{n+1}/\mu_n - \varepsilon/2 \geq d - \varepsilon.$$

By (17) we have

$$|x - p_n/q_n| \ll q_n^{-(d-\varepsilon)} \tag{18}$$

for all n and

$$|x - p_n/q_n| \ll q_n^{-(c-\varepsilon)} \tag{19}$$

for infinitely many n . (15) follows from the last inequality. Since $d > 1$, we get for $P/Q \in \{p_n/q_n\}$ ($P \in \mathbb{Z}$, $Q \in \mathbb{N}$),

$$|x - P/Q| \gg Q^{-(1+c/(d-1)+\varepsilon)}, \tag{20}$$

using Lemma 5. On the other hand, since

$$|x - p_n/q_n| \gg q_{n+1}^{-1}$$

we get by (16)

$$|x - p_n/q_n| \gg q_n^{-(c+\varepsilon)},$$

which together with (20) yields (14). The transcendency follows from Lemma 6 with (16), (18), and

$$\log q'_n / \log q_n \ll \lambda_n / \mu_n \rightarrow 0 \quad (n \rightarrow \infty).$$

This completes the proof of Theorem 2.

REMARK 4. If $c = \overline{\lim}_{n \rightarrow \infty} \mu_{n+1} / \mu_n > 2$ in Theorem 2, x is transcendental by Roth's Theorem and (19), without the condition $\underline{\lim}_{n \rightarrow \infty} \mu_{n+1} / \mu_n > 1$. In particular, $c > 2$ if $a \geq 3$.

I would like to express my thanks to Professors Iekata Shiokawa and Jun'ichi Tamura for their valuable advices concerning the paper.

References

- [1] H. Kano: *A remark on Wagner's ring of normal numbers*, Arch. Math. **60** (1993), 46–50.
- [2] H. Kano and I. Shiokawa: *Rings of normal and nonnormal numbers*, to appear in Israel J. Math.
- [3] A.N. Korobov: *Continued fractions of certain normal numbers*, Math. Notes **47** (1990) No.2, 128–132 = Mat. Zamet. **47** (1990), No. 2, 28–33.
- [4] N.M. Korobov: *Trigonometric sums with exponential functions and the distribution of signs in repeating decimals*, Math. Notes **8** (1970), No.5, 831–837. = Mat. Zamet. **8** (1970), No.5, 641–652.
- [5] N.M. Korobov: *On the distribution of digits in periodic fractions*, Math. USSR Sb. **18** (1972), No.4, 659–676 = Mat. Sb. **89** (131) (1972), No.4, 654–670.
- [6] L. Kuipers and H. Niederreiter: *Uniform Distribution of Sequences*, J. Wiley and Sons 1974.
- [7] K. Mahler: *Arithmetische Eigenschaften einer Klasse von Dezimalbrüchen*, Nederl. Akad. Wetensch. Proc., Ser. A, **40** (1937), 421–428.
- [8] Y. Nakai and I. Shiokawa: *A class of normal numbers*, Japan. J. Math. Vol. **16**, No. 1 (1990), 17–29.
- [9] I. Shiokawa: *Asymptotic distributions of digits in integers*, Colloquia Mathematica Societatis János Bolyai **51**, Number Theory, 1987, 505–525.
- [10] R.G. Stoneham: *A general arithmetic construction of transcendental non-Liouville normal numbers from rational fractions*, Acta Arith. **XVI** (1970), 239–253. Errata to the paper in Acta Arith. **XVII** (1971).
- [11] J. Tamura: *Irrationality measure of g -adic fractions with digits defined by dis-*

tributions, preprint.

- [12] G. Wagner: *On rings of normal and nonnormal numbers*, preprint.

Department of Mathematics
Faculty of Science and Technology
Keio University
Hiyoshi 3-14-1, Kohoku-ku,
Yokohama-shi, 223 Japan

