

Title	On multiply transitive groups. VIII
Author(s)	Oyama, Tuyosi
Citation	Osaka Journal of Mathematics. 1969, 6(2), p. 315-319
Version Type	VoR
URL	<a href="https://doi.org/10.18910/10490">https://doi.org/10.18910/10490</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## ON MULTIPLY TRANSITIVE GROUPS VIII

TUYOSI OYAMA

(Received May 6, 1969)

### 1. Introduction

Let  $G$  be a 4-fold transitive group on  $\Omega = \{1, 2, \dots, n\}$ , and let  $P$  be a Sylow 2-subgroup of a stabilizer of four points in  $G$ . By a theorem of M. Hall [2, Theorem 5.8.1] and a lemma of E. Witt [7, Theorem 9.4], we have that  $P$  fixes exactly four, five, six, seven or eleven points and the normalizer of  $P$  in  $G$  restricted on the set of the fixed points of  $P$  is  $S_4$ ,  $S_5$ ,  $A_6$ ,  $A_7$  or  $M_{11}$ . (cf. H. Nagao and T. Oyama [5], Lemma 1).

The purpose of this paper is to prove the following

**Theorem.** *Let  $G$  be a 4-fold transitive group. If a Sylow 2-subgroup of a stabilizer of four points in  $G$  fixes exactly six points, then  $G$  must be  $A_6$ .*

The above theorem of M. Hall is that if a stabilizer of four points in  $G$  is of odd order then  $G$  must be one of the following groups:  $S_4$ ,  $S_5$ ,  $A_6$ ,  $A_7$  or  $M_{11}$ . Therefore to prove our theorem we may assume that a Sylow 2-subgroup of a stabilizer of four points in  $G$  is not identity.

### 2. Definitions and notations

A permutation  $x$  is called semi-regular if there is no point fixed by  $x$ . A permutation group  $G$  is called semi-regular if every non-identity element of  $G$  is semi-regular on the points actually moved by  $G$ .

For a permutation group  $G$  on  $\Omega$  the subgroup of  $G$  consisting of all the elements fixing the points  $i, j, \dots, k$  of  $\Omega$  will be denoted by  $G_{i,j,\dots,k}$ , which we shall call the stabilizer of the points  $i, j, \dots, k$ . The totality of points left fixed by a subset  $X$  of  $G$  will be denoted by  $I(X)$ , and if a subset  $\Delta$  of  $\Omega$  is a fixed block of  $X$ , then the restriction of  $X$  on  $\Delta$  will be denoted by  $X^\Delta$ . A  $G$ -orbit of minimal length is called a minimal orbit of  $G$ .

For subsets  $X, Y$  of a group  $G$ ,  $\langle X, Y \rangle$  is the subgroup of  $G$  generated by the elements of  $X$  and  $Y$ , and  $N_G(X)$  is the normalizer of  $X$  in  $G$ .

### 3. Proof of the theorem

In the following two lemmas we assume that  $G$  is a 4-fold transitive group

on  $\Omega = \{1, 2, \dots, n\}$ , and  $P$  is a Sylow 2-subgroup of  $G_{1234}$ . For a point  $t$  of a minimal orbit of  $P$  in  $\Omega - I(P)$  let  $P_t = Q$ ,  $N_G(Q) = N$  and  $I(Q) = \Delta$ .

**Lemma 1.** *Let  $R$  be a Sylow 2-subgroup of  $N_{ijkl}$  for  $\{i, j, k, l\} \subseteq \Delta$ . Then  $R^\Delta$ , which is a Sylow 2-subgroup of  $(N^\Delta)_{ijkl}$ , is semi-regular and  $|I(R)| = |I(P)|$ .*

*Proof.* Let  $\varphi$  be a natural homomorphism of  $N_{ijkl}$  onto  $(N^\Delta)_{ijkl}$ . Then  $\varphi(R) = R^\Delta$ . Since  $R$  is a Sylow 2-subgroup of  $N_{ijkl}$ ,  $R^\Delta$  is a Sylow 2-subgroup of  $(N^\Delta)_{ijkl}$ .

Let  $P'$  be a Sylow 2-subgroup of  $G_{ijkl}$  containing  $R$ . Since  $Q$  is a normal 2-subgroup of  $N_{ijkl}$  and  $R$  is a Sylow 2-subgroup of  $N_{ijkl}$ ,  $P' \geq R \geq Q$  and  $I(P') \subseteq I(R) \subseteq \Delta$ . Since  $G$  is 4-fold transitive,  $P$  and  $P'$  are conjugate, and  $|I(P)| = |I(P')|$ . Therefore  $I(P')$  is a proper subset of  $\Delta$ . For any point  $r$  of  $\Delta - I(P')$ ,  $P'_r \geq Q = P_t$ . From the assumption that  $t$  belongs to a minimal  $P$ -orbit we have

$$|P' : P'_r| = |r^{P'}| \geq |t^P| = |P : P_t| = |P' : Q|.$$

Therefore  $P'_r = Q$  and  $R_r = Q$ . Thus  $R^\Delta$  is identity or semi-regular and  $I(R^\Delta) = I(P')$ . Since  $\Delta = I(Q) \supseteq I(P')$  and  $P'_\Delta = Q$ ,  $N_{P'}(Q)^\Delta$  is a non-identity 2-group of  $(N^\Delta)_{ijkl}$ . Therefore  $R^\Delta$ , which is a Sylow 2-subgroup of  $(N^\Delta)_{ijkl}$ , is semi-regular and  $|I(R)| = |I(R^\Delta)| = |I(P')| = |I(P)|$ .

**Lemma 2.** *In Lemma 1 if  $|I(P)| = 6$ , that is  $N_G(P)^{I(P)} = A_6$ , then  $R^\Delta$  is an elementary abelian group,  $N_N(R)^{I(R)} \leq A_6$  and  $|\Delta| \geq 10$ .*

*Proof.* If  $R$  has an element  $x = (i' j' k' l') \dots$ , where  $\{i', j', k', l'\} \subset \Delta$ , then  $x^\Delta$  has no 2-cycle by Lemma 1. On the other hand,  $x$  normalizes  $G_{i'j'k'l'}$  and  $Q$ . Therefore  $x$  normalizes some Sylow 2-subgroup  $P'$  of  $G_{i'j'k'l'}$  containing  $Q$ . From the assumption  $x^{I(P')} \in A_6$ . Hence  $x^{I(P')}$  has a 2-cycle. Since  $I(P') \subset \Delta$ ,  $x^\Delta$  has a 2-cycle, which is a contradiction. Thus  $R^\Delta$  is elementary abelian.

From Lemma 1  $|I(R)| = 6$ . If  $N_N(R)^{I(R)} \not\leq A_6$ , then  $N_N(R)$  has a 2-element  $x$  such that  $x^{I(R)}$  is an odd permutation. On the other hand  $x$  normalizes some Sylow 2-subgroup  $P''$  of  $G_{I(R)}$ . Since  $G_{I(R)}$  contains a Sylow 2-subgroup of  $G_{ijkl}$ ,  $P''$  is a Sylow 2-subgroup of some stabilizer of four points in  $G$ , and  $I(P'') = I(R)$ . Then  $x^{I(R)} = x^{I(P'')} \in N_G(P'')^{I(P'')} = A_6$ , which is a contradiction. Thus  $N_N(R)^{I(R)} \leq A_6$ .

Since  $|I(R)| = 6$  and  $\Delta \supseteq I(R)$ ,  $|\Delta| \geq 8$ . Suppose that  $|\Delta| = 8$ . Let  $\Delta = \{i, j, k, l, r, s, u, v\}$  and  $I(R) = \{i, j, k, l, r, s\}$ . Then  $R$  has the following 2-element

$$a = (i)(j)(k)(l)(r)(s)(uv) \dots$$

Since  $a$  normalizes  $G_{ijuv}$  and  $Q$ ,  $a$  normalizes a Sylow 2-subgroup  $P'''$  of  $G_{ijuv}$  containing  $Q$ . It follows from  $I(P''') \subset \Delta$  that  $a^{I(P''')}$  is a transposition. This is a contradiction since  $N_G(P''')^{I(P''')} = A_6$ . Therefore  $|\Delta| \geq 10$ .

From now on we consider a permutation group  $G$  on  $\Omega = \{1, 2, \dots, n\}$ , which is not necessarily 4-fold transitive. In the following two lemmas we assume that  $G$  satisfies the following condition.

- (\*) *Let  $P$  be a Sylow 2-subgroup of any stabilizer of four points in  $G$ . Then*
  - (i)  *$P$  is semi-regular and elementary abelian,*
  - (ii)  *$|\Omega| \geq 10$  and  $|I(P)| = 6$ ,*
  - (iii)  *$N_G(P)^{I(P)} \leq A_6$ .*

By Lemma 1 and Lemma 2,  $N^\Delta$  satisfies the assumption (\*). In Lemma 4 we shall show that there is no group satisfying (\*). Thus if  $G$  is a 4-fold transitive group as in Theorem, then  $G_{1234}$  is of odd order, and hence  $G$  must be  $A_6$ .

**Lemma 3.**  *$G$  is a doubly transitive group and  $N_G(P)^{I(P)}$  is  $A_6$  or  $A_6^*$ , where  $A_6^*$  is a doubly transitive group of degree 6 isomorphic to  $A_6$ .*

Proof. For any two points  $i_1$  and  $i_2$  let  $P'$  be a Sylow 2-subgroup of  $G_{i_1 i_2 i_3 i_4}$ , where  $\{i_3, i_4\} \subset \Omega - \{i_1, i_2\}$ . Then we have an involution  $a$  of  $P'$ , which has the following form

$$a = (i_1)(i_2)(i_3)(i_4)(i_5)(i_6)(j_1 j_2) \dots$$

where  $I(P') = \{i_1, i_2, \dots, i_6\}$ . Since  $a$  normalizes  $G_{i_1 i_2 j_1 j_2}$ ,  $a$  normalizes a Sylow 2-subgroup  $P''$  of  $G_{i_1 i_2 j_1 j_2}$ . From (iii)

$$a^{I(P'')} = (i_1)(i_2)(j_1 j_2)(j_3 j_4),$$

where  $I(P'') = \{i_1, i_2, j_1, j_2, j_3, j_4\}$ . Hence  $\langle a, P'' \rangle$  is a 2-group and fixes exactly two points  $i_1$  and  $i_2$ . From a lemma of D. Livingstone and A. Wagner [4, Lemma 6]  $G$  is a doubly transitive group on  $\Omega$ .

Since  $G$  is doubly transitive, for any two points  $u$  and  $v$  of  $I(P)$  there is a Sylow 2-subgroup  $S$  of  $G_{uv}$ , which contains  $P$  and fixes only two points  $u$  and  $v$ . Let  $T = N_S(P)$ . Then  $T \cong P$  and  $I(S) \subseteq I(T) \subseteq I(P)$ . Suppose that  $I(S) \subsetneq I(T)$ , then  $|I(T)| = 4$  or  $6$ . By (iii)  $|I(T)| = 6$ . Therefore  $I(T) = I(P)$ . Since  $P$  is a Sylow 2-subgroup of  $G_{I(P)}$ ,  $T = P$ , which is a contradiction. Therefore  $I(T) = I(S) = \{u, v\}$ . This shows that for any two points  $u, v$  of  $I(P)$   $N_G(P)^{I(P)}$  contains a 2-group, which fixes only two points  $u$  and  $v$ . Using also Lemma 6 of [3] we have that  $N_G(P)^{I(P)}$  is doubly transitive. Therefore, by assumption (iii),  $N_G(P)^{I(P)}$  is either  $A_6$  or  $A_6^*$ , where  $A_6^*$  is a doubly transitive group of degree 6 isomorphic to  $A_5$  (see B. Huppert [3] II, 4.7 Satz).

To prove the next lemma we need the following result in [6]: Let  $G$  be a 4-fold transitive group. If a Sylow 2-subgroup  $P$  of a stabilizer of four points in  $G$  is semi-regular and not identity, then  $|I(P)| \neq 6$ .

**Lemma 4.** *There is no group satisfying (\*).*

Proof. Assume that  $G$  satisfies (\*). We may assume that  $P$  is a Sylow

2-subgroup of  $G_{1234}$  and  $I(P)=\{1, 2, \dots, 6\}$ . In the case  $N_G(P)^{I(P)}=A_6^*$ , we may assume that  $A_6^*$  is generated by  $\{(3\ 4)(5\ 6), (1\ 2\ 3)(4\ 5\ 6), (1\ 3\ 5\ 6\ 4)\}$  (see W. Burnside [1] §166).

For any point  $i$  of  $\Omega-\{1, 2, 3\}$  let  $P'$  be a Sylow 2-subgroup of  $G_{123i}$ . By (ii)  $P'$  fixes six points, say  $1, 2, 3, i, j$  and  $k$ . Let  $Q$  be a Sylow 2-subgroup of  $G_{123}$  containing  $P'$ . Since  $|\Omega|$  is even,  $|I(Q)|$  is also even. Therefore  $Q$  fixes at least four points, and hence  $Q$  fixes six points by (ii), which are the points of  $I(P')$ . Thus  $Q=P'$  is a Sylow 2-subgroup of  $G_{123}$ , and any point in  $\Omega-\{1, 2, 3\}$  belongs to a  $G_{123}$ -orbit of odd length. On the other hand  $P$  and  $P'$  are conjugate in  $G_{123}$ , hence  $G_{123}$  has an element taking  $\{4, 5, 6\}$  into  $\{i, j, k\}$ . Thus  $G_{123}$  has exactly one or three orbits in  $\Omega-\{1, 2, 3\}$ . If  $G_{123}$  has three orbits in  $\Omega-\{1, 2, 3\}$ , then three points 4, 5 and 6 belong to different  $G_{123}$ -orbits, say  $T_4, T_5$  and  $T_6$  respectively.

Suppose that  $G_{123}$  is transitive on  $\Omega-\{1, 2, 3\}$ . Since a Sylow 2-subgroup of  $G_{12}$  fixes only two points 1 and 2,  $G_{12}$  has an element taking 3 into some point of  $\Omega-\{1, 2, 3\}$ . Therefore  $G_{12}$  is transitive on  $\Omega-\{1, 2\}$ . It follows from Lemma 3 that  $G$  is 4-fold transitive on  $\Omega$ . But this contradicts the theorem in [6].

From now on we assume that  $G_{123}$  has three orbits  $T_4, T_5$  and  $T_6$  in  $\Omega-\{1, 2, 3\}$ . Suppose that  $N_G(P)^{I(P)}=A_6$ . Then  $N_G(P)$  contains an element  $x$  of the form

$$x = (1)(2)(3)(4\ 5\ 6)\dots$$

Since  $x \in G_{123}$ , 4, 5 and 6 belong to the same  $G_{123}$ -orbit, which is a contradiction.

Thus we have that  $N_G(P')^{I(P')}=A_6^*$ , for any Sylow 2-subgroup  $P'$  of an arbitrary stabilizer of four points in  $G$ .

Suppose that  $P$  has two involutions  $x$  and  $y$ . Since  $P$  is elementary abelian by (i), we may assume by (ii) that  $x$  and  $y$  are of the following forms

$$\begin{aligned} x &= (1)(2)\dots(6)(ij)(kl)\dots, \\ y &= (1)(2)\dots(6)(ik)(jl)\dots. \end{aligned}$$

$\langle x, y \rangle$  normalizes some Sylow 2-subgroup of  $G_{ijkl}$ . Hence the restriction of  $\langle x, y \rangle$  on the set of the points fixed by this Sylow 2-subgroup is a four group and fixes two points. But a stabilizer of two points in  $A_6^*$  is of order 2, which is a contradiction. Therefore  $P$  is of order 2, and any Sylow 2-subgroup of a stabilizer of four points in  $G$  is also of order 2.

Let  $a$  be an involution of  $P$ . We may assume by (ii) that  $a$  is of the form

$$a = (1)(2)\dots(6)(ij)\dots$$

Then  $a$  normalizes a Sylow 2-subgroup of  $G_{12ij}$ , and hence  $a$  commutes with some involution  $b$  in  $G_{12ij}$ . Since  $b$  fixes only six points,  $b^{I(a)}$  is not identity. Since  $N_G(P)^{I(P)}=A_6^*$ ,  $b$  must be of the form

$$b = (1) (2) (3\ 4) (5\ 6) (i) (j) (k) (l) \dots,$$

and then  $a$  have a 2-cycle  $(k\ l)$ . If  $a$  and  $b$  have two 2-cycles  $(i'\ j')$   $(k'\ l')$  and  $(i'\ k')$   $(j'\ l')$  respectively, then  $\langle a, b \rangle$  normalizes some Sylow 2-subgroup of  $G_{i'j'k'l'}$ . Using the same argument as above we have a contradiction. Therefore if  $a$  has 2-cycles in  $\Omega - \{1, 2, \dots, 6, i, j, k, l\}$ , then  $b$  has the same 2-cycles. Since  $a$  commutes with  $b$ ,  $ab$  is also an involution, and fixes two points 1 and 2. Therefore  $|I(ab)| = 2$  or 6 by (ii). If  $|I(ab)| = 2$ , then

$$\begin{aligned} a &= (1) (2) \dots (6) (i\ j) (k\ l), \\ b &= (1) (2) (3\ 4) (5\ 6) (i) (j) (k) (l), \end{aligned}$$

and hence  $|\Omega| = 10$ . If  $|I(ab)| = 6$ , then

$$\begin{aligned} a &= (1) (2) \dots (6) (i\ j) (k\ l) (i'\ j') (k'\ l'), \\ b &= (1) (2) (3\ 4) (5\ 6) (i) (j) (k) (l) (i'\ j') (k'\ l'), \end{aligned}$$

and hence  $|\Omega| = 14$ . On the other hand from the assumption that  $A_6^*$  has an element  $(1\ 2\ 3) (4\ 5\ 6)$ , there is an element

$$c = (1\ 2\ 3) (4\ 5\ 6) \dots$$

in  $N_G(P)$ . Then  $c$  normalizes  $G_{123}$ , and since the  $G_{123}$ -orbits in  $\Omega - \{1, 2, 3\}$  are  $T_4, T_5$  and  $T_6$ ,  $c$  takes  $T_4$  into  $T_5$ ,  $T_5$  into  $T_6$  and  $T_6$  into  $T_4$ . Therefore  $T_4, T_5$  and  $T_6$  are of the same length and  $|\Omega - \{1, 2, 3\}|$  is divisible by 3. But  $|\Omega| - 3 = 7$  or 11, which is not divisible by 3. This contradiction arises from the first assumption that there is a group which satisfies the conditions (i), (ii) and (iii).

Now by Lemma 2 and 4 we have that, for a 4-fold transitive group  $G$  as in the theorem, the stabilizer of four points in  $G$  is of odd order. Therefore  $G$  must be  $A_6$ .

YAMAGUCHI UNIVERSITY

**References**

[1] W. Burnside: *Theory of Groups of Finite Order*, Second edition, Cambridge Univ. Press, 1911.  
 [2] M. Hall: *The Theory of Groups*, Macmillan, New York, 1959.  
 [3] B. Huppert: *Endliche Gruppen I*, Springer, Berlin-Heidelberg-New York, 1967.  
 [4] D. Livingstone and A. Wagner: *Transitivity of finite permutation groups on unordered sets*. *Math. Z.* **90** (1965), 393-403.  
 [5] H. Nagao and T. Oyama: *On multiply transitive groups II*, *Osaka J. Math.* **2** (1965), 129-136.  
 [6] T. Oyama: *On multiply transitive groups VII*, *Osaka J. Math.* **5** (1968), 155-164.  
 [7] H. Wielandt: *Finite Permutation Groups*, Academic Press, New York, 1964.

