

Title	The Cauchy problem and the martingale problem for integro-differential operators with non-smooth kernels
Author(s)	Abels, Helmut; Kassmann, Moritz
Citation	Osaka Journal of Mathematics. 2009, 46(3), p. 661-683
Version Type	VoR
URL	https://doi.org/10.18910/10499
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

THE CAUCHY PROBLEM AND THE MARTINGALE PROBLEM FOR INTEGRO-DIFFERENTIAL OPERATORS WITH NON-SMOOTH KERNELS

HELMUT ABELS* and MORITZ KASSMANN†

(Received September 3, 2007, revised April 15, 2008)

Abstract

We consider the linear integro-differential operator L defined by

$$Lu(x) = \int_{\mathbb{R}^n} (u(x+y) - u(x) - \mathbb{1}_{[1,2]}(\alpha) \mathbb{1}_{\{|y|\leq 2\}}(y)y \cdot \nabla u(x))k(x, y) dy.$$

Here the kernel $k(x, y)$ behaves like $|y|^{-n-\alpha}$, $\alpha \in (0, 2)$, for small y and is Hölder-continuous in the first variable, precise definitions are given below. We study the unique solvability of the Cauchy problem corresponding to L . As an application we obtain well-posedness of the martingale problem for L . Our strategy follows the classical path of Stroock-Varadhan. The assumptions allow for cases that have not been dealt with so far.

1. Introduction

A linear operator $A: C_0^2(\mathbb{R}^n) \rightarrow C(\mathbb{R}^n)$ is said to satisfy the global maximum principle if $Au(x^*) \leq 0$ for all $x^* \in \{x \in \mathbb{R}^n; u(x) \geq u(y), \forall y \in \mathbb{R}^n\}$. It is well-known that infinitesimal generators of strongly continuous contraction semi-groups on $C_0(\mathbb{R}^n)$ generating Markov processes satisfy the global maximum principle. Surprisingly, the global maximum principle implies already a certain structure of A , see [12]. More precisely, A is the sum of a possibly degenerate elliptic diffusion operator with bounded coefficients, a drift and a jump part which we call L . Since L alone generates pure jump processes which generalize Lévy processes it is sometimes called a Lévy-type operator, see [19], [5], [17] and [22] for surveys.

It is the aim of this work to study important properties of the operator L which is defined by

$$(1.1) \quad Lu(x) = \int_{\mathbb{R}^n} (u(x+y) - u(x) - \mathbb{1}_{B_2}(y)y \cdot \nabla u(x))k(x, y) dy$$

2000 Mathematics Subject Classification. Primary 47G20; Secondary 47G30, 60J75, 60J35, 60G07, 35K99, 35B65, 47A60.

*Partially supported by DFG (German Science Foundation) through SFB 611.

†Partially supported by DFG (German Science Foundation) through SFB 611.

if $1 \leq \alpha < 2$ and

$$(1.2) \quad Lu(x) = \int_{\mathbb{R}^n} (u(x+y) - u(x))k(x, y) dy$$

if $0 < \alpha < 1$. Here $k: \mathbb{R}^n \times (\mathbb{R}^n \setminus \{0\}) \rightarrow (0, \infty)$ is Hölder continuous of order $\tau \in (0, 1)$ in $x \in \mathbb{R}^n$, measurable in $y \in \mathbb{R}^n \setminus \{0\}$ and can be decomposed as $k = k_1 + k_2$ such that $k_1(x, y) = 0$ for $|y| \geq 2$, k_1 is $(n + 1)$ -times differentiable in y , and the following estimates are satisfied:

$$(1.3) \quad \|\partial_y^\beta k_1(\cdot, y)\|_{C^\tau(\mathbb{R}^n)} \leq C|y|^{-n-\alpha-|\beta|}, \quad 0 < |y| \leq 2,$$

$$(1.4) \quad k_1(x, y) \leq c|y|^{-n-\alpha}, \quad 0 < |y| \leq 1, \quad x \in \mathbb{R}^n,$$

$$(1.5) \quad \|k_2(\cdot, y)\|_{C^\tau(\mathbb{R}^n)} \leq C|y|^{-n-\alpha'}, \quad 0 < |y| \leq 1,$$

$$(1.6) \quad \int_{|y| \geq 1} \|k_2(\cdot, y)\|_{C^\tau(\mathbb{R}^n)} dy < \infty,$$

$$(1.7) \quad \lim_{|y| \rightarrow \infty} \|k_2(\cdot, y)\|_{C^\tau(\mathbb{R}^n)} = 0$$

for all $\beta \in \mathbb{N}_0^n$ with $|\beta| \leq N := n + 1$, where $0 \leq \alpha' < \alpha < 2$. Moreover, we assume $k_1(x, -y) = k_1(x, y)$ if $\alpha = 1$. There are many examples satisfying these assumptions, see the discussion below. A model case is given by $k(x, y) = c|y|^{-n-\alpha}$, $y \neq 0$, which leads to $L = -(-\Delta)^{\alpha/2}$. Other examples are given by $k(x, y) = g(x, y)|y|^{-n-\alpha}$, $y \neq 0$, if g is sufficiently smooth, positive and bounded from above and away from zero. Note that g does not need to be homogeneous in y nor in x .

Our main result concerning the Cauchy-Problem for L is given by the following theorem. In the following $C^s(\mathbb{R}^n)$, $s > 0$, denotes the Hölder-Zygmund space and $C_0^s(\mathbb{R}^n) = \overline{C_0^\infty(\mathbb{R}^n)}^{|\cdot|, \| \cdot \|_{C^s}}$. For a precise definition of the function spaces we refer to Section 2.1 below.

Theorem 1.1. *Let k satisfy (1.3)–(1.5), let L be defined as in (1.1), and let $T > 0$, $0 < s < \tau$, $0 < \theta < 1$. Then for every $f \in C^\theta([0, T]; C_0^s(\mathbb{R}^n))$ with $f(0) = 0$ there is a unique $u \in C^{1,\theta}([0, T]; C_0^s(\mathbb{R}^n)) \cap C^\theta([0, T]; C_0^{s+\alpha}(\mathbb{R}^n))$ solving*

$$(1.8) \quad \partial_t u - Lu = f \quad \text{in } (0, T) \times \mathbb{R}^n,$$

$$(1.9) \quad u(0, \cdot) = 0 \quad \text{in } \mathbb{R}^n.$$

If f is non-negative, then u is non-negative as well.

The latter theorem will be a direct consequence of the fact that L generates an analytic semi-group on $C_0^s(\mathbb{R}^n)$ with $0 < s < \tau$. In order to prove this we will construct an approximate resolvent to L using pseudodifferential operators with non-smooth symbols.

Further down we formulate and prove an important corollary to the above theorem. It involves the martingale problem which we briefly review. By $\mathcal{D}([0, \infty); \mathbb{R}^n)$ we denote the space of all càdlàg paths. A probability measure \mathbb{P}^μ on $\mathcal{D}([0, \infty); \mathbb{R}^n)$ is said to be a solution to the martingale problem for $(L, D(L))$ with domain $D(L)$ being contained in the set of bounded functions $f: \mathbb{R}^n \rightarrow \mathbb{R}$, L defined as in (1.1) and μ a probability measure on \mathbb{R}^n if, for any $\phi \in D(L)$

$$\left(\phi(\Pi_t) - \phi(\Pi_0) - \int_0^t (L\phi)(\Pi_s) ds \right)_{t \geq 0}$$

is a \mathbb{P}^μ -martingale with respect to the filtration $(\sigma(\Pi_s; s \leq t))_{t \geq 0}$ and $\mathbb{P}^\mu(\Pi_0 = \mu) = 1$. Here Π is the usual coordinate process, i.e., $\Pi: [0, \infty) \times \mathcal{D}([0, \infty); \mathbb{R}^n) \rightarrow \mathbb{R}^n$, $\Pi_t(\omega) = \omega(t)$. If for every μ there is a unique solution \mathbb{P}^μ of the martingale problem, we say that the martingale problem for $(L, D(L))$ is well-posed.

As a corollary to Theorem 1.1 we obtain the following result.

Theorem 1.2. *Let L be defined as above. Then the martingale problem for $(L, C_0^\infty(\mathbb{R}^n))$ is well-posed.*

Proof. The existence of a solution \mathbb{P}^μ for a given distribution μ on \mathbb{R}^n has been established by several authors, see Theorem 2.2 in [40], Theorem IX.2.31 in [20] and Theorem 3.2 in [15]. Note that these papers establish existence for a class which is much larger than the class for which uniqueness is shown. Because of these results, it is sufficient for us to prove uniqueness. Theorem 1.1 and more precisely, Corollary 2.17 below, provide a bounded analytic semi-group $(T_t)_{t \geq 0}$ on $C_0^s(\mathbb{R}^n)$ for any $s \in (0, \tau)$ with generator $(L, C_0^{s+\alpha}(\mathbb{R}^n))$. In particular, this implies that $R(\lambda - L) = C_0^s(\mathbb{R}^n)$ is dense in $C_0(\mathbb{R}^n)$ for all $\lambda > 0$ and the condition of the Hille-Yosida theorem for Feller semi-groups, cf. [21, Theorem 17.11] are satisfied, where the global maximum principle is easily verified. Thus L is a closable operator on $C_0(\mathbb{R}^n)$ such that \bar{L} generates a Feller semi-group. Uniqueness of the martingale problem now follows from [14, Chapter IV, Theorem 4.1]. □

Studying the existence of pure jump processes, i.e., processes without a diffusion component, together with their properties is a field of still increasing interest. We list some references dealing with the martingale problem for non-local operators such as L . In the case $k(x, y) = k(y)$ with k as in (1.1) L is a generator of a Lévy jump process, i.e., a jump process with independent stationary increments. There are different and more elegant approaches than the martingale problem to the existence of a corresponding process, see [7], [38].

The martingale problem for an operator of the form $A + L$ where A is a non-degenerate elliptic operator and L is an operator of our type has been studied first in [27], [40], [30]. Since A is a second order operator L is a lower order perturbation of

A for many questions. [28], [29] seem to be the first articles treating the martingale problem for pure jump processes generated by operators like L . The main assumptions are that $k(x, y)$ is a perturbation of $\tilde{k}(x, y) = |y|^{-d-\alpha}$, $y \neq 0$, together with quite strong regularity assumptions. More general results have been obtained in [35] using techniques from partial differential equations. In the latter article $k(x, y)$ is assumed to be twice continuously differentiable in the first variable.

Strong results on the well-posedness have been obtained in [32], [33], [34]. The authors use a setup similar to the one of the so called Calderon-Zygmund approach in the theory of partial differential equations. In [32], [33] $k(x, y)$ is assumed to be only continuous in the first variable but some additional homogeneity is assumed in the second variable. To add a personal comment, these results have been underestimated in the literature from our point of view. This is maybe due to the fact that the journal is not available easily and that the articles are written in a somewhat dense style.

Using pseudodifferential operators and anisotropic Sobolev spaces built with continuous negative definite functions [15] proves well-posedness of the martingale problem under assumptions like $x \mapsto k(x, y) \in C^{3n}(\mathbb{R}^n)$ but allowing for a more general dependence of $k(x, y)$ on y . Moreover, the extension of L to a generator of a Feller semi-group is discussed. See [10] for similar techniques in infinite dimensions and [36] for related questions. In the setting of [15] a parametrix for the pseudodifferential operator is constructed in [11]. These results do not apply to our setting since we assume only Hölder regularity of the mapping $x \mapsto k(x, y)$.

The results of [35], [32], [33], [15] and the ones in the present work do not imply one another but have a large region of intersection. The assumptions on the x -dependence of $k(x, y)$ in [35], [15], [34] are more restrictive but the assumptions on the y -dependence are partly weaker than ours. The situation is reversed when comparing our results to [32], [33]. Our techniques solving the Cauchy problem are different from [35], [15] and [32].

The authors of [13] prove solvability of the Cauchy problem for a time dependent pseudodifferential operator $L(t) = p(t, x, D_x)$ where the principal part of the symbol $p(t, x, \xi)$ is homogeneous in ξ of degree $\alpha \in [1, 2]$ and uniformly Hölder continuous in (t, x) . Their results do not apply to the uniqueness for solutions of the martingale problem since sufficient regularity of solutions to the Cauchy problem is not provided.

In the above list we do not mention results concerning what is sometimes called “stable-like” cases, i.e. when $k(x, y) \approx |y|^{-d-\alpha(x)}$, $y \neq 0$. Well-posedness of the martingale problem is proved in one spatial dimension in [4] when $\alpha(\cdot)$ is Dini-continuous. Uniqueness problems for stochastic differential equations in similar situations but including higher dimensions and also diffusion coefficients are considered in [45]. The techniques of [4] can be extended to higher dimensions and to a larger class of problems, see to a larger class of problems, see [6]. See [18], [25] for results on the question when the linear operators of type L extend to generators of Feller processes in the case when the y -singularity of $k(x, y)$ is of variable order. [16] provides such a result together with well-posedness of the martingale problem when $x \mapsto \alpha(x)$ is smooth

where $\alpha(x)$ is the order of differentiability of L .

One scope of this contribution is to present an application of the theory of pseudo-differential operators with non-smooth coefficients to jump processes. We hope to draw the attention of probabilists to this method.

2. The Cauchy problem for Lévy-type operators

2.1. Preliminaries and notation. The characteristic function of a set A is denoted by $\mathbb{1}_A$. Furthermore, we define $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$ for $\xi \in \mathbb{R}^n$. Moreover, we define $\Sigma_\delta := \{z \in \mathbb{C} \setminus \{0\} : |\arg z| < \delta\}$ for $0 < \delta \leq \pi$.

As usual, $C_0^\infty(\mathbb{R}^n)$ denotes the set of all smooth and compactly supported functions $f: \mathbb{R}^n \rightarrow \mathbb{R}$, $\mathcal{S}(\mathbb{R}^n)$ denotes the space of all smooth and rapidly decreasing functions, and $\mathcal{S}'(\mathbb{R}^n) = (\mathcal{S}(\mathbb{R}^n))'$ the space of tempered distributions. $C^k(\mathbb{R}^n)$, $k \in \mathbb{N}$, shall be the usual Banach space of continuous functions with bounded continuous derivatives up to order k . By $C_0^k(\mathbb{R}^n)$ we denote the closure of $C_0^\infty(\mathbb{R}^n)$ with respect to the norm of $C^k(\mathbb{R}^n)$. $C^s(M; X)$, where $s \in (0, 1)$, $M \subseteq \mathbb{R}^n$, M closed, and X is a Banach space, is the space of uniformly bounded Hölder continuous functions $f: M \rightarrow X$ of order s with uniformly bounded Hölder constant. Moreover, $C^s(M) = C^s(M; \mathbb{R})$ and $f \in C^{1,s}([0, T]; X)$ iff $f: [0, T] \rightarrow X$ is continuously differentiable and $(d/dt)f \in C^s([0, T]; X)$. Finally, if $f: \mathbb{R}^n \rightarrow \mathbb{R}$, we define $(\tau_h f)(x) = f(x + h)$, $x, h \in \mathbb{R}^n$, and $\Delta_h f = \tau_h f - f$.

For functions $f \in \mathcal{S}(\mathbb{R}^n)$ the Fourier transform \mathcal{F} and its inverse \mathcal{F}^{-1} are defined via

$$\mathcal{F}(f)(\xi) = \int e^{-ix \cdot \xi} f(x) dx, \quad \mathcal{F}^{-1}(f)(x) = \int e^{ix \cdot \xi} f(\xi) d\xi,$$

where $d\xi = (2\pi)^{-n} d\xi$. When there is ambiguity we use subscripts to indicate the variables with respect to which the Fourier transform is taken, i.e., $\mathcal{F}(f)$ would be written as $\mathcal{F}_{x \mapsto \xi}(f)$. Finally, $\mathcal{F}: \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ is defined by duality and $D_{x_j} := (1/i)\partial_{x_j}$, $j = 1, \dots, n$, where ∂_{x_j} is the usual partial derivative. D_x denotes the vector $(D_{x_1}, \dots, D_{x_n})$.

We use a dyadic partition of unity $\varphi_j \in C_0^\infty(\mathbb{R}^n)$, $j \in \mathbb{N}_0$, which satisfies $\text{supp } \varphi_0 \subset B_2(0)$ and $\text{supp } \varphi_j \subset \{2^{j-1} \leq |\xi| \leq 2^{j+1}\}$ for $j \in \mathbb{N}$. Then the Hölder-Zygmund space $\mathcal{C}^s(\mathbb{R}^n)$, $s > 0$, consists of all $f \in \mathcal{S}'(\mathbb{R}^n)$ satisfying

$$\|f\|_{\mathcal{C}^s} = \sup\{2^{ks} \|\varphi_k(D_x)f\|_{L^\infty} : k \in \mathbb{N}_0\} < \infty,$$

where

$$\varphi_k(D_x)f = \mathcal{F}^{-1}[\varphi_k(\xi)\mathcal{F}[f](\xi)].$$

Note that $\mathcal{C}^s(\mathbb{R}^n) = B_{\infty\infty}^s(\mathbb{R}^n)$, where $B_{pq}^s(\mathbb{R}^n)$, $s \in \mathbb{R}$, $1 \leq p, q \leq \infty$, denotes the usual Besov space. Moreover, it is well-known that $\mathcal{C}^s(\mathbb{R}^n) = C^s(\mathbb{R}^n)$ for $s \in \mathbb{R}_+ \setminus \mathbb{N}$, cf. [42, Appendix A] or Triebel [44, Section 2.7].

The closure of $C_0^\infty(\mathbb{R}^n)$ in $C^s(\mathbb{R}^n)$ is denoted by $C_0^s(\mathbb{R}^n)$. We will use the following sufficient criterion for a function to belong to $C_0^s(\mathbb{R}^n)$:

Proposition 2.1. *Let $0 < s < s' < 1$. Then every $f \in C^{s'}(\mathbb{R}^n)$ satisfying*

$$(2.1) \quad \lim_{R \rightarrow \infty} \|f\|_{C^s(\mathbb{R}^n \setminus B_R(0))} = 0$$

belongs to $C_0^s(\mathbb{R}^n)$.

Proof. Let $\varphi_\varepsilon(x) = \varepsilon^{-n} \varphi(\varepsilon^{-1}x)$, $\varphi \in C_0^\infty(\mathbb{R}^n)$ with $\int \varphi(x) dx = 1$, be a standard mollifier. Then $\varphi_\varepsilon * f \xrightarrow{\varepsilon \rightarrow 0} f$ in $C^s(\mathbb{R}^n)$ since $f \in C^{s'}(\mathbb{R}^n)$. Moreover, (2.1) implies that each $\varphi_\varepsilon * f$ can be approximated by smooth, compactly supported functions up to an arbitrarily small error in $C^s(\mathbb{R}^n)$. This proves the proposition. \square

2.2. Pseudodifferential operators with non-smooth symbols. In the following, the principal part of the Lévy-type operator will be represented as pseudodifferential operator with a symbol of the following kind:

DEFINITION 2.2. Let $n, n' \in \mathbb{N}$, $N \in \mathbb{N}_0$, $m \in \mathbb{R}$, and let $\tau \in (0, 1)$. Then a function $p: \mathbb{R}^{n'} \times \mathbb{R}^n \rightarrow \mathbb{C}$ belongs to $C^\tau S_{1,0;N}^m(\mathbb{R}^{n'}; \mathbb{R}^n)$ if $p(x, \xi)$ is Hölder continuous w.r.t. $x \in \mathbb{R}^{n'}$, N -times continuously differentiable w.r.t. $\xi \in \mathbb{R}^n$ and satisfies

$$(2.2) \quad \|\partial_\xi^\beta p(\cdot, \xi)\|_{C^\tau(\mathbb{R}^n)} \leq C \langle \xi \rangle^{m-|\beta|}$$

uniformly in $\xi \in \mathbb{R}^n$ and for all $|\beta| \leq N$. Moreover, let

$$\|p\|_{C^\tau S_{1,0;N}^m} := \sup_{\xi \in \mathbb{R}^n, |\beta| \leq N} \langle \xi \rangle^{-m+|\beta|} \|\partial_\xi^\beta p(\cdot, \xi)\|_{C^\tau(\mathbb{R}^n)}.$$

REMARK 2.3. Note that $\bigcap_{\tau > 0, N \in \mathbb{N}} C^\tau S_{1,0;N}^m(\mathbb{R}^n; \mathbb{R}^n)$ coincides with the classical symbol class $S_{1,0}^m(\mathbb{R}^n; \mathbb{R}^n)$ as defined in [23]. A first treatment of pseudodifferential symbols which are merely Hölder continuous in the space variable x and the associated operators was done by Kumano-go and Nagase [24]. Further results and many references can be found in the monographs by Taylor [42, 43].

For $a = a(x, y, \xi) \in C^\tau S_{1,0;N}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$ we define the associated *pseudodifferential operator in (x, y) -form* (formally) by

$$(2.3) \quad a(x, D_x, x)f := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(x-y)\cdot \xi} a(x, y, \xi) f(y) dy d\xi.$$

So far, it is not clear whether $a(x, D_x, x)f$ in (2.3) is well-defined even for $f \in C_0^\infty(\mathbb{R}^n)$. This will be clarified later in each particular situation we have to deal with.

REMARK 2.4. In order to underline the connection between the operator $a(x, D_x, x)$ and the corresponding symbol $a(x, y, \xi)$ we write $a(x, \xi, y)$ instead of $a(x, y, \xi)$ in the sequel.

In the special case that $a(x, \xi, y) = p(x, \xi)$, $p \in S^m_{1,0,N}(\mathbb{R}^n; \mathbb{R}^n)$, and $f \in \mathcal{S}(\mathbb{R}^n)$, the operator in (2.3) is well-defined as iterated integrals and coincides with

$$p(x, D_x)f = \int_{\mathbb{R}^n} e^{ix \cdot \xi} p(x, \xi) \hat{f}(\xi) d\xi,$$

which is a pseudodifferential operator in x -form. The adjoints of x -form pseudodifferential operators are the pseudodifferential operators in y -form, which corresponds to the case $a(x, \xi, y) = p(y, \xi)$, $p \in S^m_{1,0,N}(\mathbb{R}^n; \mathbb{R}^n)$, and is (formally) given by

$$p(D_x, x)f := \mathcal{F}^{-1} \left[\int_{\mathbb{R}^n} e^{-iy \cdot \xi} p(y, \xi) f(y) dy \right].$$

If $f \in \mathcal{S}(\mathbb{R}^n)$, the inner integral defines a bounded continuous function in $\xi \in \mathbb{R}^n$ and $p(D_x, x)$ is a well-defined operator $p(D_x, x): \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$.

REMARK 2.5. Working with non-smooth symbols it is important to distinguish between pseudodifferential operators in x -form and in y -form since the mapping properties are different, cf. Theorem 2.6 below. The principal part of the operator L will be a pseudodifferential operator in x -form; but it is important to take the approximate resolvent $Q_\lambda = q_\lambda(D_x, x) \approx (\lambda - L)^{-1}$ as an operator in y -form, not in x -form. Otherwise the mapping properties of Q_λ would not fit to $(\lambda - L)^{-1}: C^s_0(\mathbb{R}^n) \rightarrow C^{s+\alpha}_0(\mathbb{R}^n)$ for $0 < s < \tau$. This technique was already successfully applied to the resolvent equation of the Stokes operator in suitable domains with non-smooth boundary, cf. [3, 1]. An alternative way for a parametrix construction is described in [2, Section 6], where the operator is first reduced to a zero order operator and then the parametrix is constructed in x -form. The latter article deals with pseudodifferential *boundary value problems*; but the construction also applies to pseudodifferential equations on \mathbb{R}^n .

Mapping properties of pseudodifferential operators with non-smooth coefficients have been studied by several authors starting with the pioneering work of Kumano-go and Nagase [24], cf. Taylor [42, 43] and the references given there. For our purposes we will use the following theorem, which is a consequence of the results by Marschall [31].

Theorem 2.6. *Let $N > n/2$, $\tau \in (0, 1)$, and let $p \in C^\tau S_{1,0,N}^m(\mathbb{R}^n; \mathbb{R}^n)$. Then*

$$(2.4) \quad p(x, D_x): C_0^{s+m}(\mathbb{R}^n) \rightarrow C^s(\mathbb{R}^n) \quad \text{if } 0 < s < \tau, s + m > 0$$

and

$$(2.5) \quad p(D_x, x): C_0^{s+m}(\mathbb{R}^n) \rightarrow C^s(\mathbb{R}^n) \quad \text{if } s > 0, 0 < s + m < \tau$$

are bounded operators. Moreover, the operator norms can be estimated by $C \|p\|_{C^\tau S_{1,0,N}^m}$, where C is independent of $p \in C^\tau S_{1,0,N}^m(\mathbb{R}^n; \mathbb{R}^n)$.

REMARK 2.7. Note that for an operator $p(x, D_x)$ in x -form the order of the range space C^s is limited by the smoothness of the symbol in x . For the corresponding operator in y -form, $p(D_x, x)$, the order of the domain C_0^{s+m} is limited by τ .

Proof of Theorem 2.6. First of all, we note that the symbol class $C^\tau S_{1,0,N}^m(\mathbb{R}^n; \mathbb{R}^n)$ coincides with the symbol class $S_{1,0}^m(\tau, N)$ defined in [31]. Moreover, if $f \in \mathcal{S}(\mathbb{R}^n)$, then $p(x, D_x)f$ defined as above coincides with the definition in [31] as a limit of operators obtained from a symbol decomposition, cf. proof of [31, Proposition 2.4]. Hence [31, Proposition 2.4] implies that

$$\|p(x, D_x)f\|_{C^s(\mathbb{R}^n)} \leq C \|f\|_{C^{s+m}(\mathbb{R}^n)}$$

for $f \in \mathcal{S}(\mathbb{R}^n)$ provided that $0 < s < \tau$ and $s + m > 0$.

By our definition of $p(D_x, x): \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$

$$\langle p(D_x, x)f, g \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{ix \cdot \xi} p(y, \xi) f(y) dy \hat{g}(-\xi) d\xi = \int_{\mathbb{R}^n} f(x) q(x, D_x)g dx$$

for all $f, g \in \mathcal{S}(\mathbb{R}^n)$ with $q(x, \xi) = p(x, -\xi)$. Because of [31, Proposition 4.3], $q(x, D_x)^*: C^{s+m}(\mathbb{R}^n) \rightarrow C^s(\mathbb{R}^n)$ provided that $0 < s + m < \tau$ and $s > 0$.

Finally, it is easy to observe that all estimates done in the proof of [31, Proposition 4.3] are uniform for all $p \in C^\tau S_{1,0,N}^m(\mathbb{R}^n; \mathbb{R}^n)$ with $\|p\|_{C^\tau S_{1,0,N}^m} \leq 1$, which is nothing but the boundedness of the linear mapping from the symbol space $C^\tau S_{1,0,N}^m(\mathbb{R}^n; \mathbb{R}^n)$ into the corresponding space of linear operators. □

The next important ingredient are kernel estimates of the Schwartz kernel associated to a pseudodifferential operator. We follow the presentation given in [39, Chapter 6, Paragraph 4]. Given $a \in C^\tau S_{1,0,N}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$ we define for $j \in \mathbb{N}_0$

$$k_j(x, y, z) := \mathcal{F}_{\xi \mapsto z}^{-1}[a_j(x, \cdot, y)], \quad a_j(x, \xi, y) := a(x, \xi, y)\varphi_j(\xi),$$

where φ_j is the Dyadic partition of unity introduced above.

First of all, we have

Lemma 2.8. *Let $a \in C^\tau S^m_{1,0;N}(\mathbb{R}^n \times \mathbb{R}^n)$, $m \in \mathbb{R}$, $N \in \mathbb{N}_0$, $\tau \in (0, 1)$, and let $k_j(x, y, z)$ be defined as above. Then*

$$(2.6) \quad \|\partial_z^\alpha k_j(\dots, z)\|_{C^\tau(\mathbb{R}^n \times \mathbb{R}^n)} \leq C_{\alpha, M} \|a\|_{C^\tau S^m_{1,0;N}} |z|^{-M} 2^{j(n+m-M+|\alpha|)}$$

for all $\alpha \in \mathbb{N}_0^n$, $M = 0, \dots, N$, where $C_{\alpha, M}$ does not depend on $j \in \mathbb{N}_0$ and $a \in C^\tau S^m_{1,0;N}(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$.

Proof. We start with

$$z^\gamma D_z^\alpha k_j(x, y, z) = \int_{\mathbb{R}^n} e^{iz \cdot \xi} D_\xi^\gamma [\xi^\alpha a_j(x, \xi, y)] d\xi$$

for all $\alpha, \gamma \in \mathbb{N}_0^n$. We estimate the integral on the right hand side from above. Firstly, the integrand is supported in the ball $\{|\xi| \leq 2^{j+1}\}$, which has volume bounded by a multiple of 2^{nj} . Secondly, since the support is also limited by the condition $2^{j-1} \leq |\xi|$ (when $j \neq 0$) and $c2^j \leq \langle \xi \rangle \leq C2^j$ on $\{2^{j-1} \leq |\xi| \leq 2^{j+1}\}$,

$$|D_\xi^\gamma [\xi^\alpha a_j(x, \xi, y)]| \leq C_{\alpha, \gamma} \|a\|_{C^\tau S^m_{1,0;N}} 2^{j(m+|\alpha|-|\gamma|)}$$

due to the symbol estimates of $\xi^\alpha a_j(x, \xi, y) \in C^\tau S^{m+|\alpha|}_{1,0;N}(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$. Hence

$$\sup_{x, y \in \mathbb{R}^n} |z^\gamma D_z^\alpha k_j(x, y, z)| \leq C_{\alpha, \gamma} \|a\|_{C^\tau S^m_{1,0;N}} 2^{j(n+m+|\alpha|-M)}, \quad \text{whenever } |\gamma| = M.$$

Taking the supremum over all γ with $|\gamma| = M$, gives (2.6) with $C^\tau(\mathbb{R}^n \times \mathbb{R}^n)$ replaced by $C^0(\mathbb{R}^n \times \mathbb{R}^n)$. In order to get the same for $C^\tau(\mathbb{R}^n \times \mathbb{R}^n)$ one simply replaces $a_j(x, \xi, y)$ and $k_j(x, y, z)$ by $a_j(x, \xi, y) - a_j(x', \xi, y')$ and $k_j(x, y, z) - k_j(x', y', z)$, resp., in the estimates above and uses that

$$\begin{aligned} & |D_\xi^\gamma [\xi^\alpha (a_j(x, \xi, y) - a_j(x', \xi, y'))]| \\ & \leq C_{\alpha, \gamma} \|a\|_{C^\tau S^m_{1,0;N}} 2^{j(m+|\alpha|-|\gamma|)} (|x - x'| + |y - y'|)^\tau. \end{aligned}$$

This finishes the proof. □

Using the latter lemma, we are able to prove the following kernel estimate:

Theorem 2.9. *Let $a \in C^\tau S^m_{1,0;N}(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$, $\tau \in (0, 1)$, $m > -n$, and $N \in \mathbb{N}_0$ such that $N > n+m$ and let k_j be defined as above. Then for every $x, y, z \in \mathbb{R}^n$, $z \neq 0$,*

$$k(x, y, z) := \sum_{j=0}^{\infty} k_j(x, y, z)$$

exists, converges uniformly in $x, y \in \mathbb{R}^n$, $|z| \geq \varepsilon > 0$, and satisfies

$$\|\partial_z^\alpha k(\cdot, \cdot, z)\|_{C^\tau(\mathbb{R}^n \times \mathbb{R}^n)} \leq \begin{cases} C_\alpha \|a\|_{C^\tau S_{1,0;N}^m} |z|^{-n-m-|\alpha|} & \text{for } |z| \leq 1, \\ C_\alpha \|a\|_{C^\tau S_{1,0;N}^m} |z|^{-N} & \text{for } |z| \geq 1, \end{cases}$$

uniformly in $z \neq 0$ for all $\alpha \in \mathbb{N}_0$ with $|\alpha| < N - n - m$, where C is independent of $a \in C^\tau S_{1,0;N}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$.

Proof. First we consider the case when $0 < |z| \leq 1$. We brake the above sum into two parts: the first where $2^j \leq |z|^{-1}$, the second where $2^j > |z|^{-1}$. In order to estimate the first sum we use (2.6) with $M = 0$:

$$\sum_{2^j \leq |z|^{-1}} \|\partial_z^\alpha k_j(\cdot, \cdot, z)\|_{C^\tau(\mathbb{R}^n \times \mathbb{R}^n)} \leq C \|a\|_{C^\tau S_{1,0;N}^m} \sum_{2^j \leq |z|^{-1}} 2^{j(n+m+|\alpha|)},$$

where

$$\sum_{2^j \leq |z|^{-1}} 2^{j(n+m+|\alpha|)} = O(|z|^{-n-m-|\alpha|})$$

since $n + m + |\alpha| > 0$.

Next, for the second sum, we use again (2.6) with $M = N$ and get the estimate

$$\begin{aligned} \sum_{2^j > |z|^{-1}} \|\partial_z^\alpha k_j(\cdot, \cdot, z)\|_{C^\tau(\mathbb{R}^n \times \mathbb{R}^n)} &\leq C_\alpha \|a\|_{C^\tau S_{1,0;N}^m} |z|^{-M} \sum_{2^j > |z|^{-1}} 2^{j(n+m+|\alpha|-M)} \\ &\leq C'_\alpha \|a\|_{C^\tau S_{1,0;N}^m} |z|^{-n-m-|\alpha|}. \end{aligned}$$

Finally, we consider the situation $|z| \geq 1$. Since $N > n + m + |\alpha|$, (2.6) shows that

$$\begin{aligned} \sum_{j=0}^\infty \|\partial_z^\alpha k_j(\cdot, \cdot, z)\|_{C^\tau(\mathbb{R}^n \times \mathbb{R}^n)} &\leq C_\alpha |z|^{-N} \|a\|_{C^\tau S_{1,0;N}^m} \sum_{j=0}^\infty 2^{j(n+m-N+|\alpha|)} \\ &\leq C'_\alpha \|a\|_{C^\tau S_{1,0;N}^m} |z|^{-N}. \end{aligned}$$

Hence the proof is complete. □

The following corollary shows that (2.4) can be improved to $p(x, D_x): C_0^{s+m}(\mathbb{R}^n) \rightarrow C_0^s(\mathbb{R}^n)$ under the same assumptions.

Corollary 2.10. *Let $N > n + m$, $\tau \in (0, 1)$, let $p \in C^\tau S_{1,0;N}^m(\mathbb{R}^n; \mathbb{R}^n)$, and let $f \in C_0^\infty(\mathbb{R}^n)$. Then $p(x, D_x)f \in C_0^s(\mathbb{R}^n)$ for all $0 < s < \tau$ with $s + m > 0$ and $p(D_x, x)f \in C_0^s(\mathbb{R}^n)$ provided that $0 < s + m < \tau$ and $s > 0$.*

Proof. For simplicity we only treat the case of the operator in x -form. The other case is treated in the same way.

Fix $0 < s < \tau$ with $s + m > 0$ and choose $s' \in (s, \tau)$. Then $p(x, D_x)f \in \mathcal{C}^{s'}(\mathbb{R}^n)$ due to Theorem 2.6. Hence, using Proposition 2.1, it is sufficient to show (2.1). Because of Theorem 2.9 with $a(x, \xi, y) = p(x, \xi)$,

$$\begin{aligned} p(x, D_x)f &= \sum_{j=0}^{\infty} p_j(x, D_x)f = \sum_{j=0}^{\infty} \int_{\mathbb{R}^n} k_j(x, x - y)f(y) dy \\ &= \int_{\mathbb{R}^n} k(x, x - y)f(y) dy \quad \text{for all } x \notin \text{supp } f. \end{aligned}$$

The latter representation and the kernel estimate stated in Theorem 2.9 imply that for sufficiently large $R > 0$

$$\begin{aligned} &\|p(x, D_x)f\|_{\mathcal{C}^s(\mathbb{R}^n \setminus B_R(0))} \\ &\leq \sup_{z \neq 0} |z|^N \|k(\cdot, z)\|_{\mathcal{C}^s(\mathbb{R}^n)} \sup_{|x| \geq R} \left(\int_{\text{supp } f} |x - y|^{-N} dy \|f\|_{\mathcal{C}^s(\mathbb{R}^n)} \right) \\ &\leq C |\text{supp } f| \|f\|_{\infty} |R|^{-N} \rightarrow_{R \rightarrow \infty} 0. \end{aligned}$$

Hence (2.1) holds and therefore $p(x, D_x)f \in \mathcal{C}_0^s(\mathbb{R}^n)$. The statement for $p(D_x, x)f$ is proved in the same way. □

Recall that, if $a \in S_{1,0}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$ is a smooth symbol, then by the results of the classical theory of pseudodifferential operators

$$a(x, D_x, x) = p(x, D_x),$$

where $p \in S_{1,0}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$ and

$$p(x, \xi) = a(x, \xi, x) + r(x, \xi),$$

with $r \in S_{1,0}^{m-1}(\mathbb{R}^n; \mathbb{R}^n)$, see [23, Chapter 2, Section 3]. In the case $a \in C^\tau S_{1,0;N}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$, $0 \leq \tau \leq m$, the following result can be applied to

$$r(x, \xi, y) = a(x, \xi, y) - a(x, \xi, x).$$

Proposition 2.11. *Let $r \in C^\tau S_{1,0;N}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$, where $\tau \in (0, 1)$, $0 \leq m < \tau$, and $N = n + 1$. Moreover, we assume that $r(x, \xi, x) = 0$. Then*

$$r(x, D_x, x) := \sum_{j=0}^{\infty} r_j(x, D_x, x)$$

converges absolutely in $\mathcal{L}(\mathcal{C}^s(\mathbb{R}^n))$ for each $0 < s < \tau - m$ and satisfies

$$(2.7) \quad \|r(x, D_x, x)\|_{\mathcal{L}(\mathcal{C}_0^s(\mathbb{R}^n))} \leq C \|r\|_{C^\tau S_{1,0,N}^m},$$

where C does not depend on $r \in C^\tau S_{1,0,N}^m(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{R}^n)$. Moreover, $r(x, D_x, x)$ maps $\mathcal{C}_0^s(\mathbb{R}^n)$ into itself.

Proof. First we denote

$$r^M(x, D_x, x)f := \sum_{j=0}^M r_j(x, D_x, x)f.$$

Using that

$$r_j(x, D_x, x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} k_j(x, y, x - y)f(y) dy, \quad f \in \mathcal{S}(\mathbb{R}^n),$$

we have

$$r^M(x, D_x, x) = \int_{\mathbb{R}^n} k^M(x, y, x - y)f(y) dy, \quad f \in \mathcal{S}(\mathbb{R}^n),$$

with $k^M(x, y, z) := \sum_{j=0}^M k_j(x, y, z)$. Note that $k^M(x, x, z) = k_j(x, x, z) = 0$ since $r(x, \xi, x) = 0$. By the proof of Theorem 2.9 it is obvious that

$$\|k^M(\cdot, \cdot, z)\|_{C^\tau(\mathbb{R}^n \times \mathbb{R}^n)} \leq \begin{cases} C \|r\|_{C^\tau S_{1,0,N}^m} |z|^{-n-m} & \text{if } |z| \leq 1, \\ C \|r\|_{C^\tau S_{1,0,N}^m} |z|^{-n-1} & \text{if } |z| \geq 1, \end{cases}$$

uniformly in $z \neq 0$ and $M \in \mathbb{N}$. But this implies

$$(2.8) \quad \begin{aligned} |k^M(x, y, x - y)| &= |k^M(x, y, x - y) - k^M(x, x, x - y)| \\ &\leq C \|r\|_{C^\tau S_{1,0,N}^m} |x - y|^{-n-m+\tau} (1 + |x - y|)^{m-1}. \end{aligned}$$

Hence Lebesgue's theorem on dominated convergence implies that

$$r(x, D_x, x)f = \lim_{M \rightarrow \infty} r^M(x, D_x, x)f = \int_{\mathbb{R}^n} k(x, y, x - y)f(y) dy$$

exists for every $x \in \mathbb{R}^n$ and $f \in L^\infty(\mathbb{R}^n)$. Moreover, since (2.8) holds for $k(x, y, x - y)$ as well, we conclude

$$(2.9) \quad \|r(x, D_x, x)\|_{\mathcal{L}(L^\infty(\mathbb{R}^n))} \leq C \|r\|_{C^\tau S_{1,0,N}^m}.$$

In order to prove (2.7), we use the relation

$$\Delta_h r(x, D_x, x)f = r(x, D_x, x)(\Delta_h f) + \int_{\mathbb{R}^n} k_h(x, y, x - y)f(y + h) dy,$$

where $(\Delta_h f)(x) = f(x + h) - f(x)$, $h \in \mathbb{R}^n$, and

$$k_h(x, y, z) = k(x + h, y + h, z) - k(x, y, z).$$

Moreover, $k_h(x, y, z)$ is the kernel belonging to $r_h(x, D_x, x)$ with $r_h(x, \xi, y) = r(x + h, \xi, y + h) - r(x, \xi, y)$ and it is easy to prove that

$$\|r_h\|_{C^{\tau-s}S_{1,0,N}^m} \leq C|h|^s \|r_h\|_{C^\tau S_{1,0,N}^m}$$

uniformly in $h \in \mathbb{R}^n$ for each $0 < s < \tau$. Hence using (2.9) for r and r_h , we conclude that

$$\begin{aligned} \|\Delta_h r(x, D_x, x)f\|_{L^\infty} &\leq C\|r\|_{C^\tau S_{1,0,N}^m} \|\Delta_h f\|_{L^\infty} + C\|r_h\|_{C^{\tau-s}S_{1,0,N}^m} \|f\|_{L^\infty} \\ &\leq C\|r\|_{C^\tau S_{1,0,N}^m} \|f\|_{C^s(\mathbb{R}^n)} |h|^s \end{aligned}$$

for $0 < s < \tau - m$. This finishes the proof of (2.7). The last statement is proved by showing that $r(x, D_x, x)f \in C_0^s(\mathbb{R}^n)$ for $f \in C_0^\infty(\mathbb{R}^n)$. This can be done in the same way as in Corollary 2.10 using the decay of the kernel $k(x, y, z)$ as $|z| \rightarrow \infty$ and Proposition 2.1. □

2.3. Application to the resolvent equations. In this section we construct an approximate resolvent Q_λ to a Lévy-type operator L as introduced in (1.1), (1.2). Here $Q_\lambda = q_\lambda(D_x, x)$ is a pseudodifferential operator obtained by inverting the symbol of the principal part of $\lambda - L$.

More precisely, because of the assumption on the kernel, we have a decomposition

$$Lu(x) = L^1u(x) + L^2u(x), \quad u \in \mathcal{S}(\mathbb{R}^n),$$

where L^j denotes the same kind of operator with kernel k^j , $j = 1, 2$. Here L^1 can be considered as principle part and L^2 is of lower order in the following sense:

Lemma 2.12. *Let L^2 be as above. Then L^2 extends to a bounded operator $L^2: C_0^{s+\alpha''}(\mathbb{R}^n) \rightarrow C_0^s(\mathbb{R}^n)$ for any $\alpha'' > \alpha'$ and $0 < s < \tau$ provided that $s + \alpha'' > 1$ if $\alpha \geq 1$.*

Proof. First of all, if $u \in C^{s'}(\mathbb{R}^n)$ and $1 < s' < 2$, then

$$(2.10) \quad |u(x + y) - u(x) - y \cdot \nabla u(x)| \leq C\|u\|_{C^{s'}(\mathbb{R}^n)} |y|^{s'}, \quad |y| \leq 1.$$

First we assume that $1 \leq \alpha' < \alpha < 2$. Then (2.10) with $s' = \alpha''$ yields

$$(2.11) \quad \begin{aligned} & \|L^2 u\|_{L^\infty(\mathbb{R}^n)} \\ & \leq C \left(\sup_{x \in \mathbb{R}^n, |y| \leq 1} |y|^{n+\alpha'} |k_2(x, y)| + \int_{|y| \geq 1} \|k_2(\cdot, y)\|_\infty dy \right) \|u\|_{C^{\alpha''}(\mathbb{R}^n)} \end{aligned}$$

with a constant C independent of k_2 . Moreover,

$$(2.12) \quad \Delta_h(L^2 u) = L^2(\Delta_h u) + L_h^2(\tau_h u),$$

where L_h^2 is the Lévy-type operator with kernel $k_h^2(x, y) := k^2(x + h, y) - k^2(x, y)$. By the assumptions on the kernel,

$$\sup_{x \in \mathbb{R}^n, |y| \leq 1} |y|^{n+\alpha'} |k_h^2(x, y)| + \int_{|y| \geq 1} \|k_h^2(\cdot, y)\|_\infty dy \leq C|h|^s$$

uniformly in $h \in \mathbb{R}^n$. Therefore using (2.11) with L^2 replaced by holds for L_h^2 and k_2 replaced by k_h^2 we conclude

$$\|L_h^2(\tau_h u)\|_{L^\infty(\mathbb{R}^n)} \leq C|h|^s \|u\|_{C^{\alpha''}(\mathbb{R}^n)}.$$

Hence, using the inequality above, (2.12), and (2.11), we conclude

$$\|\Delta_h(L^2 u)\|_{L^\infty(\mathbb{R}^n)} \leq C(\|\Delta_h u\|_{C^{\alpha''}(\mathbb{R}^n)} + |h|^s \|u\|_{C^{\alpha''}(\mathbb{R}^n)}) \leq Ch^s \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)},$$

where we have used $\|\Delta_h u\|_{C^{\alpha''}(\mathbb{R}^n)} \leq C|h|^s \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)}$. The latter inequality can be easily proved by first proving the cases $s = 0, 1$ and then using interpolation. Hence $L^2: C^{s+\alpha''}(\mathbb{R}^n) \rightarrow C^s(\mathbb{R}^n)$.

Secondly, if $0 < \alpha < 1$, then the proof above is easily modified using

$$|u(x + y) - u(x)| \leq C \|u\|_{C^{s'}(\mathbb{R}^n)} |y|^{s'}, \quad |y| \leq 1,$$

for $u \in C^{s'}(\mathbb{R}^n)$ and $s' \in (0, 1)$ instead of (2.10).

It remains to consider the case $0 \leq \alpha' < 1 \leq \alpha$. Using (2.10) with $s' = s + \alpha'' \in (1, 2)$ we conclude as before

$$(2.13) \quad \begin{aligned} & \|L^2 u\|_{L^\infty(\mathbb{R}^n)} \\ & \leq C \left(\sup_{x \in \mathbb{R}^n, |y| \leq 1} |y|^{n+\alpha'} |k_2(x, y)| + \int_{|y| \geq 1} \|k_2(\cdot, y)\|_\infty dy \right) \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)} \end{aligned}$$

with a constant C independent of k_2 . We use again (2.12). The second term can be estimated in the same manner as before to obtain

$$\|L_h^2(\tau_h u)\|_{L^\infty(\mathbb{R}^n)} \leq C|h|^s \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)}.$$

But the first term in (2.12) has to be estimated differently: Using (2.10) with u replaced by $\Delta_h u$, we have on one hand

$$\begin{aligned} & |\Delta_h u(x+y) - \Delta_h u(x) - y \cdot \nabla \Delta_h u(x)| \\ & \leq C \|\Delta_h u\|_{C^{s+\alpha''}(\mathbb{R}^n)} |y|^{s+\alpha''} \leq C' \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)} |y|^{s+\alpha''}, \quad |y| \leq 1. \end{aligned}$$

On the other hand

$$\begin{aligned} & |\Delta_h u(x+y) - \Delta_h u(x) - y \cdot \nabla \Delta_h u(x)| \\ & \leq C \|\Delta_h u\|_{C^1(\mathbb{R}^n)} |y| \leq C' |y| |h|^{s+\alpha''-1} \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)}, \quad |y|, |h| \leq 1. \end{aligned}$$

Interpolation of both inequalities yields

$$|\Delta_h u(x+y) - \Delta_h u(x) - y \cdot \nabla \Delta_h u(x)| \leq C |h|^s |y|^{\alpha''} \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)}$$

uniformly in $|h|, |y| \leq 1$. With this inequality

$$\|L^2 \Delta_h u\|_{L^\infty(\mathbb{R}^n)} \leq C |h|^s \|u\|_{C^{s+\alpha''}(\mathbb{R}^n)}, \quad |h| \leq 1,$$

is proved in the same way as before.

Finally, if $f \in C_0^\infty(\mathbb{R}^n)$, one easily proves $L^2 f \in C_0^s(\mathbb{R}^n)$ with the aid of Proposition 2.1 and (1.7). □

For the principal part L^1 , we use

$$\begin{aligned} u(x+y) - u(x) - y \cdot \nabla u(x) &= \mathbb{F}_{\xi \mapsto x}^{-1} [(e^{iy \cdot \xi} - 1 - iy \cdot \xi) \hat{u}(\xi)], \\ u(x+y) - u(x) &= \mathbb{F}_{\xi \mapsto x}^{-1} [(e^{iy \cdot \xi} - 1) \hat{u}(\xi)]. \end{aligned}$$

Hence L^1 can be represented as a pseudodifferential operator

$$L^1 u(x) = \int_{\mathbb{R}^n} e^{ix \cdot \xi} p(x, \xi) \hat{u}(\xi) d\xi,$$

where

$$\begin{aligned} p(x, \xi) &:= \int_{\mathbb{R}^n} (e^{iy \cdot \xi} - 1 - iy \cdot \xi) k_1(x, y) dy & \text{if } \alpha \in [1, 2), \\ p(x, \xi) &:= \int_{\mathbb{R}^n} (e^{iy \cdot \xi} - 1) k_1(x, y) dy & \text{if } \alpha \in (0, 1). \end{aligned}$$

Note that in the borderline case $\alpha = 1$ we also have

$$p(x, \xi) = \int_{\mathbb{R}^n} (e^{iy \cdot \xi} - 1) k_1(x, y) dy$$

since $k_1(x, -y) = k_1(x, y)$ by the assumptions.

The following lemma shows that p is a symbol in the class studied above.

Lemma 2.13. *Let $k_1: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ be N -times differentiable w.r.t. the second variable satisfying*

$$(2.14) \quad \|\partial_y^\beta k_1(\cdot, y)\|_{C^\tau(\mathbb{R}^n)} \leq C|y|^{-n-\alpha-|\beta|}$$

for all $0 < |y| \leq 2$ and $|\beta| \leq N$ and $k_1(x, y) = 0$ for $|y| \geq 2$ and $k_1(x, -y) = k_1(x, y)$ if $\alpha = 1$. Then $p \in C^\tau S_{1,0;N}^\alpha(\mathbb{R}^n; \mathbb{R}^n)$ where p is defined as above.

Proof. We denote $f(s) = e^{is} - 1 - is$, $s \in \mathbb{R}$, if $\alpha \in [1, 2)$ and $f(s) = e^{is} - 1$, $s \in \mathbb{R}$, if $\alpha \in (0, 1)$. Let $\gamma, \beta \in \mathbb{N}_0^n$ with $m = |\gamma| = |\beta| \leq N$. Then

$$\partial_\xi^\beta (\xi^\gamma f(y \cdot \xi)) = \partial_\xi^\beta (\partial_y^\gamma F^m(y \cdot \xi)) = \partial_y^\gamma (\partial_\xi^\beta F^m(y \cdot \xi)) = \partial_y^\gamma (y^\beta f(y \cdot \xi))$$

where F^m denotes the m -th primitive of f . Therefore

$$\begin{aligned} \partial_\xi^\beta (\xi^\gamma p(x, \xi)) &= \int_{\mathbb{R}^n} \partial_y^\gamma (y^\beta f(y \cdot \xi)) k_1(x, y) dy \\ &= (-1)^m \int_{\mathbb{R}^n} y^\beta f(y \cdot \xi) \partial_y^\gamma k_1(x, y) dy \\ &= (-1)^m |\xi|^{-n-m} \int_{\mathbb{R}^n} z^\beta f\left(z \cdot \frac{\xi}{|\xi|}\right) (\partial_y^\gamma k_1)\left(x, \frac{z}{|\xi|}\right) dz. \end{aligned}$$

Hence, if $\alpha \neq 1$

$$\begin{aligned} \|\partial_\xi^\beta (\xi^\gamma p(\cdot, \xi))\|_{C^\tau(\mathbb{R}^n)} &\leq C|\xi|^{-n-m} \int_{\mathbb{R}^n} |z|^m \frac{|z|^j}{1+|z|} \left| \frac{z}{|\xi|} \right|^{-n-\alpha-m} dz \\ &\leq C' |\xi|^\alpha, \end{aligned}$$

where $j = 2$ if $\alpha \geq 1$ and $j = 1$ else. Moreover, if $\alpha = 1$, we use that

$$\begin{aligned} \partial_\xi^\beta (\xi^\gamma p(x, \xi)) &= (-1)^m |\xi|^{-n-m} \int_{|z| \leq 1} z^\beta f\left(z \cdot \frac{\xi}{|\xi|}\right) (\partial_y^\gamma k_1)\left(x, \frac{z}{|\xi|}\right) dz \\ &\quad + (-1)^m |\xi|^{-n-m} \int_{|z| > 1} z^\beta (e^{z \cdot \xi / |\xi|} - 1) (\partial_y^\gamma k_1)\left(x, \frac{z}{|\xi|}\right) dz \end{aligned}$$

since $k_1(x, -y) = k_1(x, y)$ by assumption. Therefore

$$\begin{aligned} \|\partial_\xi^\beta (\xi^\gamma p(\cdot, \xi))\|_{C^\tau(\mathbb{R}^n)} &\leq C|\xi|^{-n-m} \int_{\mathbb{R}^n} |z|^m \frac{|z|^2}{1+|z|^2} \left| \frac{z}{|\xi|} \right|^{-n-\alpha-m} dz \\ &\leq C' |\xi|^\alpha, \end{aligned}$$

also in the case $\alpha = 1$.

Since $\beta, \gamma \in \mathbb{N}_0^n$ with $|\beta| = |\gamma| \leq N$ are arbitrary, the latter estimate implies

$$\|\xi^\gamma \partial_\xi^\beta p(\cdot, \xi)\|_{C^\tau(\mathbb{R}^n)} \leq C|\xi|^\alpha$$

for all $|\beta| = |\gamma| \leq N$, which is easy to prove by induction. Hence

$$\|\partial_\xi^\beta p(\cdot, \xi)\|_{C^\tau(\mathbb{R}^n)} \leq C|\xi|^{\alpha-|\beta|}$$

since $\gamma \in \mathbb{N}_0^n$ with $|\gamma| = |\beta|$ is arbitrary. □

Hence (1.3), Lemma 2.13, Theorem 2.6, and Corollary 2.10 imply that

$$p(x, D_x): C_0^{s+\alpha}(\mathbb{R}^n) \rightarrow C_0^\alpha(\mathbb{R}^n)$$

for all $0 < s < \tau$. Moreover, (1.4) implies

$$\begin{aligned} -\operatorname{Re} p(x, \xi) &= \int_{\mathbb{R}^n} (1 - \cos y \cdot \xi) k_1(x, y) dy \\ &\geq c \int_{B_2(0)} (1 - \cos y \cdot \xi) |y|^{-n-\alpha} dy \geq C|\xi|^\alpha \end{aligned}$$

for all $|\xi| \geq 1$ and $-\operatorname{Re} p(x, \xi) \geq 0$ for all $\xi \in \mathbb{R}^n$. Since $|p(x, \xi)| \leq C|\xi|^\alpha$, we conclude that

$$\left| \frac{\operatorname{Im} p(x, \xi)}{\operatorname{Re} p(x, \xi)} \right| \leq M$$

uniformly in $|\xi| \geq 1$. Thus $p(x, \xi) \in \mathbb{C} \setminus \Sigma_\delta$ for $\delta := \pi - \arctan M > \pi/2$ and for all $|\xi| \geq 1$.

Hence, we can define

$$q_\lambda(y, \xi) := (\lambda - p(y, \xi))^{-1}, \quad y, \xi \in \mathbb{R}^n, \lambda \in \Sigma_{\delta'}, |\lambda| \geq R,$$

for $0 < \delta' < \delta$ and $R > \sup_{x \in \mathbb{R}^n, |\xi| \leq 1} |p(x, \xi)|$.

Since $p \in C^\tau S_{1,0,N}^\alpha(\mathbb{R}^n; \mathbb{R}^n)$, we have $q_\lambda \in C^\tau S_{1,0,N}^{-\alpha}(\mathbb{R}^n; \mathbb{R}^n)$. More precisely, the following lemma holds:

Lemma 2.14. *Let q_λ, δ be defined as above and $\lambda \in \Sigma_{\delta'}$ where $\delta' \in (0, \delta)$ is arbitrary. Then there is some $R > 0$ such that $q_\lambda \in C^\tau S_{1,0,N}^{-\alpha}$ for all $\lambda \in \Sigma_{\delta'}$ with $|\lambda| \geq R$. Moreover, for each $\alpha' \in [0, \alpha]$*

$$\|q_\lambda\|_{C^\tau S_{1,0,N}^{-\alpha'}} \leq C_{\delta'} (1 + |\lambda|)^{-(\alpha-\alpha')/\alpha}$$

uniformly in $\lambda \in \Sigma_{\delta'}$ with $|\lambda| \geq R$.

Proof. First of all, by a simple geometric observation

$$|\lambda - z| \geq c_{\delta'} \max\{|\lambda|, |z|\} \quad \text{if } \lambda \in \Sigma_{\delta'}, z \in \mathbb{C} \setminus \Sigma_{\delta}$$

provided that $0 < \delta' < \delta$. As seen above $p(x, \xi) \in \mathbb{C} \setminus \Sigma_{\delta}$ for $|\xi| \geq 1$ and some $\delta > \pi/2$ and $|p(x, \xi)| \geq c|\xi|^\alpha$ for $|\xi| \geq 1$. Hence

$$(2.15) \quad |\lambda - p(x, \xi)| \geq c_{\delta'} \max\{|\lambda|, |\xi|^\alpha\}$$

for all $|\xi| \geq 1$ and $\lambda \in \Sigma_{\delta'}$ with $0 < \delta' < \delta$ arbitrary. Moreover, since $|p(x, \xi)| \leq C$ for all $|\xi| \leq 1$ and $x \in \mathbb{R}^n$, we conclude that (2.15) holds for all $\xi \in \mathbb{R}^n$ and $\lambda \in \Sigma_{\delta'}$ with $|\lambda| \geq R$ for some $R > 0$ sufficiently large. Using this, $p \in C^\tau S_{1,0,N}^\alpha(\mathbb{R}^n; \mathbb{R}^n)$, and the chain rule, one derives in a straight-forward manner that

$$\|\partial_\xi^\beta q_\lambda(\cdot, \xi)\|_{C^\tau(\mathbb{R}^n)} \leq C_{\delta'} \frac{\langle \xi \rangle^{-|\beta|}}{|\lambda| + |\xi|^\alpha} \leq C_{\delta'} |\lambda|^{-(\alpha - \alpha')/\alpha} \langle \xi \rangle^{-\alpha' - |\beta|}$$

uniformly in $\xi \in \mathbb{R}^n$ and $\lambda \in \Sigma_{\delta'}$, $|\lambda| \geq R > 0$ and for all $|\beta| \leq N$, which proves the statement. □

Application of Theorem 2.6, Corollary 2.10 and the lemma above gives:

Corollary 2.15. *Let $q_\lambda, \delta, \delta'$ be as above and let $0 < s < \tau$. Then*

$$q_\lambda(D_x, x): C_0^s(\mathbb{R}^n) \rightarrow C_0^{s+\alpha}(\mathbb{R}^n)$$

is a bounded linear operator, which satisfies

$$\|q_\lambda(D_x, x)\|_{\mathcal{L}(C_0^s(\mathbb{R}^n), C_0^{s+\alpha'}(\mathbb{R}^n))} \leq C_{\delta'} |\lambda|^{-(\alpha - \alpha')/\alpha} \quad \text{for all } \lambda \in \Sigma_{\delta'}, |\lambda| \geq R,$$

for all $0 \leq \alpha' \leq \alpha$ with some sufficiently large $R > 0$.

Now we are in the position to prove the following key lemma.

Lemma 2.16. *Let $q_\lambda, \delta, \delta'$ be as above and let $0 < s < \tau$. Then*

$$(\lambda - p(x, D_x))q_\lambda(D_x, x) = I - R_\lambda$$

with

$$\|R_\lambda\|_{\mathcal{L}(C_0^s(\mathbb{R}^n))} \leq C_{\delta'} |\lambda|^{-\varepsilon}$$

uniformly in $\lambda \in \Sigma_{\delta'}$ with $|\lambda| \geq M$ for sufficiently large $M > 0$ and some $\varepsilon > 0$ depending on s, τ .

Proof. First of all, for each $f \in C_0^\infty(\mathbb{R}^n)$, $q_\lambda(D_x, x)f \in C^{s'+\alpha}(\mathbb{R}^n)$ with $s < s' < \tau$. We conclude

$$\sum_{j=0}^N \varphi_j(D_x)q_\lambda(D_x, x)f \rightarrow q_\lambda(D_x, x)f \quad \text{in } C^{s+\alpha}(\mathbb{R}^n) \quad \text{as } N \rightarrow \infty.$$

Therefore

$$q_\lambda(D_x, x)f = \sum_{j=0}^\infty \varphi_j(D_x)q_\lambda(D_x, x)f = \sum_{j=0}^\infty q_{\lambda,j}(D_x, x)f$$

where $q_{\lambda,j}(\xi, y) = q_\lambda(\xi, y)\varphi_j(\xi)$. Hence

$$\begin{aligned} (\lambda - p(x, D_x))q_\lambda(D_x, x)f &= \sum_{j=0}^\infty (\lambda - p(x, D_x))q_{\lambda,j}(D_x, x)f \\ &= f + \sum_{j=0}^\infty a_{\lambda,j}(x, D_x, x)f, \end{aligned}$$

where $a_{\lambda,j}(x, y, \xi) = a_\lambda(x, \xi, y)\varphi_j(\xi)$ and

$$a_\lambda(x, y, \xi) = \frac{\lambda - p(x, \xi)}{\lambda - p(y, \xi)} - 1 = (p(y, \xi) - p(x, \xi))q_\lambda(y, \xi).$$

Using Lemma 2.14, we conclude

$$\|a_\lambda\|_{C^\tau S_{1,0,N}^{-\alpha'}} \leq C \|p\|_{C^\tau S_{1,0,N}^\alpha} \|q_\lambda\|_{C^\tau S_{1,0,N}^{-\alpha'}} \leq C_{\delta'}(1 + |\lambda|)^{-(\alpha - \alpha')/\alpha}.$$

Since $a_\lambda(x, \xi, x) = 0$, we can use Proposition 2.11 to conclude that

$$a_\lambda(x, D_x, x) = \sum_{j=0}^\infty a_{\lambda,j}(x, D_x, x)$$

is well-defined as limit in $\mathcal{L}(C_0^s(\mathbb{R}^n))$ and satisfies

$$\|a_\lambda(x, D_x, x)\|_{\mathcal{L}(C_0^s(\mathbb{R}^n))} \leq C \|a_\lambda\|_{C^\tau S_{1,0,N}^{-\alpha'}} \leq C_{\delta'}(1 + |\lambda|)^{-(\alpha - \alpha')/\alpha}$$

for each $0 < \alpha' < \alpha$ with $\alpha - \alpha' < \tau - s$. □

Recall that an unbounded operator $A: \mathcal{D}(A) \subseteq X \rightarrow X$ generates an analytic semi-group on a Banach space X if and only if A is closed, $\mathcal{D}(A)$ is dense, and there are

some $\delta > \pi/2$, $\omega \in \mathbb{R}$, and $M \geq 1$ such that $(\lambda - A)^{-1}$ exists for all $\lambda \in \omega + \Sigma_\delta$ and satisfies

$$(2.16) \quad \|(\lambda - A)^{-1}\|_{\mathcal{L}(X)} \leq \frac{M}{|\lambda - \omega|} \quad \text{for all } \lambda \in \omega + \Sigma_\delta,$$

cf. [37].

Corollary 2.17. *Let $0 < s < \tau$. Then $p(x, D_x)$ and L generate an analytic semi-group on $\mathcal{C}_0^s(\mathbb{R}^n)$ with domains $\mathcal{D}(L) = \mathcal{D}(p(x, D_x)) = \mathcal{C}_0^{s+\alpha}(\mathbb{R}^n)$. Moreover, if $A = p(x, D_x)$ or $A = L$, then*

$$\|(\lambda - A)^{-1}\|_{\mathcal{L}(\mathcal{C}_0^s(\mathbb{R}^n), \mathcal{C}_0^{s+\alpha'}(\mathbb{R}^n))} \leq C_{\delta'} |\lambda|^{-(\alpha-\alpha')/\alpha} \quad \text{for all } \lambda \in \Sigma_{\delta'}, |\lambda| \geq R,$$

for all $0 \leq \alpha' \leq \alpha$ with some sufficiently large $R > 0$ and some $\delta' > \pi/2$.

Proof. By a standard Neumann series argument Lemma 2.16 yields that

$$(\lambda - p(x, D_x))^{-1}: \mathcal{C}_0^s(\mathbb{R}^n) \rightarrow \mathcal{C}_0^{s+\alpha}(\mathbb{R}^n)$$

exists for all $\lambda \in \Sigma_{\delta'}$ with $|\lambda| \geq R$ for some $R > 0$ and satisfies

$$\|(\lambda - p(x, D_x))^{-1}\|_{\mathcal{L}(\mathcal{C}_0^s(\mathbb{R}^n))} \leq 2 \|q_\lambda(D_x, x)\|_{\mathcal{L}(\mathcal{C}_0^s(\mathbb{R}^n))} \leq C |\lambda|^{-1}.$$

This implies (2.16) for a suitable choice of ω . Hence $p(x, D_x)$ generates an analytic semi-group on $\mathcal{C}_0^s(\mathbb{R}^n)$ with domain $\mathcal{D}(p(x, D_x)) = \mathcal{C}_0^{s+\alpha}(\mathbb{R}^n)$.

Similarly,

$$(\lambda - L)q_\lambda(D_x, x) = I - R_\lambda + L^2 q_\lambda(D_x, x),$$

where

$$\|L^2 q_\lambda(D_x, x)\|_{\mathcal{L}(\mathcal{C}_0^s(\mathbb{R}^n))} \leq C \|q_\lambda(D_x, x)\|_{\mathcal{L}(\mathcal{C}_0^s(\mathbb{R}^n), \mathcal{C}_0^{s+\alpha''}(\mathbb{R}^n))} \leq C_{\delta, \delta', \alpha''} |\lambda|^{-(\alpha-\alpha'')/\alpha}$$

uniformly in $\lambda \in \Sigma_{\delta'}$, $|\lambda| \geq R$, with arbitrary $\alpha' < \alpha'' < \alpha$. Thus the same arguments as before show that L generates an analytic semi-group.

Finally, the uniform estimate of $(\lambda - A)^{-1}$ easily follows from Corollary 2.15. \square

Proof of Theorem 1.1. Because of Corollary 2.17, well-known results from semi-group theory imply the existence of a unique classical solution $u \in \mathcal{C}^{1,\theta}([0, T]; \mathcal{C}_0^s(\mathbb{R}^n)) \cap \mathcal{C}^\theta([0, T]; \mathcal{D}(L))$ of (1.8)–(1.9), cf. [37, Chapter 4, Theorem 3.5]. Finally, since $(\lambda - L)^{-1}: \mathcal{C}_0^s(\mathbb{R}^n) \rightarrow \mathcal{C}_0^{s+\alpha}(\mathbb{R}^n)$ is a bounded operator for $\lambda = R$, the graph norm on $\mathcal{D}(L)$, i.e., $\|u\|_{\mathcal{C}^s} + \|Lu\|_{\mathcal{C}^s}$, is equivalent to the norm of $\mathcal{C}^{s+\alpha}(\mathbb{R}^n)$. That u inherits the non-negativity from f is easily established using the maximum principle. \square

References

- [1] H. Abels: *Bounded imaginary powers and H_∞ -calculus of the Stokes operator in unbounded domains*; in *Nonlinear Elliptic and Parabolic Problems*, Progr. Nonlinear Differential Equations Appl. **64**, Birkhäuser, Basel, 2005, 1–15.
- [2] H. Abels: *Pseudodifferential boundary value problems with non-smooth coefficients*, *Comm. Partial Differential Equations* **30** (2005), 1463–1503.
- [3] H. Abels: *Reduced and generalized Stokes resolvent equations in asymptotically flat layers, part II: H_∞ -calculus*, *J. Math. Fluid Mech.* **7** (2005), 223–260.
- [4] R.F. Bass: *Uniqueness in law for pure jump Markov processes*, *Probab. Theory Related Fields* **79** (1988), 271–287.
- [5] R.F. Bass: *Stochastic differential equations with jumps*, *Probab. Surv.* **1** (2004), 1–19 (electronic).
- [6] R.F. Bass and H. Tang: *The martingale problem for a class of stable-like processes*, *Stochastic Process. Appl.* **119** (2009), 1144–1167.
- [7] J. Bertoin: *Lévy Processes*, Cambridge Tracts in Mathematics **121**, Cambridge Univ. Press, Cambridge, 1996.
- [8] P. Billingsley: *Convergence of Probability Measures*, Wiley, New York, 1968.
- [9] P. Billingsley: *Convergence of Probability Measures*, Wiley Series in Probability and Statistics: Probability and Statistics, Second edition, Wiley, New York, 1999.
- [10] V. Bogachev, P. Lescot and M. Röckner: *The martingale problem for pseudo-differential operators on infinite-dimensional spaces*, *Nagoya Math. J.* **153** (1999), 101–118.
- [11] B. Böttcher: *A parametrix construction for the fundamental solution of the evolution equation associated with a pseudo-differential operator generating a Markov process*, *Math. Nachr.* **278** (2005), 1235–1241.
- [12] Ph. Courège: *Sur la forme intégro-différentielle des opérateurs de c_k^∞ dans c satisfaisant au principe du maximum*, *Séminaire Brelot-Choquet-Deny, Théorie du potentiel* **10**, 1–38, 1965–1966.
- [13] S.D. Eidelman, S.D. Ivasyshen and A.N. Kochubei: *Analytic Methods in the Theory of Differential and Pseudo-Differential Equations of Parabolic Type*, *Operator Theory: Advances and Applications* **152**, Birkhäuser, Basel, 2004.
- [14] S.N. Ethier and T.G. Kurtz: *Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics, Markov Processes*, Wiley, New York, 1986.
- [15] W. Hoh: *The martingale problem for a class of pseudo-differential operators*, *Math. Ann.* **300** (1994), 121–147.
- [16] W. Hoh: *Pseudo differential operators with negative definite symbols of variable order*, *Rev. Mat. Iberoamericana* **16** (2000), 219–241.
- [17] N. Jacob: *Pseudo Differential Operators and Markov Processes, III*, Imp. Coll. Press, London, 2005.
- [18] N. Jacob and H.-G. Leopold: *Pseudo-differential operators with variable order of differentiation generating Feller semigroups*, *Integral Equations Operator Theory* **17** (1993), 544–553.
- [19] N. Jacob and R.L. Schilling: *Lévy-type processes and pseudodifferential operators*; in *Lévy Processes*, Birkhäuser, Boston, Boston, MA, 2001, 139–168.
- [20] J. Jacod and A.N. Shiryaev: *Limit Theorems for Stochastic Processes*, second edition, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]* **288**, Springer, Berlin, 2003.
- [21] O. Kallenberg: *Foundations of Modern Probability*, Springer, New York, 1997.
- [22] M. Kassmann: *\mathcal{L} -harmonische Funktionen und Sprungprozesse*, *Mitt. Dtsch. Math.-Ver.* **14** (2006), 80–87.
- [23] H. Kumano-go: *Pseudodifferential Operators*, Translated from the Japanese by the author, Rémi Vaillancourt and Michihiro Nagase, MIT Press, Cambridge, Mass., 1981.
- [24] H. Kumano-go and M. Nagase: *Pseudo-differential operators with non-regular symbols and applications*, *Funkcial. Ekvac.* **21** (1978), 151–192.
- [25] K. Kikuchi and A. Negoro: *On Markov process generated by pseudodifferential operator of variable order*, *Osaka J. Math.* **34** (1997), 319–335.
- [26] V. Kolokoltsov: *Symmetric stable laws and stable-like jump-diffusions*, *Proc. London Math.*

- Soc. (3) **80** (2000), 725–768.
- [27] T. Komatsu: *Markov processes associated with certain integro-differential operators*, Osaka J. Math. **10** (1973), 271–303.
- [28] T. Komatsu: *On the martingale problem for generators of stable processes with perturbations*, Osaka J. Math. **21** (1984), 113–132.
- [29] T. Komatsu: *Pseudodifferential operators and Markov processes*, J. Math. Soc. Japan **36** (1984), 387–418.
- [30] J.-P. Lepeltier and B. Marchal: *Problème des martingales et équations différentielles stochastiques associées à un opérateur intégro-différentiel*, Ann. Inst. H. Poincaré Sect. B (N.S.) **12** (1976), 43–103.
- [31] J. Marschall: *Pseudodifferential operators with nonregular symbols of the class $S_{\rho, \delta}^m$* , Comm. Partial Differential Equations **12** (1987), 921–965.
- [32] R. Mikulevičius and H. Pragarauskas: *On the Cauchy problem for certain integro-differential operators in Sobolev and Hölder spaces*, Liet. Mat. Rink. **32** (1992), 299–331.
- [33] R. Mikulyavichyus and G. Pragarauskas: *The martingale problem related to nondegenerate Lévy operators*, Liet. Mat. Rink. **32** (1992), 377–396.
- [34] R. Mikulyavichyus and G. Pragarauskas: *On the uniqueness of solutions of the martingale problem that is associated with degenerate Lévy operators*, Liet. Mat. Rink. **33** (1993), 455–475.
- [35] A. Negoro and M. Tsuchiya: *Stochastic processes and semigroups associated with degenerate Lévy generating operators*, Stochastics Stochastics Rep. **26** (1989), 29–61.
- [36] O. Okitaloshima and J.A. van Casteren: *On the uniqueness of the martingale problem*, Internat. J. Math. **7** (1996), 775–810.
- [37] A. Pazy: *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer, New York, 1983.
- [38] K. Sato: *Lévy Processes and Infinitely Divisible Distributions*, Cambridge Studies in Advanced Mathematics **68**, Translated from the 1990 Japanese original; Revised by the author, Cambridge Univ. Press, Cambridge, 1999.
- [39] E.M. Stein: *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, with the assistance of T.S. Murphy; Monographs in Harmonic Analysis, III, Princeton Mathematical Series **43**, Princeton Univ. Press, Princeton, NJ, 1993.
- [40] D.W. Stroock: *Diffusion processes associated with Lévy generators*, Z. Wahrsch. Verw. Gebiete **32** (1975), 209–244.
- [41] D.W. Stroock and S.R.S. Varadhan: *Multidimensional Diffusion Processes*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences] **233**, Springer, Berlin, 1979.
- [42] M.E. Taylor: *Pseudodifferential Operators and Nonlinear PDE*, Birkhäuser Boston, Boston, MA, 1991.
- [43] M.E. Taylor: *Tools for PDE*, Mathematical Surveys and Monographs, Amer. Math. Soc., Providence, RI, 2000.
- [44] H. Triebel: *Interpolation Theory, Function Spaces, Differential Operators*, North-Holland, Amsterdam, 1978.
- [45] M. Tsuchiya: *Lévy measure with generalized polar decomposition and the associated SDE with jumps*, Stochastics Stochastics Rep. **38** (1992), 95–117.

Helmut Abels
NWF I - Mathematik
Universität Regensburg
D-93040 Regensburg
Germany

Moritz Kassmann
Fakultät für Mathematik
Universität Bielefeld
Postfach 100131
D-33501 Bielefeld
Germany
e-mail: moritz.kassmann@uni-bielefeld.de