



Title	On a construction of exceptional pseudosymmetric sets
Author(s)	Nobusawa, Nobuo
Citation	Osaka Journal of Mathematics. 1984, 21(4), p. 701–706
Version Type	VoR
URL	<a href="https://doi.org/10.18910/10716">https://doi.org/10.18910/10716</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Nobusawa, N.  
Osaka J. Math.  
21 (1984), 701-706

## ON A CONSTRUCTION OF EXCEPTIONAL PSEUDOSYMMETRIC SETS

NOBUO NOBUSAWA

(Received January 17, 1984)

### 1. Introduction

If we denote the conjugation  $b^{-1}ab$  by  $a \circ b$  or  $a^{\sigma(b)}$  in a group, it satisfies (i)  $\sigma(b)$  is a permutation on the group, (ii)  $a \circ a = a$ , and (iii)  $(a \circ b) \circ c = (a \circ c) \circ (b \circ c)$ . Under the condition (i), (iii) is equivalent with a fundamental identity:  $\sigma[b^{\sigma(c)}] = \sigma(c)^{-1} \sigma(b) \sigma(c)$ . The present author called a binary system satisfying these conditions a pseudosymmetric set in [2] and [3]. D. Joyce called it a quandle in [1]. When every  $\sigma(b)$  is of order 2, a pseudosymmetric set is called a symmetric set (or an involutive quandle in [1], or a symmetric groupoid in [4].) A pseudosymmetric set is called *special* if it is isomorphic with a pseudosymmetric subset of a group (=a pseudosymmetric set in the above sense). It is easily seen that a special pseudosymmetric set is isomorphic with a union of conjugacy classes of some group. Its structure is determined to some extent by its enveloping group ([3]). A pseudosymmetric set which is not special is called *exceptional* by analogy with the theory of Jordan algebras. The simplest exceptional pseudosymmetric set is a set  $\{a, b, c\}$  with  $\sigma(a) = \sigma(b) = 1$  and  $\sigma(c) \neq 1$  ([4], p. 72). The structure of general pseudosymmetric sets seems to be more complicate, and our consideration will be restricted to the transitive one; a pseudosymmetric set is called transitive when the group of permutations generated by all  $\sigma(b)$  is a transitive permutation group on the set. The object of this paper is to show a construction method of transitive exceptional pseudosymmetric sets. In this direction, S. Doro discussed about the existence of transitive exceptional symmetric sets in his unpublished paper. (See the remark in the last part of this paper.) For the construction, we need an extension theory of pseudosymmetric sets, which will be given in 2. Let  $S$  be a pseudosymmetric set and  $A$  a set (a trivial pseudosymmetric set). We intend to make  $A \times S$  a pseudosymmetric set by introducing a suitable composition. For  $(a, s), (b, t) \in A \times S$ , we consider a composition  $(a, s) \circ (b, t) = (c, s \circ t)$ , where generally  $c$  depends on  $a, b, s$  and  $t$ , satisfying the three conditions of a pseudosymmetric set. But in this paper we consider a simple but important case that  $c$  depends only on  $a, s$  and  $t$ . Denote  $c = a^{\pi(s, t)}$ .  $\pi(s, t)$  is a permutation

on  $A$ . In order that  $A \times S$  becomes a pseudosymmetric set by this composition, it must satisfy two conditions (1) and (2) given in 2. In this case, we call the mapping  $\pi$  of  $S \times S$  into  $P(A)$ —the group of permutations on  $A$ —a *factor set*. The pseudosymmetric set  $A \times S$  given in this way is called a null extension (of  $A$  by  $S$ ). (“null” because  $c$  does not depend on  $b$ .) We can develop some formalism on factor sets similar to that of group extension theory. This may be useful in developing the structure theory, but for the purpose of this paper we are rather interested in a special factor set. When  $\varphi$  is a homomorphism of  $S$  into  $P(A)$ , let  $\pi(s, t) = \varphi(s)^{-1}\varphi(t)$ . It can be shown that this  $\pi$  is a factor set. Especially, if  $S$  is a pseudosymmetric subset of  $P(A)$ , we can take the imbedding mapping as  $\varphi$ . In 4, we specialize  $A = \{1, 2, 3, 4, 5, 6\}$  and  $S =$  the subset of  $P(A)$  consisting of all permutations of order 2 that move every element of  $A$ . We shall show that the null extension  $A \times S$  is a transitive exceptional pseudosymmetric set of order 90. To show that it is not special, we need a criterion for a transitive null extension to be special, which will be given in 3. Let  $X = A \times S$  be a null extension. For  $s \in S$ , let  $G_s = \{\tau \in G(X) \mid (A, s)^{\tau} = (A, s)\}$ , where  $G(X)$  is the group generated by  $\sigma(x)$  ( $x \in X$ ). If  $\tau \in G_s$ , it induces a permutation  $\bar{\tau}$  on  $A$  from  $(a, s)^{\tau} = (a^{\bar{\tau}}, s)$ . Then,  $F_s = \{\bar{\tau} \mid \tau \in G_s\}$  is a group of permutations on  $A$ . The criterion we are looking for is that  $F_s$  is a regular permutation group if  $X$  is special and if  $S$  is an effective pseudosymmetric set, i.e.,  $\sigma(s) \neq \sigma(t)$  whenever  $s \neq t$ . This example can be generalized. In place of 6, we take an even number  $m \geq 6$ , and let  $A = \{1, 2, \dots, m\}$ .  $S$  is the set of permutations of order 2 on  $A$  that move every element of  $A$ . The null extension  $A \times S$  constructed as above can be shown to be a transitive exceptional pseudosymmetric set. Thus, we can construct infinitely many finite transitive exceptional pseudosymmetric sets.

## 2. Null extensions and factor sets

Let  $S = \{s, t, \dots\}$  be a pseudosymmetric set and  $A = \{a, b, \dots\}$  a set. Let  $\pi$  be a mapping of  $S \times S$  into  $P(A)$ , the group of permutations on  $A$ . When  $\pi$  satisfies the following two conditions (1) and (2), we say that  $\pi$  is a factor set (with respect to  $A$  and  $S$ ).

$$(1) \quad \pi(s, s) = 1,$$

$$(2) \quad \pi(s, t)\pi(s \circ t, u) = \pi(s, u)\pi(s \circ u, t \circ u).$$

Let  $\pi$  be a factor set. We define a composition on  $A \times S = \{(a, s) \mid a \in A \text{ and } s \in S\}$  by  $(a, s) \circ (b, t) = (a^{\pi(s, t)}, s \circ t)$ . It can be checked that  $A \times S$  satisfies the three conditions of a pseudosymmetric set. We call it a null extension (of  $A$  by  $S$ ) with the factor set  $\pi$  and denote it by  $[A \times S; \pi]$ . We have an example: Let  $\varphi$  be a homomorphism of  $S$  into  $P(A)$ . Define  $\pi$  by  $\pi(s, t) = \varphi(s)^{-1}\varphi(t)$ .

$\pi$  is a factor set. Since  $\varphi$  is a homomorphism, we have  $\varphi(s \circ t) = \varphi(s) \circ \varphi(t) = \varphi(t)^{-1} \varphi(s) \varphi(t)$ . Using it, we can check that both terms of (2) coincide with  $\varphi(s)^{-2} \varphi(t) \varphi(u)$ .

We derive some formalism on factor sets. Let  $[A \times S; \pi_1]$  and  $[A \times S; \pi_2]$  be nullextensions with factor sets  $\pi_1$  and  $\pi_2$  respectively. Denote elements of  $[A \times S; \pi_i]$  by  $(a, s)_i$ ,  $i=1, 2$ , and also let  $(A, s)_i = \{(a, s)_i \mid a \in A\}$ . When there is an isomorphism  $f$  of  $[A \times S; \pi_1]$  onto  $[A \times S; \pi_2]$  such that  $f[(A, s)_1] = (A, s)_2$  for every  $s$ , we say that  $[A \times S; \pi_1]$  and  $[A \times S; \pi_2]$  are equivalent, or  $\pi_1 \sim \pi_2$ . In this case,  $f[(a, s)_1] = (a^{\theta(s)}, s)_2$ , where  $\theta(s) \in P(A)$ . From  $f[(a, s)_1 \circ (b, t)_1] = f[(a, s)_1] \circ f[(b, t)_1]$ , we can obtain  $\pi_1(s, t) \theta(s \circ t) = \theta(s) \pi_2(s, t)$ , or

$$(3) \quad \pi_2(s, t) = \theta(s)^{-1} \pi_1(s, t) \theta(s \circ t).$$

Conversely, if (3) is satisfied for a mapping  $\theta$  of  $S$  into  $P(A)$ , then  $\pi_1 \sim \pi_2$  as we can see easily. We can also see that " $\sim$ " is an equivalent relation and hence all factor sets with respect to  $A$  and  $S$  can be classified to factor set classes. Also note that if a factor set  $\pi_1$  is given, define  $\pi_2$  by (3), and we can check that  $\pi_2$  is a factor set. Let  $\varepsilon(s, t) = 1$  for any  $s$  and  $t$ .  $\varepsilon$  is a factor set. When  $\pi \sim \varepsilon$ , we say that  $\pi$  is trivial or that  $[A \times S; \pi]$  is splitting. The condition for  $\pi \sim \varepsilon$  is given by

$$(4) \quad \pi(s, t) = \theta(s)^{-1} \theta(s \circ t)$$

for some  $\theta$ . Especially, for a homomorphism  $\varphi$  of  $S$  into  $P(A)$ , put  $\pi(s, t) = \varphi(s)^{-1} \varphi(s \circ t)$ .  $\pi$  is a trivial factor set. Since  $\varphi(s \circ t) = \varphi(s) \circ \varphi(t) = \varphi(t)^{-1} \varphi(s) \varphi(t)$ , we have  $\pi(s, t) = [\varphi(s), \varphi(t)]$  (a commutator). It is interesting to note that if we use this  $\pi$  in (2), we get an identity in a group:  $[x, y][x^y, z] = [x, z][x^z, y^z]$ . Lastly, we define  $F_s$  for  $A \times S$ . Let  $X = A \times S$ , a null extension, and  $G(X)$  the group generated by all  $\sigma(x)$ ,  $x \in X$ . If  $\tau \in G(X)$  and  $(A, s)^\tau = (A, s)$ , then  $\tau$  induces a permutation  $\bar{\tau}$  on  $A$  by  $(a, s)^\tau = (a^{\bar{\tau}}, s)$ . Let  $F_s$  be the set of all such  $\tau$ . It is a permutation group on  $A$ . Suppose that  $\pi_1 \sim \pi_2$  via (3). Let  $F_s^{(i)}$  be  $F_s$  for  $[A \times S; \pi_i]$ ,  $i=1, 2$ . We show that  $F_s^{(2)} = \theta(s)^{-1} F_s^{(1)} \theta(s)$ . Let  $X_i = [A \times S; \pi_i]$ . The isomorphism  $f$  of  $X_1$  onto  $X_2$  induces in a natural sense an isomorphism of  $G(X_1)$  onto  $G(X_2)$ , which we denote by  $g$ , i.e.,  $f(x^\tau) = f(x)^{g(\tau)}$  for  $x \in X_1$  and  $\tau \in G(X_1)$ . Let  $\tau \in G(X_1)$  such that  $(A, s)_1^\tau = (A, s)_1$ . Then  $(a^{\bar{\tau}}, s)_1 = (a, s)_1$ . So,  $f[(a^{\bar{\tau}}, s)_1] = f[(a, s)_1]^{g(\tau)}$ , and hence  $(a^{\bar{\tau}\theta(s)}, s)_2 = (a^{\theta(s)}, s)_2^{g(\tau)} = (a^{\theta(s)\bar{\tau}}, s)_2$ , where  $\bar{\tau} = g(\tau) \in F_s^{(2)}$ . Thus,  $\bar{\tau} = \theta(s)^{-1} \tau \theta(s)$ , or  $F_s^{(2)} = \theta(s)^{-1} F_s^{(1)} \theta(s)$  as we asserted.

### 3. Transitive pseudosymmetric sets

Let  $X = \{x, y, \dots\}$  be a transitive pseudosymmetric set, and let  $S = \{\sigma(x) \mid x \in X\} = \sigma(X)$ .  $S$  is a special pseudosymmetric set and  $\sigma$  is considered as a homomorphism of  $X$  onto  $S$ . Corresponding to the homomorphism  $\sigma$ , we

have a coset-decomposition (or a normal decomposition in [3]) of  $X$ ;  $X = \bigcup X_i$ , where  $X_i$  is the set of inverse images of an element in  $S$  by  $\sigma$ . From now on,  $x_i$  indicate representatives of  $X_i$ . So,  $X_i = \{x \in X \mid \sigma(x) = \sigma(x_i)\}$ . Every element of  $G(X)$  induces a permutation on  $\{X_i\}$ , the set of cosets. That  $X$  is transitive implies that all  $X_i$  are bijective each other. We choose a set  $A$  to which all  $X_i$  are bijective. Then,  $X$  is bijective to  $A \times S$ , where  $X_i$  is mapped to  $(A, s_i)$  for  $s_i = \sigma(x_i) \in S$ . Through this bijection, we can make  $A \times S$  a pseudosymmetric set isomorphic with  $X$ . Then,  $A \times S$  is a null extension. The different choice of bijections in the above only leads to an equivalent null extension. Now suppose that  $X$  is special, and let  $K$  be its enveloping group (the group generated by  $X$ ). We show that  $A$  is a subgroup of  $C(K)$ , the center of  $K$ . If  $\sigma(x) = \sigma(y)$ , then  $x^{-1}zx = y^{-1}zy$  for every  $z$  in  $X$  hence in  $K$ . Thus,  $xy^{-1} \in C(K)$ . So, every element  $y$  of  $X_1$  is expressed as  $y = ux_1$  with  $u \in C(K)$ . Let  $N = \{u \in C(K) \mid ux_1 \in X_1\}$ .  $X_1 = Nx_1$ . We show that  $N$  is a subgroup. Let  $u$  and  $v \in N$ . Take  $\tau$  in  $G(X)$  such that  $x_1^\tau = ux_1$ . Then,  $(vx_1)^\tau = v(x_1^\tau)$ . Note that  $G(X)$  is the inner automorphism group of  $K$ . So,  $(vx_1)^\tau = vux_1 \in X_1$ , and hence  $vu \in N$ . In the above,  $x_1^{\tau^{-1}} = u^{-1}x_1$ , and hence  $u^{-1} \in N$ . Thus,  $N$  is a subgroup. We can also show  $X_i = Nx_i$  in the same way. We can take  $A = N$ . Let  $s = \sigma(x_i)$ . We consider  $F_s$  for our  $N \times S$ . The effect of elements of  $F_s$  on  $N$  is determined as follows. Let  $G_{X_1} = \{\tau \in G(X) \mid X_1^\tau = X_1\}$ . If  $\tau \in G_{X_1}$ , then  $\tau$  induces a permutation  $\bar{\tau}$  on  $N$  defined from  $(ux_1)^\tau = u^\tau x_1$ .  $F_s = \{\bar{\tau} \mid \tau \in G_{X_1}\}$ . For  $\tau \in G_{X_1}$ , let  $x_1^\tau = wx_1$  with  $w$  in  $N$ . Then,  $u^\tau = uw$ . Conversely, for any element  $w$  in  $N$ , choose  $\tau$  in  $G(X)$  such that  $x_1^\tau = wx_1$ . Then  $\bar{\tau} \in F_s$ . So,  $F_s$  is isomorphic with  $N$  and is of course a regular permutation group on  $N$ . We have obtained

**Theorem.** *Let  $X$  be a transitive special pseudosymmetric set. Then,  $X$  is isomorphic with a null extension  $N \times S$ , where  $N$  is a subgroup of  $C(K)$  and  $S = \sigma(X)$ . In this case,  $F_s$  is isomorphic with  $N$  and is a regular permutation group on  $N$ .*

In the above, we found that every transitive pseudosymmetric set  $X$  is isomorphic with a null extension  $[A \times S; \pi]$ . However, even if  $X$  is a null extension itself, i.e.,  $X = [A' \times S'; \pi']$ , we can not say that  $S'$  is isomorphic with  $S$  and  $A'$  is bijective to  $A$ . But if  $S'$  is effective, i.e.,  $S' \xrightarrow{\sigma} \sigma(S')$ , we can conclude it. For, let  $X = [A' \times S'; \pi']$  be transitive and suppose  $S'$  is effective. Then,  $\sigma(X)$  is isomorphic with  $S'$ . As a matter of fact, the mapping  $\sigma[(a', s')] \rightarrow s'$  gives the isomorphism, where  $a' \in A'$  and  $s' \in S'$ . On the other hand,  $S = \sigma(X)$ . So,  $S' \cong S$ . Also,  $(A', s')$  corresponds to  $(A, s)$  in the isomorphism  $X \cong [A \times S; \pi]$ , where  $s'$  corresponds to  $s$  in the above isomorphism. So,  $A'$  is bijective to  $A$ . We may replace  $S'$  by  $S$  and  $A'$  by  $A$ , and we have  $X = [A \times S; \pi']$ . The isomorphism  $X \cong [A \times S; \pi]$  gives  $\pi' \sim \pi$ . Especially, if

$X$  is special, we have  $X=[N \times S; \pi'] \cong [N \times S; \pi]$  where  $\pi' \sim \pi$ . Using the last discussion of the previous section, we can conclude that  $F_s$  for  $[N \times S; \pi']$  is a regular permutation group on  $N$ . We obtained

**Corollary.** *Suppose that a null extension  $[A \times S; \pi]$  is transitive and special. Suppose also that  $S$  is effective. Then,  $F_s$  is a regular permutation group on  $A$  for every  $s$ .*

#### 4. An exceptional pseudosymmetric set

Suppose that  $S$  is a pseudosymmetric subset of  $P(A)$  for a set  $A$ . Define a mapping  $\mu$  of  $S \times S$  into  $P(A)$  by  $\mu(s, t)=s^{-1}t$ . We have shown in 2 that  $[A \times S; \mu]$  is a null extension. Denote  $(s \circ t) \circ u$  by  $s \circ t \circ u$ , etc. It can be shown by induction that, in  $[A \times S; \mu]$ , we have  $(a, s) \circ (b, t_1) \circ \dots \circ (c, t_n) = (a^o, s \circ t_1 \circ \dots \circ t_n)$ , where  $\rho=s^{-n}t_1 \dots t_n$ . Now we specialize  $A$  and  $S$  as follows. Let  $A=\{1, 2, 3, 4, 5, 6\}$  and  $S$  = the set of all permutations of order 2 that move every element of  $A$ .  $S$  consists of all conjugates of  $s=(12)(34)(56)$  and contains 15 elements. So, the order of  $A \times S$  is 90. Denote this  $[A \times S; \mu]$  by  $X_6$ . We show that  $X_6$  is transitive and exceptional. First, note that  $S$  is transitive and effective. To show that  $X_6$  is transitive, it is enough to show that  $F_s$  is transitive. Let  $t_1=(13)(24)(56)$ ,  $t_2=(14)(23)(56)$ ,  $t_3=(12)(35)(46)$  and  $t_4=(12)(36)(45)$ . Then,  $(1, s) \circ (1, t_1) \circ (1, t_2) = (1^o, s \circ t_1 \circ t_2) = (2, s)$ , where  $\rho=s^{-2}t_1t_2=t_1t_2$  and  $1^o=2$ . We have also  $(1, s) \circ (1, t_1) \circ (1, t_3) \circ (1, s) \circ (1, t_4) = (3, s)$ . In this way, we can show that 1 is mapped to any elements by elements of  $F_s$  and  $F_s$  is transitive. Next, we show that  $X_6$  is exceptional. Due to Corollary, it is enough to show that  $F_s$  is not regular. We have already shown that  $(1, s) \circ (1, t_1) \circ (1, t_2) = (2, s)$ . On the other hand,  $(5, s) \circ (1, t_1) \circ (1, t_2) = (5, s)$  as we can check. This implies that there is an element  $\bar{\tau}$  in  $F_s$  such that  $\bar{\tau} \neq 1$  and  $5^{\bar{\tau}}=5$ , and hence  $F_s$  is not regular. Therefore,  $X_6$  is transitive and exceptional. It is not hard to generalize this example. In place of 6 in the above, we take an even number  $m \geq 6$ , and construct  $X_m$  in the similar way. We can show that  $X_m$  is transitive and exceptional.

**REMARK.**  $X_6$  is actually a symmetric set. However, we do not call it an exceptional symmetric set. We did not define an exceptional symmetric set in this paper. Usually, a group is considered as a symmetric set via the composition  $a \circ b = ba^{-1}b$ . So, the structure of a group considered as a pseudosymmetric set is different from that considered as a symmetric set. For the latter, see [5].

**References**

- [1] D. Joyce: *Simple quandles*, J. Algebra **79** (1982), 307–318.
- [2] N. Nobusawa: *A remark on conjugacy classes in simple groups*, Osaka J. Math. **18** (1981), 749–754.
- [3] \_\_\_\_\_: *Some structure theorems on pseudo-symmetric sets*, Osaka J. Math. **20** (1983), 727–734.
- [4] R.S. Pierce: *Symmetric groupoids*, Osaka J. Math. **15** (1978), 51–76.
- [5] N. Umaya: *On symmetric structure of a group*, Proc. Japan Acad. **52** (1976), 174–176.

Department of Mathematics  
University of Hawaii  
Honolulu, Hawaii 96822  
U.S.A.