



Title	On nonsingular QF-3' rings with injective dimension ≤ 1
Author(s)	Sumioka, Takeshi
Citation	Osaka Journal of Mathematics. 1978, 15(1), p. 1-11
Version Type	VoR
URL	https://doi.org/10.18910/10756
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

ON NON-SINGULAR QF-3' RINGS WITH INJECTIVE DIMENSION ≤ 1

TAKESHI SUMIOKA

(Received November 26, 1976)

Let R be a ring and E an injective hull of R_R . We call R right QF-3' if every finitely generated submodule of E is torsionless. The dimension of a right R -module M is defined as the superior of the lengths n of chains $M_0 \subset M_1 \subset \cdots \subset M_n$ of submodules of M such that each factor module M_{i+1}/M_i is not torsion (under the Lambek torsion theory).

We shall characterize a finite dimensional (i.e. the dimensions of R_R and ${}_R R$ are finite) right and left QF-3' ring using the dimension of modules with some properties. Next, let R be a noetherian ring. Jans [4] proved that every finitely generated torsionless right R -module is reflexive if and only if R has injective dimension ≤ 1 as a left R -module. When R is (not necessarily noetherian) a commutative integral domain, results analogous to above one were proved by Matlis [6], where he further dealt with some properties on torsion modules. These properties, in case of noetherian QF-3' ring, were investigated by Zaks [11] and Sato [8]. In this note, we shall give characterizations of non-singular noetherian (or artinian) QF-3' ring with injective dimension ≤ 1 using the Lambek torsion theory.

Throughout we assume that if a ring R is said to be noetherian or QF-3', etc., we mean right and left noetherian or right and left QF-3', etc.. Moreover we assume that "every R -module" means "every right R -module and every left R -module," and " R has injective dimension ≤ 1 " does " R has injective dimension ≤ 1 as right and left R -modules," etc..

In this note, "torsion theory" means the Lambek torsion theory, which is cogenerated by an injective hull of $R_R({}_R R)$. We denote its torsion radical by t . Let R be a ring and M a right R -module. A chain

$$M_1 \subset M_2 \subset M_3 \subset \cdots (\text{resp. } M_1 \supset M_2 \supset M_3 \supset \cdots)$$

of submodules of M is called *t-chain* (of M) if M_{i+1}/M_i (resp. M_i/M_{i+1}) is not torsion for each i . A module M is called *finite dimensional* if any ascending *t-chain* and any descending *t-chain* of M terminate. A ring R is called *right finite dimensional* if R_R is finite dimensional, (refer Goldman [3] for these defini-

tions and their properties). The *dimension* of M is said to be *equal to or larger than* n and we denote it by $\dim M \geq n$, if M has a t -chain of length n . The *dimension* of M is said to be *equal to* n if $\dim M \geq n$ and $\dim M \not\geq n+1$, and in particular, $\dim M = 0$ if M is torsion, and $\dim M = \infty$ if $\dim M \geq n$ for any positive integer n .

The following lemma is immediate from the fact that if $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is a short exact sequence of right R -modules, B is torsion if and only if A and C are torsion.

Lemma 1. *Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be an exact sequence of right R -modules. Then we have $\dim M = \dim L + \dim N$.*

By Lemma 1, it is easy to see that a right R -module M is finite dimensional if and only if $\dim M = n$ for some integer $n \geq 0$. Let R be a right non-singular ring and M a torsion-free right R -module. Then we should note that this dimension of M is equal to the Goldie dimension of M . A right R -module M is called *finitely imbedded* (more briefly *FI*), if M is imbedded in some finitely generated right R -module. A ring R is called *right QF-3'* if R satisfies the following equivalent properties:

- (1) Any finitely generated submodule of E is torsionless, where E is an injective hull of R_R .
- (2) Every FI torsion-free right R -module is torsionless.
- (3) The maximal right quotient ring Q of R is a left quotient ring of R , and every finitely generated submodule of an injective hull $E(Q_Q)$ of Q_Q is torsionless.

The equivalence of these was proved by Masaike [5].

If M is a right R -module, we denote $\text{Hom}_R(M, R)$ and $\text{Ext}_R^1(M, R)$ by M^* and M_* , respectively, and these are naturally regarded as left R -modules.

Lemma 2. *Let R be a right QF-3' ring and M an FI right R -module. Then M is torsion if and only if $M^* = 0$.*

Proof. This is immediate from the definition of FI modules.

Lemma 3. *Let R be a finite dimensional right QF-3' ring. Then the following statements for a right R -module M are equivalent :*

- (1) M is finite dimensional torsionless.
- (2) M is FI torsionless.
- (3) M is FI torsion-free.
- (4) M is imbedded in a finitely generated free right R -module.

Proof. We shall only show that (1) implies (4), for $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$ are clear. Assume (1). It follows from the definition of "finite dimensional" that there is a finitely generated submodule N of M such that M/N is torsion. Hence

M^* is imbedded in N^* . Since N^* is imbedded in a finitely generated free left R -module, so is M^* , and in particular M^* is also finite dimensional. Therefore, using a similar discussion for M^* it follows that M^{**} is imbedded in a finitely generated free right R -module. Thus (4) is satisfied since the canonical map $M \rightarrow M^{**}$ is monomorphic.

Proposition 1. *Let R be a right QF-3' ring. Then the following statements are equivalent:*

- (1) *R is right finite dimensional.*
- (2) *R satisfies the ascending chain conditions on annihilator right ideals and annihilator left ideals.*

Proof. First we note that the condition (2) is clearly equivalent to a condition that R satisfies both the chain conditions on annihilator right ideals. If I is a right ideal of R , I is right annihilator if and only if R/I is torsionless. Hence, (1) implies (2). Assume (2), and let I and I' be right ideals of R such that $I \subset I'$ and I'/I is not torsion. For a subset A of R , we denote by $r(A)$ (resp. $l(A)$) the right (resp. left) annihilator ideal of A . Put $J = r l(I)$ and $J' = r l(I')$, and let K/I be the torsion submodule of R/I . Then, since R/J is torsion-free, J/I contains K/I , i.e., $J \supset K$. By the assumption, a cyclic torsion-free right R -module R/K is torsionless, and hence K is an annihilator right ideal of R . Therefore we have $K=J$ from $I \subset K \subset I$, and so J/I is torsion. Since I'/I is not torsion, so is J'/J , and in particular $J \subsetneq J'$. This implies by the condition (2) that any ascending t -chain and any descending t -chain of R_R terminate. Thus the condition (1) is satisfied.

REMARKS (i) In Proposition 1, (1) implies (2) without the assumption that R is right QF-3'. Hence, for a semi-prime ring R , R is right finite dimensional if and only if R is right Goldie.

(ii) From right-left symmetry of the condition (2) in Proposition 1, it follows that a left QF-3' and right finite dimensional ring is also left finite dimensional.

Lemma 4. *Let R be a right QF-3' ring, M an FI right R -module and N a submodule of M . Then the monomorphism $(M/N)^* \rightarrow M^*$ derived by the canonical epimorphism $M \rightarrow M/N$ is isomorphic if and only if N is torsion.*

Proof. "If" part is trivial, and we shall show "only if" part. Assume that the map $(M/N)^* \rightarrow M^*$ is isomorphic. In case M is torsionless, we easily see $N=0$. Now let M be an FI right R -module and M' the torsion submodule of M . Then, we have a following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \rightarrow & M' \cap N & \longrightarrow & M' & \longrightarrow & M' + N/N \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & N & \longrightarrow & M & \longrightarrow & M/N \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \rightarrow & N/M' \cap N & \rightarrow & M/M' & \rightarrow & M/M' + N \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0.
\end{array}$$

Moreover, from this we obtain the following commutative diagram with exact rows and isomorphic columns:

$$\begin{array}{ccccccc}
0 & \rightarrow & (M/M' + N)^* & \rightarrow & (M/M')^* & \rightarrow & (N/M' \cap N)^* \\
& & \wr & & \wr & & \wr \\
0 & \longrightarrow & (M/N)^* & \longrightarrow & M^* & \longrightarrow & N^*.
\end{array}$$

Since R is right QF-3', M/M' is torsionless, and so we have $N/M' \cap N = 0$ by the above case, and hence $N \subset M'$. Thus N is torsion.

Lemma 5. *Let R be a right finite dimensional right QF-3' ring. If M is an FI right R -module, we have $\dim M_R \leq \dim_R (M^*)$.*

Proof. In case $\dim M = 0$, the assertion is trivial. In case $\dim M = 1$, we have $\dim M^* \geq 1$, since M^* is non-zero torsionless by Lemma 2. Assume that the assertion is satisfied for $\dim M \leq n-1$, and let the dimension of M be equal to n . Then there is a submodule L of M such that $\dim L = 1$ and so $\dim M/L = n-1$. Then the map $(M/L)^* \rightarrow M^*$ is not isomorphic by Lemma 4, and hence we have an exact sequence $0 \rightarrow (M/L)^* \rightarrow M^* \rightarrow K \rightarrow 0$ with non-zero torsion-free left R -module K . By inductual assumption, we have $\dim (M/L)^* \geq n-1$. Therefore $\dim M^* \geq \dim M$.

Let M and N be right R -modules. For a homomorphism $f: M \rightarrow N$, we denote its dual map by $f^*: N^* \rightarrow M^*$. On the other hand, by $\varphi_M: M \rightarrow M^{**}$ we denote the canonical homomorphism of M to M^{**} .

Theorem 1. *Let R be a right finite dimensional ring. Then the following conditions are equivalent:*

- (1) R is QF-3'.
- (2) $\dim M = \dim M^*$ for every FI module M .
- (3) $\dim M = \dim M^*$ for every FI torsion-free module M .

Proof. (1) implies (2). Let $\dim R_R = n$. Since, by Lemma 5 and Remark (ii), $\dim R_R \leq \dim_R (R^*) \leq \dim (R^{**})_R$, we have $\dim R_R = \dim_R R$. Therefore we only show that $\dim M = \dim M^*$ for every FI right R -module M , because the same arguments hold for FI left R -modules. If $\dim M = 0$, this assertion is

trivial. Suppose that M is a cyclic right R -module and N is the kernel of an epimorphism $R \rightarrow M$. If the dimension of M is n , we have $M_R \simeq R_R$, and so $\dim M = \dim M^*$. If $\dim M = n-1$, and so $\dim N = 1$, then we have $\dim M^* = n-1$ by Lemma 4 and Lemma 5. Suppose $\dim M = n-2$ and $\dim N = 2$. Let L be a submodule N such that $\dim L = 1$. Since R/L is a cyclic right R -module with dimension $n-1$, we have $\dim (R/L)^* = n-1$ as the above case. Therefore, by Lemma 4 and Lemma 5, $\dim M^* = n-2$. Iterating this discussion we have $\dim M = \dim M^*$ for any cyclic right R -module M . Next suppose that M is a finitely generated right R -module. Then, using induction on the number of generators of M , the assertion is easily showed by Lemma 5. In the case where M is an FI right R -module, there is a finitely generated submodule N of M such that M/N is torsion, i.e., $\dim M = \dim N$. Then, from the exact sequence $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$, we have the monomorphism $M^* \rightarrow N^*$ and so $\dim M^* \leq \dim N^*$. Since N is finitely generated, $\dim N = \dim N^*$. Thus we have $\dim M = \dim M^*$ by Lemma 5.

(2) implies (3). This is trivial.

(3) implies (1). Since $\dim {}_R R = \dim R_R$, R is left finite dimensional. Let L be a finitely generated torsion-free left R -module, and let $0 \rightarrow K \rightarrow G \rightarrow L \rightarrow 0$ be an exact sequence with a finitely generated free left R -module G . Then we have an exact sequence $0 \rightarrow L^* \rightarrow G^* \rightarrow K^* \rightarrow L_* \rightarrow 0$, which implies $\dim L_* = 0$ by the assumption. Next let M be a finitely generated torsion-free right R -module. Then we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & J & \rightarrow & F & \rightarrow & M \rightarrow 0 \\ & & \downarrow & & \downarrow \varphi_F & & \downarrow \varphi_M \\ 0 & \rightarrow & J' & \rightarrow & F^{**} & \rightarrow & M^{**} \rightarrow (F^*/J')_* \rightarrow 0, \end{array}$$

where F is a finitely generated free right R -module and J' is the annihilator of J in F^* . Since F^*/J' is torsionless, $\dim (F^*/J')_* = 0$, so $\dim \text{Im } \varphi_M = \dim M$. This implies that φ_M is monomorphic. Thus R is right QF-3' and similarly left QF-3'.

Corollary 1. *Let R be a finite dimensional QF-3' ring. Then we have:*

- (1) *If M is a finite dimensional right R -module, M^* is reflexive.*
- (2) *Let M be a finitely generated right R -module, and let $f: L \rightarrow M$ be a monomorphism. Then M_* and $\text{Coker } f^*$ are torsion.*

Proof. (1) This follows from a fact that the composition map of natural maps $\varphi_M^*: M^* \rightarrow M^{***}$ and $(\varphi_M)^*: M^{***} \rightarrow M^*$ is identity, and Theorem 1. (2) The assertion for M_* is immediate from the equivalence of (1) and (2) in Theorem 1. Hence $(M/L)_*$ is also torsion, and so the left R -module $\text{Coker } f^*$ imbedded in $(M/L)_*$ is torsion.

Proposition 2. *Let R be a finite dimensional QF-3' ring. Then, for a finite dimensional torsion-free right R -module M , $\dim M = \dim M^*$ if and only if M is torsionless.*

Proof. Suppose $\dim M = \dim M^*$ and $\dim M = n$. We shall show by induction on n that the canonical map $M \rightarrow M^{**}$ is monomorphic. The result holds for $n=1$, since $M^* \neq 0$. Assume the result for positive integer $\leq n-1$. There exists a submodule N of M such that $\dim N=1$ and M/N is a torsion-free right R -module with $\dim M/N = n-1$. Then an exact sequence $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$ derives the exact sequence $0 \rightarrow (M/N)^* \rightarrow M^* \rightarrow L \rightarrow 0$, where L is the image of the derived map $M^* \rightarrow N^*$. Now, let A be a finite dimensional right R -module. Then there is a finitely generated submodule B of A such that A/B is torsion, and so $\dim A^* \leq \dim B^*$. Hence we have $\dim A^* \leq \dim A$, since $\dim B = \dim B^*$ by Theorem 1. Therefore in particular, $\dim N^* \leq 1$ and $\dim (M/N)^* \leq n-1$. On the other hand, $\dim M^* = n$. Consequently $\dim N^* = 1$ and $\dim (M/N)^* = n-1$, and so N and M/N are torsionless by inductive assumption. Moreover $\dim L = 1$, so the dual map $\iota^*: N^{**} \rightarrow L^*$ of the inclusion $\iota: L \rightarrow N^*$ is monomorphic. Consider a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & N & \rightarrow & M & \rightarrow & M/N \rightarrow 0 \\ & & \downarrow & & \downarrow \varphi_M & & \downarrow \varphi_{M/N} \\ 0 & \rightarrow & L^* & \rightarrow & M^{**} & \rightarrow & (M/N)^{**} \end{array}$$

with exact rows, where $N \rightarrow L^*$ is the composition of the maps $\varphi_N: N \rightarrow N^{**}$ and $\iota^*: N^{**} \rightarrow L^*$. Then the middle map is monomorphic, since so are both the out sides. Thus M is torsionless. The converse is followed from Lemma 3 and Theorem 1.

The following lemma is a slight extension of Matlis [6, Lemma, p. 19], and this extension is essentially necessary in the proof of Theorem 3.

Lemma 6. *Let R be a ring, and let $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ be an exact sequence of right R -modules. Suppose that N is reflexive, M is torsion-free and L , L_* and L_{**} are simple torsion. Then M is reflexive.*

Proof. By the sequence

$$0 \rightarrow N \xrightarrow{f} M \xrightarrow{g} L \rightarrow 0 \quad \dots\dots\dots (A),$$

an exact sequence $0 \rightarrow M^* \xrightarrow{f^*} N^* \xrightarrow{\delta} L_*$ is obtained. Suppose that f^* is isomorphic. Then its dual map f^{**} is also isomorphic. Since, in a commutative diagram

$$\begin{array}{ccc} N & \xrightarrow{f} & M \\ \downarrow \varphi_N & f^{**} & \downarrow \varphi_M \\ N^{**} & \xrightarrow{\quad} & M^{**}, \end{array}$$

φ_N and f^{**} are isomorphic, the exact sequence (A) splits. This is a contradiction because M is torsion-free and L is torsion. Thus f^* is not isomorphic, that is $\delta \neq 0$, which implies that δ is epimorphic, since L_* is simple. Thus we have an exact sequence $0 \rightarrow N^{**} \xrightarrow{f^{**}} M^{**} \xrightarrow{\delta'} L_{**}$. δ' is epimorphism, since f^{**} is not isomorphic. Therefore we have a map $\eta: L \rightarrow L_{**}$ and the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & N & \xrightarrow{f} & M & \xrightarrow{g} & L \rightarrow 0 \\ & & \downarrow \varphi_N & & \downarrow \varphi_M & & \downarrow \eta \\ 0 & \rightarrow & N^{**} & \xrightarrow{f^{**}} & M^{**} & \xrightarrow{\delta'} & L_{**} \rightarrow 0. \end{array}$$

Assume that φ_M is not epimorphic. Put $M_0 = \text{Im } \varphi_M$. Since φ_N is isomorphic, $\text{Im } f^{**} \subset M_0$. Therefore, since L_{**} is simple, $M_0 = \text{Im } f^{**}$, which implies (A) splits. This is a contradiction. Thus φ_M is an epimorphism. Therefore η is isomorphic, and consequently φ_M is isomorphic.

Let R be a ring, and let M be a finitely generated torsion right R -module such that M_* is torsion. Consider an exact sequence $0 \rightarrow L \rightarrow P \rightarrow M \rightarrow 0$ with finitely generated projective right R -module P . Then we have an exact sequence $0 \rightarrow P^* \rightarrow L^* \rightarrow M^* \rightarrow 0$ derived from the above sequence. Therefore, since M^* is torsion, we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & L & \rightarrow & P & \rightarrow & M \rightarrow 0 \\ & & \downarrow \varphi_L & & \downarrow \varphi_P & & \downarrow \\ 0 & \rightarrow & L^{**} & \rightarrow & P^{**} & \rightarrow & M_{**} \end{array}$$

with exact rows, where $M \rightarrow M_{**}$ is the map induced by the left side square:

$$\begin{array}{ccc} L & \longrightarrow & P \\ \downarrow & & \downarrow \\ L^{**} & \longrightarrow & P^{**}. \end{array}$$

We denote this map by $\mu_M: M \rightarrow M_{**}$. Clearly μ_M does not depend on selection of the finitely generated projective right R -module P .

Theorem 2. *Let R be a noetherian non-singular QF-3' ring with the maximal quotient ring Q . Then the following statements are equivalent:*

- (1) *For every cyclic torsion R -module M , μ_M is isomorphic.*
- (2) *Any finitely generated submodule of Q is reflexive.*
- (3) *R has injective dimension ≤ 1 .*

Proof. (3) implies (1) and (2). These follows from Sato [8] and Jans [4]. (2) implies (3). Let I be an essential right ideal of R . Since any essential

right ideal is reflexive and I^* is a noetherian left R -module, R/I has a composition series. Hence any cyclic torsion right R -module, and consequently, any finitely generated torsion right R -module has a composition series. Next, we have an isomorphism $\text{Hom}_R(R, Q) \simeq \text{Hom}_R(I, Q)$, since R/I is torsion and Q is injective. On the other hand, there is a monomorphism $\text{Hom}_R(I, R) \rightarrow \text{Hom}_R(I, Q)$. Thus I^* is imbedded in ${}_R Q$, and so every submodule of I^* is reflexive by the assumption. Therefore, since we have an exact sequence $0 \rightarrow R^* \rightarrow I^* \rightarrow (R/I)_* \rightarrow 0$, we easily see that if R/I is simple torsion, so is $(R/I)_*$. Similarly, if S is simple torsion as a left R -module, so is S_* as a right R -module. Now let M be an essential submodule of a finitely generated free right R -module F . Then there is an essential submodule N of M such that N is isomorphic to a finite direct sum of right ideals of R , and so N is reflexive. Since F/N is finitely generated torsion, F/N and consequently M/N are artinian. Therefore, from Lemma 6, it follows by induction on the length of composition series of M/N that M is reflexive. Thus, by Jans [4], we have $\text{inj. dim. } {}_R R \leq 1$. Similarly, we can show $\text{inj. dim. } R_R \leq 1$.

(1) implies (3). Let I be an essential right ideal of R . Then, we have a commutative diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & I & \longrightarrow & R & \longrightarrow & R/I \longrightarrow 0 \\ & & \downarrow \varphi_I & & \downarrow \varphi_R & & \downarrow \mu_{R/I} \\ 0 & \rightarrow & I^{**} & \rightarrow & R^{**} & \rightarrow & (R/I)^{**} \rightarrow (I^*)_* \rightarrow 0. \end{array}$$

Since $\mu_{R/I}$ is isomorphic, I is reflexive and $(I^*)_* = 0$. Therefore, it is easy to see that S_* is simple torsion for every simple torsion module S . Thus the assertion is showed as the proof of (2) \Rightarrow (3).

REMARKS (i) From the above commutative diagram, we obtain the following exact sequence:

$$0 \rightarrow I \rightarrow I^{**} \rightarrow R/I \rightarrow (R/I)^{**} \rightarrow \text{Ext}_R^1(I^*, R) \rightarrow 0.$$

If R is an integral domain, this may be identified with the exact sequence of Matlis [6, Theorem 1.2].

(ii) Let R be a noetherian integral domain with quotient field Q . If A is a finitely generated submodule of Q , A is isomorphic to an ideal of R . Therefore, the equivalence of (2) and (3) in Theorem 2 is an extension of that of (2) and (4) in Matlis [6, Theorem 3.8].

Proposition 3. *Let R be a non-singular right finite dimensional ring. Suppose that for every finitely generated torsion R -module A , A_* is torsion. Then R is QF -3'.*

Proof. From the assumption, it is easily showed that $\dim {}_R R = \dim R_R$, and

$\dim I^* = 1$ for every uniform right ideal I of R , and in particular R is left finite dimensional. Now, let L be a finitely generated torsion-free right R -module. Since R_R is non-singular, there is a finite number of uniform right ideals whose direct sum is isomorphic to an essential submodule of L . Consequently, we have $\dim L^* = \dim L$ from the assumption. Next, let N be an FI torsion-free right R -module. Then N is clearly imbedded in a finitely generated torsion-free right R -module M . Since M is finite dimensional, $N \oplus K$ is essential in M with some finitely generated right R -module K . From the above case, we have $\dim M^* = \dim M$ and $\dim K^* = \dim K$, and therefore $\dim N^* = \dim N$. Similar arguments also hold for FI torsion-free left R -modules. Consequently, R is QF-3' by Theorem 1.

Lemma 7. *Let R be a noetherian ring. Then the following conditions are equivalent :*

- (1) *If M is a finitely generated torsion module, then so is M_* , and M_{**} is isomorphic to M .*
- (2) *If M is a finitely generated torsion module, then so is M_* , and μ_M is isomorphic.*

Proof. It is trivial that (2) implies (1). Assume (1). Then it is clear that if S is a simple torsion module, S_* is non-zero. First we show that if S is simple torsion as a right R -module, so is S_* as a left R -module. Let N be a maximal submodule of S_* , and $0 \rightarrow N \rightarrow S_* \rightarrow S_*/N \rightarrow 0$ the natural exact sequence. Then we have the derived exact sequence $0 \rightarrow (S_*/N)_* \rightarrow S_{**}$, since N is torsion. By the assumption, S_{**} is isomorphic to S , and so S_{**} is simple. This implies $(S_*/N)_* \simeq S_{**} \simeq S$. Therefore $S_* \simeq (S_*N)_{**} \simeq S_*/N$, and so S_* is simple. If an R -module M has a composition series, we denote its length by $l(M_*)$. Let M be a torsion R -module with $l(M) = n$. Then, by induction on n , we can easily show $l(M_*) = n$. Now, let M be a finitely generated torsion right R -module. Then there are a finitely generated free right R -module F and its submodule K such that $F/K \simeq M$. In order to show that M is artinian, let

$$F \supset K_1 \supset K_2 \supset \cdots \supset K \quad \text{.....(A)}$$

by a chain of submodules of F such that $l(F/K_i) = i$ for each i . Since F/K is a finitely generated torsion right R -module, we may assume that we have a chain

$$F^* \subset K_1^* \subset K_2^* \subset \cdots \subset K^* \quad \text{.....(B)}$$

of submodules of K^* . Then we have $K^*/F^* \simeq (F/K)_*$ and $K_i^*/F^* \simeq (F/K_i)_*$, and so K^*/F^* is also finitely generated torsion and $l(K_i^*/F^*) = i$. On the other hand K^* is noetherian, which shows that the chain (B) and consequently (A) terminate. Now, we show that the canonical inclusion $\varphi_K: K \rightarrow K^{**}$ is an

isomorphism. Since K^*/F^* is torsion and $F \simeq F^{**}$, we may assume that $K \subset K^{**} \subset F$. Then we have $F^* \subset K^{***} \subset K^*$. Therefore $(\varphi_K)^*: K^{***} \rightarrow K^*$ is isomorphic, since $(\varphi_K)^*$ is always epimorphic. Which implies $l(F/K) = l(F/K^{**})$, and so φ_K is isomorphic. We have, however, $l(M) = l(M^{**})$. Thus μ_M is isomorphic.

The following Corollary is immediate from Proposition 3, Lemma 7 and Sato [8]. (It seems that an isomorphism $M \simeq \text{Ext}_R^1(\text{Ext}_R^1(M, R), R)$ in Sato [8, Theorem 2.3] means μ_M .)

Corollary 2. *Let R be a noetherian non-singular ring. Then the following statements are equivalent :*

- (1) *If M is a finitely generated torsion module, so is M_* , and $M_{**} \simeq M$.*
- (2) *R is a QF-3' ring with injective dimension ≤ 1 .*

A ring R is called *right QF-3* if R has a minimal faithful right R -module. The following lemma is a slight extension of Rutter [7, Corollary 3].

Lemma 8. *Let R be a right perfect ring satisfying the ascending chain condition on annihilator right ideals. If R is right QF-3', then R is QF-3.*

Proof. By Faith [2] R is semi-primary, and it follows from the proof of Proposition 1 that R is right finite dimensional. Let M be a finitely generated submodule of an injective hull $E(R_R)$ of R_R . Then $\bigcap_{f \in M^*} \text{Ker } f = 0$ since M is torsionless. For every subset A of M^* , $M / \bigcap_{f \in A} \text{Ker } f$ is torsionless. But M is finite dimensional, which implies that there exist f_1, \dots, f_n in M^* such that $\bigcap_{i=1}^n \text{Ker } f_i = 0$, and so M is imbedded in a free right R -module. Therefore $E(R_R)$ is projective by Rutter [7], and R is right QF-3. Thus, by Colby-Rutter [1], R is QF-3.

Corollary 3. *Let R be an artinian ring. Then the following statements are equivalent :*

- (1) *R is a non-singular QF-3' ring with injective dimension ≤ 1 .*
- (2) *R is hereditary QF-3.*

Proof. Assume (1). By Lemma 8 (or Rutter [7]), R is QF-3. Since R is non-singular and artinian, R has the semi-simple maximal quotient ring. Therefore, by Sumioka [9], R is hereditary. The converse is clear since any QF-3 ring is QF-3' (see Tachikawa [10], p. 47).

Theorem 3. *Let R be a non-singular artinian ring. Then the following conditions are equivalent :*

- (1) *If S is a simple torsion module, then so is S_* .*

(2) R is hereditary QF-3.

Proof. It follows from Sato [8] that (2) implies (1). Assume (1). Let M be a finitely generated torsion-free right R -module with $l(M)=n$, where $l(M)$ expresses the length of a composition series of M . By induction on n , we shall show that M is reflexive. Suppose $l(M)=1$. Then, since M is a non-singular simple module, M is projective and in particular reflexive. Let $l(M)=n$, and assume that the result holds for $n-1$. Let N be a maximal submodule of M , and consider the exact sequence $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$. By inductive assumption, N is reflexive. If M/N is simple torsion-free, then M/N is projective, and so the above sequence splits, and consequently M is reflexive. If M/N is simple torsion, then the result holds, by Lemma 6. Thus every finitely generated torsion-free right R module is reflexive. The similar statement on left R -module is also true. Therefore R is hereditary QF-3 by Jans [4] and Corollary 3.

OSAKA CITY UNIVERSITY

References

- [1] R.R. Colby and E.A. Rutter: *Generalization of QF-3 Algebras*, Trans, Amer. Math. Soc. **153** (1971), 371–386.
- [2] C. Faith: *Rings with ascending condition on annihilators*, Nagoya Math. J. **27** (1966), 179–191.
- [3] O. Goldman: *Elements of non commutative arithmetic I*, J. Algebra **35** (1975), 308–341.
- [4] J.P. Jans: *Duality in Noetherian rings*, Proc. Amer. Math. Soc. **12** (1961), 829–835.
- [5] K. Masaike: *On quotient rings and torsionless modules*, Sci. Rep. Tokyo Kyoiku Daigaku A. **11** (1971), 26–30.
- [6] E. Matlis: *Reflexive domains*, J. Algebra **8** (1968), 1–33.
- [7] E.A. Rutter: *A characterization of QF-3 rings*, Pacific J. Math. **51** (1974), 533–536.
- [8] H. Sato: *A duality of torsion modules over a QF-3 one-dimensional Gorenstein ring*, Sci. Rep. Tokyo Kyoiku Daigaku A. **13** (1975), 28–36.
- [9] T. Sumioka: *A characterization of the triangular matrix rings over QF rings*, Osaka J. Math. **12** (1975), 449–456.
- [10] H. Tachikawa: *Quasi-Frobenius rings and generalization*, Lecture Note in Math. 351, Springer, Berlin, 1973.
- [11] A. Zaks: *Hereditary noetherian rings*, J. Algebra **29** (1974), 513–527.

