OUKA
The University of Osaka
Institutional Knowledge Archive

Title	Z/kZ-finiteness for certain S^1-spaces
Author(s)	Sumi, Toshio
Citation	Osaka Journal of Mathematics. 1992, 29(3), p. 607-615
Version Type	VoR
URL	https://doi.org/10.18910/10890
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Sumi, T.
Osaka J. Math.
29 (1992), 607-615

Z/kZ-FINITENESS FOR CERTAIN S^1-SPACES

TOSHIO SUMI

(Received July 25, 1991)
(Revised September 6, 1991)

Introduction

Let G-$\mathcal{FDCW}$ denote the category of G-spaces having the G-homotopy type of a finitely dominated G-CW complex for a compact Lie group G. Lück [8] has introduced a functor Wa^G from G-$\mathcal{FDCW}$ into the category of abelian groups and has realized the equivariant finiteness obstriction as the element $w^G(X)$ in $Wa^G(X)$. That is, a finitely dominated G-CW complex X is G-homotopy equivalent to a finite G-CW complex if and only if $w^G(X)=0$. When G is the trivial group, there is an isomorphism from $Wa^G(X)$ to the reduced projective group $\tilde{K}_0(\boldsymbol{Z}[\pi_1(X)])$ which sends the element $w^G(X)$ to the Wall's finiteness obstruction ([14]).

Anderson [1] and Ehrlich [4] have studied a sufficient condition for $w^{\{1\}}(E)=0$ for some fibration $E\to B$ with fiber S^1. Munkholm, Pedresen [11], Lück [6, 7, 9] and others have studied the transfer map $\tilde{K}_0(\boldsymbol{Z}[\pi_1(B)])\to\tilde{K}_0(\boldsymbol{Z}[\pi_1(E)])$. The purpose of this paper is to get a sufficient condition for $w^L(X)=0$ for a S^1-space X and a finite cyclic group L.

We call G-maps $f_0\colon Y_0\to X$ and $f_0\colon Y_4\to X$ equivalent if there exists a commutative diagram



such that (Y_1, Y_0) and (Y_3, Y_4) are relatively finite G-CW complexes, and $Y_1\to Y_2$ and $Y_3\to Y_2$ are G-homotopy equivalences. The group $Wa^G(X)$ consists of equivalence classes $[f\colon Y\to X]$ of the set of G-maps $f\colon Y\to X$ with Y finitely dominated and $w^G(X)$ is the equivalence class containing the identity 1_X of X. The additive structure on $Wa^G(X)$ is given by a disjoint sum:

$$[f\colon Y\to X]+[g\colon Z\to X]=[f\amalg g\colon Y\amalg Z\to X]$$

Let K be a closed subgroup of G. For a K-space X, we define $\mathrm{ind}_K^G X$ as the orbit space $G\times_K X$ of the product space $G\times X$ with respect to the K-action $k\cdot(g,x)=(gk^{-1},kx)$. For a K-map $f: X\to Y$, we have an induced map $1\times_K f$: $G\times_K X\to G\times_K Y$, denoted by $\mathrm{ind}_K^G f$. The induction functor ind_K^G induces a transformation Ind_K^G: $(Wa^K, w^K)\to(Wa^G, w^G)$. To consider G-maps as K-maps implies a transformation Res_K^G: $(Wa^G, w^G)\to(Wa^K, w^K)$.

Throughout this paper we denote by ν the restriction of the G-action of X to $G\times\{x_0\}$. If $G=S^1$ and X is connected, the order of the image of $H_1(\nu;\boldsymbol{Z})$ is independent of taking a point x_0 of X.

Our main results are as follows:

Theorem A. *Let X be a connected S^1-CW complex which has finitely many orbit types. Suppose that the S^1-map $\nu: S^1\to X$ defined as above induces a monomorphism $H_1(\nu;\boldsymbol{Z})$ between 1-dimensional homology groups. Then there exist a proper subgroup K of S^1 and a K-CW complex Y such that $S^1\times_K Y$ is S^1-homotopy equivalent to X.*

Theorem B. *Let $G=S^1$ and let X be a connected G-space. If the above defined G-map $\nu: G\to X$ induces an injective homomorphism $H_1(\nu;\boldsymbol{Z})$, then the restriction homomorphism $\mathrm{Res}_H^G(X)$: $Wa^G(X)\to Wa^H(X)$ is trivial for any proper subgroup H of G.*

This paper is organized as follows. Let $F\to E\xrightarrow{p}B$ be a G-fibration with fibre F ([15]). In [13], we have constructed a transfer $p^!$: $Wa^G(B)\to Wa^G(E)$. But this homomorphism does not always send $w^G(B)$ to $w^G(E)$. It is originated from that (Wa^G, w^G) is not a functorial additive invariant for G-$\mathcal{TDCW}$. In section 1 we study a K-CW structure on $p^{-1}(1K)$ for a G-CW complex E which has a G-map $E\to G/K$. In section 2 we show that if $G\times_K X$ has the G-homotopy type of a finite G-CW complex then X has the K-homotopy type of a finite K-CW complex. In section 3, we prove Theorem A in the case where X is free. We use the fact that $\pi_1(X/K)$ has the subgroup $\pi_1(S^1/K)$ as a direct summand for some closed subgroup K of G. The last section consists of the proof of the main theorems. The proof of Theorem A is obtained from applying the free case.

1. K-CW structure on X of a G-CW complex $G\times_K X$

Let G be a compact Lie group. We study a space $p^{-1}(\{pt\})$ for a G-map p from a G-space onto an orbit space G/K of G. We note that it has a canonical K-action.

Proposition 1.1. (cf.[12]) *A G-map $p: E\to B$ is a G-fibration if and only if $p^K: E^K\to B^K$ is a fibration for any closed subgroup K of G.*

Proof. This follows essentially from Theorem 4.1 in [2].

Since $Y\to G\times_K Y\to G/K$ is a G-fibration ([15, 13]), we have that $Y^L\to (G\times_K Y)^L\to (G/K)^L$ is a fibration for any closed subgroup L of G.

We symbolize 1 as the identity element of G. For any G-map $p: X\to G/K$ it is a G-fibration with fibre $p^{-1}(1K)$. The following lemma is a key to show the main theorems. It implies that a G-map $p: X\to G/K$ with X a G-CW complex is G-homotopy equivalent to a G-fibration whose fibre is a K-CW complex.

Lemma 1.2. *Let X be a G-CW complex which has a G-map $p: X\to G/K$. A K-CW complex can be constructed from the G-CW structure of X such that the K-space $V=p^{-1}(1K)$ is homotopy equivalent to it. In particular it is a finite K-CW complex if X is a finite G-CW complex.*

Proof. Clearly we have $G\times_K V$ and X are G-homeomorphic. Then we construct a K-CW complex W and a K-homotopy equivalence $W\to V$ by induction on the dimension of cells of X. By the existence of the G-map p, we obtain that L is subconjugate to K for any isotropy subgroup L of G in X. We can regard a 0-cell $G/L\times e^0$ of X as $G\times_K K/aLa^{-1}\times e^0$ for $a\in G$ with $aLa^{-1}\leq K$. Suppose that $X=G\times_K Y\cup_\phi G/L\times e^n$ for some K-CW complex Y. Let C be a connected component of $(G/K)^L$ which contains $p^L\circ\phi(1L\times \dot{e}^n)$. Take $aK\in C$ and let $\psi: G/aLa^{-1}\to G/L$ be the canonical G-map. Then the pushout of

$$\begin{array}{ccc} G/aLa^{-1}\times \dot{e}^n & \hookrightarrow & G/aLa^{-1}\times e^n \\ \downarrow & & \\ G\times_K Y & & \end{array}$$

is G-homotopy equivalent to X. Then we can assume that $L\leq K$ and $p^L\circ\phi(1L\times \dot{e}^n)$ is contained in the connected component of $1K$.

$$\begin{array}{ccccc} 1L\times e^n & \hookleftarrow & 1L\times \dot{e}^n & & \\ & & \downarrow \phi|_{1L\times \dot{e}^n} & & \\ Y^L & \to & (G\times_K Y)^L & \xrightarrow{p^L} & (G/K)^L \end{array}$$

Since the map $p^L\circ\phi|_{1L\times \dot{e}^n}$ is homotopic to a constant map, there is a map $\sigma: 1L\times e^n\to (G/K)^L$ such that σ coincides with $p^L\circ\phi$ over $1L\times \dot{e}^n$ and $\sigma(0)=1K$. We define a map $\tau: 1L\times e^n\times I\to (G/K)^L$ as $\tau(s, t)=\sigma((1-t)s)$. Since $(G\times_K Y)^L\to (G/K)^L$ is a fibration with fibre Y^L, there exists a homotopy $F: 1L\times \dot{e}^n\times I\to (G\times_K Y)^L$ such that $F_0=\phi|_{1L\times \dot{e}^n}$ and $F_1(1L\times \dot{e}^n)\subseteq Y^L$. This map can be canonically extended to a G-map Φ from $G/L\times \dot{e}^n\times I$ to $G\times_K Y$. Let W be a K-CW complex obtained from the following pushout.

$$\begin{array}{ccc} K/L\times\dot{e}^n & \hookrightarrow & K/L\times e^n \\ \downarrow & & \downarrow\Phi_1 \\ Y & & \end{array}$$

By the property of pushout, we get a K-map $k\colon W\to V$.

$$\begin{array}{ccccc} W & \hookrightarrow & G\times_K W & \longrightarrow & G/K \\ \downarrow k & & \downarrow \mathrm{ind}_K^G k & & \| \\ V & \hookrightarrow & X=G\times_K V & \xrightarrow{p} & G/K \end{array}$$

Since the G-map $\mathrm{ind}_K^G k$ is a G-homotopy equivalence, we have the K-map k is a K-homotopy equivalence.

Theorem 1.3. *Let $f\colon Y\to X$ be a K-map between K-CW complexes and $(V, G\times_K Y)$ be a G-CW pair. If there is a G-map $g\colon V\to G\times_K X$ which is an extension of the G-map $1\times_K f$, then there exists a K-map $k\colon W\to X$ unique up to K-homotopy equivalence which fullfills the following conditions.*

(1) *(W, Y) is a K-CW pair.*

(2) *There is a G-homotopy equivalence $h\colon (G\times_K W, G\times_K Y)\to(V, G\times_K Y)$ such that $\mathrm{ind}_K^G k$ and $g\circ h$ are G-homotopic.*

(3) *The number of the relative cells of (W, Y) equals that of $(V, G\times_K Y)$.*

2. Induction homomorphism

Let $D^G(X)$ be the set of equivalence classes of the set of G-maps $f\colon Y\to X$ where Y has the G-homotopy type of a G-CW complex. Here the equivalence relation is defined as in introduction. For a G-map $f\colon Y\to X$, we denote by $[f\colon Y\to X]$ its represented element of $D^G(X)$. The additive structure on $D^G(X)$ is given as the one of $Wa^G(X)$. A G-map from a finite G-CW complex to X represents the zero element of $D^G(X)$. Then $D^G(X)$ is a semigroup and we obtain a map $Wa^G(X)\to D^G(X)$ which preserves the abelian structures.

Lemma 2.1. *The element of $D^G(X)$ represented by the identity map of X is invertible if and only if X is finitely dominated.*

Proof. The "if" part is trivial and then we show the "only if" part. There is the commutative diagram

$$\begin{array}{ccccccc} X\amalg Y_1 & \xrightarrow{i} & Y_2 & \xleftarrow{h} & Y_3 & & \\ & \searrow & \downarrow & \swarrow_r & & & \\ & & X & & & & \end{array}$$

such that $(Y_2, X \coprod Y_1)$ is a relatively finite G-CW complex, Y_3 is a finite G-CW complex, and h is a G-homotopy equivalence. Let $h^{-1}: Y_2 \to Y_3$ be the G-homotopy inverse of h. Then r is a domination with section $h^{-1} \circ i$.

Proposition 2.2. *Let K be a closed subgroup of G. A K-space X is a finitely dominated K-space if and only if $G \times_K X$ is a finitely dominated G-space.*

Proof. Suppose $G \times_K X$ is dominated by a finite G-CW complex Y_3. There is a commutative diagram such that f_3 is a domination with section $h^{-1}ij$.

$$\begin{array}{ccccccc} G \times_K X & \overset{j}{\hookrightarrow} & (G \times_K X) \coprod Y_1 & \overset{i}{\hookrightarrow} & Y_2 & \overset{h}{\leftarrow} & Y_3 \\ & & {\scriptstyle id \coprod f_1}\searrow & & \downarrow{\scriptstyle f_2} & \swarrow{\scriptstyle f_3} & \\ & & & & G \times_K X & & \end{array}$$

We let $Z_1 = (pf_1)^{-1}(1K)$, $Z_2 = (pf_2)^{-1}(1K)$, and $Z_3 = (pf_3)^{-1}(1K)$ for short, where $p: G \times_K X \to X$ is the canonical projection. We have G-homeomorphisms h_l $(l = 1, \cdots, 3)$ such that the following diagram is commutative.

$$\begin{array}{ccccc} X \coprod Z_1 & \xrightarrow{\bar{i}} & Z_2 & \xleftarrow{\bar{h}} & Z_3 \\ \downarrow & & \downarrow & & \downarrow \\ (G \times_K X) \coprod (G \times_K Z_1) & \hookrightarrow & G \times_K Z_2 & \leftarrow & G \times_K Z_3 \\ \| h_1 & & \| h_2 & & \| h_3 \\ (G \times_K X) \coprod Y_1 & \xrightarrow{i} & Y_2 & \xleftarrow{h} & Y_3 \end{array}$$

By taking the Z_l's, the G-maps f_l induce K-maps $\bar{f}_l: Z_l \to X$ $(l = 1, \cdots, 3)$. Since $\bar{h}$ is a K-homotopy equivalence and the diagram

$$\begin{array}{ccccc} X \coprod Z_1 & \overset{\bar{i}}{\hookrightarrow} & Z_2 & \overset{\bar{h}}{\leftarrow} & Z_3 \\ & {\scriptstyle id \coprod \bar{f}_1}\searrow & \downarrow{\scriptstyle \bar{f}_2} & \swarrow{\scriptstyle \bar{f}_3} & \\ & & X & & \end{array}$$

commutes, Z_3 dominates X. By Lemma 1.2, Z_3 has the K-homotopy type of a finite K-CW complex. This completes the proof.

Let $\Phi: D^G(G \times_K X) \to D^K(X)$ be a homomorphism induced by a mapping assigning $k: W \to X$, described as in Theorem 1.3, to any G-map $g: V \to G \times_K X$. It is an inverse isomorphism of a homomorphism $D^K(X) \to D^G(G \times_K X)$ induced by ind_K^G. Since $G \times_K W \underset{G}{\simeq} V$, it follows from Proposition 2.2 that $\Phi(Wa^G(G \times_K X)) \subset Wa^K(X)$. Then we have:

Theorem 2.3. *Let K be any closed subgroup of G and let X be a K-space. The induction homomorphism* $\mathrm{Ind}_K^G(X)$: $Wa^K(X)\to Wa^G(G\times_K X)$ *is an isomorphism. In particular $G\times_K X$ has the G-homotopy type of a finite G-CW complex if and only if X has the K-homotopy type of a finite K-CW complex.*

3. Free S^1-spaces

In this section we study Theorem A for free S^1-spaces. We denote a group S^1 by G and let X be a connected free G-CW complex such that $H_1(\nu;\boldsymbol{Z})$ is injective. If the projection $X\to X/G$ is a principal G-bundle and the fundamental group of X is abelian, Anderson [1] has shown that the universal cover of X is $\pi_1(X/G)$-homeomorphic to the product of the universal cover of X/G and the real space $\boldsymbol{R}$ with some $\pi_1(X/G)$-action on $\boldsymbol{R}$. We show that for some $K\leq G$ and some CW complex V, the G-space X/K is G-homotopy equivalent to $G/K\times_{(1K)}V=G/K\times V$.

Lemma 3.1. *Let X be a connected free G-space such that the G-map ν: $G\to X$ induces a monomorphism $H_1(\nu;\boldsymbol{Z})$. There is a finite subgroup K of G such that $\pi_1(X/K)$ isomorphic to $\pi_1(G/K)\oplus\pi_1(X/G)$.*

Proof. For any $K\leq G$, we have a short exact sequence:

$$1\to\pi_1(G/K)\to\pi_1(X/K)\to\pi_1(X/G)\to 1$$

We construct a splitting $\pi_1(X/K)\to\pi_1(G/K)$ for some $K<G$. By the assumption, there is an epimorphism μ: $\pi_1(X)\to\boldsymbol{Z}$ such that the following diagram commutes:

$$\begin{array}{ccc}\pi_1(G) & \overset{\nu}{\rightarrowtail} & \pi_1(X)\\ & {\scriptstyle n}\searrow & \downarrow{\scriptstyle \mu}\\ & & \boldsymbol{Z}\end{array}$$

Here n is multiplication by $n\geq 0$. Let K be a subgroup of G with order n.

$$\begin{array}{ccccc}\pi_1(G) & \xrightarrow{\nu} & \pi_1(X) & \xrightarrow{\mu} & \boldsymbol{Z}\\ \downarrow{\scriptstyle n} & & \downarrow{\scriptstyle p} & \nearrow{\scriptstyle \overline{\mu}} & \\ \pi_1(G/K) & \xrightarrow{\bar{\nu}} & \pi_1(X/K) & & \\ \downarrow{\scriptstyle \partial} & & \downarrow{\scriptstyle \partial} & & \\ K & = & K & & \end{array}$$

By a chasing method, the equation $\bar{\nu}(m)p(y)=\bar{\nu}(m')p(y')$ implies that there is

$z \in \pi_1(G)$ satisfying $m=m'+nz$ and $y=\nu(z^{-1})y'$. Then we have

$$m+\mu(y) = m'+nz+\mu(\nu(z^{-1}))+\mu(y') = m'+\mu(y') .$$

For any $x=\bar{\nu}(m)p(y) \in \pi_1(X/K)$ we define as $\bar{\mu}(x)=m+\mu(y)$. Then the map $\bar{\mu}: \pi_1(X/K) \to \boldsymbol{Z}$ is a homomorphism with $\bar{\mu}\circ p=\mu$, since the image of $\bar{\nu}$ is a subgroup of the center of $\pi_1(X/K)$. Since both $\mu\circ\nu$ and n are multiplication by n, we have $\bar{\mu}\circ\bar{\nu}=1$ and $\bar{\mu}$ is the required splitting.

Propoistion 3.2. *Let X be as in Theorem A. If X is free, then there are a proper subgroup K of G and a CW complex V such that $G/K\times V$ and X/K are G-homotopy equivalent.*

Proof. Let K be a subgroup of G such that $\pi_1(X/K) \cong \pi_1(G/K) \oplus \pi_1(X/G)$. We denote by $p: V \to X/K$ the covering space corresponding to $\pi_1(X/G) \leq \pi_1(X/K)$. The G-map $G/K\times V \to X/K$, sending (gK, v) to $g\cdot p(v)$, induces an isomorphism of homotopy groups. By a Whitehead theorem of the equivariant version [10], it is a G-homotopy equivalence.

By Lemma 1.2 there is a K-CW complex Y such that $G\times_K Y$ and X is G-homotopy equivalent.

REMARK. Let Y be a K-space obtained from the G-homotopy pullback of the G-map p through the covering map $V \to X/K$. Then the G-map $G\times_K Y \to X$ induced by the given K-map $Y \to X$ is a G-homotopy equivalence.

4. Proof of Theorems A and B

In this section, we also denote S^1 by G.

Proposition 4.1. *Let X be as in Thoerem A and let H be a finite subgroup of G. Then $H_1(p\circ\nu; \boldsymbol{Z})$ is monic for the projection $p: X \to X/H$.*

Proof. As the H-action on X comes from a G-action by restriction, H acts trivially on $H_1(X; \boldsymbol{Z})$. Applying Theorem 2.4 [3, p. 120], we obtain that the projection induces an isomorphicm $H_1(p; \boldsymbol{Q}): H_1(X; \boldsymbol{Q}) \to H_1(X/H; \boldsymbol{Q})$. Then $H_1(p; \boldsymbol{Z})$ is injective on any free abelian subgroup of rank one in $H_1(X; \boldsymbol{Z})$.

We note that $\pi_1(\nu)$ is monic does not imply that $\pi_1(p\circ\nu)$ is injective.

Proof of Theorem A. Let H be a cyclic subgroup of which order is a common multiple of order of all isotropy subgroups in X. Clearly X/H is a connected free G/H-CW complex. By the argument in the previous section, there are a free K/H-CW complex V and a G/H-homotopy equivalence $h: G/H\times_{K/H} V \to X/H$. We see canonically V as a K-CW complex. Then h induces a G-homotopy equivalence $h': G\times_K V \to X/H$. The G-space W obtained from a

G-homotopy pullback of h' through the projection $X \to X/H$ is G-homotopy equivalent to X.

$$\begin{array}{ccc} W & \longrightarrow & X \\ \downarrow & & \downarrow \\ G\times_K V & \xrightarrow[G]{\simeq} & X/H \end{array}$$

On the other hand, by Lemma 1.2, there is a K-CW complex Y such that $G\times_K Y$ and W are G-homotopy equivalent. This completes the proof.

To prove Theorem B we may show the following:

Proposition 4.3. *Let X be a connected finitely dominated G-space which fulfills that $H_1(v; \boldsymbol{Z})$ is injective. Then the K-space X has the K-homotopy type of a finite K-CW complex for any finite subgroup K of G.*

Proof. X has the G-homotopy type of a G-CW complex with finitely many orbit types [5, Theorem 1.4]. By Theorem A and Proposition 2.2, there is a finitely dominated L-space Y such that $X \underset{G}{\simeq} G\times_L Y$. Since $G\times_L Y \to G/L$ is a K-fibration, we have the result. (See Theorem 3.6 [13].)

Theorem 4.4. *Let G be any compact Lie group and let K be a subgroup of G. Let X be a finitely dominated G-space with X/G connected. If the rank of the image of $H_1(W_GK \to X/K; \boldsymbol{Z})$ is not zero, then* $\mathrm{Res}^G_K(X)\colon Wa^G(X) \to Wa^K(X)$ *is a zero map.*

Proof. Let T be a maximal torus of W_GK. Since $H_1(T \to W_GK; \boldsymbol{Z})$ is epic, there is a proper subgroup $\bar{C}$ of W_GK such that $\bar{C}$ is isomorphic to S^1 and $H_1(\bar{C} \to X/K; \boldsymbol{Z})$ is injective. Then there is a $\bar{C}$-map $f\colon X/K \to \bar{C}/\bar{L}$ for some finite subgroup $\bar{L}$ of $\bar{C}$. Let C (resp. L) be the preimage of $\bar{C}$ (resp. $\bar{L}$) under the projection $N_GK \to W_GK$. Clearly $h\colon \bar{C}/\bar{L} \cong C/L$. Since K is a normal subgroup of C, the projection $p\colon X \to X/K$ is a C-map. Then $h\circ f\circ p\colon X \to C/L$ is an equivariant C-fibration. The K-space C/L has a trivial K-action and its Euler characteristic is zero. If we apply Theorem 2.6 [13] to the equivariant K-fibration $h\circ f\circ p$, we conclude the proof.

References

[1] D.R. Anderson: *The Wall invariant of certain S^1-bundles*, Proc. Amer. Math. Soc. **31** (1972), 529–535.

[2] G.E. Bredon: Equivariant cohomology theories, Lecture Notes in Math. 34, Springer-Verlag, Berlin and New York, 1967,

[3] G.E. Bredon: Introduction to compact transformation groups, Acadeimc Press,

New York, 1972.

[4] K. Ehrlich: *Fibrations and a transfer map in algebraic K-theory,* J. Pure Appl. Algebra **14**(1979), 131–136.

[5] K. Iizuka: *Finiteness condition for G-CW complexes,* Jap. J. Math. **10**(1984), 55–69.

[6] W. Lück: *The transfer maps induced in the algebraic K_0-and K_1-groups by a fibration I,* Math. Scand. **59**(1986), 93–121.

[7] W. Lück: *The transfer maps induced in the algebraic K_0-and K_1-groups by a fibration II,* J. Pure Appl. Algebra **45**(1987), 143–169.

[8] W. Lück: *The geometric finiteness obstruction,* Proc. London Math. Soc. **54**(1987), 367–384.

[9] W. Lück: Transformation groups and algebraic K-theory, Lecture Notes in Math. 1408, Springer-Verlag, 1989.

[10] T. Matumoto: *On G-CW complexes and a theorem of J.H.C. Whitehead,* J. Fac. Univ. Tokyo Sect. I A Math. **18**(1971), 363–374.

[11] H.J. Munkholm and E.K. Pedersen: *On the Wall finiteness obtruction for the total space of certain fibrations.* Trans. Amer. Math. Soc. **261**(1980), 529–545.

[12] R.L. Rubinsztein: *On the equivariant parallelizable manifolds,* Math. Scand. **62**(1988), 217–245.

[13] T. Sumi: *Finiteness obstructions of equivariant fibrations,* Publ. Res. Inst. Math. Sci, **27** (1991), 627–637

[14] C.T.C. Wall: *Finiteness conditions for CW complexes,* Ann. of Math. **81**(1965), 56–69.

[15] S. Waner: *Equivariant fibrations and transfer,* Trans. Amer. Math. Soc. **258** (1980), 369–384.

Department of Mathematics
Kyushu University 33
Fukuoka 812, Japan
E-mail: d75190a@kyu-cc.cc.kyushu-u.ac.jp,
sumi@math sci. kyushu-u.ac.jp