



Title	Crossing change and exceptional Dehn surgery
Author(s)	Miyazaki, Katura; Motegi, Kimihiko
Citation	Osaka Journal of Mathematics. 2002, 39(4), p. 773-777
Version Type	VoR
URL	https://doi.org/10.18910/10957
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Miyazaki, K. and Motegi, K.
Osaka J. Math.
39 (2002), 773–777

CROSSING CHANGE AND EXCEPTIONAL DEHN SURGERY

KATURA MIYAZAKI and KIMIHIKO MOTEKI*

(Received November 27, 2000)

1. Introduction

Thurston's hyperbolic Dehn surgery theorem [11], [12] asserts that if a knot K in the 3-sphere S^3 is hyperbolic (i.e., $S^3 - K$ admits a complete hyperbolic structure of finite volume), then all but finitely many Dehn surgeries on K yield hyperbolic 3-manifolds. By an *exceptional surgery* on a hyperbolic knot we mean a nontrivial Dehn surgery producing a non-hyperbolic manifold. Refer to [3], [6] for a survey on Dehn surgery on knots. We empirically know that 'most' knots are hyperbolic and 'most' hyperbolic knots have no exceptional surgeries. In this paper, we demonstrate the abundance of hyperbolic knots with no exceptional surgeries by showing that every knot is 'close' to infinitely many such hyperbolic knots in terms of crossing change.

We regard that two knots are the same if they are isotopic in S^3 . For a knot K in S^3 , let $B_n(K)$ be the set of knots each of which is obtained by changing at most n crossings in a diagram of K .

Theorem 1.1. *For every knot K in S^3 , $B_1(K)$ contains infinitely many hyperbolic knots each of which has no exceptional surgeries. In particular, an arbitrary knot can be deformed into a hyperbolic knot with no exceptional surgeries by a single crossing change.*

In Section 3, we raise some questions on the distribution of hyperbolic knots (with no exceptional surgeries).

2. Proofs

A (2-string) *tangle* is a pair (B, t) where B is a 3-ball and t is a pair of disjoint arcs properly embedded in B . We call (B, t) a *trivial tangle* if there is a homeomorphism from (B, t) to $(D \times I, \{x, y\} \times I)$, where D is a disk containing x and y in its interior. A tangle (B, t) is said to be *atoroidal* if $B - \text{int } N(t)$ contains no incompressible tori. A tangle (B, t) is said to be ∂ -*irreducible* if $B - \text{int } N(t)$ is ∂ -irreducible.

2000 Mathematics Subject Classification : Primary 57M25.

*Supported in part by Grant-in-Aid for Encouragement of Young Scientists (No. 11740051), The Ministry of Education, Science, Sports and Culture.




Fig. 2.




Fig. 3.

all nontrivial Dehn surgeries on K_n produce hyperbolic, Haken manifolds. Since every non-hyperbolic knot yields non-hyperbolic manifolds after infinitely many Dehn surgeries, K_n is a hyperbolic knot.

It remains to show that $\{K_n\}_{n \in \mathbb{Z}}$ contains infinitely many distinct knots. Note that K_n is obtained from K_0 by twisting n times along the disk D in Fig. 3.

CLAIM 2.2. The circle ∂D does not bound a disk D' in S^3 which intersects K_0 in at most one point.

Proof. Since the algebraic intersection number of K_0 and D is zero, it is sufficient to show that there is no disk D' satisfying $\partial D' = \partial D$ and $D' \cap K_0 = \emptyset$. Suppose for a contradiction that we had such a disk D' . Let A be an obvious annulus in $B_1 - B_1 \cap K_0$ connecting ∂D and an essential simple loop in $\partial B_1 - \text{int } N(B_1 \cap K_0)$. Then the existence of a (possibly singular) disk $A \cup D'$ would imply the compressibility of $F = \partial B_1 - \text{int } N(B_1 \cap K_0)$ in $E(K_0)$. This is a contradiction. \square

By applying [5, Theorem 3.2] we see that $\{K_n\}_{n \in \mathbb{Z}}$ consists of infinitely many distinct knots. This completes the proof of Theorem 1.1. \square

REMARKS. (1) In Theorem 1.1 we can choose a crossing change of K so that the resulting hyperbolic knot is concordant to K and has the same Alexander invariant as K . In fact, by [9, Lemma 3.3] the crossing change given in the proof of Theorem 1.1 does not change knot concordance class and Alexander invariant. (2) Kawauchi [4] proved that $B_1(K)$ contains infinitely many hyperbolic knots as an application of the imitation theory. In [4, Theorem 3.1], let L'' be the knot obtained from L by nugatory crossing change (so that $L \cong L''$), then Theorem 3.1(1), (2) implies that an imitation L^* of L is a hyperbolic knot and obtained from L by a single crossing change.

3. Questions

On the distribution of hyperbolic knots in the knot table, Adams [1] conjectures that the proportion of hyperbolic knots among all prime knots with minimal crossing number less than n approaches 1 as $n \rightarrow \infty$. For any knot projection G of n crossings, we can obtain 2^n knot diagrams by indicating over-under relation at each crossing point in G . Weeks [13] asks if “under reasonable conditions” most knot diagrams obtained from a given knot projection represent hyperbolic knots; we paraphrase this question as follows.

Question 1. Let $f(G)$ be the proportion of diagrams representing hyperbolic knots among all the knot diagrams obtained from a knot projection G . Let \mathcal{D}_n be the set of knot projections G in S^2 of n crossings such that if C is a simple closed curve in S^2 meeting G transversely in two non-crossing points, then a component of $G - C$ is an embedded arc in S^2 . We denote by $g(r, n)$ the proportion of knot projections $G \in \mathcal{D}_n$ satisfying $f(G) > r$. Then for each r with $0 < r < 1$, does $g(r, n)$ approach 1 as $n \rightarrow \infty$?

On the distribution of hyperbolic knots with no exceptional surgeries we raise the following questions.

Question 2. (1) Is there a hyperbolic knot K such that each hyperbolic knot in $B_1(K)$ has no exceptional surgeries?

(2) If K is a trivial knot, then $B_1(K)$ contains hyperbolic, twist knots, each admitting exceptional surgeries [2]. Is there a nontrivial knot K such that $B_1(K)$ contains infinitely many hyperbolic knots with exceptional surgeries?

ACKNOWLEDGEMENTS. We would like to thank the referee for some very helpful comments.

References

- [1] C.C. Adams: *The knot book*, W.H. Freeman and Company, 1994.
- [2] M. Brittenham and Y.Q. Wu: *The classification of Dehn surgeries on 2-bridge knots*, Comm. Anal. Geom. **9** (2001), 97–113.
- [3] C.McA. Gordon and Dehn Filling: a survey, Knot theory (Warsaw, 1995), 129–144, Banach Center Publ., 42, Polish Acad. Sci., Warsaw, 1998.
- [4] A. Kawauchi: Almost identical link imitations and the skein polynomial, 465–476, Knots 90, Walter de Gruyter, 1992.
- [5] M. Kouno, K. Motegi and T. Shibuya: *Twisting and knot types*, J. Math. Soc. Japan, **44** (1992), 199–216.
- [6] J. Luecke: *Dehn surgery on knots in the 3-sphere*, Proc. International Congress of Mathematicians, Zürich, Switzerland 1994, Birkhäuser Verlag (1995), 585–594.
- [7] J. Morgan and H. Bass (eds.): *The Smith conjecture*, Academic Press, 1984.
- [8] R. Myers: *Excellent 1-manifolds in compact 3-manifolds*, Topology Appl. **49** (1993), 115–127.
- [9] Y. Nakanishi: *Primeness of links*, Math. Sem. Notes, Kobe Univ. **9** (1981), 415–440.
- [10] T. Soma: *Simple links and tangles*, Tokyo J. Math. **6** (1983), 65–73.
- [11] W.P. Thurston: The geometry and topology of 3-manifolds, Lecture notes, Princeton University, 1979.
- [12] W.P. Thurston: *Three dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. **6** (1982), 357–381.
- [13] J. Weeks: private correspondence.
- [14] Y-Q. Wu: *Dehn surgery on arborescent knots*, J. Diff. Geom. **43** (1996), 171–197.

Katura Miyazaki
 Faculty of Engineering
 Tokyo Denki University
 2-2 Kanda-Nishikicho
 Tokyo 101-8457, Japan
 e-mail: miyazaki@cck.dendai.ac.jp

Kimihiko Motegi
 Department of Mathematics
 Nihon University
 3-25-40 Sakurajosui, Setagaya-ku
 Tokyo 156-8550, Japan
 e-mail: motegi@math.chs.nihon-u.ac.jp