

Title	A note on the relation of $Z_2$ -graded complex cobordism to complex K-theory
Author(s)	Yosimura, Zen-ichi
Citation	Osaka Journal of Mathematics. 1975, 12(3), p. 583-595
Version Type	VoR
URL	<a href="https://doi.org/10.18910/11134">https://doi.org/10.18910/11134</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## A NOTE ON THE RELATION OF $Z_2$ -GRADED COMPLEX COBORDISM TO COMPLEX K-THEORY

ZEN-ICHI YOSIMURA

(Received December 16, 1974)

Let  $MU^*( )$  and  $K^*( )$  denote the  $Z_2$ -graded complex cobordism theory and the complex  $K$ -theory respectively. The Thom homomorphism  $\mu_*: \pi_0(MU) \rightarrow \pi_0(K)$  on coefficient groups is identified (up to sign) with the classical Todd genus  $Td: \Lambda \rightarrow Z$ . We denote by  $I$  the ideal of  $\Lambda$  to be the kernel of  $Td: \Lambda \rightarrow Z$ . Wolff [7] proved that the decreasing filtration  $\{I^q MU^*( )\}$  of  $MU^*( )$  consists of cohomology theories defined on the category of based finite  $CW$ -complexes, and the associated quotients  $I^q MU^*( )/I^{q+1} MU^*( )$  are determined by the complex  $K$ -theories  $KG_q^*( )$  with coefficients  $G_q = I^q/I^{q+1}$ .

The purpose of this note is to extend the Wolff's result to the category of based  $CW$ -complexes. Let  $F_q MU$  be the  $CW$ -spectrum associated with the cohomology theory  $I^q MU^*( )$ , i.e.,  $\{Y, F_q MU\}^* \cong I^q MU^*(Y)$  for any based finite  $CW$ -complex (or finite  $CW$ -spectrum). We show that  $\{F_q MU^*( )\}$  is a decreasing filtration of  $MU^*( )$  consisting of  $\Lambda$ -modules so that the associated quotients are equal to  $KG_q^*( )$ , and in addition that  $F_{q+1} MU^*( )$  is a direct summand of  $F_q MU^*( )$ .

Moreover we give a tower

$$MU \rightarrow \dots \rightarrow Q_q MU \rightarrow Q_{q-1} MU \rightarrow \dots \rightarrow Q_0 MU = K$$

of  $MU$ -module spectra such that  $KG_q \rightarrow Q_q MU \rightarrow Q_{q-1} MU$  is a cofiber sequence of  $MU$ -module spectra, which factorizes the Thom map  $\mu_*: MU \rightarrow K$ .

Baas [3] constructed a tower of  $CW$ -spectra

$$MU \rightarrow \dots \rightarrow MU\langle n \rangle \rightarrow MU\langle n-1 \rangle \rightarrow \dots \rightarrow MU\langle 0 \rangle = H$$

factorizing the Thom map  $\mu: MU \rightarrow H$ . In appendix we show that the tower is of  $MU$ -module spectra and the sequence  $\Sigma^{2n} MU\langle n \rangle \xrightarrow{m_{x_n}} MU\langle n \rangle \rightarrow MU\langle n-1 \rangle$  is a cofiber sequence where  $m_{x_n}$  is the multiplication by  $x_n$  a ring generator of  $\Lambda$  with degree  $2n$ .

### 1. Decreasing filtration of $MU_*( )$

1.1. A pair  $(E, \rho)$  is called a  $Z_2$ -graded  $CW$ -spectrum if  $E$  is a  $CW$ -spectrum

and  $\rho: \Sigma^2 E \rightarrow E$  is a homotopy equivalence. Such a pair  $(E, \rho)$  gives rise to natural isomorphisms

$$\rho_*: E_*(X) \rightarrow E_{*+2}(X), \quad \rho^*: E^{*+2}(X) \rightarrow E^*(X)$$

for any  $CW$ -spectrum  $X$ . So we can define  $Z_2$ -graded homology and cohomology theories  $E_*( )$ ,  $E^*( )$  by putting

$$E_*(X) = E_0(X) \oplus E_1(X), \quad E^*(X) = E^0(X) \oplus E^1(X).$$

For a  $CW$ -spectrum  $E$  we put

$$E = \bigvee_n \Sigma^{2n} E, \quad \bar{E} = \prod_n \Sigma^{2n} E.$$

Taking the canonical identifications  $\rho: \Sigma^2 E \rightarrow E$  and  $\bar{\rho}: \Sigma^2 \bar{E} \rightarrow \bar{E}$  as structure morphisms  $E$  and  $\bar{E}$  admit structures of  $Z_2$ -graded  $CW$ -spectra respectively. From definition it follows that

$$E_0(X) \cong \sum_n E_{2n}(X), \quad E_1(X) \cong \sum_n E_{2n+1}(X),$$

$$\bar{E}^0(X) \cong \prod_n E^{2n}(X), \quad \bar{E}^1(X) \cong \prod_n E^{2n+1}(X)$$

for all  $CW$ -spectra  $X$ . In particular, the canonical morphism  $H \rightarrow \bar{H}$  becomes a homotopy equivalence for the Eilenberg-MacLane spectrum  $H$ .

The  $BU$ -spectrum  $K$  may be regarded as a  $Z_2$ -graded  $CW$ -spectrum because it possesses the Bott map  $\beta: \Sigma^2 K \rightarrow K$  which is a homotopy equivalence.

Denote by  $F_n$  the direct sum of  $n$ -copies of the integers  $Z$  and by  $F$  the direct limit of  $F_n$ , i.e.,  $F$  is a free abelian group with countably many factors. Putting

$$BU_{F_n} = BU \times \cdots \times BU, \text{ the product of } n\text{-copies of } BU,$$

$$BU_F = \bigcup_n BU_{F_n}, \text{ the union of } BU_{F_n},$$

we obtain

**Proposition 1.** *There exists a natural isomorphism*

$$[X, BU_F] \rightarrow KF^0(X)$$

for any based connected  $CW$ -complex  $X$ .

*Proof.* Let  $Y$  be a based connected finite  $CW$ -complex. Then we have a sequence of natural isomorphisms

$$[Y, BU_F] \leftarrow \varinjlim [Y, BU_{F_n}] \leftarrow \varinjlim [Y, BU] \otimes F_n \rightarrow \varinjlim K^0(Y) \otimes F_n$$

$$\rightarrow K^0(Y) \otimes F \rightarrow KF^0(Y).$$

Therefore the contravariant functor  $KF^0$  defined on the category of based connected  $CW$ -complexes is represented by  $BU_F$  (use [1, Addendum 1.5]).

Proposition 1 implies that  $BU_F$  is homotopy equivalent to  $\Omega_0^2 BU_F$  where  $\Omega_0^2$  means the component of the base point in the double loop space. Hence we have

(1.1) *in the  $BU$ -spectrum  $KF$  with the coefficients  $F$  every even term is the based  $CW$ -complex  $BU_F$ .*

1.2. Let us denote by  $MU$  the unitary Thom spectrum and by  $\mu_c: MU \rightarrow K$  the Thom map which is a ring morphism. The composition

$$\mu_c: MU \rightarrow K \rightarrow K$$

of  $\vee \Sigma^{2n} \mu_c$  and  $\vee \beta^n$  is a morphism of  $Z_2$ -graded ring-spectra, called the  $Z_2$ -graded Thom map. As is well known, it is characterized by the coefficient homomorphism  $\mu_{c\sharp}: \pi_{\sharp}(MU) \rightarrow \pi_{\sharp}(K)$  which coincides (up to sign) with the classical Todd genus  $Td$ . Putting  $\Lambda = \pi_{\sharp}(MU)$ ,  $\pi_{\sharp}(K) = Z$  is viewed as a  $Z_2$ -graded  $\Lambda$ -module via  $\mu_{c\sharp} = Td$  and it is written  $Z_{Td}$  for emphasis.

Using the kernel  $I$  of  $Td: \Lambda \rightarrow Z$  we define a decreasing filtration  $\{I^q\}_{q \geq 0}$  consisting of ideals of  $\Lambda$ . Denoting by  $G_q$  the associated  $Z_2$ -graded  $\Lambda$ -module  $I^q/I^{q+1}$ , we see easily [7, Satz 3.8] that

(1.2)  $G_0 \cong Z_{Td}$  and  $G_q$  is a free abelian group with countably many factors for  $q \geq 1$ .

For a  $Z_2$ -graded  $\Lambda$ -module  $A$  we have a decreasing filtration  $\{I^q A\}_{q \geq 0}$  consisting of submodules of  $A$ , whose associated  $Z_2$ -graded  $\Lambda$ -module  $I^q A/I^{q+1} A$  is written  $G_q(A)$ . Applying the commutative diagram

$$\begin{array}{ccccccc} 0 \rightarrow \text{Tor}_1^{\Lambda}(Z_{Td}, A) & \rightarrow & I \otimes A & \rightarrow & A & \rightarrow & Z_{Td} \otimes A \rightarrow 0 \\ & & \downarrow \cong & & \parallel & & \downarrow \cong \\ 0 \rightarrow IA & \rightarrow & A & \rightarrow & G_0(A) & \rightarrow & 0 \end{array}$$

with exact rows, we get an isomorphism

$$(1.3) \quad G_q \otimes_{\Lambda} A \xrightarrow{\cong} G_q \otimes_{\mathbb{Z}} (Z_{Td} \otimes_{\Lambda} A) \xrightarrow{\cong} G_q \otimes_{\mathbb{Z}} G_0(A)$$

by means of ‘‘4 lemma’’.

**Proposition 2.** *Let  $A$  be a  $\Lambda$ -module with  $\text{Tor}_k^{\Lambda}(Z_{Td}, A) = 0$  for all  $k \geq 1$ . Then, for every  $q \geq 0$  both  $I^q \otimes_{\Lambda} A \rightarrow I^q A$  and  $G_q \otimes_{\Lambda} A \rightarrow G_q(A)$  are isomorphisms and  $\text{Tor}_k^{\Lambda}(I^q, A) = \text{Tor}_k^{\Lambda}(G_q, A) = 0$  for all  $k \geq 1$ .*

Proof. Choose a free  $\Lambda$ -module  $F$  such that  $A$  is isomorphic to a quotient  $F/B$ . By induction on  $q$  we shall show that the sequences

$$0 \rightarrow I^q B \rightarrow I^q F \rightarrow I^q A \rightarrow 0, \quad 0 \rightarrow G_q(B) \rightarrow G_q(F) \rightarrow G_q(A) \rightarrow 0$$

are exact. The  $q=0$  case is evident because of (1.3). Applying induction

hypotesis and “3×3 lemma” we find easily that  $0 \rightarrow I^q B \rightarrow I^q F \rightarrow I^q A \rightarrow 0$  is exact. So we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & G_q \otimes_{\mathbb{Z}} G_0(B) & \rightarrow & G_q \otimes_{\mathbb{Z}} G_0(F) & \rightarrow & G_q \otimes_{\mathbb{Z}} G_0(A) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & G_0(I^q B) & \rightarrow & G_0(I^q F) & \rightarrow & G_0(I^q A) \rightarrow 0 \end{array}$$

with exact rows. Since all vertical arrows are epimorphisms and in particular the central one is an isomorphism, all vertical arrows become isomorphisms. Consequently we get that  $0 \rightarrow G_q(B) \rightarrow G_q(F) \rightarrow G_q(A) \rightarrow 0$  is exact.

Next, we consider the commutative diagrams

$$\begin{array}{ccccccc} 0 & \rightarrow & \text{Tor}_1^\Lambda(I^q, A) & \rightarrow & I^q \otimes B & \rightarrow & I^q \otimes F \rightarrow I^q \otimes A \rightarrow 0 \\ & & & & \downarrow & & \downarrow \\ 0 & \rightarrow & I^q B & \rightarrow & I^q F & \rightarrow & I^q A \rightarrow 0 \\ \\ 0 & \rightarrow & \text{Tor}_1^\Lambda(G_q, A) & \rightarrow & G_q \otimes B & \rightarrow & G_q \otimes F \rightarrow G_q \otimes A \rightarrow 0 \\ & & & & \downarrow & & \downarrow \\ 0 & \rightarrow & G_q \otimes_{\mathbb{Z}} G_0(B) & \rightarrow & G_q \otimes_{\mathbb{Z}} G_0(F) & \rightarrow & G_q \otimes_{\mathbb{Z}} G_0(A) \rightarrow 0 \end{array}$$

with exact rows. Remark that  $\text{Tor}_k^\Lambda(Z_{Td}, B) = 0$  for all  $k \geq 1$ . By use of “4 lemma” and (1.3) we see that all vertical arrows are isomorphisms, and hence we obtain the required results.

For a  $\Lambda$ -module  $A$  we put  $J_q(A) = A/I^{q+1}A$  and abbreviate  $J_q = J_q(\Lambda)$  when  $A = \Lambda$ . As an immediate corollary of Proposition 2 we have

**Corollary 3.** *Let  $A$  be a  $\Lambda$ -module with  $\text{Tor}_k^\Lambda(Z_{Td}, A) = 0$  for all  $k \geq 1$ . Then  $J_q \otimes A \rightarrow J_q(A)$  is an isomorphism and  $\text{Tor}_k^\Lambda(J_q, A) = 0$  for all  $k \geq 1$ .*

**1.3.** Let  $\mathcal{M}\mathcal{U}$  denote the category of comodules over  $MU_*(MU)$  which are finitely presented as  $\Lambda$ -modules. Notice that  $\mathcal{M}\mathcal{U}$  is an abelian category which has enough projectives, and also that  $MU_*(Y)$  lies in the category  $\mathcal{M}\mathcal{U}$  whenever  $Y$  is a finite  $CW$ -spectrum. Since the functor  $M \rightarrow Z_{Td} \otimes_{\Lambda} M$  is exact on  $\mathcal{M}\mathcal{U}$  [5, Example 3.3] it follows immediately that

$$(1.4) \quad \text{Tor}_k^\Lambda(Z_{Td}, M) = 0 \quad \text{for all } k \geq 1 \text{ if } M \text{ lies in } \mathcal{M}\mathcal{U}.$$

Proposition 1 and Corollary 2 combined with (1.4) say that

$$(1.5) \quad \text{the functors } M \rightarrow I^q M \cong I^q \otimes_{\Lambda} M, M \rightarrow G_q(M) \cong G_q \otimes_{\Lambda} M \text{ and } M \rightarrow J_q(M) \cong J_q \otimes_{\Lambda} M \text{ on } \mathcal{M}\mathcal{U} \text{ are exact.}$$

**Theorem 1** (Wolff [7]). *i) Both  $I^q MU_*( )$  and  $MU_*( )/I^{q+1}MU_*( )$  are*

homology theories defined on the category of  $CW$ -spectra, so that  $I^q \otimes_{\Lambda} \mathbf{MU}_*(X) \rightarrow I^q \mathbf{MU}_*(X)$  and  $\Lambda/I^{q+1} \otimes_{\Lambda} \mathbf{MU}_*(X) \rightarrow \mathbf{MU}_*(X)/I^{q+1} \mathbf{MU}_*(X)$  are natural isomorphisms for all  $CW$ -spectra  $X$ .

ii)  $I^q \mathbf{MU}_*( )/I^{q+1} \mathbf{MU}_*( )$  is a homology theory defined on the category of  $CW$ -spectra such that there exists a natural isomorphism  $I^q \mathbf{MU}_*(X)/I^{q+1} \mathbf{MU}_*(X) \rightarrow KG_{q^*}(X)$  for any  $CW$ -spectrum  $X$  which is induced by the  $Z_2$ -graded Thom map  $\mu_c$ .

Proof. i) and the first half of ii) are immediate from (1.5). The latter half of ii) is also valid because we have a natural isomorphism

$$G_q(\mathbf{MU}_*(X)) \xleftarrow{\cong} G_q \otimes_{\Lambda} \mathbf{MU}_*(X) \xrightarrow{\cong} G_q \otimes_{\mathbb{Z}} (Z_{Td} \otimes_{\Lambda} \mathbf{MU}_*(X)) \xrightarrow{\cong} G_q \otimes_{\mathbb{Z}} K_*(X) \xrightarrow{\cong} KG_{q^*}(X).$$

Let  $\phi: E_*(X) \rightarrow F_*(X)$  be a natural transformation for any  $CW$ -spectrum  $X$ . According to [1, Addendum 1.5] there exists a morphism  $f: E \rightarrow F$  inducing  $\phi$ , and it is unique up to weak homotopy. The proof in [1] is actually given for the category of based connected  $CW$ -complexes, but it is easily extended to that of  $CW$ -spectra. Such a morphism  $f$  is uniquely chosen (up to homotopy) under the assumption that  $F^0(E)$  is Hausdorff.

Let  $E_*( )$  be a  $Z_2$ -graded homology theory defined on the category of  $CW$ -spectra, i.e., a homology theory equipped with a natural isomorphism  $E_*(X) \rightarrow E_{*+2}(X)$  for any  $CW$ -spectrum  $X$ . Then it gives  $E$  a structure of  $Z_2$ -graded  $CW$ -spectrum. In particular, the induced structure is unique if  $E^0(E)$  is Hausdorff.

Recall that the  $Z_2$ -graded  $CW$ -spectrum  $\mathbf{MU}$  is equipped with the canonical identification  $\rho: \Sigma^2 \mathbf{MU} \rightarrow \mathbf{MU}$  as structure morphism. Since  $\mathbf{MU}^*(\mathbf{MU})$  is Hausdorff (use Proposition 6 below), the  $Z_2$ -graded  $CW$ -spectrum  $(\mathbf{MU}, \rho)$  is characterized only by the  $Z_2$ -graded homology theory  $\mathbf{MU}_*( )$ .

Denote by  $F_q \mathbf{MU}$  and  $Q_q \mathbf{MU}$  the representing spectra of the new homology theories  $I^q \mathbf{MU}_*( )$  and  $\mathbf{MU}_*( )/I^{q+1} \mathbf{MU}_*( )$  respectively, i.e.,

$$I^q \mathbf{MU}_*(X) \cong \{ \Sigma^*, F_q \mathbf{MU} \wedge X \}, \quad \mathbf{MU}_*(X)/I^{q+1} \mathbf{MU}_*(X) \cong \{ \Sigma^*, Q_q \mathbf{MU} \wedge X \}$$

for any  $CW$ -spectrum  $X$ . Of course, they are both  $Z_2$ -graded  $CW$ -spectra. Then there exist morphisms

$$i_q: F_{q+1} \mathbf{MU} \rightarrow F_q \mathbf{MU}, \quad j_q: Q_q \mathbf{MU} \rightarrow Q_{q-1} \mathbf{MU}, \\ \iota_q: F_{q+1} \mathbf{MU} \rightarrow \mathbf{MU}, \quad \pi_q: \mathbf{MU} \rightarrow Q_q \mathbf{MU}$$

which induce the canonical morphisms in homology groups, and moreover we have morphisms

$$\mu_q: F_q\mathbf{MU} \rightarrow KG_q, \quad \nu_q: KG_q \rightarrow Q_q\mathbf{MU}$$

such that  $\mu_{q*}: F_q\mathbf{MU}_*(X) \rightarrow KG_{q*}(X)$  and  $\nu_{q*}: KG_{q*}(X) \rightarrow Q_q\mathbf{MU}_*(X)$  in homology groups are natural homomorphisms induced by the  $Z_2$ -graded Thom map  $\mu_c$ .

**Lemma 4.** *Let  $E \xrightarrow{f} F \xrightarrow{g} G$  be a sequence which satisfies the property that  $0 \rightarrow E_*(X) \rightarrow F_*(X) \rightarrow G_*(X) \rightarrow 0$  is a short exact sequence for every CW-spectrum  $X$ . Then it is a cofiber sequence.*

*Proof.* Let  $C_f$  be the mapping cone of  $f$ , i.e.,  $E \rightarrow F \rightarrow C_f$  a cofiber sequence. Then for any CW-spectrum  $X$  we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & E_*(X) & \rightarrow & F_*(X) & \rightarrow & C_{f*}(X) \rightarrow 0 \\ & & \parallel & & \parallel & & \downarrow \phi = h_* \\ 0 & \rightarrow & E_*(X) & \rightarrow & F_*(X) & \rightarrow & G_*(X) \rightarrow 0 \end{array}$$

with exact rows. Clearly  $h: C_f \rightarrow G$  which induces  $\phi$  is a homotopy equivalence.

By virtue of Lemma 4 we verify that

$$(1.6) \quad F_{q+1}\mathbf{MU} \rightarrow \mathbf{MU} \rightarrow Q_q\mathbf{MU}, \quad F_{q+1}\mathbf{MU} \rightarrow F_q\mathbf{MU} \rightarrow KG_q \text{ and } KG_q \rightarrow Q_q\mathbf{MU} \rightarrow Q_{q-1}\mathbf{MU} \text{ are all cofiber sequences.}$$

## 2. $Z_2$ -graded $\mathbf{MU}$ -module spectra

**2.1.** The inclusion  $Z \subset Q$  induces a natural transformation  $ch: E^*(X) \rightarrow EQ^*(X)$  for any CW-spectrum  $X$ , called the Chern-Dold character.

**Proposition 5.** *If  $ch: E^*(X) \rightarrow EQ^*(X)$  is a monomorphism, then  $E^*(X)$  is Hausdorff.*

*Proof.* Since  $EQ^*(X)$  is always Hausdorff [8, Proposition 4], the result is immediate.

Let  $W$  be a connective CW-spectrum with  $H_*(W)$  free and assume that  $\pi_*(E)$  is torsion free. Then  $H^*(W; \pi_*(E)) \rightarrow H^*(W; \pi_*(E) \otimes Q)$  is a monomorphism, and hence the Atiyah-Hirzebruch spectral sequences for  $E^*(W)$  and  $EQ^*(W)$  collapse. Therefore we get that

$$(2.1) \quad ch: E^*(W) \rightarrow EQ^*(W) \text{ is a monomorphism.} \quad (\text{Cf., [8, Lemma 11]}).$$

Applying Proposition 5 we obtain

**Proposition 6.** *Let  $W$  be a connective CW-spectrum with  $H_*(W)$  free. If  $\pi_*(E)$  is torsion free, then  $E^*(W)$  is Hausdorff.*

By means of Proposition 5 we get the following lemmas.

**Lemma 7.** *Assume that  $\pi_0(E)$  is torsion free and  $\pi_1(E)=0$ . Then  $E^0(KG_q \wedge MU \wedge \cdots \wedge MU)$  and  $E^0(Q_q MU \wedge MU \wedge \cdots \wedge MU)$  are Hausdorff and  $E^1(F_q MU \wedge MU \wedge \cdots \wedge MU)=0$ .*

Proof. Since  $E^{2n-1}(BU_{G_q} \wedge MU \wedge \cdots \wedge MU) = EQ^{2n-1}(BU_{G_q} \wedge MU \wedge \cdots \wedge MU) = 0$  we have a commutative square

$$\begin{array}{ccc} E^0(KG_q \wedge MU \wedge \cdots \wedge MU) & \rightarrow & \lim_{\leftarrow} E^{2n}(BU_{G_q} \wedge MU \wedge \cdots \wedge MU) \\ \downarrow & & \downarrow \\ EQ^0(KG_q \wedge MU \wedge \cdots \wedge MU) & \rightarrow & \lim_{\leftarrow} EQ^{2n}(BU_{G_q} \wedge MU \wedge \cdots \wedge MU) \end{array}$$

such that the horizontal arrows are isomorphisms. The left arrow is a monomorphism because so is the right one by use of (2.1). Since Theorem 1 implies that  $0 \rightarrow EQ^*(Q_{q-1} MU \wedge MU \wedge \cdots \wedge MU) \rightarrow EQ^*(Q_q MU \wedge MU \wedge \cdots \wedge MU) \rightarrow EQ^*(KG_q \wedge MU \wedge \cdots \wedge MU) \rightarrow 0$  is exact, an induction on  $q$  involving "4 lemma" shows that  $ch: E^0(Q_q MU \wedge MU \wedge \cdots \wedge MU) \rightarrow EQ^0(Q_q MU \wedge MU \wedge \cdots \wedge MU)$  is a monomorphism. Then we find that  $E^0(Q_q MU \wedge MU \wedge \cdots \wedge MU) \rightarrow E^0(MU \wedge \cdots \wedge MU)$  is a monomorphism because so is  $EQ^0(Q_q MU \wedge MU \wedge \cdots \wedge MU) \rightarrow EQ^0(MU \wedge \cdots \wedge MU)$ . Therefore  $E^1(MU \wedge \cdots \wedge MU) \rightarrow E^1(F_{q+1} MU \wedge MU \wedge \cdots \wedge MU)$  is an epimorphism, and hence  $E^1(F_{q+1} MU \wedge MU \wedge \cdots \wedge MU) = 0$ .

**Lemma 8.**  *$KG_p^0(F_q MU \wedge MU \wedge \cdots \wedge MU)$  and  $Q_p MU^0(F_q MU \wedge MU \wedge \cdots \wedge MU)$  are Hausdorff and  $KG_p^1(F_q MU \wedge MU \wedge \cdots \wedge MU) = Q_p MU^1(F_q MU \wedge MU \wedge \cdots \wedge MU) = 0$ .*

Proof. Putting  $X = F_q MU \wedge MU \wedge \cdots \wedge MU$ , we note by Theorem 1 i) that  $K_0(X)$  is free and  $K_1(X) = 0$ . Applying the universal coefficient sequence

$$0 \rightarrow \text{Ext}(K_{\#-1}(X), G_p) \rightarrow KG_p^{\#}(X) \rightarrow \text{Hom}(K_{\#}(X), G_p) \rightarrow 0$$

for  $K$  (see [9, (3.1)]) we get immediately that  $ch: KG_p^0(X) \rightarrow KG_p \otimes Q^0(X)$  is a monomorphism and  $KG_p^1(X) = 0$ . By induction on  $p$  we obtain that  $ch: Q_p MU^0(X) \rightarrow Q_p MU Q^0(X)$  is a monomorphism and  $Q_p MU^1(X) = 0$  because  $0 \rightarrow KG_p \otimes Q^*(X) \rightarrow Q_p MU Q^*(X) \rightarrow Q_{p-1} MU Q^*(X) \rightarrow 0$  is exact.

Assume that  $\pi_*(E)$  is free and of finite type and put again  $X = F_q MU \wedge MU \wedge \cdots \wedge MU$ . Using the universal coefficient sequence for  $E$  [9, (1.8)] we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & \text{Ext}(\hat{E}_{*-1}(X), Z) & \rightarrow & E^*(X) & \rightarrow & \text{Hom}(\hat{E}_*(X), Z) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & \text{Ext}(\hat{E}_{*-1}(X), Q) & \rightarrow & EQ^*(X) & \rightarrow & \text{Hom}(\hat{E}_*(X), Q) \rightarrow 0 \end{array}$$

with exact rows where  $\hat{E}$  is the dual of  $E$  constructed in [9]. Note that  $\pi_*(\hat{E})$  is free and hence so is  $\hat{E}_*(X)$ . Then the central arrow becomes a monomorphism. Considering the commutative square



$$\begin{array}{ccc} \bar{E}^*(X) & \rightarrow & \prod E^n(X) \\ \downarrow & & \downarrow \\ \bar{E}Q^*(X) & \rightarrow & \prod EQ^n(X) \end{array}$$

in which the upper arrow is an isomorphism, we find that the left one is a monomorphism. Thus we get that

(2.2)  $\bar{E}^*(F_q\mathbf{MU} \wedge \mathbf{MU} \wedge \cdots \wedge \mathbf{MU})$  is Hausdorff.

Let  $\widetilde{\mathbf{MU}}$  denote the mapping cone of the canonical morphism  $\mathbf{MU} \rightarrow \overline{\mathbf{MU}}$ . Since  $\prod Z/\sum Z \rightarrow \prod Q/\sum Q$  is a monomorphism we remark that

(2.3)  $\pi_0(\widetilde{\mathbf{MU}}) = \prod_n \pi_{2n}(\mathbf{MU})/\sum_n \pi_{2n}(\mathbf{MU})$  is torsion free and  $\pi_1(\widetilde{\mathbf{MU}}) = 0$ ,

(see [4, Exercise IV 20]). Then  $\widetilde{\mathbf{MU}}$  has a unique structure of  $Z_2$ -graded CW-spectrum so that the cofiber sequence  $\mathbf{MU} \rightarrow \overline{\mathbf{MU}} \rightarrow \widetilde{\mathbf{MU}}$  is of  $Z_2$ -graded CW-spectra.

**Lemma 9.**  $F_p\mathbf{MU}^0(F_q\mathbf{MU} \wedge \mathbf{MU} \wedge \cdots \wedge \mathbf{MU})$  is Hausdorff and  $F_p\mathbf{MU}^1(F_q\mathbf{MU} \wedge \mathbf{MU} \wedge \cdots \wedge \mathbf{MU})=0$

Proof. We put  $X=F_q\mathbf{MU} \wedge \mathbf{MU} \wedge \cdots \wedge \mathbf{MU}$ . From Lemma 7 it follows that  $F_p\mathbf{MU}^1(X)=0$ . In the sequence

$$F_p\mathbf{MU}^0(X) \rightarrow \mathbf{MU}^0(X) \rightarrow \overline{\mathbf{MU}}^0(X)$$

the former arrow is a monomorphism because of Lemma 8 and the latter one is so by means of (2.3) and Lemma 7. Thus the above composition is a monomorphism. On the other hand, (2.2) says that  $\overline{\mathbf{MU}}^0(X)$  is Hausdorff. So we get the remaining result.

**2.2.** Since  $F_q\mathbf{MU}^0(F_q\mathbf{MU})$  and  $Q_q\mathbf{MU}^0(Q_q\mathbf{MU})$  are both Hausdorff, we verify that


(2.4) the  $Z_2$ -graded homology theories  $I^q\mathbf{MU}_*( )$  and  $\mathbf{MU}_*( )/I^{q+1}\mathbf{MU}_*( )$  give  $F_q\mathbf{MU}$  and  $Q_q\mathbf{MU}$  unique structures of  $Z_2$ -graded CW-spectra respectively.

Moreover, by virtue of Lemmas 7, 8 and 9 we see that

(2.5)  $i_q: F_{q+1}\mathbf{MU} \rightarrow F_q\mathbf{MU}, \quad \iota_q: F_{q+1}\mathbf{MU} \rightarrow \mathbf{MU}, \quad \mu_q: F_q\mathbf{MU} \rightarrow KG_q$   
 $j_q: Q_q\mathbf{MU} \rightarrow Q_{q-1}\mathbf{MU}, \quad \pi_q: \mathbf{MU} \rightarrow Q_q\mathbf{MU}, \quad \nu_q: KG_q \rightarrow Q_q\mathbf{MU}$

are uniquely determined (up to homotopy), which induce the canonical morphisms in homology groups. In particular, the composition  $\iota_{q-1} \cdot i_q$  is homotopic to  $\iota_q$  and  $j_q \cdot \pi_q$  is so to  $\pi_{q-1}$ .

Consider the diagram



consisting of cofiber sequences. With an application of Verdier's lemma (see [2, Lemma 6.8]) we get a cofiber sequence  $KG_q \rightarrow Q_qMU \rightarrow Q_{q-1}MU \rightarrow \Sigma KG_q$  (denoted by dotted arrows in the above diagram) which makes the diagram homotopy commutative. Clearly this yields the canonical exact sequence  $0 \rightarrow KG_{q*}(X) \rightarrow Q_qMU_*(X) \rightarrow Q_{q-1}MU_*(X) \rightarrow 0$ . By uniqueness of  $v_q, j_q$  the above cofiber sequence coincides with  $KG_q \xrightarrow{v_q} Q_qMU \xrightarrow{j_q} Q_{q-1}MU \rightarrow \Sigma KG_q$ .

The multiplication  $\phi: MU \wedge MU \rightarrow MU$  gives rise to natural  $Z_2$ -graded homomorphisms

$$\begin{aligned}
 m_q: F_qMU_*(X) \otimes MU_*(Y) &\rightarrow F_qMU_*(X \wedge Y) \\
 \bar{m}_q: Q_qMU_*(X) \otimes MU_*(Y) &\rightarrow Q_qMU_*(X \wedge Y)
 \end{aligned}$$

for all  $CW$ -spectra  $X$  and  $Y$ . By use of Lemmas 7 and 9 there exist unique pairings

$$\phi_q: F_qMU \wedge MU \rightarrow F_qMU, \quad \bar{\phi}_q: Q_qMU \wedge MU \rightarrow Q_qMU$$

which induce the above  $m_q$  and  $\bar{m}_q$  respectively. Then it follows that

(2.6) both  $F_qMU$  and  $Q_qMU$  are (associative)  $Z_2$ -graded  $MU$ -module spectra.

**Proposition 10.** Let  $M$  be a  $Z_2$ -graded ring spectrum,  $E, F$  and  $G$   $Z_2$ -graded  $M$ -module spectra and  $E \rightarrow F \rightarrow G$  a cofiber sequence. Assume that  $E^0(E), E^0(E \wedge M), G^0(F)$  and  $G^0(F \wedge M)$  are Hausdorff, or that  $F^0(E), F^0(E \wedge M), G^0(G)$  and  $G^0(G \wedge M)$  are Hausdorff. If for any  $CW$ -spectrum  $X$   $0 \rightarrow E_*(X) \rightarrow F_*(X) \rightarrow G_*(X) \rightarrow 0$  is a short exact sequence of  $Z_2$ -graded  $M_*(\ )$ -modules, then the cofiber sequence  $E \rightarrow F \rightarrow G$  is of  $Z_2$ -graded  $M$ -module spectra.

Proof. Assuming that  $F^0(E), F^0(E \wedge M), G^0(G)$  and  $G^0(G \wedge M)$  are Hausdorff, we consider the diagrams

$$\begin{array}{ccccccc}
 \Sigma^2 E & \rightarrow & \Sigma^2 F & \rightarrow & \Sigma^2 G & \rightarrow & \Sigma^3 E & & E \wedge M & \rightarrow & F \wedge M & \rightarrow & G \wedge M & \rightarrow & \Sigma E \wedge M \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 E & \rightarrow & F & \rightarrow & G & \rightarrow & \Sigma E & & E & \rightarrow & F & \rightarrow & G & \rightarrow & \Sigma E
 \end{array}$$

with cofiber sequences. Under the first two assumptions two left squares become homotopy commutative because they induce the  $Z_2$ -graded homomorphism  $E_{\sharp}(\ ) \rightarrow F_{\sharp}(\ )$  of  $M_{\sharp}(\ )$ -modules. Therefore there exist morphisms  $\Sigma^2 G \rightarrow G$  and  $G \wedge M \rightarrow G$  which make the above diagrams into morphisms of cofiber sequences. As is easily checked, they give  $G_{\sharp}(\ )$  a structure of  $Z_2$ -graded  $M_{\sharp}(\ )$ -module, which coincides with the original one. So, using the remaining assumptions again we see that the above morphisms are homotopic to the given ones respectively. Consequently the cofiber sequence  $E \rightarrow F \rightarrow G$  becomes the required one.

Another case is similarly proved.

The ring spectrum  $K$  may be regarded as a  $Z_2$ -graded  $MU$ -module spectrum via the  $Z_2$ -graded Thom map  $\mu_c: MU \rightarrow K$ .

Applying Proposition 10 to three cofiber sequences of (1.6) we get

**Theorem 2.** *The sequences  $F_{q+1}MU \rightarrow MU \rightarrow Q_qMU$ ,  $F_{q+1}MU \rightarrow F_qMU \rightarrow KG_q$  and  $KG_q \rightarrow Q_qMU \rightarrow Q_{q-1}MU$  are cofiber sequences of  $Z_2$ -graded  $MU$ -module spectra.*

Proof. The assumptions needed in Proposition 10 are satisfied by Lemmas 7,8 and 9.

As a result we have a tower

$$(2.7) \quad MU \rightarrow \dots \rightarrow Q_qMU \rightarrow Q_{q-1}MU \rightarrow \dots \rightarrow Q_0MU = K$$

of  $Z_2$ -graded  $MU$ -module spectra such that  $KG_q \rightarrow Q_qMU \rightarrow Q_{q-1}MU$  is a cofiber sequence, which factorizes the  $Z_2$ -graded Thom map  $\mu_c: MU \rightarrow K$ .

2.3. Here we extend the Wolff's result to the case of based CW-complexes.

**Proposition 11.** *There exists an (unstable) natural homomorphism*

$$\Phi_q: KG_q^*(X) \rightarrow F_qMU^*(X)$$

for any based CW-complex  $X$ , which satisfies the equality that  $\mu_{q^*} \cdot \Phi_q = \text{id}$ .

Proof. We may assume that  $X$  is connected. Let  $i: BU_{G_q} \rightarrow KG_q$  be the inclusion. Then we can choose a morphism  $c_q: BU_{G_q} \rightarrow F_qMU$  such that  $i$  is homotopic to the composition  $\mu_{q^*} \cdot c_q$ , because  $F_{q+1}MU^1(BU_{G_q}) = 0$ . In the commutative diagram

$$\begin{array}{ccc} [X, BU_{G_q}] \xrightarrow{J_0} \{X, BU_{G_q}\} & \xrightarrow{i^*} & \{X, KG_q\} = KG_q^0(X) \\ & \searrow c_{q^*} & \uparrow \mu_{q^*} \\ & & \{X, F_qMU\} = F_qMU^0(X) \end{array}$$

the composition  $i_* \cdot J_0$  is an isomorphism because of Proposition 2 (see [6, Theorem 14.5]). So we put that  $\Phi_q = c_{q*} \cdot J_0 \cdot (i_* \cdot J_0)^{-1}$ .

REMARK. If  $\Phi_q$  is stable, then we have a natural split exact sequence

$$0 \rightarrow F_{q+1}MU^*(X) \rightarrow F_qMU^*(X) \rightarrow KG_q^*(X) \rightarrow 0$$

for every  $CW$ -spectrum  $X$ . Therefore  $F_qMU$  becomes homotopy equivalent to the wedge  $F_{q+1}MU \vee KG_q$ . However  $H_*(F_qMU)$  is a free abelian group and  $H_*(KG_q)$  is a  $Q$ -module. This is a contradiction.

We now obtain our main result.

**Theorem 3.** *For any based  $CW$ -complex  $X$  the natural sequences*

$$\begin{aligned} 0 \rightarrow F_{q+1}MU^*(X) \rightarrow F_qMU^*(X) \rightarrow KG_q^*(X) \rightarrow 0 \\ 0 \rightarrow KG_q^*(X) \rightarrow Q_qMU^*(X) \rightarrow Q_{q-1}MU^*(X) \rightarrow 0 \end{aligned}$$

of  $Z_2$ -graded  $\Lambda$ -modules are split exact.

Proof. The first case is immediate from Proposition 11. On the other hand, a diagram chase shows that the second sequence is exact for any based  $CW$ -complex  $X$  and hence it is split.

## Appendix

Recall that  $MU$  is a ring spectrum with coefficients  $\Lambda_* = Z[x_1, \dots, x_n, \dots]$  where  $\deg x_n = 2n$ . By killing certain bordism classes Baas [3] constructed homology theories  $MU\langle n \rangle_*( )$  with coefficient  $\pi_*(MU\langle n \rangle) = \Lambda_*(x_{n+1}, \dots)$ , whose representing spectrum we denote by  $MU\langle n \rangle$ .  $MU\langle n \rangle_*( )$  is an (associative)  $MU_*( )$ -module, thus there exists a natural homomorphism

$$m_n: MU_*(X) \otimes MU\langle n \rangle_*(Y) \rightarrow MU\langle n \rangle_*(X \wedge Y)$$

for any  $CW$ -spectra  $X$  and  $Y$ . This gives us a pairing

$$\phi\langle n \rangle: MU \wedge MU\langle n \rangle \rightarrow MU\langle n \rangle$$

by which the above  $m_n$  is induced.

An easy computation shows that  $MU\langle n \rangle \hat{Z}/Z^{2k-1}(MU \wedge \dots \wedge MU \wedge MU\langle n \rangle) = 0$  because  $\pi_{2l+1}(MU\langle n \rangle) = 0$  for all  $l$ . Then [8, Theorem 1] says that

(A.1)  $MU\langle n \rangle^{2k}(MU \wedge \dots \wedge MU \wedge MU\langle m \rangle)$  is Hausdorff (see also [9, Corollary 13]).

Hence  $\phi\langle n \rangle$  is uniquely determined (up to homotopy) and moreover

(A.2)  $MU\langle n \rangle$  is an (associative)  $MU$ -module spectrum.

An important relationship between  $MU\langle n \rangle_*( )$  and  $MU\langle n-1 \rangle_*( )$  is given in the form of a natural exact sequence

$$\rightarrow MU\langle n \rangle_{*-2n}(X) \xrightarrow{\cdot x_n} MU\langle n \rangle_*(X) \xrightarrow{t_n} MU\langle n-1 \rangle_*(X) \rightarrow MU\langle n \rangle_{*-2n-1}(X) \rightarrow$$

of  $MU_*( )$ -modules where  $\cdot x_n$  denotes the multiplication by  $x_n$ . Because of (A.1) there exists a unique morphism  $\tau_n: MU\langle n \rangle \rightarrow MU\langle n-1 \rangle$  of  $MU$ -module spectra whose induced homomorphism is the above  $t_n$ . On the other hand, the composition

$$m_{x_n}: \Sigma^{2n} MU\langle n \rangle \xrightarrow{x_n \wedge 1} MU \wedge MU\langle n \rangle \xrightarrow{\phi\langle n \rangle} MU\langle n \rangle$$

is characterized by the above multiplication  $\cdot x_n$ .

**Lemma A.** *Let  $E \xrightarrow{f} F \xrightarrow{g} G$  be a sequence of CW-spectra such that the composition  $g \cdot f$  is homotopic to the zero. If  $0 \rightarrow \pi_*(E) \rightarrow \pi_*(F) \rightarrow \pi_*(G) \rightarrow 0$  is exact, then  $E \rightarrow F \rightarrow G$  is a cofiber sequence. (Cf., Lemma 4).*

*Proof.* Let  $C_f$  be the mapping cone of  $f: E \rightarrow F$ . Then  $g: F \rightarrow G$  admits a factorization  $F \rightarrow C_f \xrightarrow{h} G$ . Considering the commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & \pi_*(E) & \rightarrow & \pi_*(F) & \rightarrow & \pi_*(C_f) \rightarrow 0 \\ & & \parallel & & \parallel & & \downarrow h_* \\ 0 & \rightarrow & \pi_*(E) & \rightarrow & \pi_*(F) & \rightarrow & \pi_*(G) \rightarrow 0 \end{array}$$

with exact rows, we see easily that  $h: C_f \rightarrow G$  is a homotopy equivalence.

Using (A.1) the composition  $\tau_n \cdot m_{x_n}$  becomes homotopic to the zero. We get therefore that

$$(A.3) \quad \Sigma^{2n} MU\langle n \rangle \xrightarrow{m_{x_n}} MU\langle n \rangle \xrightarrow{\tau_n} MU\langle n-1 \rangle \text{ is a cofiber sequence.}$$

OSAKA CITY UNIVERSITY

**References**

- [1] J.F. Adams: *A variant of E.H. Brown's representability theorem*, *Topology* **10** (1971), 185-198.
- [2] J.F. Adams: *Stable Homotopy and Generalized Homology*, Chicago Lecture in Math., Univ. of Chicago Press.
- [3] N.A. Baas: *On bordism theory of manifolds with singularities*, *Math. Scand.* **33** (1973), 279-302.

- [4] P.A. Griffith: *Infinite Abelian Group Theory*, Chicago Lecture in Math., Univ. of Chicago Press.
- [5] P.S. Landweber: *Homological properties of comodules over  $MU_*(MU)$  and  $BP_*$  ( $BP$ )*, to appear.
- [6] R. Vogt: *Boardman's Stable Homotopy Category*, Lecture notes series 21, Aarhus Univ. (1970).
- [7] G. Wolff: *Der einfluss von  $K^*(-)$  auf  $U^*(-)$* , *Manuscripta Math.* **10** (1973), 141–161.
- [8] Z. Yosimura: *On cohomology theories of infinite CW-complexes*, III, *Publ. Res. Inst. Math. Sci.* **9** (1974), 683–706.
- [9] Z. Yosimura: *Universal coefficient sequences for cohomology theories of CW-spectra*, *Osaka J. Math.* **12** (1975), 305–323.

