



Title	Stiefel manifolds as framed boundaries
Author(s)	Minami, Haruo
Citation	Osaka Journal of Mathematics. 1990, 27(1), p. 185-189
Version Type	VoR
URL	https://doi.org/10.18910/11170
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

STIEFEL MANIFOLDS AS FRAMED BOUNDARIES

HARUO MINAMI

(Received February 8, 1989)

Let $V_{n,q}$ denote the Stiefel manifold of orthogonal q -frames in F^n where $F = \mathbf{R}, \mathbf{C}, \mathbf{H}$. We regard this space as the homogeneous space G_n/G_{n-q} of right cosets modulo $1_q \times G_{n-q}$ where G_k denotes the relevant group $SO(k)$, $SU(k)$ or $Sp(k)$. Then this space obtains a framing in a canonical way as mentioned below [3]. We denote this framing ambiguously by \mathcal{F} and we write $[V_{n,q}, \mathcal{F}]$ for an element in π_*^S defined by the pair $(V_{n,q}, \mathcal{F})$ via the Thom-Pontrjagin construction. In this note we prove the following

Theorem. *Let $1 \leq q \leq n-1$, $n-1$ or n according as $F = \mathbf{R}, \mathbf{C}$ or \mathbf{H} . Then*

$$[V_{n,q}, \mathcal{F}] = 0.$$

We denote by \mathcal{R} the right invariant framing of G_n and by \mathcal{R}^α the framing obtained by twisting \mathcal{R} by a representation α [5]. Also we write $[G_n, \alpha]$ for $[G_n, \mathcal{R}^\alpha]$. Let

$$\rho_n: G_n \subset GL(dn, \mathbf{R})$$

be the standard real representation of G_n where $d = \dim_{\mathbf{R}} F$. Then by the theorem we have

Corollary ([1], [5]).

$$[SO(n), (n-1)\rho_n] = 0, [SU(n), (n-1)\rho_n] = 0 \quad \text{and} \quad [Sp(n), n\rho_n] = 0.$$

REMARK. By taking $G_k = U(k)$ instead of $SU(k)$ we get $[U(n), n\rho_n] = 0$ analogously.

The proof of the theorem uses the arguments parallel to [4]. Actually we construct a bounding manifold for $V_{n,q} = G_n/G_{n-q}$.

Let V_k denote the representation space of ρ_k . There is then the real vector bundle

$$\xi_k: G_n \times_{G_k} V_k \rightarrow G_n/G_k$$

for $k < n$. If $S(V_k)$ denote the unit sphere of V_k , then we have a canonical G_k -equivariant diffeomorphism.

$$G_k/G_{k-1} \approx S(V_k).$$

This and $G_n/G_{n-q} = G_n \times_{G_{n-q+1}} G_{n-q+1}/G_{n-q}$ imply that the homogeneous fibre bundle

$$(1) \quad G_{n-q+1}/G_{n-q} \rightarrow G_n/G_{n-q} \xrightarrow{\pi} G_n/G_{n-q+1}$$

is isomorphic to the sphere bundle of ξ_{n-q+1} . Hence we have

$$G_n/G_{n-q} \approx S(\xi_{n-q+1})$$

where $S(\xi_k)$ denotes the total space of the sphere bundle of ξ_k . Denote also by $D(\xi_k)$ the total space of the disc bundle of ξ_k . Evidently we then have

$$\partial D(\xi_{n-q+1}) \approx V_{n,q}.$$

To prove the theorem it therefore suffices to show that the framing \mathcal{F} of $V_{n,q}$ extends over $D(\xi_{n-q+1})$.

So we first recall the framing of [3]. Let G be a compact connected Lie group and H a closed subgroup of G . Let $\tau(G/H)$ denote the tangent bundle of G/H . Consider the principal H -bundle

$$H \rightarrow G \xrightarrow{\pi} G/H.$$

Then we have a decomposition of the tangent bundle of G

$$\tau(G) \cong \pi^* \tau(G/H) \oplus \tau_H(G)$$

where $\tau_H(G)$ is the bundle of tangents along the fibres. This isomorphism is compatible with the right action of H , so that we obtain an isomorphism of vector bundles over G/H

$$\tau(G)/H \cong \tau(G/H) \oplus \tau_H(G)/H.$$

Let $\tau_g(G)$ denote the tangent space at $g \in G$ and $R_{g^{-1}}: \tau_g(G) \rightarrow \tau_e(G)$ denote the differential of right multiplication by g^{-1} where e is the identity element of G . Then the right invariant framing of G

$$\mathcal{R}: \tau(G) \cong G \times \tau_e(G)$$

is given by $\mathcal{R}(v) = (g, R_{g^{-1}}(v))$ where $v \in \tau_g(G)$. This gives

$$\tau(G)/H \cong G/H \times \tau_e(G).$$

as vector bundles over G/H .

By ad_H we denote the adjoint representation of H on $\tau_e(H)$. We consider the differential $L_{a^{-1}}: \tau_a(gH) \rightarrow \tau_e(H)$ induced by the left multiplication by a^{-1} where $a \in gH$. Then similarly we have

$$\tau_H(G)/H \cong G \times_H \tau_e(H)$$

as vector bundles over G/H where H acts on $\tau_e(H)$ via ad_H .

Combining these three bundle equations we have

$$(2) \quad G/H \times \tau_e(G) \cong \tau(G/H) \oplus G \times_{\text{ad}_H} \tau_e(H)$$

as vector bundles over G/H . So we find that if ad_H is contained in the image of the restriction map $RO(G) \rightarrow RO(H)$ of real representation rings, then formula (2) gives rise to a framing of G/H .

Here we return to the framing of $V_{n,q}$. We consider the restrictions of ad_{G_n} and ρ_n to G_k for $k < n$. Now we write $\text{ad}_k = \text{ad}_{G_k}$ briefly. Then we have

$$\begin{aligned} \text{Lemma 1.} \quad & (i) \quad \rho_n|_{G_{n-1}} \cong \rho_{n-1} \oplus d \cdot 1, \\ & (ii) \quad \text{ad}_n|_{G_{n-1}} \cong \text{ad}_{n-1} \oplus \rho_{n-1} \oplus (d-1) \cdot 1 \end{aligned}$$

where 1 denotes the trivial 1-dimensional real representation.

Proof. (i) is obvious. By observing the maximal root of G_n [2] we see that $c(\text{ad}_n) = \lambda^2 \tilde{\rho}_n, (\lambda^1 \tilde{\rho}_n)(\lambda^{n-1} \tilde{\rho}_n) - 1$ or $(\lambda^1 \tilde{\rho}_n)^2 - \lambda^2 \tilde{\rho}_n$ where $\tilde{\rho}_n$ denotes the canonical complex representation $G_n \subset GL(n, \mathbf{C}), GL(n, \mathbf{C})$ or $GL(2n, \mathbf{C})$ according as $F = \mathbf{R}, \mathbf{C}$ or \mathbf{H} . Here c is the complexification and $\lambda^i \tilde{\rho}_n$ is the i -th exterior power of $\tilde{\rho}_n$. For this fact, however, we refer to [6]. So we can readily obtain (ii) using (i).

From Lemma 1 it follows that for $k < n$

$$(3) \quad \begin{aligned} \text{ad}_n|_{G_k} &\cong \text{ad}_k \oplus (n-k) \rho_k \oplus ((n-k)(dn-dk+d-2)/2) \cdot 1, \\ \rho_n|_{G_k} &\cong \rho_k \oplus d(n-k) \cdot 1. \end{aligned}$$

Hence we get

$$(4) \quad \text{ad}_n|_{G_k} \oplus s(n, k) \cdot 1 \cong \text{ad}_k \oplus (n-k) \rho_n|_{G_k}$$

for $k < n$ where $s(n, k) = (n-k)(dn-dk-d+2)/2$.

Denote by W_k the representation space of ad_k . Then using (2) when $G = G_n, H = G_{n-q}$ and (4) we obtain a framing of $V_{n,q}$

$$(5) \quad \mathcal{F}: \tau(V_{n,q}) \oplus (V_{n,q} \times W_n) \oplus s(n, n-q) \cdot 1 \cong V_{n,q} \times (\tau_e(G_n) \oplus qV_n)$$

where 1 denotes the trivial real line bundle.

We now give this framing \mathcal{F} more directly. Let W'_k be a direct summand of $W_n|_{G_k}$ such that $W_n|_{G_k} \cong W'_k \oplus W_k$ for $k < n$. Then we have

$$\text{Lemma 2 ([1]).} \quad \tau(V_{n,n-k}) \cong G_n \times_{G_k} W'_k \quad \text{for } k < n.$$

Proof. There is an obvious isomorphism of vector bundles over $V_{n,n-k}$

$$G_n \times_{G_k} (W_n|_{G_k}) \cong V_{n,n-k} \times \tau_e(G_n).$$

So consider the composite with the isomorphism of (2) when $G=G_n$, $H=G_k$

$$G_n \times_{G_k} (W_n|_{G_k}) \cong \tau(V_{n,n-k}) \oplus G_n \times_{G_k} W_k.$$

Then we see that this map sends identically the direct summand $G_n \times_{G_k} W_k$ to that on the right-hand side and isomorphically another direct summand $G_n \times_{G_k} W'_k$ to $\tau(V_{n,n-k})$, and so the result follows.

Using Lemma 2 we can interpret \mathcal{F} of (5) as follows. By (4) we have

$$(6) \quad W'_{k-q} \oplus (W_n|_{G_{n-q}}) \oplus s(n, n-q) \cdot 1 \cong (W_n \oplus qV_n)|_{G_{n-q}}.$$

This gives rise to an isomorphism of real vector bundles associated with the principal G_{n-q} -bundle $G_n \rightarrow V_{n,q}$ with modules on both sides as fibres. It is easily seen that this isomorphism induces \mathcal{F} of (5) precisely.

Proof of Theorem. By the second formula of (3) it follows that

$$\xi_{n-q+1} \oplus d(q-1) \cdot 1 \cong V_{n,q-1} \times V_n.$$

Taking the sum of this and the framing \mathcal{F} of $V_{n,q-1}$ we have

$$(7) \quad \begin{aligned} &\tau(V_{n,q-1}) \oplus \xi_{n-q+1} \oplus (V_{n,q-1} \times W_n) \oplus (s(n, n-q) - 1) \cdot 1 \\ &\cong V_{n,q-1} \times (\tau_e(G_n) \oplus qV_n) \end{aligned}$$

since $s(n, n-q) = s(n, n-q+1) + d(q-1) + 1$. Now from the above arguments about the fibre bundle of (1) it is clear that

$$\tau(V_{n,q}) \oplus 1 \cong \pi^*(\tau(V_{n,q-1}) \oplus \xi_{n-q+1})$$

where π is the projection map of (1). Therefore by pulling the isomorphism of (7) back along π we obtain another framing of $V_{n,q}$

$$\mathcal{F}': \tau(V_{n,q}) \oplus (V_{n,q} \times W_n) \oplus s(n, n-q) \cdot 1 \cong V_{n,q} \times (\tau_e(G_n) \oplus qV_n).$$

Denote by $\tilde{\pi}$ the canonical projection map $D(\xi_{n-q+1}) \rightarrow V_{n,q-1}$. Moreover we then have

$$\tau(D(\xi_{n-q+1})) \cong \tilde{\pi}^*(\tau(V_{n,q-1}) \oplus \xi_{n-q+1}),$$

so that by pulling the isomorphism of (7) back along $\tilde{\pi}$ again we obtain a framing of $D(\xi_{n-q+1})$. Identifying $V_{n,q}$ with $S(\xi_{n-q+1})$, this framing is obviously an extension of \mathcal{F}' over $D(\xi_{n-q+1})$ since $\tilde{\pi}|_{V_{n,q}} = \pi$. Hence it follows $[V_{n,q}, \mathcal{F}'] = 0$ and so it suffices to show that \mathcal{F} agrees up to sign with \mathcal{F}' .

By (3) we have

$$W'_{n-q} \oplus 1 \cong (W'_{n-q+1} \oplus V_{n-q+1})|_{G_{n-q}}.$$

Using this and Lemma 2 we can verify that formula (6) also gives rise to either

\mathcal{F}' or $-\mathcal{F}'$ in the same way as the case of \mathcal{F} . Therefore this proves the theorem.

Proof of Corollary. Since $V_{n,n-1}=SO(n)$, $SU(n)$ and $V_{n,n}=Sp(n)$, we set $V_{n,q}=G_n$ in (5) and so in defining \mathcal{F} we consider $\text{ad}_{n-q}=0$. Hence by definition it follows that \mathcal{F} is just the framing twisting \mathcal{R} of G_n by $\text{ad}_n-q\rho_n$, so that by the theorem it follows $[G_n, \text{ad}_n-q\rho_n]=0$. Therefore we have $[G_n, q\rho_n]=0$ by [5].

References

- [1] L. Astey, M.A. Guest and G. Pastor: *Lie groups as framed boundaries*, Osaka J. Math. **25** (1988), 891–907.
- [2] N. Bourbaki: *Groupes et algèbres de Lie*, Hermann, Paris, 1968.
- [3] P. Löffler and L. Smith: *Line bundles over framed manifolds*, Math. Z. **138** (1974), 35–52.
- [4] H.V. Pittie and L. Smith: *Generalized flag manifolds as framed boundaries*, Math. Z. **142** (1975), 191–193.
- [5] B. Steer: *Orbits and the homotopy class of a compactification of a classical map*, Topology **15** (1976), 383–393.
- [6] I. Yokota: *Groups and representations* (in Japanese), Shôkabô, Tokyo, 1973.

Department of Mathematics
Naruto University of Education
Naruto, Tokushima 772
Japan

