



Title	Totally geodesic hypersurfaces of naturally reductive homogeneous spaces
Author(s)	Tsukada, Kazumi
Citation	Osaka Journal of Mathematics. 1996, 33(3), p. 697-707
Version Type	VoR
URL	<a href="https://doi.org/10.18910/11213">https://doi.org/10.18910/11213</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Tsukada, K.  
Osaka J. Math.  
33 (1996), 697-707

## **TOTALLY GEODESIC HYPERSURFACES OF NATURALLY REDUCTIVE HOMOGENEOUS SPACES**

KAZUMI TSUKADA

(Received June 8, 1995)

### **1. Introduction**

Totally geodesic submanifolds of Riemannian symmetric spaces have been well investigated and it has been shown that they have beautiful and fruitful properties. In particular, due to the  $(M_+, M_-)$ -theory by B.Y. Chen and T. Nagano [1] this subject has made great progress. Naturally reductive homogeneous spaces are known as a natural generalization of Riemannian symmetric spaces. K. Tojo [6] investigated totally geodesic submanifolds of naturally reductive homogeneous spaces and obtained a necessary and sufficient condition of their existence. We will recall his result in section 3. Moreover he implicitly made the following conjecture.

**Conjecture.** *If a simply connected irreducible naturally reductive homogeneous space  $M$  admits a totally geodesic hypersurface, then  $M$  has constant sectional curvature.*

The conjecture is regarded as a generalization of the result which was shown in the case of Riemannian symmetric spaces by B.Y. Chen and T. Nagano [1]. K. Tojo gave an affirmative answer to the conjecture in the case that  $\dim M = 3, 4$  and  $5$  [6] and in the case that  $M$  is a normal homogeneous space [7]. We shall prove that the conjecture above is true.

**Main Theorem.** *If a simply connected irreducible (as a Riemannian manifold) naturally reductive homogeneous space  $M$  admits a totally geodesic hypersurface, then  $M$  has constant sectional curvature.*

We shall discuss the irreducibility of naturally reductive homogeneous spaces in Section 2 and prove the main theorem in Section 3.

### **2. Irreducibility of naturally reductive homogeneous spaces**

We first recall basic definitions and properties of naturally reductive

homogeneous spaces, following J.E. D'Atri and W. Ziller [2] and S. Kobayashi and K. Nomizu [3]. See also O. Kowalski and L. Vanhecke [4], [5]. Let  $(M, g)$  be a homogeneous Riemannian manifold. Let  $K$  be a connected Lie group of isometries which acts transitively and almost effectively on  $M$  and let  $H$  be the isotropy subgroup at a point  $o \in M$ . Let  $\mathfrak{k}$  be the Lie algebra of  $K$  and  $\mathfrak{h}$  the subalgebra corresponding to  $H$ . Let  $\mathfrak{m}$  be an  $Ad(H)$ -invariant subspace which is complementary to  $\mathfrak{h}$  in  $\mathfrak{k}$ . We denote by  $x_{\mathfrak{h}}$  and  $x_{\mathfrak{m}}$  the  $\mathfrak{h}$ -component and the  $\mathfrak{m}$ -component of  $x \in \mathfrak{k}$ , respectively. As usual we identify  $\mathfrak{m}$  with the tangent space  $T_o M$  at  $o$  and denote by  $\langle \cdot, \cdot \rangle$  the inner product on  $\mathfrak{m}$  induced from the metric  $g_o$  on  $T_o M$ .

**DEFINITION 2.1.** A homogeneous Riemannian manifold  $(M, g)$  is said to be a *naturally reductive homogeneous space* if there exist  $K$  and  $\mathfrak{m}$  as above such that

$$(2.1) \quad \langle [x, y]_{\mathfrak{m}}, z \rangle + \langle y, [x, z]_{\mathfrak{m}} \rangle = 0 \quad \text{for any } x, y, z \in \mathfrak{m}.$$

From now on we assume that  $(M, g)$  is a naturally reductive homogeneous space. Then by a theorem of Kostant we may assume that  $\mathfrak{k} = \mathfrak{m} + [\mathfrak{m}, \mathfrak{m}]$ . Let  $\Lambda_{\mathfrak{m}} : \mathfrak{m} \rightarrow \mathfrak{so}(\mathfrak{m})$  be a linear mapping which corresponds to the Riemannian connection  $\nabla$  (see [3] Chapter X), where  $\mathfrak{so}(\mathfrak{m})$  denotes the Lie algebra consisting of skew symmetric endomorphisms of  $(\mathfrak{m}, \langle \cdot, \cdot \rangle)$ . Then  $\Lambda_{\mathfrak{m}}$  is given by

$$(2.2) \quad \Lambda_{\mathfrak{m}}(x)(y) = \frac{1}{2} [x, y]_{\mathfrak{m}} \quad \text{for } x, y \in \mathfrak{m}$$

(cf. Theorem 3.3 p.201 in [3]),

**DEFINITION 2.2.** A subspace  $V$  of  $\mathfrak{m}$  is said to be  $\Lambda_{\mathfrak{m}}$ -*invariant* if it satisfies  $\Lambda_{\mathfrak{m}}(x)(V) \subset V$  for any  $x \in \mathfrak{m}$ . Moreover a  $\Lambda_{\mathfrak{m}}$ -invariant subspace  $V$  is  $\Lambda_{\mathfrak{m}}$ -*irreducible* if  $V$  has only trivial  $\Lambda_{\mathfrak{m}}$ -invariant subspaces.

We set  $\mathfrak{m}_0 = \{v \in \mathfrak{m} \mid \Lambda_{\mathfrak{m}}(x)(v) = 0 \text{ for any } x \in \mathfrak{m}\}$ . Then we evidently have the following orthogonal decomposition into  $\Lambda_{\mathfrak{m}}$ -invariant subspaces:

$$(2.3) \quad \mathfrak{m} = \mathfrak{m}_0 \oplus \mathfrak{m}_1 \oplus \cdots \oplus \mathfrak{m}_r,$$

where for each  $i$  ( $1 \leq i \leq r$ )  $\mathfrak{m}_i$  is  $\Lambda_{\mathfrak{m}}$ -irreducible and  $\Lambda_{\mathfrak{m}}(x)|_{\mathfrak{m}_i} \neq 0$  for some  $x \in \mathfrak{m}$ .

**Theorem 2.3.** Let  $M = K/H$  be a naturally reductive homogeneous space with  $Ad(H)$ -invariant decomposition  $\mathfrak{k} = \mathfrak{h} \oplus \mathfrak{m}$ . We assume that  $\mathfrak{k} = \mathfrak{m} + [\mathfrak{m}, \mathfrak{m}]$ . Let

$$\mathfrak{m} = \mathfrak{m}_0 \oplus \mathfrak{m}_1 \oplus \cdots \oplus \mathfrak{m}_r$$

be the decomposition of  $\mathfrak{m}$  which satisfies (2.3). If we set

$$\mathfrak{k}_i = \mathfrak{m}_i + [\mathfrak{m}_i, \mathfrak{m}_i] \quad (i = 0, 1, \dots, r)$$

$$\mathfrak{h}_i = \mathfrak{k}_i \cap \mathfrak{h} \quad (i=0,1,\dots,r),$$

then we have  $\mathfrak{k} = \mathfrak{k}_0 \oplus \mathfrak{k}_1 \oplus \dots \oplus \mathfrak{k}_r$ , and  $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1 \oplus \dots \oplus \mathfrak{h}_r$  as direct sums of Lie algebras.

Proof. We first show the following identity.

**Lemma 2.4.** *Let  $M = K/H$  be a homogeneous space with  $Ad(H)$ -invariant decomposition  $\mathfrak{k} = \mathfrak{h} \oplus \mathfrak{m}$ . Then the following holds:*

$$[[x,y]_{\mathfrak{m}},z]_{\mathfrak{h}} + [[y,z]_{\mathfrak{m}},x]_{\mathfrak{h}} + [[z,x]_{\mathfrak{m}},y]_{\mathfrak{h}} = 0$$

for  $x,y,z \in \mathfrak{m}$ .

Proof of Lemma 2.4. By the Jacobi's identity, we have

$$\begin{aligned} 0 &= [[x,y],z] + [[y,z],x] + [[z,x],y] \\ &= [[x,y]_{\mathfrak{h}},z] + [[y,z]_{\mathfrak{h}},x] + [[z,x]_{\mathfrak{h}},y] \\ &\quad + [[x,y]_{\mathfrak{m}},z] + [[y,z]_{\mathfrak{m}},x] + [[z,x]_{\mathfrak{m}},y] \end{aligned}$$

for  $x,y,z \in \mathfrak{m}$ .

Comparing the  $\mathfrak{h}$ -components of both sides, we obtain the identity in Lemma 2.4.  $\square$

By (2.2) and (2.3), we have  $[\mathfrak{m},\mathfrak{m}_i]_{\mathfrak{m}} \subset \mathfrak{m}_i$ . In particular,

$$(2.4) \quad [\mathfrak{m}_i, \mathfrak{m}_j]_{\mathfrak{m}} = 0 \quad \text{for } i \neq j,$$

$$(2.5) \quad [\mathfrak{m}_i, \mathfrak{m}_i]_{\mathfrak{m}} = \mathfrak{m}_i \quad \text{for } i \geq 1.$$

**Lemma 2.5.** *The following relations hold:*

- (1)  $[\mathfrak{m}_i, \mathfrak{m}_j] = 0$  for  $i \neq j$ .
- (2)  $[[\mathfrak{m}_i, \mathfrak{m}_i], \mathfrak{m}_j] = 0$  for  $i \neq j$ .
- (3)  $[[\mathfrak{m}_i, \mathfrak{m}_i]_{\mathfrak{h}}, \mathfrak{m}_i] \subset \mathfrak{m}_i$ .
- (4)  $[[\mathfrak{m}_i, \mathfrak{m}_i], \mathfrak{m}_i] \subset \mathfrak{m}_i + [\mathfrak{m}_i, \mathfrak{m}_i]$ .

Proof of Lemma 2.5. (1) It is sufficient to prove that  $[\mathfrak{m}_i, \mathfrak{m}_j]_{\mathfrak{h}} = 0$  for  $i \neq j$ . We may assume that  $i \geq 1$ . By Lemma 2.4, we have for  $x, y \in \mathfrak{m}_i$  and  $z \in \mathfrak{m}_j$ ,

$$[[x,y]_{\mathfrak{m}},z]_{\mathfrak{h}} = -[[y,z]_{\mathfrak{m}},x]_{\mathfrak{h}} - [[z,x]_{\mathfrak{m}},y]_{\mathfrak{h}} = 0.$$

Since  $[\mathfrak{m}_i, \mathfrak{m}_i]_{\mathfrak{m}} = \mathfrak{m}_i$  for  $i \geq 1$ , we have  $[\mathfrak{m}_i, \mathfrak{m}_j]_{\mathfrak{h}} = 0$ .

(2) From the Jacobi's identity and (1), it follows that for  $x, y \in \mathfrak{m}_i$ ,  $z \in \mathfrak{m}_j$

$$[[x,y],z] = -[[y,z],x] - [[z,x],y] = 0.$$

(3) By (1) and (2), we obtain  $[[x, y]_{\mathfrak{h}}, z] = 0$  for  $x, y \in \mathfrak{m}_i$ ,  $z \in \mathfrak{m}_j$  ( $i \neq j$ ). Therefore for  $x, y, v \in \mathfrak{m}_i$ ,  $z \in \mathfrak{m}_j$  ( $i \neq j$ )

$$\langle [[x, y]_{\mathfrak{h}}, v], z \rangle = -\langle v, [[x, y]_{\mathfrak{h}}, z] \rangle = 0,$$

that is,  $[[x, y]_{\mathfrak{h}}, \mathfrak{m}_i] \subset \mathfrak{m}_i$ .

(4) By (3) and (2.5), we obtain (4).  $\square$

Proof of Theorem 2.3. We first prove that each  $\mathfrak{k}_i$  is an ideal of  $\mathfrak{k}$ . In fact applying the relations in Lemma 2.5, we obtain the following:

$$\begin{aligned} [\mathfrak{m}, \mathfrak{m}_i] &\subset [\mathfrak{m}_i, \mathfrak{m}_i], \\ [\mathfrak{m}, [\mathfrak{m}_i, \mathfrak{m}_i]] &\subset [\mathfrak{m}_i, [\mathfrak{m}_i, \mathfrak{m}_i]] \subset \mathfrak{m}_i + [\mathfrak{m}_i, \mathfrak{m}_i], \\ [[\mathfrak{m}, \mathfrak{m}], \mathfrak{m}_i] &\subset \left[ \sum_{j=0}^r [[\mathfrak{m}_j, \mathfrak{m}_j], \mathfrak{m}_i], \right. \\ &\quad \left. \subset [[\mathfrak{m}_i, \mathfrak{m}_i], \mathfrak{m}_i] \subset \mathfrak{m}_i + [\mathfrak{m}_i, \mathfrak{m}_i], \right. \\ [[\mathfrak{m}, \mathfrak{m}], [\mathfrak{m}_i, \mathfrak{m}_i]] &\subset [[[m, m], \mathfrak{m}_i], \mathfrak{m}_i] \\ &\subset [\mathfrak{m}_i + [\mathfrak{m}_i, \mathfrak{m}_i], \mathfrak{m}_i] \subset \mathfrak{m}_i + [\mathfrak{m}_i, \mathfrak{m}_i]. \end{aligned}$$

Since  $[\mathfrak{m}_i, \mathfrak{m}_i]_{\mathfrak{m}} \subset \mathfrak{m}_i$  ( $i = 0, 1, \dots, r$ ), we have  $\mathfrak{h}_i = [\mathfrak{m}_i, \mathfrak{m}_i]_{\mathfrak{h}}$  and hence  $\mathfrak{k}_i = \mathfrak{m}_i \oplus \mathfrak{h}_i$  (direct sum). Finally we shall show that  $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1 \oplus \dots \oplus \mathfrak{h}_r$  as a direct sum of vector spaces. Let  $x$  be a vector of  $(\mathfrak{h}_0 + \dots + \mathfrak{h}_i) \cap \mathfrak{h}_{i+1}$ . Since  $x \in \mathfrak{h}_{i+1}$  by (1) and (2), it follows  $[x, v] = 0$  for any  $v \in \mathfrak{m}_0 + \dots + \mathfrak{m}_i + \mathfrak{m}_{i+2} + \dots + \mathfrak{m}_r$ . On the other hand since  $x \in \mathfrak{h}_0 + \dots + \mathfrak{h}_i$  again by (1) and (2), it follows  $[x, v] = 0$  for any  $v \in \mathfrak{m}_{i+1}$ . These imply  $[x, v] = 0$  for any  $v \in \mathfrak{m}$ . Since  $K$  acts almost effectively on  $M$ , we have  $x = 0$ . Hence  $(\mathfrak{h}_0 + \dots + \mathfrak{h}_i) \cap \mathfrak{h}_{i+1} = 0$ . Since  $[\mathfrak{m}, \mathfrak{m}]_{\mathfrak{h}} = \mathfrak{h}$ , we have  $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1 \oplus \dots \oplus \mathfrak{h}_r$ . Noticing that  $\mathfrak{k}_i$  are ideals of  $\mathfrak{k}$ , we have  $\mathfrak{k} = \mathfrak{k}_0 \oplus \mathfrak{k}_1 \oplus \dots \oplus \mathfrak{k}_r$  and  $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1 \oplus \dots \oplus \mathfrak{h}_r$  as direct sums of Lie algebras.  $\square$

**Corollary 2.6.** *Let  $M = K/H$  be a simply connected irreducible (as a Riemannian manifold) naturally reductive homogeneous space. If  $\Lambda_{\mathfrak{m}} \neq 0$ ,  $\mathfrak{m}$  is  $\Lambda_{\mathfrak{m}}$ -irreducible.*

**Proof.** Let  $\mathfrak{m} = \mathfrak{m}_0 \oplus \mathfrak{m}_1 \oplus \dots \oplus \mathfrak{m}_r$  be the decomposition of  $\mathfrak{m}$  which satisfies (2.3). By Theorem 2.3, we see that each  $\mathfrak{m}_i$  is an invariant subspace by the holonomy algebra of the Riemannian connection (cf. see [3] Chapter X §4). Therefore the above decomposition has the only one factor. Since  $\Lambda_{\mathfrak{m}} \neq 0$ ,  $\mathfrak{m} \neq \mathfrak{m}_0$  and thus  $\mathfrak{m}$  is  $\Lambda_{\mathfrak{m}}$ -irreducible.  $\square$

### 3. Proof of the Main Theorem

We first recall a theorem of K. Tojo ([6]). Let  $M = K/H$  be a naturally

reductive homogeneous space with  $Ad(H)$ -invariant decomposition  $\mathfrak{k} = \mathfrak{h} \oplus \mathfrak{m}$ . According to [6], we put  $\varphi_x = \Lambda_{\mathfrak{m}}(x)$  for simplicity. Since  $\varphi_x$  is a skew symmetric endomorphism on  $(\mathfrak{m}, \langle \cdot, \cdot \rangle)$ ,  $e^{\varphi_x}$  is defined as a linear isometry on  $(\mathfrak{m}, \langle \cdot, \cdot \rangle)$ . Then K. Tojo showed the following (Theorem 3.2 in [6]).

**Theorem 3.1.** *Let  $V$  be a subspace of  $\mathfrak{m}$  (which is canonically identified with  $T_0 M$ ). Then there exists a totally geodesic submanifold of  $M$  through  $o$  whose tangent space at  $o$  is  $V$  if and only if the following holds:*

$$R(e^{\varphi_x}(V), e^{\varphi_x}(V))e^{\varphi_x}(V) \subset e^{\varphi_x}(V) \quad \text{for any } x \in V,$$

where  $R$  denotes the Riemannian curvature tensor of  $M$ .

The above theorem is considered as a generalization of the Lie triple system in Riemannian symmetric spaces due to E. Cartan.

Now we shall prove Main Theorem. Let  $M$  be as in Main Theorem. If  $\Lambda_{\mathfrak{m}} = 0$ , then  $M$  is a simply connected irreducible Riemannian symmetric space. In this case, our theorem has been proved by B.Y. Chen and T. Nagano [1]. Therefore we assume that  $\Lambda_{\mathfrak{m}} \neq 0$ . By Corollary 2.6, it follows that  $\mathfrak{m}$  is  $\Lambda_{\mathfrak{m}}$ -irreducible. Let  $S$  be a totally geodesic hypersurface of  $M$ . Since  $M$  is a homogeneous Riemannian manifold, we may assume that  $S$  is through  $o$ . Let  $V$  be a hyperplane (i.e., a subspace with codimension 1) of  $\mathfrak{m}$  which is a tangent space of  $S$  at  $o$ . We denote by  $\xi$  the unit vector of  $\mathfrak{m}$  which is orthogonal to  $V$ . We set

$$V_1 = \{\varphi_{\xi}x \mid x \in \mathfrak{m}\} = \{\varphi_{\xi}x \mid x \in V\}.$$

Then  $V_1$  is a subspace of  $V$ . In fact for any  $x \in \mathfrak{m}$ ,  $\langle \varphi_{\xi}x, \xi \rangle = -\langle x, \varphi_{\xi}\xi \rangle = 0$ . Since  $\mathfrak{m}$  is  $\Lambda_{\mathfrak{m}}$ -irreducible,  $V_1 \neq 0$ . We set  $O_1 = R\xi \oplus V_1$ .

**Lemma 3.2.** *The following equations hold:*

- (1)  $\langle R(x, y)z, \xi \rangle = 0$ .
- (2)  $\langle \varphi_{\xi}x, y \rangle \langle R(z, \xi)\xi, w \rangle - \langle \varphi_{\xi}x, z \rangle \langle R(y, \xi)\xi, w \rangle = \langle R(y, z)w, \varphi_{\xi}x \rangle$

for  $x, y, z \in V, w \in \mathfrak{m}$ .

**Proof of Lemma 3.2.** Applying Theorem 3.1, we obtain

$$(3.1) \quad \langle R(e^{t\varphi_x}y, e^{t\varphi_x}z)e^{t\varphi_x}w, e^{t\varphi_x}\xi \rangle = 0$$

for  $x, y, z, w \in V, t \in \mathbb{R}$ .

Putting  $t = 0$  in (3.1), we obtain (1). Differentiating (3.1) with respect to  $t$  at  $t = 0$ ,

$$(3.2) \quad \begin{aligned} & \langle R(\varphi_x y, z)w, \xi \rangle + \langle R(y, \varphi_x z)w, \xi \rangle \\ & + \langle R(y, z)\varphi_x w, \xi \rangle + \langle R(y, z)w, \varphi_x \xi \rangle = 0. \end{aligned}$$

We put  $\varphi_x y = \langle \varphi_x y, \xi \rangle \xi + v$ , where  $v \in V$ . Then by the equation (1) in this lemma

$$\begin{aligned}\langle R(\varphi_x y, z) w, \xi \rangle &= \langle \varphi_x y, \xi \rangle \langle R(\xi, z) w, \xi \rangle + \langle R(v, z) w, \xi \rangle \\ &= \langle \varphi_x y, \xi \rangle \langle R(z, \xi) \xi, w \rangle.\end{aligned}$$

Similarly we have

$$\begin{aligned}\langle R(y, \varphi_x z) w, \xi \rangle &= -\langle \varphi_x z, y \rangle \langle R(y, \xi) \xi, w \rangle \\ \langle R(y, z) \varphi_x w, \xi \rangle &= \langle \varphi_x z, w \rangle \langle R(y, z) \xi, \xi \rangle = 0.\end{aligned}$$

Substituting them in (3.2), we obtain (2) for  $w \in V$ . If  $w = \xi$ , the both sides of (2) are equal to 0. Therefore the equation (2) holds for all  $w \in \mathfrak{m}$ .  $\square$

By Lemma 3.2 (2), it follows that

$$(3.3) \quad \langle v, y \rangle \langle R(z, \xi) \xi, w \rangle - \langle v, z \rangle \langle R(y, \xi) \xi, w \rangle = -\langle R(y, z) v, w \rangle$$

for  $v \in V_1$ ,  $y, z \in V$ ,  $w \in \mathfrak{m}$ .

For  $x \in \mathfrak{m}$ , we define a symmetric endomorphism  $R_x : \mathfrak{m} \rightarrow \mathfrak{m}$  by  $R_x y = R(y, x) x$ .

**Lemma 3.3.** *There exists a constant  $c$  such that  $R_\xi x = cx$  for any  $x \in V_1$ .*

Proof of Lemma 3.3. Let  $x$  be an arbitrary non-zero vector of  $V_1$  and  $y$  be a vector of  $V$  which is orthogonal to  $x$ . Putting  $v = z = w = x$  in (3.3), we have  $\langle R(x, \xi) \xi, y \rangle = 0$ . On the other hand, clearly  $\langle R(x, \xi) \xi, \xi \rangle = 0$ . This implies that  $V_1$  is a subspace of some eigenspace with respect to  $R_\xi$ . We may take its eigenvalue as the constant  $c$ .  $\square$

**Lemma 3.4.** *For any  $v \in O_1$ , the following relations hold:*

- (1)  $R(y, z)v = 0$  for any  $y, z \in v^\perp$ ,
- (2)  $R_v x = c\{\langle v, v \rangle x - \langle x, v \rangle v\}$  for  $x \in O_1$ ,
- (3)  $R_v x = \langle v, v \rangle R_\xi x$  for  $x \in O_1^\perp$ ,

where  $v^\perp$  and  $O_1^\perp$  denote the orthogonal complements in  $\mathfrak{m}$  of  $v$  and  $O_1$ , respectively and the constant  $c$  in (2) is given in Lemma 3.3.

Proof of Lemma 3.4. We consider the following three cases for  $v \in O_1$ :

Case 1.  $v = \xi$ ;

Case 2.  $v$  is a unit vector of  $V_1$ . In this case we denote  $e$  by such a  $v$ ;

Case 3.  $v$  is an arbitrary unit vector of  $O_1$ .

Case 1. By Lemma 3.2 (1),  $R(y, z)\xi = 0$  for any  $y, z \in V$ . By Lemma 3.3

$$R_\xi x = c\{x - \langle x, \xi \rangle \xi\} \quad \text{for } x \in O_1.$$

Therefore (1), (2), and (3) in Lemma 3.4 hold for this case.

Case 2. Let  $y, z$  be vectors of  $e^\perp \cap V$ . Putting  $v=e$  in (3.3), we have  $R(y, z)e=0$ . Moreover it holds that  $R(y, \xi)e=0$ . In fact, for  $w \in V$ ,

$$\langle R(y, \xi)e, w \rangle = \langle R(e, w)y, \xi \rangle = 0$$

and

$$\langle R(y, \xi)e, \xi \rangle = -\langle R(e, \xi)\xi, y \rangle = -\langle R_\xi e, y \rangle = -c\langle e, y \rangle = 0.$$

From these, we see that (1) holds. Applying (3.3) for  $v=z=e$  and  $y \in e^\perp \cap V$ , we obtain  $R_e y = R_\xi y$ . Hence (2) and (3) hold.

Case 3. It is easily seen that the following relations hold:

$$R(y, e)\xi = -c\langle y, \xi \rangle e$$

$$R(y, \xi)e = -c\langle y, e \rangle \xi$$

for a unit vector  $e \in V_1$  and any  $y \in \mathfrak{m}$ .

We put  $v = \cos \theta e + \sin \theta \xi$  for some unit vector  $e \in V_1$  and some  $\theta \in \mathbb{R}$ . For  $y, z \in e^\perp \cap V$ , we have

$$R(y, z)v = \cos \theta R(y, z)e + \sin \theta R(y, z)\xi = 0,$$

$$\begin{aligned} R(y, -\sin \theta e + \cos \theta \xi)v \\ = -\sin \theta \cos \theta R(y, e)e - \sin^2 \theta R(y, e)\xi \\ + \cos^2 \theta R(y, \xi)e + \sin \theta \cos \theta R(y, \xi)\xi \\ = \sin \theta \cos \theta \{R_\xi y - R_e y\} = 0. \end{aligned}$$

Hence in this case (1) holds.

For  $x \in \mathfrak{m}$ , we have

$$\begin{aligned} (3.4) \quad R_v x &= \cos^2 \theta R_e x + \sin^2 \theta R_\xi x + \sin \theta \cos \theta \{R(x, e)\xi + R(x, \xi)e\} \\ &= \cos^2 \theta R_e x + \sin^2 \theta R_\xi x - c \sin \theta \cos \theta \{\langle x, \xi \rangle e + \langle x, e \rangle \xi\}. \end{aligned}$$

For  $x \in O_1$ , (3.4) implies

$$\begin{aligned} R_v x &= c \cos^2 \theta \{x - \langle x, e \rangle e\} + c \sin^2 \theta \{x - \langle x, \xi \rangle \xi\} \\ &\quad - c \sin \theta \cos \theta \{\langle x, \xi \rangle e + \langle x, e \rangle \xi\} \\ &= c \{x - \langle x, v \rangle v\}. \end{aligned}$$

For  $x \in O_1^\perp$ , (3.4) implies  $R_v x = R_\xi x$ . □

**Lemma 3.5.** *The following identity holds:*

$$\mathfrak{S}_{x,y,z} \{ \varphi_x(R(y,z)w) - R(\varphi_x y, z)w - R(y, \varphi_x z)w - R(y, z)\varphi_x w \} = 0$$

for  $x, y, z, w \in \mathfrak{m}$ .

Here the symbol  $\mathfrak{S}$  denotes the cyclic sum with respect to the indicated variables.

**Proof of Lemma 3.5.** It is known that the covariant derivative  $\nabla R$  of  $R$  is given as follows

$$\begin{aligned} (\nabla_x R)(y, z)w &= (\varphi_x \cdot R)(y, z)w \\ &= \varphi_x(R(y, z)w) - R(\varphi_x y, z)w - R(y, \varphi_x z)w - R(y, z)\varphi_x w. \end{aligned}$$

By this and Bianchi's 2nd identity of  $\nabla R$ , we have the identity in this lemma.  $\square$

We consider the symmetric endomorphism  $R_\xi: \mathfrak{m} \rightarrow \mathfrak{m}$ . Evidently we have  $R_\xi(V) \subset V$ . Then  $V$  is decomposed into the eigenspaces of  $R_\xi$ :

$$V = \mathfrak{p}_1 \oplus \cdots \oplus \mathfrak{p}_l,$$

where each  $\mathfrak{p}_i$  ( $i=1, \dots, l$ ) is the eigenspace of  $R_\xi$  with eigenvalue  $\lambda_i$ . Here we set  $\lambda_1=c$ , where the constant  $c$  has been given in Lemma 3.3. By Lemma 3.3, it follows that  $V_1 \subset \mathfrak{p}_1$ .

**Lemma 3.6.** (1) For  $x, y \in \mathfrak{p}_1$ ,  $\varphi_x y \in R_\xi \mathfrak{p}_1 \oplus \mathfrak{p}_1$ .  
 (2) For  $x \in \mathfrak{p}_i$ ,  $y \in \mathfrak{p}_j$  ( $j \neq 1$ ),  $\varphi_x y$  is contained in the eigenspace of  $R_\xi$  with eigenvalue  $\frac{\lambda_i + \lambda_j}{2}$ .

**Proof of Lemma 3.6.** By Lemma 3.5, we have for  $x \in \mathfrak{p}_i$ ,  $y \in \mathfrak{p}_j$

$$\begin{aligned} 0 &= \varphi_\xi(R(x, y)\xi) - R(\varphi_\xi x, y)\xi - R(x, \varphi_\xi y)\xi - R(x, y)\varphi_\xi \xi \\ &\quad + \varphi_x(R(y, \xi)\xi) - R(\varphi_x y, \xi)\xi - R(y, \varphi_x \xi)\xi - R(y, \xi)\varphi_x \xi \\ &\quad + \varphi_y(R(\xi, x)\xi) - R(\varphi_y \xi, x)\xi - R(\xi, \varphi_y x)\xi - R(\xi, x)\varphi_y \xi \\ &= \lambda_j \varphi_x y - 2R_\xi(\varphi_x y) - R(y, \xi)\varphi_x \xi - \lambda_i \varphi_y x + R(x, \xi)\varphi_y \xi \\ &= (\lambda_i + \lambda_j) \varphi_x y - 2R_\xi(\varphi_x y) + 2c \langle \varphi_x \xi, y \rangle \xi. \end{aligned}$$

Hence

$$(3.5) \quad 2R_\xi(\varphi_x y) = (\lambda_i + \lambda_j) \varphi_x y + 2c \langle \varphi_x \xi, y \rangle \xi.$$

If  $i=j=1$ , then (3.5) implies  $R_\xi(\varphi_x y) = c\{\varphi_x y - \langle \varphi_x y, \xi \rangle \xi\}$ . Therefore (1) in this lemma holds. If  $j \neq 1$ , (3.5) implies  $R_\xi(\varphi_x y) = \frac{\lambda_i + \lambda_j}{2} \varphi_x y$ . Therefore (2) in this lemma

holds.  $\square$

**Lemma 3.7.** *If  $x \in \mathfrak{p}_i$ ,  $y \in \mathfrak{p}_j$  ( $i \neq j$ ), then we have  $\varphi_x y = 0$ .*

**Proof of Lemma 3.7.** We assume that  $j \neq 1$  and that  $\varphi_x y \neq 0$ . We set  $\varphi_x y = z$ . Then by Lemma 3.6 (2),  $z$  is an eigenvector of  $R_\xi$  with eigenvalue  $\frac{\lambda_i + \lambda_j}{2}$ . Since  $0 \neq \langle \varphi_x y, z \rangle = -\langle y, \varphi_x z \rangle$ ,  $y$  and  $\varphi_x z$  are eigenvectors of  $R_\xi$  with same eigenvalue. Therefore we have  $\lambda_j = \frac{1}{2}(\frac{\lambda_i + \lambda_j}{2} + \lambda_i)$  and hence  $\lambda_i = \lambda_j$ , that is,  $i = j$ . It is contrary to our assumption  $i \neq j$ . Therefore we have  $\varphi_x y = 0$ .  $\square$

Since  $V_1 \subset \mathfrak{p}_1$ , together with Lemmas 3.6 and 3.7, we see that  $\mathbf{R}\xi \oplus \mathfrak{p}_1, \mathfrak{p}_2, \dots, \mathfrak{p}_l$  are  $\Lambda_m$ -invariant subspaces. By  $\Lambda_m$ -irreducibility, we have  $\mathfrak{m} = \mathbf{R}\xi \oplus V_1$ . By this and Lemma 3.4, it holds that

$$R(v, x)y = c\{\langle x, y \rangle v - \langle v, y \rangle x\} \quad \text{for } v \in O_1 = \mathbf{R}\xi \oplus V_1 \quad \text{and } x, y \in \mathfrak{m}.$$

We define a tensor  $R_0$  of type (1,3) by

$$R_0(u, v)w = \langle v, w \rangle u - \langle u, w \rangle v$$

and define a subspace  $\mathfrak{n}$  of  $\mathfrak{m}$  by

$$\mathfrak{n} = \{x \in \mathfrak{m} \mid i(x)(R - cR_0) = 0\}.$$

The preceding result means that  $O_1 \subset \mathfrak{n}$ . Now we note that the curvature tensor  $R$  is given as follows (cf [3] p.202):

$$\begin{aligned} R(x, y)z &= -[[x, y]_{\mathfrak{h}}, z] \\ &\quad + \frac{1}{4}[x, [y, z]_{\mathfrak{m}}]_{\mathfrak{m}} - \frac{1}{4}[y, [x, z]_{\mathfrak{m}}]_{\mathfrak{m}} - \frac{1}{2}[[x, y]_{\mathfrak{m}}, z]_{\mathfrak{m}} \\ &= -[[x, y]_{\mathfrak{h}}, z] + \varphi_x \varphi_y z - \varphi_y \varphi_x z - \varphi_{(\varphi_x y - \varphi_y x)} z \end{aligned}$$

for  $x, y, z \in \mathfrak{m}$ .

Since  $R$  and  $R_0$  are invariant by the action of  $\mathfrak{h}$ , the subspace  $\mathfrak{n}$  is invariant by the action of  $\mathfrak{h}$ . In particular we see that  $[[y, z]_{\mathfrak{h}}, v] \in \mathfrak{n}$  for  $v \in \mathfrak{n}$  and  $y, z \in \mathfrak{m}$ .

We first assume that  $c \neq 0$ . For an arbitrary vector  $x \in V$ , we have

$$R(x, \xi)\xi = -[[x, \xi]_{\mathfrak{h}}, \xi] - \varphi_{\varphi_x \xi} \xi.$$

Hence  $\xi$  and  $\varphi_{\varphi_x \xi} \xi$  are contained in  $\mathfrak{n}$ . By the preceding remark, it follows that  $[[x, \xi]_{\mathfrak{h}}, \xi] \in \mathfrak{n}$ . Hence  $R(x, \xi)\xi \in \mathfrak{n}$ . On the other hand, since  $V = \mathfrak{p}_1$ ,  $R(x, \xi)\xi = cx$ . Since  $c \neq 0$ , we have  $x \in \mathfrak{n}$ . Therefore we see that  $\mathfrak{n} = \mathfrak{m}$ , that is,  $R$  has constant sectional curvature  $c$ .

We secondly assume that  $c = 0$ . We define subspaces  $V_i$  ( $i = 0, 1, 2, \dots$ ) inductively

as follows. Set  $V_0 = R\xi$ . We define  $V_{i+1}$  by a subspace linearly spanned by  $\varphi_x z$  for  $x \in \mathfrak{m}$ ,  $z \in V_i$ . We remark that  $V_1$  coincides with the subspace defined at the beginning in this section.

**Lemma 3.8.** *For each  $i$ ,  $V_i \subset \mathfrak{n} = \{x \in \mathfrak{m} \mid i(x)R = 0\}$ .*

**Proof of Lemma 3.8.** We shall prove our assertion by the induction with respect to  $i$ . It is already shown that  $V_0 \subset \mathfrak{n}$  and  $V_1 \subset \mathfrak{n}$ . Suppose that our assertion holds for  $0, 1, \dots, i$  ( $i \geq 1$ ). Then we shall prove that  $V_{i+1} \subset \mathfrak{n}$ , that is,  $\varphi_x z \in \mathfrak{n}$  for  $x \in \mathfrak{m}$ ,  $z \in V_i$ . We consider the following three cases.

Case 1.  $x \in V_j$ ,  $0 \leq j \leq i-1$ ;

Case 2.  $x \in V_i$ ;

Case 3.  $x \in (V_0 + V_1 + \dots + V_i)^\perp$ .

Case 1. Since  $\varphi_x z = -\varphi_z x \in V_{j+1}$  and  $j+1 \leq i$ ,  $\varphi_x z \in \mathfrak{n}$ .

Case 2. By Lemma 3.5, we have for  $u, v \in \mathfrak{m}$

$$\begin{aligned} 0 &= \varphi_x(R(z, u)v) - R(\varphi_x z, u)v - R(z, \varphi_x u)v - R(z, u)\varphi_x v \\ &\quad + \varphi_z(R(u, x)v) - R(\varphi_z u, x)v - R(u, \varphi_z x)v - R(u, x)\varphi_z v \\ &\quad + \varphi_u(R(x, z)v) - R(\varphi_u x, z)v - R(x, \varphi_u z)v - R(x, z)\varphi_u v \\ &= -2R(\varphi_x z, u)v. \end{aligned}$$

Therefore we have  $\varphi_x z \in \mathfrak{n}$ .

Case 3. It is sufficient to prove our assertion when  $z = \varphi_u v$  for  $u \in \mathfrak{m}$ ,  $v \in V_{i-1}$ . We first remark that  $\varphi_x v = 0$ . In fact, for any  $w \in \mathfrak{m}$ ,  $\langle \varphi_x v, w \rangle = -\langle \varphi_w v, x \rangle$  and since  $\varphi_w v \in V_i$  and  $x \in (V_0 + V_1 + \dots + V_i)^\perp$ , we have  $\langle \varphi_x v, w \rangle = 0$ . It follows that

$$\begin{aligned} R(x, u)v &= -[[x, u]_{\mathfrak{h}}, v] + \varphi_x \varphi_u v - \varphi_u \varphi_x v - 2\varphi_{\varphi_x u} v \\ &= -[[x, u]_{\mathfrak{h}}, v] + \varphi_x z - 2\varphi_{\varphi_x u} v. \end{aligned}$$

On other hand,  $R(x, u)v = -R(u, v)x - R(v, x)u = 0$  by the assumption of induction. Then we have  $\varphi_x z = [[x, u]_{\mathfrak{h}}, v] + 2\varphi_{\varphi_x u} v$ . Since the right hand side is contained in  $\mathfrak{n}$ , so is  $\varphi_x z$ .  $\square$

We set  $O_i = V_0 + V_1 + \dots + V_i$ . Evidently we have  $O_0 \subseteq O_1 \subseteq \dots \subseteq O_i \subseteq O_{i+1} \subseteq \dots$ . Therefore there exists an integer  $i$  such that  $O_i = O_{i+1}$ . Then  $O_i$  is an invariant subspace with respect to  $\Lambda_{\mathfrak{m}}$ . Since  $O_i \neq 0$ , we have  $O_i = \mathfrak{m}$ . By Lemma 3.8, it follows that  $\mathfrak{n} = \mathfrak{m}$ , that is, the curvature tensor  $R$  vanishes. Thus our theorem has been completely proved.

## References

- [1] B.Y. Chen and T. Nagano: *Totally geodesic submanifolds of symmetric spaces* II, Duke Math. J. **45** (1978), 405–425.
- [2] J.E. D'Atri and W. Ziller: *Naturally reductive metrics and Einstein metrics on compact Lie groups*, Mem. Amer. Math. Soc. **215** (1979).
- [3] S. Kobayashi and K. Nomizu: *Foundations of differential geometry* II, Interscience publishers, 1969.
- [4] O. Kowalski and L. Vanhecke: *Four-dimensional naturally reductive homogeneous spaces*, Rend. Sem. Mat. Univ. Politec. Torino (1983), 223–232.
- [5] O. Kowalski and L. Vanhecke: *Classification of five-dimensional naturally reductive spaces*, Math. Proc. Camb. Phil. Soc. **97** (1985), 445–463.
- [6] K. Tojo: *Totally geodesic submanifolds of naturally reductive homogeneous spaces* (to appear).
- [7] K. Tojo: *Totally geodesic hypersurfaces of normal homogeneous spaces* (to appear).

Department of Mathematics  
Ochanomizu University  
Tokyo 112, Japan

