



Title	The warping degree of a link diagram
Author(s)	Shimizu, Ayaka
Citation	Osaka Journal of Mathematics. 2011, 48(1), p. 209-231
Version Type	VoR
URL	https://doi.org/10.18910/11237
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

THE WARPING DEGREE OF A LINK DIAGRAM

AYAKA SHIMIZU

(Received July 28, 2009, revised October 30, 2009)

Abstract

For an oriented link diagram D , the warping degree $d(D)$ is the smallest number of crossing changes which are needed to obtain a monotone diagram from D . We show that $d(D) + d(-D) + sr(D)$ is less than or equal to the crossing number of D , where $-D$ denotes the inverse of D and $sr(D)$ denotes the number of components which have at least one self-crossing. Moreover, we give a necessary and sufficient condition for the equality. We also consider the minimal $d(D) + d(-D) + sr(D)$ for all diagrams D . For the warping degree and linking warping degree, we show some relations to the linking number, unknotting number, and the splitting number.

1. Introduction

The warping degree and a monotone diagram is defined by Kawauchi for an oriented diagram of a knot, a link [7] or a spatial graph [8]. The warping degree represents such a complexity of a diagram, and depends on the orientation of the diagram. For an oriented link diagram D , we say that D is monotone if we meet every crossing point as an over-crossing first when we travel along all components of the oriented diagram with an order by starting from each base point. This notion is earlier used by Hoste [5] and by Lickorish–Millett [9] in computing polynomial invariants of knots and links. The warping degree $d(D)$ of an oriented link diagram D is the smallest number of crossing changes which are needed to obtain a monotone diagram from D in the usual way. We give the precise definitions of the warping degree and a monotone diagram in Section 2. Let $-D$ be the diagram D with orientations reversed for all components, and we call $-D$ the inverse of D . Let $c(D)$ be the crossing number of D . We have the following theorem in [13] which is for a knot diagram:

Theorem 1.1 ([13]). *Let D be an oriented knot diagram which has at least one crossing point. Then we have*

$$d(D) + d(-D) + 1 \leq c(D).$$

Further, the equality holds if and only if D is an alternating diagram.




Fig. 1.

Let D be a diagram of an r -component link ($r \geq 1$). Let D^i be a diagram on a knot component L^i of L , and we call D^i a component of D . We define a property of a link diagram as follows:

DEFINITION 1.2. A link diagram D has *property C* if every component D^i of D is alternating, and the number of over-crossings of D^i is equal to the number of under-crossings of D^i in every subdiagram $D^i \cup D^j$ for each $i \neq j$.

Note that a diagram D has property C if D is an alternating diagram in the case that $r = 1$. We generalize Theorem 1.1 to a link diagram:

Theorem 1.3. *Let D be an oriented link diagram, and $sr(D)$ the number of components D^i such that D^i has at least one self-crossing. Then we have*

$$d(D) + d(-D) + sr(D) \leq c(D).$$

Further, the equality holds if and only if D has property C.

For example, the link diagram D in Fig. 1 has $d(D) + d(-D) + sr(D) = 3 + 3 + 2 = 8 = c(D)$. Let D be a diagram of a link. Let $u(D)$ be the unlinking number of D . As a lower bound for the value $d(D) + d(-D) + sr(D)$, we have the following inequality:

Theorem 1.4. *We have*

$$2u(D) + sr(D) \leq d(D) + d(-D) + sr(D).$$

The rest of this paper is organized as follows. In Section 2, we define the warping degree $d(D)$ of an oriented link diagram D . In Section 3, we define the linking warping degree $ld(D)$, and consider the value $d(D) + d(-D)$ to prove Theorem 1.3. In Section 4, we show relations of the linking warping degree and the linking number. In Section 5, we apply the warping degree to a link itself. In Section 6, we study relations to unknotting number and crossing number. In Section 7, we define the splitting number and consider relations between the warping degree and the splitting number. In




Fig. 2.

Section 8, we show methods for calculating the warping degree and the linking warping degree.

2. The warping degree of an oriented link diagram

Let L be an r -component link, and D a diagram of L . We take a sequence \mathbf{a} of base points a_i ($i = 1, 2, \dots, r$), where every component has just one base point except at crossing points. Then $D_{\mathbf{a}}$, the pair of D and \mathbf{a} , is represented by $D_{\mathbf{a}} = D_{a_1}^1 \cup D_{a_2}^2 \cup \dots \cup D_{a_r}^r$ with the order of \mathbf{a} . A self-crossing point p of $D_{a_i}^i$ is a *warping crossing point of $D_{a_i}^i$* if we meet the point first at the under-crossing when we go along the oriented diagram $D_{a_i}^i$ by starting from a_i ($i = 1, 2, \dots, r$). A crossing point p of $D_{a_i}^i$ and $D_{a_j}^j$ is a *warping crossing point between $D_{a_i}^i$ and $D_{a_j}^j$* if p is the under-crossing of $D_{a_i}^i$ ($1 \leq i < j \leq r$). A crossing point p of $D_{\mathbf{a}}$ is a *warping crossing point of $D_{\mathbf{a}}$* if p is a warping crossing point of $D_{a_i}^i$ or a warping crossing point between $D_{a_i}^i$ and $D_{a_j}^j$ [7].

For example in Fig. 2, p is a warping crossing point of $D_{a_1}^1$, and q is a warping crossing point between $D_{a_1}^1$ and $D_{a_2}^2$. We define the warping degree for an oriented link diagram [7]. The *warping degree of $D_{\mathbf{a}}$* , denoted by $d(D_{\mathbf{a}})$, is the number of warping crossing points of $D_{\mathbf{a}}$. The *warping degree of D* , denoted by $d(D)$, is the minimal warping degree $d(D_{\mathbf{a}})$ for all base point sequences \mathbf{a} of D . Ozawa showed that a non-trivial link which has a diagram D with $d(D) = 1$ is a split union of a twist knot or the Hopf link and r trivial knots ($r \geq 0$) [11]. Fung also showed that a non-trivial knot which has a diagram D with $d(D) = 1$ is a twist knot [14].

For an oriented link diagram and its base point sequence $D_{\mathbf{a}} = D_{a_1}^1 \cup D_{a_2}^2 \cup \dots \cup D_{a_r}^r$, we denote by $d(D_{a_i}^i)$ the number of warping crossing points of $D_{a_i}^i$. We denote by $d(D_{a_i}^i, D_{a_j}^j)$ the number of warping crossing points between $D_{a_i}^i$ and $D_{a_j}^j$. By definition, we have that

$$d(D_{\mathbf{a}}) = \sum_{i=1}^r d(D_{a_i}^i) + \sum_{i < j} d(D_{a_i}^i, D_{a_j}^j).$$




Fig. 3.

Thus, the set of the warping crossing points of $D_{\mathbf{a}}$ is divided into two types in the sense that the warping crossing point is self-crossing or not. The pair $D_{\mathbf{a}}$ is *monotone* if $d(D_{\mathbf{a}}) = 0$. For example, $D_{\mathbf{a}}$ depicted in Fig. 3 is monotone. Note that a monotone diagram is a diagram of a trivial link. Hence we have $u(D) \leq d(D)$, where $u(D)$ is the unlinking number of D ([10], [15]).

3. Proof of Theorem 1.3

In this section, we prove Theorem 1.3. We first define the linking warping degree, which is like a restricted warping degree and which has relations to the crossing number and the linking number (see also Section 4). The number of non-self warping crossing points does not depend on the orientation. We define the *linking warping degree* of $D_{\mathbf{a}}$, denoted by $ld(D_{\mathbf{a}})$, by the following formula:

$$ld(D_{\mathbf{a}}) = \sum_{i < j} d(D_{a_i}^i, D_{a_j}^j) = d(D_{\mathbf{a}}) - \sum_{i=1}^r d(D_{a_i}^i),$$

where $D_{a_i}^i, D_{a_j}^j$ are components of $D_{\mathbf{a}}$ ($1 \leq i < j \leq r$). The *linking warping degree* of D , denoted by $ld(D)$, is the minimal $ld(D_{\mathbf{a}})$ for all base point sequences \mathbf{a} . It does not depend on any choices of orientations of components. For example, the diagram D in Fig. 4 has $ld(D) = 2$. A pair $D_{\mathbf{a}}$ is *stacked* if $ld(D_{\mathbf{a}}) = 0$. A diagram D is *stacked* if $ld(D) = 0$. For example, the diagram E in Fig. 4 is a stacked diagram. We remark that a similar notion is mentioned in [5]. Note that a monotone diagram is a stacked diagram. A link L is *completely splittable* if L has a diagram D without non-self crossings. Notice that a completely splittable link has some stacked diagrams.

The *linking crossing number* of D , denoted by $lc(D)$, is the number of non-self crossing points of D . Remark that $lc(D)$ is always even. For an unordered diagram D , we assume that D^i and $D^i \cup D^j$ denote subdiagrams of D with an order. We have the following relation of linking warping degree and linking crossing number.




Fig. 4.

Lemma 3.1. *We have*

$$ld(D) \leq \frac{lc(D)}{2}.$$

Further, the equality holds if and only if the number of over-crossings of D^i is equal to the number of under-crossings of D^i in every subdiagram $D^i \cup D^j$ for every $i \neq j$.

Proof. Let \mathbf{a} be a base point sequence of D , and $\tilde{\mathbf{a}}$ the base point sequence \mathbf{a} with the order reversed. We call $\tilde{\mathbf{a}}$ the reverse of \mathbf{a} . Since we have that $ld(D_{\mathbf{a}}) + ld(D_{\tilde{\mathbf{a}}}) = lc(D)$, we have the inequality $ld(D) \leq lc(D)/2$. Let D be a link diagram such that the number of over-crossings of D^i is equal to the number of under-crossings of D^i in every subdiagram $D^i \cup D^j$ for each $i \neq j$. Then we have $ld(D_{\mathbf{a}}) = lc(D)/2$ for every base point sequence \mathbf{a} . Hence we have $ld(D) = lc(D)/2$. On the other hand, we consider the case the equality $2ld(D) = lc(D)$ holds. For an arbitrary base point sequence \mathbf{a} of D and its reverse $\tilde{\mathbf{a}}$, we have

$$ld(D_{\mathbf{a}}) \geq ld(D) = lc(D) - ld(D) \geq lc(D) - ld(D_{\tilde{\mathbf{a}}}) = ld(D_{\tilde{\mathbf{a}}}) \geq ld(D).$$

Then we have $lc(D) - ld(D_{\mathbf{a}}) = ld(D)$. Hence we have $ld(D_{\mathbf{a}}) = ld(D)$ for every base point sequence \mathbf{a} . Let $\mathbf{a}' = (a_1, a_2, \dots, a_{k-1}, a_{k+1}, a_k, a_{k+2}, \dots, a_r)$ be the base point sequence which is obtained from $\mathbf{a} = (a_1, a_2, \dots, a_k, a_{k+1}, \dots, a_r)$ by exchanging a_k and a_{k+1} ($k = 1, 2, \dots, r-1$). Then, the number of over-crossings of D^k is equal to the number of under-crossings of D^k in the subdiagram $D^k \cup D^{k+1}$ of $D_{\mathbf{a}}$ because we have $ld(D_{\mathbf{a}}) = ld(D_{\mathbf{a}'})$. This completes the proof. \square

We next consider the value $d(D) + d(-D)$ for an oriented link diagram D and the inverse $-D$. We have the following proposition:

Proposition 3.2. *Let D be an oriented link diagram. The value $d(D) + d(-D)$ does not depend on the orientation of D .*

Proof. Let D' be D with the same order and another orientation. Since we have $d(D^{i'}) = d(D^i)$ or $d(D^{i'}) = d(-D^i)$, we have $d(D^{i'}) + d(-D^{i'}) = d(D^i) + d(-D^i)$ for

each D^i and $D^{i'}$. Then we have

$$\begin{aligned}
 d(D') + d(-D') &= \sum_{i=1}^r d(D^{i'}) + ld(D') + \sum_{i=1}^r d(-D^{i'}) + ld(-D') \\
 &= \sum_{i=1}^r \{d(D^{i'}) + d(-D^{i'})\} + 2 ld(D') \\
 &= \sum_{i=1}^r \{d(D^i) + d(-D^i)\} + 2 ld(D) \\
 &= \sum_{i=1}^r d(D^i) + ld(D) + \sum_{i=1}^r d(-D^i) + ld(-D) \\
 &= d(D) + d(-D). \quad \square
 \end{aligned}$$

A link diagram is a *self-crossing diagram* if every component of D has at least one self-crossing. In other words, a diagram D of an r -component link L is a self-crossing diagram if $sr(D) = r$. We have the following lemma:

Lemma 3.3. *Let D be a self-crossing diagram of an r -component link. Then we have*

$$d(D) + d(-D) + r \leq c(D).$$

Further, the equality holds if and only if D has property C.

Proof. We have

$$\begin{aligned}
 d(D) + d(-D) + r &= \sum_{i=1}^r d(D^i) + ld(D) + \sum_{i=1}^r d(-D^i) + ld(-D) + r \\
 &= \sum_{i=1}^r \{d(D^i) + d(-D^i) + 1\} + 2 ld(D) \\
 &\leq \sum_{i=1}^r c(D^i) + 2 ld(D) \\
 &\leq \sum_{i=1}^r c(D^i) + lc(D) \\
 &= c(D),
 \end{aligned}$$

where the first inequality is obtained by Theorem 1.1, and the second inequality is obtained by Lemma 3.1. Hence we have the inequality. The equality holds if and only if D has property C which is obtained by Theorem 1.1 and Lemma 3.1. \square

We give an example of Lemma 3.3.




Fig. 5.




Fig. 6.

EXAMPLE 3.4. In Fig. 5, there are three diagrams with 12 crossings. The diagram D is a diagram such that any component is alternating and has 3 over-non-self crossings and 3 under-non-self crossings. Then we have $d(D) + d(-D) + r = 12 = c(D)$. The diagram D' is a diagram which has a non-alternating component diagram. Then we have $d(D') + d(-D') + r = 10 < c(D')$. The diagram D'' is a diagram such that a component has 2 over-non-self crossings and 4 under-non-self crossings. Then we have $d(D'') + d(-D'') + r = 10 < c(D'')$.

Lemma 3.3 is only for self-crossing link diagrams. We prove Theorem 1.3 which is for every link diagram.

Proof of Theorem 1.3. For every component D^i such that D^i has no self-crossings, we apply a Reidemeister move of type I as shown in Fig. 6. Then we obtain the diagram $D^{i'}$ from D^i , and $D^{i'}$ satisfies $d(D^{i'}) = d(-D^{i'}) = 0 = d(D^i) = d(-D^i)$ and $c(D^{i'}) = 1 = c(D^i) + 1$. For example the base points a_i, b_i in Fig. 6 satisfy $d(D^i_{a_i}) = d(D^i) = 0$, $d(-D^i_{b_i}) = d(-D^i) = 0$. We remark that every D^i and $D^{i'}$ are alternating. We denote by D' the diagram obtained from D by this procedure. Since every component has at least one self-crossing, we apply Lemma 3.3 to D' . Then we have

$$d(D') + d(-D') + r \leq c(D').$$

And we obtain

$$d(D) + d(-D) + r \leq c(D) + (r - sr(D)).$$

Hence we have

$$d(D) + d(-D) + sr(D) \leq c(D).$$

The equality holds if and only if D has property C . □

4. The linking warping degree and linking number

In this section, we consider the relation of the linking warping degree and the linking number. For a crossing point p of an oriented diagram, $\varepsilon(p)$ denotes the sign of p , namely $\varepsilon(p) = +1$ if p is a positive crossing, and $\varepsilon(p) = -1$ if p is a negative crossing. For an oriented subdiagram $D^i \cup D^j$, the *linking number* of D^i with D^j is defined to be

$$\text{Link}(D^i, D^j) = \frac{1}{2} \sum_{p \in D^i \cap D^j} \varepsilon(p).$$

The linking number of D^i with D^j is independent of the diagram (cf. [3], [7]). We have a relation of the linking warping degree and the linking number of a link diagram in the following proposition:

Proposition 4.1. *For a link diagram D , we have the following (i) and (ii).*

(i): *We have*

$$\sum_{i < j} |\text{Link}(D^i, D^j)| \leq ld(D).$$

Further, the equality holds if and only if non-self under-crossings of D^i in $D^i \cup D^j$ are all positive or all negative with an orientation for every subdiagram $D^i \cup D^j$ ($i < j$).

(ii): *We have*

$$(1) \quad \sum_{i < j} |\text{Link}(D^i, D^j)| \equiv ld(D) \pmod{2}.$$

Proof. (i): For a subdiagram $D^i \cup D^j$ ($i < j$) with $d(D^i, D^j) = m$, we show that

$$|\text{Link}(D^i, D^j)| \leq d(D^i, D^j).$$

Let p_1, p_2, \dots, p_m be the warping crossing points between D^i and D^j , and $\varepsilon(p_1), \varepsilon(p_2), \dots, \varepsilon(p_m)$ the signs of them. Since a stacked diagram is a diagram of a completely splittable link, we have

$$(2) \quad \text{Link}(D^i, D^j) - (\varepsilon(p_1) + \varepsilon(p_2) + \dots + \varepsilon(p_m)) = 0$$

by applying crossing changes at p_1, p_2, \dots, p_m for $D^i \cup D^j$. Then we have

$$|\text{Link}(D^i, D^j)| = |\varepsilon(p_1) + \varepsilon(p_2) + \dots + \varepsilon(p_m)| \leq m = d(D^i, D^j).$$




Fig. 7.

Hence we obtain

$$\sum_{i < j} |\text{Link}(D^i, D^j)| \leq ld(D).$$

The equality holds if and only if non-self under-crossings of D^i in $D^i \cup D^j$ are all positive or all negative with an orientation for every subdiagram $D^i \cup D^j$ ($i < j$).

(ii): By the above equality (2), we observe that $\text{Link}(D^i, D^j) = \varepsilon(p_1) + \varepsilon(p_2) + \cdots + \varepsilon(p_m) = \varepsilon(q_1) + \varepsilon(q_2) + \cdots + \varepsilon(q_n)$, where p_k (resp. q_k) is a non-self under-crossing (resp. over-crossing) of D^i in $D^i \cup D^j$, $ld(D^i \cup D^j) = m$ and $lc(D^i \cup D^j) = m + n$. A similar fact is also mentioned in [12]. We have

$$\begin{aligned} \text{Link}(D^i, D^j) &= \varepsilon(p_1) + \varepsilon(p_2) + \cdots + \varepsilon(p_m) \\ &\equiv m \pmod{2} \\ &= d(D^i, D^j). \end{aligned}$$

Hence we have the modular equality

$$\sum_{i < j} |\text{Link}(D^i, D^j)| \equiv ld(D) \pmod{2}. \quad \square$$

EXAMPLE 4.2. In Fig. 7, D has $(0, 2, 3)$, E has $(0, 2, 2)$, and F has $(4, 4, 4)$, where (l, m, n) of D denotes that $\sum_{i < j} |\text{Link}(D^i, D^j)| = l$, $ld(D) = m$, and $lc(D)/2 = n$.

The *total linking number* of an oriented link L is defined to be $\sum_{i < j} \text{Link}(D^i, D^j)$ with a diagram and an order. We have the following corollary:

Corollary 4.3. *We have*

$$\sum_{i < j} \text{Link}(D^i, D^j) = \sum_{k=1}^r \{\varepsilon(p_k) \mid p_k: \text{a non-self warping crossing point of } D_{\mathbf{a}}\},$$

where \mathbf{a} is a base point sequence of D .




Fig. 8.

Corollary 4.3 is useful in calculating the total linking number of a diagram. For example in Fig. 8, the diagram D with 4 components and 11 crossing points has $ld(D) = 4$. We have that the total linking number of D is 0 by summing the signs of only 4 crossing points.

5. To a link invariant

In this section, we consider the minimal $d(D) + d(-D)$ for minimal crossing diagrams D of L in the following formula:

$$e(L) = \min\{d(D) + d(-D) \mid D: \text{a diagram of } L \text{ with } c(D) = c(L)\},$$

where $c(L)$ denotes the crossing number of L . In the case where K is a non-trivial knot, we have

$$(3) \quad e(K) + 1 \leq c(K).$$

Further, the equality holds if and only if K is a prime alternating knot [13]. Note that the condition for the equality of (3) requires that D is a minimal crossing diagram in the definition of $e(L)$. We next define $c^*(L)$ and $e^*(L)$ as follows:

$$c^*(L) = \min\{c(D) \mid D: \text{a self-crossing diagram of } L\},$$

$$e^*(L) = \min\{d(D) + d(-D) \mid D: \text{a self-crossing diagram of } L \text{ with } c(D) = c^*(L)\}.$$

As a generalization of the above inequality (3), we have the following theorem:

Theorem 5.1. *For an r -component link L , we have*

$$e^*(L) + r \leq c^*(L).$$

Further, the equality holds if and only if every self-crossing diagram D of L with $c(D) = c^(L)$ has property C.*




Fig. 9.

Proof. Let D be a self-crossing diagram of L with $c(D) = c^*(D)$. We assume that D satisfies the equality $d(D) + d(-D) = e^*(L)$. Then we have

$$\begin{aligned}
 e^*(L) + r &= d(D) + d(-D) + r \\
 &= \sum_{i=1}^r d(D^i) + ld(D) + \sum_{i=1}^r d(-D^i) + ld(-D) + r \\
 &= \sum_{i=1}^r \{d(D^i) + d(-D^i) + 1\} + 2ld(D) \\
 &\leq \sum_{i=1}^r c(D^i) + 2ld(D) \\
 &\leq \sum_{i=1}^r c(D^i) + lc(D) \\
 &= c(D) = c^*(L),
 \end{aligned}$$

where the first inequality is obtained by Theorem 1.1, and the second inequality is obtained by Lemma 3.1. If D has a non-alternating component D^i , or D has a diagram $D^i \cup D^j$ such that the number of over-crossings of D^i is not equal to the number of under-crossings of D^i , then we have $e^*(L) + r < c^*(L)$. On the other hand, the equality holds if D has property C . \square

We have the following example:

EXAMPLE 5.2. For non-trivial prime alternating knots L^1, L^2, \dots, L^r ($r \geq 2$), we have a non-splittable link L by performing n_i -full twists for every L^i and L^{i+1} ($i = 1, 2, \dots, r$) with $L^{r+1} = L^1$ as shown in Fig. 9, where we assume that n_1 and n_r have the same sign. Note that we do not change the type of knot components L^i . Let D be a diagram of L with $c(D) = c(L)$. Then we notice that D is a self-crossing diagram with $c(D) = c^*(L)$. We also notice that D has property C because $lc(D^i \cup D^j) = 2|n_i|$ and $\text{Link}(D^i, D^j) = n_i$, and $lc(D^1 \cup D^r) = 2|n_1 + n_r|$ and $\text{Link}(D^1, D^r) = n_1 + n_r$ in

the case where $r = 2$. Hence we have $e^*(L) + r = c^*(L)$ in this case.

We have the following corollary:

Corollary 5.3. *Let L be an r -component link whose all components are non-trivial. Then we have*

$$e(L) + r \leq c(L).$$

Further, the equality holds if and only if every diagram D of L with $c(D) = c(L)$ has property C.

Proof. Since every diagram D of L is a self-crossing diagram, we have $e(L) = e^*(L)$ and $c(L) = c^*(L)$. \square

We also consider the minimal $d(D) + d(-D) + sr(D)$ and the minimal $sr(D)$ for diagrams D of L in the following formulas:

$$f(L) = \min\{d(D) + d(-D) + sr(D) \mid D: \text{a diagram of } L\},$$

$$sr(L) = \min\{sr(D) \mid D: \text{a diagram of } L\}.$$

Note that the value $f(L)$ and $sr(L)$ also do not depend on the orientation of L . Jin and Lee mentioned in [6] that every link has a diagram which restricts to a minimal crossing diagram for each component. Then we have the following proposition:

Proposition 5.4. *The value $sr(L)$ is equal to the number of non-trivial knot components of L .*

The following corollary is directly obtained from Theorem 1.3.

Corollary 5.5. *We have*

$$f(L) \leq c(L).$$

Proof. For a diagram D with $c(D) = c(L)$, we have

$$f(L) \leq d(D) + d(-D) + sr(D) \leq c(D) = c(L),$$

where the second inequality is obtained by Theorem 1.3. \square

We have the following question:

QUESTION 5.6. When does the equality $f(L) = c(L)$ hold?




Fig. 10.

EXAMPLE 5.7. In Fig. 10, there are two link diagrams D and E . We assume that D (resp. E) is a diagram of a link L (resp. M). We have $f(L) = c(L) = 5$ because we have $d(D) + d(-D) + sr(D) = 2 + 2 + 1$ and we know $d(D^i) \geq u(3_1) = 1$, $ld(D) \geq 1$, and $sr(D) \geq sr(L) = 1$, where D^i is any diagram of 3_1 . On the other hand, we have that $f(M) < c(M)$ because $f(M) \leq d(E) + d(-E) + sr(E) = 3 + 3 + 1 = 7 < 10 = c(M)$.

6. Relations of warping degree, unknotting number, and crossing number

In this section, we enumerate several relations of the warping degree, the unknotting number or unlinking number, and the crossing number. Let $|D|$ be D with orientation forgotten. We define the minimal warping degree of D for all orientations as follows:

$$d(|D|) := \min\{d(D) \mid D: |D| \text{ with an orientation}\}.$$

Note that the minimal $d(|D|)$ for all diagrams D of L is equal to the ascending number $a(L)$ [11]:

$$a(L) = \min\{d(|D|) \mid D: \text{a diagram of } L\}.$$

Let E be a knot diagram, and D a diagram of an r -component link. We review the relation of the unknotting number $u(E)$ (resp. the unlinking number $u(D)$) and the crossing number $c(E)$ (resp. $c(D)$) of E (resp. D). The following inequalities are well-known [10]:

$$(4) \quad u(E) \leq \frac{c(E) - 1}{2},$$

$$(5) \quad u(D) \leq \frac{c(D)}{2}.$$

Moreover, Taniyama mentioned the following conditions [15]: The necessary condition for the equality of (4) is that E is a reduced alternating diagram of some $(2, p)$ -torus knot, or E is a diagram with $c(E) = 1$. The necessary condition for the equality of (5) is that every D^i is a simple closed curve on \mathbb{S}^2 and every subdiagram $D^i \cup D^j$ is an alternating diagram.

Hanaki and Kanadome characterized the link diagrams D which satisfy $u(D) = (c(D) - 1)/2$ as follows [4]: Let $D = D^1 \cup D^2 \cup \cdots \cup D^r$ be a diagram of an r -component link. Then we have

$$u(D) = \frac{c(D) - 1}{2}$$

if and only if exactly one of D^1, D^2, \dots, D^r is a reduced alternating diagram of a $(2, p)$ -torus knot, the other components are simple closed curves on \mathbb{S}^2 , and the non-self crossings of the subdiagram $D^i \cup D^j$ are all positive, all negative, or empty for each $i \neq j$. In addition, they showed that any minimal crossing diagram D of a link L with $u(L) = (c(L) - 1)/2$ satisfies $u(D) = (c(D) - 1)/2$.

Abe and Higa study the knot diagrams D which satisfy

$$u(D) = \frac{c(D) - 2}{2}.$$

Let D be a knot diagram with $u(D) = (c(D) - 2)/2$. They showed in [1] that for any crossing point p of D , one of the components of D_p is a reduced alternating diagram of a $(2, p)$ -torus knot and the other component of D_p has no self-crossings, where D_p is the diagram obtained from D by smoothing at p . In addition, they showed that any minimal crossing diagram D of a knot K with $u(K) = (c(K) - 2)/2$ satisfies the above condition.

By adding to (4), we have the following corollary:

Corollary 6.1. *For a knot diagram E , we have*

$$u(E) \leq d(|E|) \leq \frac{c(E) - 1}{2}.$$

Further, if we have

$$u(E) = d(|E|) = \frac{c(E) - 1}{2},$$

then E is a reduced alternating diagram of some $(2, p)$ -torus knot, or E is a diagram with $c(E) = 1$.

By adding to (5), we have the following corollary.

Corollary 6.2. (i): *For an r -component link diagram D , we have*

$$u(D) \leq d(|D|) \leq \frac{c(D)}{2}.$$

(ii): We have

$$u(D) \leq d(|D|) = \frac{c(D)}{2}$$

if and only if every D^i is a simple closed curve on \mathbb{S}^2 and the number of over-crossings of D^i is equal to the number of under-crossings of D^i in every subdiagram $D^i \cup D^j$ for each $i \neq j$.

(iii): If we have

$$u(D) = d(|D|) = \frac{c(D)}{2},$$

then every D^i is a simple closed curve on \mathbb{S}^2 and for each pair i, j , the subdiagram $D^i \cup D^j$ is an alternating diagram.

Proof. (i): The equality $u(D) \leq d(|D|)$ holds because $u(D) \leq d(D)$ holds for every oriented diagram. We show that $d(|D|) \leq c(D)/2$. Let D be an oriented diagram which satisfies

$$d(D) = \sum_{i=1}^r d(D^i) + ld(D) = d(|D|).$$

Then D also satisfies

$$(6) \quad d(D^i) \leq \frac{c(D^i)}{2}$$

for every component D^i because of the orientation of D . By Lemma 3.1, we have

$$(7) \quad ld(D) \leq \frac{lc(D)}{2}.$$

Then we have

$$\sum_{i=1}^r d(D^i) + ld(D) \leq \sum_{i=1}^r \frac{c(D^i)}{2} + \frac{lc(D)}{2}$$

by (6) and (7). Hence we obtain the inequality

$$d(|D|) \leq \frac{c(D)}{2}.$$

(ii): Suppose that the equality $d(|D|) = c(D)/2$ holds. Then the equalities

$$(8) \quad d(D^i) = \frac{c(D^i)}{2}$$

and

$$(9) \quad ld(D) = \frac{lc(D)}{2}$$

hold by (6) and (7), where D has an orientation such that $d(D) = d(|D|)$. The equality (8) is equivalent to that $c(D^i) = 0$ for every D^i . We prove this by an indirect proof. We assume that $c(D^i) > 0$ for a component D^i . In this case, we have the inequality

$$(10) \quad d(D^i) + d(-D^i) + 1 \leq c(D^i)$$

by Theorem 1.1 since D^i has a self-crossing. We also have

$$(11) \quad d(D^i) = d(-D^i) = \frac{c(D^i)}{2}$$

because $d(D^i) \leq d(-D^i)$ and (8). By substituting (11) for (10), we have

$$c(D^i) + 1 \leq c(D^i).$$

This implies that the assumption $c(D^i) > 0$ is contradiction. Therefore every D^i is a simple closed curve. The inequality (9) is equivalent to that the number of over-crossings of D^i is equal to the number of under-crossings of D^i in every subdiagram $D^i \cup D^j$ for each $i \neq j$ by Lemma 3.1. On the other hand, suppose that every D^i is a simple closed curve, and the number of over-crossings of D^i is equal to the number of under-crossings of D^i in every subdiagram $D^i \cup D^j$ for each $i \neq j$, then we have

$$d(|D|) = ld(D) = \frac{lc(D)}{2} = \frac{c(D)}{2}.$$

(iii): This holds by Corollary 6.2 (i) and above Taniyama's condition. \square

Let K be a knot, and L an r -component link. Let $u(K)$ be the unknotting number of K , and $u(L)$ be the unlinking number of L . The following inequalities are also well-known [10]:

$$(12) \quad u(K) \leq \frac{c(K) - 1}{2},$$

$$(13) \quad u(L) \leq \frac{c(L)}{2}.$$

The following conditions are mentioned by Taniyama [15]: The necessary condition for the equality of (12) is that K is a $(2, p)$ -torus knot (p : odd, $\neq \pm 1$). The necessary condition for the equality of (13) is that L has a diagram D such that every D^i is a simple closed curve on \mathbb{S}^2 and every subdiagram $D^i \cup D^j$ is an alternating diagram.

By adding to (12) and (3), we have the following corollary:

Corollary 6.3. (i): *We have*

$$u(K) \leq \frac{e(K)}{2} \leq \frac{c(K) - 1}{2}.$$

(ii): *We have*

$$u(K) \leq \frac{e(K)}{2} = \frac{c(K) - 1}{2}$$

if and only if K is a prime alternating knot.

(iii): *If we have*

$$u(K) = \frac{e(K)}{2} = \frac{c(K) - 1}{2},$$

then K is a $(2, p)$ -torus knot (p : odd, $\neq \pm 1$).

By adding to (13), we have the following corollary:

Corollary 6.4. *For a diagram of an unoriented r -component link, we have*

$$u(L) \leq \frac{e(L)}{2} \leq \frac{c(L)}{2}.$$

Further, if the equality $u(L) = e(L)/2 = c(L)/2$ holds, then L has a diagram $D = D^1 \cup D^2 \cup \dots \cup D^r$ such that every D^i is a simple closed curve on \mathbb{S}^2 and for each pair i, j , the subdiagram $D^i \cup D^j$ is an alternating diagram.

Proof. We prove the inequality $u(L) \leq e(L)/2$. Let D be a minimal crossing diagram of L which satisfies $e(L) = d(D) + d(-D)$. Then we obtain

$$e(L) = d(D) + d(-D) \geq 2u(D) \geq 2u(L).$$

The condition for the equality is due to above Taniyama's condition. □

7. Splitting number

In this section, we define the splitting number and enumerate relations of the warping degree and the complete splitting number. The *splitting number* (resp. *complete*




Fig. 11.




Fig. 12.

splitting number) of D , denoted by $Split(D)$ (resp. $split(D)$), is the smallest number of crossing changes which are needed to obtain a diagram of a splittable (resp. completely splittable) link from D . The splitting number of a link which is the minimal $Split(D)$ for all diagrams D is defined by Adams [2]. The *linking splitting number* (resp. *linking complete splitting number*) of D , denoted by $lSplit(D)$ (resp. $lsplit(D)$), is the smallest number of non-self-crossing changes which are needed to obtain a diagram of a splittable (resp. completely splittable) link from D . We have the following propositions:

Proposition 7.1. (i): *We have*

$$split(D) \leq d(|D|).$$

(ii): *We have*

$$split(D) \leq lsplit(D) \leq ld(D) \leq \frac{lc(D)}{2} \leq \frac{c(D)}{2}.$$

We give examples of Proposition 7.1.

EXAMPLE 7.2. The diagram D in Fig. 11 has $split(D) = 2 < d(|D|) = 3$. The diagram E in Fig. 11 has $split(E) = d(|E|) = 3$.

EXAMPLE 7.3. The diagram D in Fig. 12 has $split(D) = 1 < lsplit(D) = 2$. The diagram E in Fig. 12 has $split(E) = lsplit(E) = 2$.




Fig. 13.

EXAMPLE 7.4. The diagram D in Fig. 13 has $lsplit(D) = 3 < ld(D) = 5$. The diagram E in Fig. 13 has $lsplit(E) = ld(E) = 5$.

We raise the following question:

QUESTION 7.5. When does the equality

$$split(D) = d(|D|),$$

$$split(D) = lsplit(D)$$

or

$$lsplit(D) = ld(D)$$

hold?

8. Calculation of warping degree

In this section, we show methods for calculating the warping degree and linking warping degree by using matrices. First, we give a method for calculating the warping degree $d(D)$ of an oriented knot diagram D . Let a be a base point of D . We can obtain the warping degree $d(D_a)$ of D_a by counting the warping crossing points easily. Let $[D_a]$ be a sequence of some “ o ” and “ u ”, which is obtained as follows. When we go along the oriented diagram D from a , we write down “ o ” (resp. “ u ”) if we reach a crossing point as an over-crossing (resp. under-crossing) in numerical order. We next perform normalization to $[D_a]$, by deleting the subsequence “ ou ” repeatedly, to obtain the normalized sequence $\lfloor D_a \rfloor$. Then we have

$$d(D) = d(D_a) - \frac{1}{2} \# \lfloor D_a \rfloor,$$

where $\# \lfloor D_a \rfloor$ denotes the number of entries in $\lfloor D_a \rfloor$. Thus, we obtain the warping degree $d(D)$ of D . In the following example, we find the warping degree of a knot diagram by using the above algorithm.




Fig. 14.




Fig. 15.

EXAMPLE 8.1. For the oriented knot diagram D and the base point a in Fig. 14, we have $d(D_a) = 4$ and $[D_a] = [ouuuouuuououououou]$. By normalizing $[D_a]$, we obtain $[D_a] = [uuoo]$. Hence we find the warping degree of D as follows:

$$d(D) = 4 - \frac{1}{2} \times 4 = 2.$$

For some types of knot diagram, this algorithm is useful in formulating the warping degree or looking into its properties. We enumerate the properties of an oriented diagram of a pretzel knot of odd type in the following example:

EXAMPLE 8.2. Let $D = P(\varepsilon_1 n_1, \varepsilon_2 n_2, \dots, \varepsilon_m n_m)$ be an oriented pretzel knot diagram of odd type ($\varepsilon_i \in +1, -1$, n_i, m : odd > 0), where the orientation is given as shown in Fig. 15. We take base points a, b in Fig. 15. Then we have

$$d(D_a) = d(-D_b) = \frac{c(D)}{2} + \sum_i \frac{(-1)^{i+1} \varepsilon_i}{2}$$

and

$$\#[D_a] = \#[-D_b].$$

Hence we have $d(D) = d(-D)$ in this case. In particular, if D is alternating i.e. $\varepsilon_1 =$




Fig. 16.

$\varepsilon_2 = \cdots = \varepsilon_m = \pm 1$, then we have that

$$d(D) = \frac{c(D)}{2} - \frac{1}{2}.$$

We next explore how to calculate the linking warping degree $ld(D)$ by using matrices. For a link diagram D and a base point sequence \mathbf{a} of D , we define an r -square matrix $M(D_{\mathbf{a}}) = (m_{ij})$ by the following rule:

- For $i \neq j$, m_{ij} is the number of crossings of D^i and D^j which are under-crossings of D^i .
- For $i = j$, $m_{ij} = d(D^i)$.

We show an example.

EXAMPLE 8.3. For $D_{\mathbf{a}}$ and $D_{\mathbf{b}}$ in Fig. 16, we have

$$M(D_{\mathbf{a}}) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 2 & 2 & 0 \end{pmatrix}, \quad M(D_{\mathbf{b}}) = \begin{pmatrix} 0 & 2 & 2 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

We note that $ld(D_{\mathbf{a}})$ is obtained by summing the upper triangular entries of $M(D_{\mathbf{a}})$, that is

$$ld(D_{\mathbf{a}}) = \sum_{i < j} m_{ij},$$

and we notice that

$$d(D_{\mathbf{a}}) = \sum_{i \leq j} m_{ij},$$

where m_{ij} is an entry of $M(D_{\mathbf{a}})$ ($i, j = 1, 2, \dots, r$). For the base point sequence $\mathbf{a}' = (a_1, a_2, \dots, a_{k+1}, a_k, \dots, a_r)$ which is obtained from a base point sequence \mathbf{a} by

exchanging a_k and a_{k+1} ($k = 1, 2, \dots, r-1$), the matrix $M(D_{\mathbf{a}'})$ is obtained as follows:

$$M(D_{\mathbf{a}'}) = P_k M(D_{\mathbf{a}}) P_k^{-1},$$

where

$$P_k = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & 0 & 1 & \\ & & 1 & 0 & \\ & & & & \ddots & \\ & & & & & 1 \end{pmatrix}; \quad m_{ij} = \begin{cases} 1 & \text{for } (i, j) = (k, k+1), (k+1, k) \\ & \text{and } (i, j) = (i, i) (i \neq k, k+1), \\ 0 & \text{otherwise.} \end{cases}$$

With respect to the linking warping degree, we have

$$ld(D_{\mathbf{a}'}) = ld(D_{\mathbf{a}}) - m_{k\ k+1} + m_{k+1\ k},$$

where $m_{k\ k+1}, m_{k+1\ k}$ are entries of $M(D_{\mathbf{a}})$. To enumerate the permutation of the order of $\mathbf{a} = (a_1, a_2, \dots, a_r)$, we consider a matrix $Q = P^{r-1} P^{r-2} \dots P^2 P^1$, where P^n denotes $P_n P_{n+1} \dots P_{k_n}$ ($n \leq k_n \leq r-1$) or the identity matrix E_r . Since Q depends on the choices of k_n ($n = 1, 2, \dots, r-1$), we also denote Q by $Q_{\mathbf{k}}$, where $\mathbf{k} = (k_1, k_2, \dots, k_{r-1})$ ($n \leq k_n \leq r$) and we regard $P^n = E_r$ in the case $k_n = r$. Hence we obtain the following formula:

$$ld(D) = \min_{\mathbf{k}} \left\{ \sum_{i < j} m_{ij} \mid m_{ij}: \text{an entry of } Q_{\mathbf{k}} M(D_{\mathbf{a}}) Q_{\mathbf{k}}^{-1} \right\}.$$

Thus, we obtain the warping degree of an oriented link diagram by summing the warping degrees $d(D^i)$ ($i = 1, 2, \dots, r$) and the linking warping degree $ld(D)$.

ACKNOWLEDGMENTS. The author would like to thank the referee for helpful remarks. She would also like to thank Ryo Hanaki for giving her valuable information about unknotting number and ascending number. She is grateful to Takefumi Nosaka for his helpful advices about the manner of writing. She is also grateful to Reiko Shinjo for pointing out Adams' paper. She is deeply grateful to the members of Friday Seminar on Knot Theory in Osaka City University, including Professor Taizo Kanenobu, Tetsuya Abe, In Dae Jong, and Kengo Kishimoto, who assisted her in uncountable comments and the tender encouragement. She especially wants to thank her advisor, Professor Akio Kawauchi, whose support and encouragement inspired her to greater efforts throughout this research project.

References

- [1] T. Abe, R. Hanaki and R. Higa: *The unknotting number and band-unknotting number of a knot*, to appear in Osaka J. Math.
- [2] C.C. Adams: *Splitting versus unlinking*, J. Knot Theory Ramifications **5** (1996), 295–299.
- [3] P.R. Cromwell: *Knots and Links*, Cambridge Univ. Press, Cambridge, 2004.
- [4] R. Hanaki and J. Kanadome: *On an inequality between unknotting number and crossing number of links*, to appear in J. Knot Theory Ramifications.
- [5] J. Hoste: *A polynomial invariant of knots and links*, Pacific J. Math. **124** (1986), 295–320.
- [6] J.H. Lee and G.T. Jin: *Link diagrams realizing prescribed subdiagram partitions*, Kobe J. Math. **18** (2001), 199–202.
- [7] A. Kawauchi: *Lectures on Knot Theory*, Kyoritsu shuppan, 2007, (in Japanese).
- [8] A. Kawauchi: *On a complexity of a spatial graph*; in *Knots and Soft-Matter Physics, Topology of Polymers and Related Topics in Physics, Mathematics and Biology*, Bussei Kenkyu **92**, 2009, 16–19.
- [9] W.B.R. Lickorish and K.C. Millett: *A polynomial invariant of oriented links*, Topology **26** (1987), 107–141.
- [10] Y. Nakanishi: *A note on unknotting number*, Math. Sem. Notes Kobe Univ. **9** (1981), 99–108.
- [11] M. Ozawa: *Ascending number of knots and links*, to appear in J. Knot Theory Ramifications.
- [12] D. Rolfsen: *Knots and Links*, Publish or Perish, Berkeley, CA, 1976.
- [13] A. Shimizu: *The warping degree of a knot diagram*, to appear in J. Knot Theory Ramifications.
- [14] A. Stoimenow: *Gauß diagram sums on almost positive knots*, Compos. Math. **140** (2004), 228–254.
- [15] K. Taniyama: *Unknotting numbers of diagrams of a given nontrivial knot are unbounded*, J. Knot Theory Ramifications **18** (2009), 1049–1063.

Graduate School of Science
 Osaka City University
 Osaka 558–8585
 Japan
 e-mail: shimizu1984@gmail.com