

Title	Note on locally compact groups
Author(s)	Yamabe, Hidehiko
Citation	Osaka Mathematical Journal. 1951, 3(1), p. 77-82
Version Type	VoR
URL	https://doi.org/10.18910/11390
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Note on Locally Compact Groups

By Hidehiko YAMABE

§1. The purpose of this note is to study the problem proposed by C. Chevalley: Is it true that a locally compact group which has no arbitrarily small¹⁾ subgroup is a Lie group?

Concerning the above problem two theorems will be proved in this note. One of them is:

Theorem 1. *A locally euclidean group G which has a neighbourhood of the identity containing no non-trivial subgroup, has a neighbourhood \tilde{U} of the identity, through any point of which there exists one and only one one-parameter subgroup²⁾.*

The other is:

Theorem 2. *If $(U_n)^n$ is contained in \tilde{U} , then G is a Lie group, where U_n denotes the aggregate of the n -th roots of elements in a neighbourhood U .*

§2. For an element x of a neighbourhood U of the identity e we denote by $\delta_v(x)$ the smallest number n such that $x^{2^n} \in U$. The group G is said to have the property (S) if there exists a neighbourhood U of e such that $\delta_v(x) < \infty$ for every x in $U - \{e\}$. According to Kuranishi³⁾ a locally euclidean group G which has the property (S), has a neighbourhood of the identity e , through any point of which one can draw one and only one one-parameter subgroup. Therefore we have only to show that a locally compact group which has no arbitrarily small subgroup has the property (S).

In order to prove this we shall need the following Lemma.

Lemma 1. Let W be a neighbourhood of the identity e which contains no non-trivial subgroup in it. For an arbitrarily small neigh-

1) A small subgroup means a subgroup contained in a sufficiently small neighbourhood of the identity.

2) This theorem was proved with the co-operation of Dr. Gotô. Cf. the forthcoming Nagoya Math. Journal.

3) See Kuranishi: *Differentiability of locally compact groups*, Nagoya Math. Journal Vol. 1, 1950, 71-81.

neighbourhood U of e there exists a neighbourhood U^* of e such that if $x, x^i \in U^*$ and $x^i \in W$ for all $i \leq k$, then $x^k \in U$ for all $i \leq k$.

Proof.

Put

$$X = \{x^k; x^k \in U, x^i \in W \text{ for all } i \leq k, x^j \in U \text{ for some } j \leq k.\}$$

If $U - X$ does not contain any neighbourhood of e , then there exists a sequence $\{a_n\}$ such that

$$a_n^{k_n} \rightarrow e$$

and

$$a_n^{j_n} \in U,$$

with

$$j_n \leq k_n.$$

We may suppose that $a_n^{j_n}$ converges to $\bar{a} \in \bar{W}$.

For an integer r we can easily find integers r_n such that

$$r_n \equiv r j_n \pmod{k_n}$$

$$0 < r_n \leq k_n.$$

Then $\{a_n^{r_n}\}$ converges to \bar{a}^r because

$$a_n^{r_n} = a_n^{r j_n} \cdot a_n^{p_n k_n},$$

where p_n are integers whose absolute values are less than r .

Let us denote by A the aggregate of limit points \bar{a}^r . A is clearly a non-trivial subgroup of G contained in W , because $A \ni \bar{a} \neq e$. This contradicts the hypothesis and whence we complete the proof of Lemma 1.

Now we shall have the

Theorem 1'. *A locally compact group which has no arbitrarily small subgroup, has the property (S).*

Proof. Let us take a neighbourhood U of e such that $U^2 \subset W$. Let V be a neighbourhood of e contained in U^* . If $x, x^2, x^3, \dots, x^{2^j} \in U$ and $x^{2^{j+1}} \in V$, then clearly $x^i \in W$ for all $i \leq 2^{j+1}$ and by Lemma 1 $x^i \in U$ for all $i \leq 2^{j+1}$. Therefore for a large j , $x^{2^j} \in V$, which shows that G has the property (S).

§3. Concerning Theorem 2 we shall also use the Kuranishi's⁴⁾ results. He proved that if $(x^{1/n} y^{1/n})^n$ converge uniformly to a continuous function over $U \times U$, then G is a Lie group.

We shall need some preparatory lemmas.

4) l. c. 3).

Lemma 2. Under the assumption of Theorem 2 a metric $\rho(x, y)$ can be defined so that $\rho(x^\lambda, e)$ may be differentiable at $\lambda = 0$.

Proof. At first we must define the metric. Without loss of generality \tilde{U} may be taken as a symmetric one.

Let y be an element of the boundary $Bd(\tilde{U})$ of \tilde{U} ⁵⁾, and let U be a neighbourhood of e such that

$$U^2 \subset \tilde{U}.$$

Let us define a metric $\rho(x, y)$ in U such that

$$\rho(x, e) = \inf_{(y)} \sum |\lambda_i|$$

$$\rho(x, z) = \rho(z^{-1}x, e)$$

where $\inf(y)$ means the infimum of $\sum |\lambda_i|$ when we take an arbitrary decomposition

$$x = y_1^{\lambda_1} y_2^{\lambda_2} \dots y_n^{\lambda_n}$$

for a suitable y_i 's $\in Bd(\tilde{U})$ and real λ_i 's.

It is clear that

$$\rho(x, y) = \rho(y, x)$$

$$\rho(x, y) + \rho(y, z) \geq \rho(x, z).$$

This metric $\rho(x, y)$ satisfies the metric conditions when we prove that $\rho(x, y) = 0$ implies $x = y$. We may suppose that $y = e$ and G is connected.

For a sufficiently large s

$$x^s \in \tilde{U}.$$

We denote by $t(n)$ the smallest integer such that

$$(U_n)^{t(n)} \ni x.$$

Then

$$st(n) > n,$$

$$t(n)/n > 1/s.$$

From this inequality we shall have easily

$$\rho(x, e) > 0.$$

Hence we have proved that $\rho(x, y)$ satisfies the metric conditions. From the definition we can easily see that

5) We may assume that $Bd(\tilde{U})$ intersects in only one point with any one-parameter semi-group. See H. Whitney: *On regular family of curves*, Bull. Amer. Math. Soc. 47, 145-147 (1941).

$$\rho(x^\lambda, e) \leq \lambda$$

for $x \in U$.

Then $\rho(x^\lambda, e)$ is differentiable because $\rho(x^\lambda, e)$ is subadditive and $\rho(x^\lambda, e)/\lambda \leq 1$ ⁶⁾.

§ 4. Now let us study some properties of this metric $\rho(x, y)$.

Lemma 3. The metric $\rho(x, y)$ has the following properties:

- i) $\rho(x, y)$ is left invariant,
- ii) $K_2\rho(y, e) \leq \rho(yx, x) \leq K_1\rho(y, e)$, ⁷⁾
- iii) if $\rho(x, e) = O(\mu)$, then

$$\frac{\rho(x^{-1}yx, e)}{\rho(y, e)} = 1 + O(\mu).$$

Proof. i) is evident. From the definition of this metric we have for some y_i 's $\in Bd(\bar{U})$, real λ_i 's and an arbitrarily small ε ,

$$\left| \rho(y, e) - \sum |\lambda_i| \right| \leq \varepsilon$$

and

$$y = y_1^{\lambda_1} y_2^{\lambda_2} \dots y_n^{\lambda_n}.$$

Let us consider a real number $s(y)$ with

$$x^{-1}y^{s(y)}x \in Bd(\bar{U}).$$

Then for every i

$$0 < K_2 \leq s(y_i) \leq K_1 < \infty.$$

Hence by simple calculations we obtain

$$K_2\rho(y, e) \leq \rho(x^{-1}yx, e) \leq K_1\rho(y, e).$$

Thus the proposition ii) is proved.

In case $\rho(x, e) = O(\mu)$,

$$s(y) = 1 + O(\mu),$$

because

$$\frac{\rho(x^{-1}yx, e)}{\rho(y, e)} - 1 \leq \frac{\rho(x, e) + \rho(y^{-1}xy, e)}{\rho(y, e)} \leq \frac{O(\mu)}{\rho(y, e)}.$$

Hence the proposition iii) is proved and we complete the proof of Lemma 3.

6) See, Einar Hille: *Functional Analysis and Semigroups*, Amer. Math. Soc. Coll. Pub. p. 143.

7) In this note K_i 's are all absolute constants. They could be taken near to 1 except K_4 .

§5. From the following relations

$$x(xaxa^{-1})x^{-1}(xaxa^{-1}) = x^2ax^2a^{-1},$$

we have

$$\rho(xaxa^{-1}, e)(1+C(x)) \geq \rho(x^2ax^2a^{-1}, e),$$

where $C(x)$ denotes

$$\sup_y \frac{\rho(x^{-1}yx, e)}{\rho(y, e)}$$

If $\delta_{\tilde{v}}(a)$ and $\delta_{\tilde{v}}(x)$ are $\leq n$, then

$$\begin{aligned} \rho(xaxa^{-1}, e) &\prod_{i=1}^n (1+C(x^{2^i})) \\ &\geq \rho(x^{2^n}ax^{2^n}a^{-1}, e) \\ &\geq \rho(x^{2^{n+1}}, e) + O(1/2^n) \\ &\geq 2^{n+1}K_3\rho(x, e). \end{aligned}$$

It is possible to take our neighbourhood \tilde{U} so small as to make K_3 sufficiently near to 1.

On the other hand

$$\prod_{i=1}^n (1 + C(x^{2^i}))/2^n$$

can be taken sufficiently near to 1 too.

Then we have

$$\rho(xaxa^{-1}, e) \geq K_4\rho(x, e)$$

with $K_4 > 1$.

Put

$$\begin{array}{ll} x_1 = x & a_1 = a \\ \vdots & \vdots \\ x_i = x_{i-1}a_{i-1}x_{i-1}a_{i-1}^{-1}, & a_i = a^{2^i} \\ \vdots & \vdots \end{array}$$

Then by simple calculations

$$\rho(x_p, e) \geq K_4^p\rho(x, e)$$

or

$$\rho((xa)^{2^p}a^{-2^p}, e) \geq K_4^p\rho(x, e).$$

Now take $x^{1/2^q}y^{1/2^p}x^{-1/2^q}y^{-1/2^p}$ for x and $y^{1/2^p}$ for a in the above inequality.

Then

$$\begin{aligned} &\rho(x^{1/2^q}yx^{-1/2^q}y^{-1}, e) \\ &\geq K_4^p\rho(x^{1/2^q}y^{1/2^p}x^{-1/2^q}y^{-1/2^p}, e). \end{aligned}$$

This means that

$$2^q \rho(x^{1/2} y^{1/2} x^{-1/2} y^{-1/2}, e)$$

converge to zero when p and q increase to ∞ .

Consequently

$$\rho(x^\mu y^\lambda x^{-\mu} y^{-\lambda}, e) / \mu$$

converge to zero when λ and μ decrease to zero.

§ 6. Proof of Theorem 2.

Put

$$F_n(x, y) = (x^{1/n} y^{1/n})^n.$$

Then

$$\begin{aligned} & \rho(F_n(x, y), F_{np}(x, y)) \\ &= \rho((x^{1/n} y^{1/n})^n, (x^{1/np} y^{1/np})^{np}) \\ &\leq K_1 n \rho(x^{1/n} y^{1/n}, (x^{1/np} y^{1/np})^p) \\ &\leq K_1 n \sum_{i=0}^{p-1} \rho(y^{1/np}, x^{-i/np} y^{1/np} x^{i/np}) \end{aligned}$$

By the inequality obtained in the last chapter

$$\begin{aligned} &\leq K_1 np/np \cdot \varepsilon_n \\ &= K_1 \varepsilon_n, \end{aligned}$$

where ε_n converges to zero when $n \rightarrow \infty$.

Hence we proved that $F_n(x, y)$ converge uniformly in $U \times U$, which completes the proof of Theorem 2.

To conclude this note, the author wishes to thank Dr. Gotô and Dr. Kuranishi for their valuable suggestions and advices.

(Received January 11, 1951)

Added in proof. Theorem 1 is true when the group is locally connected. The proof will be given in the next number of this journal.