

Title	On J_R -homomorphisms
Author(s)	Minami, Haruo
Citation	Osaka Journal of Mathematics. 1979, 16(3), p. 581-588
Version Type	VoR
URL	https://doi.org/10.18910/11553
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ON J_R -HOMOMORPHISMS

HARUO MINAMI

(Received October 13, 1978)

1. Introduction

In [8] Snaith proved the Adams conjecture for suspension spaces. In this paper we shall prove an analogous result to Snaith's theorem ([8], Corollary 5.2) for the Real Adams operation ψ^3 and a Real J -map J_R (see §2). This is proved by using the results of Seymour [7]. And as an application we shall determine an undecided order in the theorem of [6].

Here we shall inherit the notations and terminologies in [2], §1 and [6].

2. Homomorphism J_R

In [6] we defined the homomorphisms $J_{R,n}$ and J_R for doubly indexed suspension spaces $\Sigma^{p,q}X$, $p \geq 0$ and $q \geq 1$. Clearly, these definitions are also valid for any finite pointed τ -complex. But the natural map obtained in this manner

$$J_R: \widetilde{KR}^{-1}(X) \rightarrow \pi_s^{0,0}(X)$$

is not a homomorphism in general. As in the usual case we see that this map satisfies the following formula:

$$J_R(\alpha + \beta) = J_R(\alpha) + J_R(\beta) + J_R(\alpha)J_R(\beta) \quad \alpha, \beta \in \widetilde{KR}^{-1}(X)$$

where ab ($a, b \in \pi_s^{0,0}(X)$) denotes the product of a and b induced by the loop composition in $\Omega^{n,n}\Sigma^{n,n}$ (cf. [9], p. 314).

3. Adams operation ψ^3 in KR -theory

In this section we recall the construction of the Real Adams operation ψ_R^3 described in [7], §4.

Let S_3 be the symmetric group with two generators a, b satisfying

$$a^3 = b^2 = 1, \quad bab = a^2$$

and let Z_3 be the cyclic subgroup of S_3 generated by a . From the above relations we see that $\tau(a) = a^2$, $\tau(b) = b$ induces an automorphic involution τ on

S_3 . Z_3 is closed under the involution τ . Therefore S_3 (resp. Z_3) is regarded as a Real group with the involution τ (resp. its restriction to Z_3) in the sense of Atiyah-Segal [4].

We know that all simple S_3 - and Z_3 -modules over \mathbf{C} are as follows:

$$(3.1) \quad \begin{aligned} S_3: \tilde{\mathbf{1}} &= \{\mathbf{C} \mid a = 1, b = 1\}, \tilde{M} = \{\mathbf{C} \mid a = 1, b = -1\}, \\ \tilde{M}_1 &= \{\mathbf{C}^2 \mid av = Av, bv = Bv, v \in \mathbf{C}^2\} \\ Z_3: \mathbf{1} &= \{\mathbf{C} \mid a = 1\}, M_1 = \{\mathbf{C} \mid av = \zeta v, v \in \mathbf{C}\}, \\ M_2 &= \{\mathbf{C} \mid av = \zeta^2 v, v \in \mathbf{C}\} \end{aligned}$$

where $A = \begin{bmatrix} \zeta & 0 \\ 0 & \zeta^2 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \in GL(2, \mathbf{C})$ and $\zeta = \exp(2\pi i/3)$ (see, e.g., [5], §32).

Let G denote either S_3 or Z_3 . Clearly each G -module listed above is a Real G -module with the conjugate linear involution induced by complex conjugation. This fact shows that the forgetful map $R_R(G) \rightarrow R(G)$, which is injective in general, is surjective where $R_R(G)$ is the Grothendieck group of Real G -modules and $R(G)$ is the complex representation ring of G .

Let X be a Real space with trivial G -action and $F \rightarrow X$ be a Real G -vector bundle in the sense of [4], §6. Then we see easily that the decomposition of F as a complex G -vector bundle ([4], §8)

$$(3.2) \quad \bigoplus_M \text{Hom}^G(M, F) \otimes M \xrightarrow{\cong} F$$

becomes an isomorphism of Real G -vector bundles. Here M runs through the simple G -modules over \mathbf{C} and \underline{M} denotes the product bundle $M \times X$ over X . And so we see that (3.2) induces a natural isomorphism $KR_G(X) \cong KR(X) \otimes R(G)$.

Let $E \rightarrow X$ be a Real vector bundle over X with the involution $\tau_E: E \rightarrow E$. We define a Real structure $\tilde{\tau}_E$ on $E^{\otimes 3}$ by $\tilde{\tau}_E = (1 \otimes t)\tau_E^{\otimes 3}$ where $t: E^{\otimes 2} \rightarrow E^{\otimes 2}$ is the switching map. Then $E^{\otimes 3}$ becomes a Real S_3 -vector bundle with the S_3 -action permuting the factors.

Applying (3.2) to $E^{\otimes 3}$ we have an isomorphism of Real S_3 -vector bundles

$$(3.3) \quad \begin{aligned} E^{\otimes 3} \cong & \text{Hom}^{S_3}(\tilde{\mathbf{1}}, E^{\otimes 3}) \otimes \tilde{\mathbf{1}} \oplus \text{Hom}^{S_3}(\tilde{M}, E^{\otimes 3}) \otimes \tilde{M} \\ & \oplus \text{Hom}^{S_3}(\tilde{M}_1, E^{\otimes 3}) \otimes \tilde{M}_1 \end{aligned}$$

with the notations of (3.1). And by (3.3), as a Real Z_3 -vector bundle we obtain

$$(3.4) \quad \begin{aligned} E^{\otimes 3} \cong & (\text{Hom}^{S_3}(\tilde{\mathbf{1}}, E^{\otimes 3}) \oplus \text{Hom}^{S_3}(\tilde{M}, E^{\otimes 3})) \otimes \mathbf{1} \\ & \oplus \text{Hom}^{S_3}(\tilde{M}_1, E^{\otimes 3}) \otimes (\underline{M}_1 \oplus \underline{M}_2). \end{aligned}$$

Put

$$\begin{aligned} V_0 &= \text{Hom}^{S_3}(\tilde{\mathbf{1}}, E^{\otimes 3}) \oplus \text{Hom}^{S_3}(\tilde{M}, E^{\otimes 3}), \\ V_1 &= \text{Hom}^{S_3}(\tilde{M}_1, E^{\otimes 3}) \end{aligned}$$

and

$$N = 1 \oplus M_1 \oplus M_2, \text{ the regular representation of } Z_3,$$

then by (3.4)

$$(3.5) \quad E^{\otimes 3} \cong V_0 \otimes \underline{1} \oplus V_1 \otimes (\underline{M}_1 \oplus \underline{M}_2)$$

as a Real Z_3 -vector bundle and so

$$[E^{\otimes 3}] = ([V_0] - [V_1]) \otimes 1 + [V_1] \otimes N$$

in $KR_{Z_3}(X) = KR(X) \otimes R(Z_3)$ where $[A]$ denotes the isomorphism class of A .

Here we define ϕ_R^3 by

$$(3.6) \quad \phi_R^3([E]) = [V_0] - [V_1].$$

Then we can easily check that ϕ_R^3 satisfies the properties of Adams operation. And moreover by [3], Proposition 2.5 we see that forgetting the Real structure, ϕ_R^3 is reduced to the complex Adams operation ψ_U^3 .

4. Real Adams conjecture for ϕ_R^3

The purpose of this section is to prove the following theorem.

Theorem 4.1. *Let X be a finite pointed τ -complex. Then*

$$J_R(\phi_R^3(x)) = J_R(x) \quad \text{for any } x \in \widetilde{KR}^{-1}(X)$$

in $\pi_s^{0,0}(X) \left[\frac{1}{3} \right]$.

Let Y be a τ -space with trivial Z_3 -action. As in §3 we assume here that $E^{\otimes 3}$ has the twisted Real structure for a Real vector bundle E over Y . We have the following lemmas as in [7], §1.

Lemma 4.2 (cf. [7], Proposition 1.2). *There is a natural isomorphism of Real Z_3 -vector bundles*

$$(E \oplus F)^{\otimes 3} \cong E^{\otimes 3} \oplus F^{\otimes 3} \oplus (U'(E, F) \otimes N)$$

for Real vector bundles E and F over Y .

Lemma 4.3 (cf. [7], p.399). *For the trivial Real vector bundle \underline{n} of dimension n over Y there is a canonical isomorphism of Real Z_3 -vector bundles*

$$\theta_n: \underline{n}^{\otimes 3} \rightarrow \underline{n} \oplus (\underline{n}' \otimes \underline{N})$$

such that

$$\pi_n \theta_n((\sum_{i=1}^n z_i e_i)^{\otimes 3}, x) = (\sum_{i=1}^n z_i^3 e_i, x) \quad (z_i \in \mathbf{C}, x \in X)$$

where let π_n denote the projection of $\underline{n} \oplus (\underline{n}' \otimes \underline{N})$ onto \underline{n} and let e_1, \dots, e_n denote

the standard basis of \mathbf{C}^n .

Let $f_k: \underline{n}^{\otimes 3} \rightarrow \underline{n} \oplus (\underline{n}' \otimes \underline{N})$ ($k=1,2$) be isomorphisms of Real Z_3 -vector bundles. Consider the direct sum

$$f_1 \oplus f_2: 2\underline{n}^{\otimes 3} \rightarrow 2\underline{n} \oplus (2\underline{n}' \otimes \underline{N}).$$

By Lemma 4.2, adding $U'(\underline{n}, \underline{n}) \otimes \underline{N}$ to the above isomorphism we have an isomorphism of Real Z_3 -vector bundles

$$(2\underline{n})^{\otimes 3} \rightarrow 2\underline{n} \oplus ((2\underline{n})' \otimes \underline{N})$$

for which we write $f_1 + f_2$.

By modifying the proof of [7], Proposition 2.5 we get the following

Lemma 4.4 (cf. [7], Proposition 2.5). *Given an isomorphism of Real Z_3 -vector bundles $f: \underline{n}^{\otimes 3} \rightarrow \underline{n} \oplus (\underline{n}' \otimes \underline{N})$, there is an isomorphism of Real Z_3 -vector bundles $g: \underline{n}^{\otimes 3} \rightarrow \underline{n} \oplus (\underline{n}' \otimes \underline{N})$ such that $f+g$ is homotopic to θ_{2n} through Real Z_3 -isomorphism.*

Define a map $\delta: \mathbf{C}^n \rightarrow \mathbf{C}^n$ by $\delta(z_1, \dots, z_n) = (z_1^3, \dots, z_n^3)$ ($z_i \in \mathbf{C}$). Then δ induces a base-point-preserving τ -map of $\Sigma^{n,n}$ into itself which we denote by the same letter δ . Now, according to [2], Theorem 12.5

$$\pi_{n,n}(\Sigma^{n,n}) = Z[\rho]/(1-\rho^2)$$

for $n \geq 1$. We observe $[\delta]^\tau \in \pi_{n,n}(\Sigma^{n,n})$, the τ -homotopy class of δ .

Lemma 4.5. *With the above notations, we have*

$$[\delta]^\tau = \frac{1+3^n}{2} + \frac{1-3^n}{2}\rho \quad (n \geq 1)$$

in $\pi_{n,n}(\Sigma^{n,n})$.

Proof. We have

$$\psi(1) = 1, \phi(1) = 1, \psi(\rho) = -1 \text{ and } \phi(\rho) = 1$$

where ψ and ϕ are the forgetful and fixed-point homomorphisms respectively. So putting $[\delta]^\tau = x + y\rho$ ($x, y \in Z$) we have

$$x = \frac{1+3^n}{2} \text{ and } y = \frac{1-3^n}{2}$$

since $\psi([\delta]^\tau) = 3^n$ and $\phi([\delta]^\tau) = 1$ by the definition. q.e.d.

For a τ -map σ of $\Sigma^{l,l}$ into itself we define a τ -map $t_\sigma: \Omega^{m,m} \Sigma^{m,m} \rightarrow \Omega^{l+m,l+m} \Sigma^{l+m,l+m}$ by $t_\sigma(\eta) = \sigma \wedge \eta$ ($\eta \in \Omega^{m,m} \Sigma^{m,m}$) where ' \wedge ' denotes the smash product upon one point compactification. Let ε be a τ -map of $\Sigma^{n,n}$ into itself such that

$$[\varepsilon]^\tau = \frac{1+3^n}{2} - \frac{1-3^n}{2}\rho$$

in $\pi_{n,n}(\Sigma^{n,n})$. Then $[\varepsilon\delta]^\tau = 3^n$ for δ as in Lemma 4.5. Hence we have

$$\varepsilon \wedge \delta \simeq_\tau \varepsilon\delta \wedge 1 \simeq_\tau 3^n: \Sigma^{2n,2n} \rightarrow \Sigma^{2n,2n}$$

where 1 is the identity map of $\Sigma^{n,n}$.

For a τ -map $h: X \rightarrow GL(n, \mathbf{C})$ we define a τ -map $\tilde{h}: X \rightarrow \Omega_0^{3n,3n} \Sigma^{3n,3n}$ to be the composition

$$\begin{aligned} X &\xrightarrow{h} GL(n, \mathbf{C}) \xrightarrow{i} \Omega^{n,n} \Sigma^{n,n} \xrightarrow{t_\delta} \Omega^{2n,2n} \Sigma^{2n,2n} \\ &\xrightarrow{t_\varepsilon} \Omega^{3n,3n} \Sigma^{3n,3n} \xrightarrow{\tilde{t}} \Omega^{3n,3n} \Sigma^{3n,3n} . \end{aligned}$$

Here i is the canonical inclusion map and \tilde{t} is the map given by adding a fixed map of degree (-3^n) to the elements of $\Omega^{3n,3n} \Sigma^{3n,3n}$ with respect to the loop addition along fixed coordinates of $\Sigma^{3n,3n}$. By adh we denote the adjoint of \tilde{h} . Then, by the definition of $J_{R,3n}$ we have

Lemma 4.6. *With the above notations*

$$[\text{ad } h]^\tau = 3^n J_{R,3n}([\tilde{j}h]^\tau)$$

where j is a canonical inclusion map of $GL(n, \mathbf{C})$ into $GL(3n, \mathbf{C})$.

As we note in [6] we have

$$\widetilde{KR}^{-1}(X) \cong \widetilde{KR}(\Sigma^{0,1}X) \cong [X, GL(\infty, \mathbf{C})]^\tau .$$

So we see that any Real vector bundle over $\Sigma^{0,1}X$ is obtained from the clutching of the trivial bundles $E_1 = \mathbf{C}^m \times \Sigma_+^{0,1}X$ and $E_2 = \mathbf{C}^m \times \Sigma_-^{0,1}X$ by a base-point-preserving τ -map from X to $GL(m, \mathbf{C})$. Here,

$$\begin{aligned} \Sigma_+^{0,1}X &= \{t \wedge x \in SX \mid t \geq 0\}, \quad \Sigma_-^{0,1}X = \{t \wedge x \in SX \mid t \leq 0\}, \\ X &= \Sigma_+^{0,1}X \cup \Sigma_-^{0,1}X \end{aligned}$$

and we consider that \mathbf{C}^m has the natural Real structure, i.e., $\mathbf{C}^m = \mathbf{R}^{m,m}$.

Proof of Theorem 4.1. Denote by E_α the associated vector bundle with a base-point-preserving τ -map $\alpha: X \rightarrow GL(m, \mathbf{C})$. From (3.5) we have a decomposition

$$E_\alpha^{\otimes 3} \cong V_0 \oplus V_1 \otimes (M_1 \oplus M_2)$$

as a Real Z_3 -vector bundle over $\Sigma^{0,1}X$ where ‘ $\otimes \mathbb{1}$ ’ is omitted for the simplicity. Also we have a vector bundle V_1^* over $\Sigma^{0,1}X$ such that $V_1 \oplus V_1^* \cong 2s$ where let $\dim V_1 = s$. Adding $V_1 \oplus V_1^*$ to the above isomorphism we obtain an isomorphism

$$E_{\alpha}^{\otimes 3} \oplus 2s \cong (V_0 \oplus V_1^*) \oplus (V_1 \otimes N).$$

By Lemmas 4.2 and 4.3, adding $((2s)' \oplus U'(E_{\alpha}, 2s)) \otimes N$ we obtain an isomorphism

$$(4.1) \quad \beta: (E_{\alpha} \oplus 2s)^{\otimes 3} \xrightarrow{\cong} (V_0 \oplus V_1^*) \oplus ((2s)' \oplus U'(E_{\alpha}, 2s) \oplus V_1) \otimes N.$$

And by (3.6) we have

$$(4.2) \quad [V_0 \oplus V_1^*] \cong \psi_R^3([E_{\alpha}]) + 2s.$$

Observe the restrictions of (4.1) over $\Sigma_+^{0,1}X$ and $\Sigma_-^{0,1}X$, then β yields an isomorphism of trivial bundles over each space since $\Sigma_{\pm}^{0,1}X$ are contractible. Therefore we have a homotopy commutative diagram

$$(4.3) \quad \begin{array}{ccc} m^{\otimes 3} & \xrightarrow{\beta_+} & \underline{m} \oplus (m' \otimes N) \\ \downarrow f'^{\otimes 3} & & \downarrow g' \oplus (g'' \otimes 1) \\ m^{\otimes 3} & \xrightarrow{\beta_-} & \underline{m} \oplus (m' \otimes N). \end{array}$$

Here the dotted arrows denote isomorphisms which are defined only over X and β_{\pm} are given by $\beta_{\pm}(v, x) = (\beta(v, *), x)$ ($x \in \mathbf{C}^{m^3}$, $x \in \Sigma_{\pm}^{0,1}X$) respectively where $*$ is the base-point of $\Sigma^{0,1}X$.

Applying Lemma 4.4 to the horizontal isomorphisms in (4.3) we obtain the following homotopy commutative diagram

$$(4.4) \quad \begin{array}{ccc} (2m)^{\otimes 3} & \xrightarrow{\theta_{2m}} & 2m \oplus ((2m)' \otimes N) \\ \downarrow \tilde{f}^{\otimes 3} & & \downarrow \tilde{g} \oplus (\tilde{g}' \otimes 1) \\ (2m)^{\otimes 3} & \xrightarrow{\theta_{2m}} & 2m \otimes ((2m)' \otimes N) \end{array}$$

where \tilde{f} , \tilde{g} and \tilde{g}' are isomorphisms over X which are naturally induced from f' , g' and g'' respectively.

Put $n=2m$ in (4.4). By Lemma 4.3 we see that the composition

$$\underline{n} \xrightarrow{\Delta_n} \underline{n}^{\otimes 3} \xrightarrow{\theta_n} \underline{n} \oplus (n' \otimes N) \xrightarrow{\pi_n} \underline{n}$$

induces a constant τ -map

$$\gamma: X \rightarrow \Omega^{n, n} \Sigma^{n, n}$$

given by $\gamma(x) = \delta(x \in X)$ where $\Delta_n(u) = u^{\otimes 3}$ and δ is as in Lemma 4.5. Besides we see that \tilde{f} and \tilde{g} induce τ -maps

$$f: X \rightarrow GL(n, \mathbf{C}) \text{ and } g: X \rightarrow GL(n, \mathbf{C})$$

in the natural way. By the commutativity of (4.4), we have

$$(ig) \circ \gamma \simeq_{\tau} \gamma \circ (if): X \rightarrow \Omega^{n,n} \Sigma^{n,n}$$

where $i: GL(n, \mathbf{C}) \subset \Omega^{n,n} \Sigma^{n,n}$ denotes the inclusion map and $f \circ h$ is given by $(f \circ h)(x)(z) = f(x)(h(x)(z))$ ($x \in X, z \in \Sigma^{n,n}$) for τ -maps $f, h: X \rightarrow \Omega^{n,n} \Sigma^{n,n}$. Therefore we obtain

$$(4.5) \quad \gamma \wedge ig \simeq_{\tau} \gamma \wedge if: X \rightarrow \Omega^{2n,2n} \Sigma^{2n,2n}$$

where $f \wedge h$ is given by $(f \wedge h)(x)(z_1 \wedge z_2) = f(x)(z_1) \wedge h(x)(z_2)$ ($x \in X, z_1, z_2 \in \Sigma^{n,n}$) for τ -maps $f, h: X \rightarrow \Omega^{n,n} \Sigma^{n,n}$. Therefore, by (4.5) and Lemma 4.6 we obtain

$$3^n J_{R,3n}([jf]^\tau) = 3^n J_{R,3n}([jg]^\tau).$$

This shows

$$J_{R,3n}([jf]^\tau) = J_{R,3n}([jg]^\tau)$$

in $[\Sigma^{3n,3n} X, \Sigma^{3n,3n}]^\tau \left[\frac{1}{3} \right]$. Consequently, passing the direct limit we have

$$J_R(\{E_\alpha\}) = J_R(\psi_R^3(\{E_\alpha\}))$$

in $\pi_s^{0,0}(X) \left[\frac{1}{3} \right]$ where $\{A\}$ denotes the stable isomorphism class of A , because the vector bundles associated with f and α are stably equivariant and g represents $\psi_R^3(\{E_\alpha\})$ stably by (4.2). This completes the proof.

5. $J_R(\pi_{m,n}(GL(\infty, \mathbf{C})))$

In [6] we showed that if p is odd and k is even then the image $J_R(\pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C})))$ ($p > k \geq 0$) is a cyclic group and its order is either $m(2p)$ or $2m(2p)$. Here we prove the following

Theorem 5.1. *The image $J_R(\pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C})))$ is a cyclic group of order $m(2p)$ for $p > k \geq 0$, p odd and k even.*

Proof. Consider the following diagram

$$\begin{array}{ccc} \pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C})) & \xrightarrow{c_1} & \pi_{4p-1}(GL(\infty, \mathbf{C})) \\ \uparrow \cong & & \uparrow \cong \\ \pi_{0, 4k-1}(GL(\infty, \mathbf{C})) = \pi_{4k-1}(GL(\infty, \mathbf{R})) & \xrightarrow{c_2} & \pi_{4k-1}(GL(\infty, \mathbf{C})) \end{array}$$

where the isomorphisms are the complex and Real Thom isomorphisms, and c_1 and c_2 are the natural complexification homomorphisms. Then we see easily that this diagram is commutative and c_2 is an isomorphism since k is even. Therefore c_1 becomes an isomorphism. Let g be a generator of $\pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C})) = Z$. Then we have

$$\psi_R^3(g) = 3^{2p}g$$

because $c_1\psi_R^3 = \psi_U^3c_1$ and $\psi_U^3(c_1(g)) = 3^{2p}c_1(g)$. Moreover we have $\nu_2(3^{2p}-1) = \nu_2(m(2p))$ by [1], Lemma 2.12 (ii). When we denote by G the quotient module of $\pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C}))$ by $(\psi_R^3-1)(\pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C})))$ we obtain

$$G_{(2)} \cong Z_2^{\nu_2(m(2p))}$$

by the above arguments where $G_{(2)}$ denotes the module obtained from G by localizing at the prime ideal (2). Now Theorem 4.1 yields the following 2-local factrization:

$$\begin{array}{ccc} \pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C}))_{(2)} & \xrightarrow{J_{R(2)}} & \pi_{2p-2k, 2p+2k-1(2)}^s \\ & \searrow & \nearrow \\ & G_{(2)} & \end{array}$$

This result and the theorem of [6] show that the order of $J_R(\pi_{2p-2k, 2p+2k-1}(GL(\infty, \mathbf{C})))$ is equal to $m(2p)$.

OSAKA CITY UNIVERSITY

References

- [1] J.F. Adams: *On the groups J(X)* II, *Topology* **3** (1965), 137-171.
- [2] S. Araki and M. Murayama: *τ -Cohomology theories*, *Japan. J. Math.* **4** (1978), 363-416.
- [3] M.F. Atiyah: *Power operation in K-theory*, *Quart. J. Math., Oxford*, **17** (1966), 165-193.
- [4] M.F. Atiyah and G.B. Segal: *Equivariant K-theory and completion*, *J. Differential Geometry* **3** (1969), 1-18.
- [5] C.W. Curtis and I. Reiner: *Representation theory of finite groups and associative algebras*, *Pure and Applied Mathematics vol. XI*, J. Wiley and Sons, Inc., 1962.
- [6] H. Minami: *On Real J-homomorphisms*, *Osaka J. Math.* **16** (1979), 529-537.
- [7] R.M. Seymour: *Vector bundles invariant under the Adams operations*, *Quart. J. Math., Oxford*, **25** (1974), 395-414.
- [8] V. Snaith: *The complex J-homomorphism I*, *Proc. London Math. Soc.* **34** (1977), 269-302.
- [9] R.M.W. Wood: *Framing the exceptional Lie group G₂*, *Topology* **15** (1976), 303-320.