



Title	On separable algebras over a commutative ring
Author(s)	Endo, Shizuo; Watanabe, Yutaka
Citation	Osaka Journal of Mathematics. 1967, 4(2), p. 233-242
Version Type	VoR
URL	https://doi.org/10.18910/11833
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

ON SEPARABLE ALGEBRAS OVER A COMMUTATIVE RING*

SHIZUO ENDO AND YUTAKA WATANABE

(Received August 19, 1967)

Introduction. The notion of a separable algebra over a commutative ring was introduced in Auslander-Goldman [2], which coincides with that of a maximally central algebra in Azumaya [3] for a central algebra over a local ring. The basic properties of separable algebras were shown in [2] and [3].

The purpose of this paper is to define the reduced trace and norm of a central separable algebra over a commutative ring and to prove that a separable algebra over a commutative ring is a symmetric algebra.

Let Λ be a central separable algebra over a commutative ring R and let S be a commutative R -algebra such that $S \otimes_R \Lambda \cong \text{Hom}_S(P, P)$ for some finitely generated, faithful, projective S -module P . Then S is called, according to [2], a *splitting ring* of Λ , and especially, if $R \subseteq S$, it is called a *proper splitting ring* of Λ . It was proved in [2] that a central separable algebra over a Noetherian local ring R has a proper splitting ring which is a Galois extension of R . However, for a general commutative ring R , it is an open problem whether any central separable R -algebra has a proper (Galois) splitting ring. Therefore, our method, which will be used to defining the reduced trace and norm of a central separable R -algebra, is different from the usual one in the classical case (cf. [4]).

In § 1 we shall show that a separable algebra over a general commutative ring is extended from a separable algebra over a Noetherian commutative ring, and, in § 2, we shall prove that, in case R is a commutative ring included in a semi-local ring, a central separable R -algebra has a proper splitting ring.

§ 3 is devoted to defining the reduced trace of a central separable R -algebra Λ . If Λ has a proper splitting ring, we can define the reduced characteristic polynomial, trace and norm of Λ by using the characteristic polynomial, trace and norm of a projective module in [7], and we shall also show that there exist the analogous relations to the classical case between these and the characteristic polynomial, trace and norm of an R -algebra Λ . In the general case, we define the reduced trace of Λ , by using the above-mentioned result in § 1.

* This work was supported by the Matsunaga Science Foundation.

An algebra Λ over a commutative ring R , which is a finitely generated, faithful, projective R -module, is called, according to [6], a *symmetric R -algebra*, if $\text{Hom}_R(\Lambda, R)$ is Λ^e -isomorphic to Λ . In the classical theory, it is well known that any semi-simple algebra over a field is symmetric. However, for a general commutative ring R , it is an open problem whether a semi-simple R -algebra is symmetric or not.

In §4 we shall prove, as a partial answer to this, that a separable algebra over a commutative ring is symmetric. This includes the results in Müller [10] and DeMeyer [5].

Throughout this paper a ring means a ring with a unit element, and a (semi-) local ring means a commutative (semi-) local ring which is not always Noetherian.

1. Basic results

First we shall prove, as a generalization of (4.5) and (4.7) in [2],

Proposition 1.1. *Let Λ be an algebra over a (not always Noetherian) commutative ring R , which is a finitely generated R -module. Then the following conditions are equivalent:*

- (1) Λ is a separable R -algebra.
- (2) For any maximal ideal \mathfrak{m} of R , $\Lambda_{\mathfrak{m}}$ is a separable $R_{\mathfrak{m}}$ -algebra.
- (3) For any maximal ideal \mathfrak{m} of R , $\Lambda/\mathfrak{m}\Lambda$ is a separable R/\mathfrak{m} -algebra.

Proof. The implications (1) \Rightarrow (2) \Rightarrow (3) are obvious.

(2) \Rightarrow (1): We have $\text{w.dim}_{\Lambda^e} \Lambda = \sup_{\mathfrak{m}} \text{w.dim}_{\Lambda_{\mathfrak{m}}^e} \Lambda_{\mathfrak{m}}$ where \mathfrak{m} runs over all maximal ideals of R . If each $\Lambda_{\mathfrak{m}}$ is $R_{\mathfrak{m}}$ -separable, then we have $\text{w.dim}_{\Lambda_{\mathfrak{m}}^e} \Lambda_{\mathfrak{m}} = 0$ and so $\text{w.dim}_{\Lambda^e} \Lambda = 0$. As Λ is Λ^e -finitely presented, this shows that Λ is Λ^e -projective.

(3) \Rightarrow (2): Without loss of generality we may assume that R is a local ring with a maximal ideal \mathfrak{m} . Now suppose that $\Lambda/\mathfrak{m}\Lambda$ is R/\mathfrak{m} -separable. Let \hat{R} be the Henselization of R and put $\hat{\Lambda} = \hat{R} \otimes_R \Lambda$. Then we have $\hat{R}/\mathfrak{m}\hat{R} = R/\mathfrak{m}$ and $\hat{\Lambda}/\mathfrak{m}\hat{\Lambda} = \Lambda/\mathfrak{m}\Lambda$. Since \hat{R} is R -faithfully flat, we have $\text{w.dim}_{\Lambda^e} \Lambda = \text{w.dim}_{\hat{\Lambda}^e} \hat{\Lambda}$ and so Λ is Λ^e -projective if and only if $\hat{\Lambda}$ is $\hat{\Lambda}^e$ -projective. Hence we may further assume that R is Henselian. Then, for the projective $\Lambda^e/\mathfrak{m}\Lambda^e$ -module $\Lambda/\mathfrak{m}\Lambda$, there is a finitely generated projective Λ^e -module P such that $\bar{f}: P/\mathfrak{m}P \cong \Lambda/\mathfrak{m}\Lambda$ as Λ^e -modules. Since R is local and P, Λ^e are Λ^e -projective, there exist Λ^e -epimorphisms $f: P \rightarrow \Lambda$, which induces \bar{f} on $P/\mathfrak{m}P$, and $g: \Lambda^e \rightarrow P$ such that $f \circ g$ is the natural epimorphism of Λ^e onto Λ . The homomorphism $f \circ g$ is R -split and so f is also R -split. From this it follows directly that f is an isomorphism. Thus Λ is Λ^e -projective, which completes our proof.

It is remarked that, by (1.1), we can omit the assumption that R is Noetherian from almost all of results in [2].

The following proposition will play an important part in § 3.

Proposition 1.2. *Let Λ be a separable R -algebra, which is a finitely generated, faithful, projective R -module. Then there exist a Noetherian subring R' of R and a separable R' -subalgebra Λ' of Λ , which is a finitely generated, faithful, projective R' -module, such that $\Lambda = R \otimes_{R'} \Lambda'$.*

Proof. Let $\{\lambda_0=1, \lambda_1, \dots, \lambda_t\}$ be a set of generators of Λ over R . Let F be a free R -module with a basis $\{u_0, u_1, \dots, u_t\}$, and define the R -epimorphism $f: F \rightarrow \Lambda$ by putting $f(u_i) = \lambda_i$ for each i . Since Λ is R -projective, we have an R -homomorphism $g: \Lambda \rightarrow F$ such that $f \circ g = 1_\Lambda$. Now we put $g(\lambda_i) = \sum_{j=1}^t r_{ij} u_j$, $r_{ij} \in R$. Let R_0 be the prime ring of R and R_1 the polynomial ring over R generated by $\{r_{ij}\}$. Then the module generated by $\lambda_0, \lambda_1, \dots, \lambda_t$ over R_1 is R_1 -projective. As Λ is R -separable, defining the Λ^e -epimorphism $\varphi: \Lambda^e \rightarrow \Lambda$ by putting $\varphi(\lambda_i \otimes_R \lambda_j) = \lambda_i \lambda_j$, there is a Λ^e -homomorphism $\psi: \Lambda \rightarrow \Lambda^e$ such that $\varphi \psi = 1$. Put $\psi(\lambda_i) = \sum_{j,k} s_{ijk} (\lambda_j \otimes_R \lambda_k)$, $s_{ijk} \in R$ and $\lambda_i \lambda_j = \sum_k t_{ijk} \lambda_k$, $t_{ijk} \in R$. Furthermore let R' be the polynomial ring over R_0 generated by $\{r_{ij}\}$, $\{s_{ijk}\}$ and $\{t_{ijk}\}$, and denote by Λ' the module over R' generated by $\lambda_0, \lambda_1, \dots, \lambda_t$. Then R' is Noetherian, and Λ' is an R' -algebra which is a finitely generated, faithful, projective R' -module, as R' includes all of $\{r_{ij}\}$ and $\{t_{ijk}\}$. If we define a Λ'^e -epimorphism $\varphi': \Lambda'^e \rightarrow \Lambda'$ by putting $\varphi'(\lambda_i \otimes_{R'} \lambda_j) = \lambda_i \lambda_j$ and we put $\psi'(\lambda_i) = \sum_{j,k} s_{ijk} (\lambda_j \otimes_{R'} \lambda_k)$ for any i , then, from the fact that Λ is R -finitely generated projective, we see easily that ψ' is the well-defined Λ'^e -homomorphism of Λ' into Λ'^e such that $\varphi' \circ \psi' = 1_{\Lambda'}$. Therefore Λ' is a separable R' -algebra. Let α be the R -algebra epimorphism of $R \otimes_{R'} \Lambda'$ onto Λ which is defined by $\alpha(r \otimes_{R'} \lambda_i) = r \lambda_i$, for any $r \in R$. Let \mathfrak{m} be a maximal ideal of R and put $\mathfrak{p}' = \mathfrak{m} \cap R'$. Then we have $(R \otimes_{R'} \Lambda')_{\mathfrak{m}} = R_{\mathfrak{m}} \otimes_{R'_{\mathfrak{p}'}} \Lambda'_{\mathfrak{p}'}$ and so α induces naturally an $R_{\mathfrak{m}}$ -algebra epimorphism $\alpha_{\mathfrak{m}}: R_{\mathfrak{m}} \otimes_{R'_{\mathfrak{p}'}} \Lambda'_{\mathfrak{p}'} \rightarrow \Lambda_{\mathfrak{m}}$. Since $\Lambda'_{\mathfrak{p}'}$ is $R'_{\mathfrak{p}'}$ -free, $\alpha_{\mathfrak{m}}$ must be an isomorphism. From this it follows immediately that α is an isomorphism. Thus our proof is completed.

2. Central separable algebras with proper splitting rings

Let Λ be a central separable R -algebra and S a commutative R -algebra. If there exists a finitely generated faithful projective S -module P such that $S \otimes_R \Lambda \cong \text{Hom}_S(P, P)$ as S -algebras, then S is called, according to [2], the *splitting ring* of Λ . Especially, when $S \supseteq R$, S is called the *proper splitting ring* of Λ .

First we give, as a slight generalization of [2], (6.3),

Proposition 2.1. *Let R be a local ring with a maximal ideal \mathfrak{m} and Λ a central separable R -algebra. Then Λ has a proper splitting ring S which is a*

separable R -algebra and a finitely generated free R -module. Especially, if R is Henselian, then we can choose as S a local ring with a maximal ideal \mathfrak{m}_S .

Proof. By using (1.1) and the Henselization instead of the completion, this can be proved along the same line as in [2], (6.3).

For a central separable algebra over a general commutative ring R , we can not assure the existence of the proper splitting ring which is R -separable and R -finitely generated, projective. In this section, we shall consider only the existence of proper splitting rings. However, we could not prove the existence of a proper splitting ring for a central separable algebra over a general coefficient ring.

Proposition 2.2. *Let R be a commutative ring which is contained in a semi-local ring. Then any central separable R -algebra has a proper splitting ring. Especially, this assumption for R is satisfied by a Noetherian ring or an integral domain.*

Proof. It suffices to prove this proposition in case R is itself a semi-local ring. Let R be a semi-local ring with maximal ideals $\mathfrak{m}_1, \mathfrak{m}_2, \dots, \mathfrak{m}_t$ and put $R' = R_{\mathfrak{m}_1} \oplus R_{\mathfrak{m}_2} \oplus \dots \oplus R_{\mathfrak{m}_t}$. Then $R \subseteq R'$ and $R' \otimes_R \Lambda = \Lambda_{\mathfrak{m}_1} \oplus \Lambda_{\mathfrak{m}_2} \oplus \dots \oplus \Lambda_{\mathfrak{m}_t}$. Accordingly to (2.1), there exists a proper splitting ring S_i of $\Lambda_{\mathfrak{m}_i}$ for any i . If we put $S = S_1 \oplus S_2 \oplus \dots \oplus S_t$, then we have $R \subseteq R' \subseteq S$ and S is a proper splitting ring of Λ , as is required.

As another case, which is not included in (2.2), we have

Proposition 2.3. *Let R be a commutative ring with the total quotient ring K such that any prime ideal of K is maximal. Then any central separable R -algebra has a proper splitting ring.*

Proof. We may assume $R = K$. If we denote by \mathfrak{n} the nil radical of R , then R/\mathfrak{n} is, by our assumption, a regular ring (in the Neumann's sense). Therefore we may further assume that Λ is a finitely generated free R -module. Let $\{u_1, u_2, \dots, u_t\}$ be an R -basis of Λ with $u_1 = 1$, and put $u_i u_j = \sum_{k=1}^t r_{ijk} u_k$, $r_{ijk} \in R$. Let R_0 be the prime ring of R , and put $R' = R_0[\{r_{ijk}\}]$ and $\Omega' = \{r'_1 u_1 + \dots + r'_t u_t \mid r'_i \in R'\}$. Then Ω' is a central R' -algebra with an R' -basis $\{u_1, \dots, u_t\}$, and we have $R \otimes_{R'} \Omega' = \Lambda$. Furthermore let \tilde{R} be the integral closure of R' in R . Since R/\mathfrak{n} is regular, any non-zero divisor of \tilde{R} is a unit in R , and therefore the total quotient ring \tilde{K} of \tilde{R} can be regarded as a subring of R . From the fact that \tilde{R} is integral over R' , we see that the total quotient ring K' of R' is included in R . Since R' is Noetherian and $\tilde{K}/\mathfrak{n} \cap \tilde{K}$ is regular, $K'/\mathfrak{n}K'$ is Artinian, and so K' is itself Artinian. If we put $\Lambda' = K' \otimes_{R'} \Omega'$, then $R' \otimes_{K'} \Lambda' = \Lambda$ and, as K' is Artinian, we can easily see that Λ is a central separable K' -algebra. According to (2.1), there

exists a proper splitting ring F of Λ' which is a finitely generated projective K' -module. Now put $S = F \otimes_{K'} R$. Then $S \supseteq F$, R and $S \otimes_R \Lambda = S \otimes_R R \otimes_{K'} \Lambda' = F \otimes_{K'} R \otimes_{K'} \Lambda' = (F \otimes_{K'} R) \otimes_F F \otimes_{K'} \Lambda'$. Consequently, S is a proper splitting ring of Λ , which completes our proof.

3. The trace and norm of a central separable algebra

1. Let R be a commutative ring and P a finitely generated projective R -module. First suppose that P has (constant) rank n . Then there exists a commutative ring $S \supseteq R$ such that $S \otimes_R P$ is a free S -module of rank n . Let $\{u_1, \dots, u_n\}$ be a S -basis of $S \otimes_R P$. If $f \in \text{Hom}_R(P, P)$, then f can be regarded as an element of $\text{Hom}_S(S \otimes_R P, S \otimes_R P)$, and we can put $f(u_j) = \sum_{i=1}^n u_i s_{ij}'$ for some $s_{ij}' \in S$. Now put $\text{Pc}_P(f: X) = |s_{ij}' - X \delta_{ij}|$, $\text{T}_P(f) = \text{traces}(s_{ij}')$ and $\text{N}_P(f) = |s_{ij}'|$ where X denotes an indeterminate. It can easily be shown by using the localization at any maximal ideal of R that $\text{Pc}_P(f, X) \in R[X]$ and $\text{T}_P(f), \text{N}_P(f) \in R$ and that these are determined without depending on S and $\{u_1, \dots, u_n\}$. If P has not constant rank, there is, by [7], § 2, a unique decomposition $R = R_1 \oplus \dots \oplus R_t$ such that any $R_i \otimes_R P$ has rank n_i over R_i where $n_1 < n_2 < \dots < n_t$, and we have $\text{Hom}_R(P, P) = \sum_{i=1}^t \oplus \text{Hom}_{R_i}(R_i \otimes_R P, R_i \otimes_R P)$. Let f be an element of $\text{Hom}_R(P, P)$ and f_i the i -th component of f . Then we put $\text{Pc}_P(f: X) = \sum_{i=1}^t \oplus \text{Pc}_{R_i \otimes_R P}(f_i: X)$, $\text{T}_P(f) = \sum_{i=1}^t \oplus \text{T}_{R_i \otimes_R P}(f_i)$ and $\text{N}_P(f) = \sum_{i=1}^t \oplus \text{N}_{R_i \otimes_R P}(f_i)$ and we call them the characteristic polynomial, trace and norm of f . It can be easily shown that our definitions coincide with those in [7].

If Λ is an R -algebra which is a finitely generated projective R -module, then we use $\text{Pc}_{\Lambda/R}(f: X)$, $\text{T}_{\Lambda/R}(f)$ and $\text{N}_{\Lambda/R}(f)$ instead of $\text{Pc}_\Lambda(f: X)$, $\text{T}_\Lambda(f)$ and $\text{N}_\Lambda(f)$.

2. Now we shall define the reduced characteristic polynomial, trace and norm for a central separable algebra with a proper splitting ring.

Let Λ be a central separable R -algebra with a proper splitting ring S . Then there exists a S -algebra isomorphism $h_S: S \otimes_R \Lambda \cong \text{Hom}_S(P^{(S)}, P^{(S)})$ for some finitely generated projective S -module $P^{(S)}$.

Proposition 3.1 *For any element λ of Λ , $\text{Pc}_{P^{(S)}}(h_S(\lambda): X)$ is a polynomial of $R[X]$ which does not depend on S , $P^{(S)}$ and h_S .*

Proof. First suppose that R is a local ring. Then Λ is a projective R -module of constant rank, and so $P^{(S)}$ is also a projective S -module of constant rank. By replacing S by any extension ring S' of it and by replacing h_S by $1 \otimes h_S$, $\text{Pc}_{P^{(S)}}(h_S(\lambda): X)$ is invariant, and therefore we may further assume that

$P^{(S)}$ is S -free. Then h_S induces a S -algebra isomorphism $k_S: S \otimes_R \Lambda \cong M_n(S)$ such that $\text{Pc}_{P^{(S)}}(h_S(\lambda): X) = |XE_n - k_S(\lambda)|$. On the other hand, according to (2.1), there exists a proper splitting semi-local ring T of Λ which is R -free. For T we can define, similarly, h_T , $P^{(T)}$ and k_T . Since T is R -free, we have $R \otimes_R R = S \otimes_R R \cap R \otimes_R T$ in $S \otimes_R T$, and so we may suppose that there is a commutative ring U containing both S and T and $S \cap T = R$ in U . Now the algebra isomorphisms $k_S: S \otimes_R \Lambda \cong M_n(S)$ and $k_T: T \otimes_R \Lambda \cong M_n(T)$ can, naturally, be extended to the U -algebra isomorphisms $k_S^*: k_T^*: U \otimes_R \Lambda \cong M_n(U)$. Then $k_S^* \circ k_T^*$ is an U -algebra automorphism of $M_n(U)$ and it induces an U_m -algebra automorphism of $M_n(U_m)$ for any maximal ideal of U . As U_m is a local ring, it is inner, and so we have $|XE_n - k_S^*(\lambda^*)| = |XE_n - k_T^*(\lambda^*)|$ in $U_m[X]$ for any $\lambda^* \in U \otimes_R \Lambda$. Hence we have $\text{Pc}_{P^{(S)}}(h_S(\lambda): X) = |XE_n - k_S(\lambda)| = |XE_n - k_S^*(\lambda)| = |XE_n - k_T^*(\lambda)| = \text{Pc}_{P^{(T)}}(h_T(\lambda): X)$ in $U[X]$. However, as $\text{Pc}_{P^{(S)}}(h_S(\lambda): X) \in S[X]$ and $\text{Pc}_{P^{(T)}}(h_T(\lambda): X) \in T[X]$, we obtain $\text{Pc}_{P^{(S)}}(h_S(\lambda): X) = \text{Pc}_{P^{(T)}}(h_T(\lambda): X) \in R[X] = S[X] \cap T[X]$. Thus $\text{Pc}_{P^{(S)}}(h_S(\lambda): X)$ is a polynomial of $R[X]$. It is obvious from the above proof that this does not depend on S , $P^{(S)}$ and h_S , which completes our proof for a local ring R .

Let R be a general commutative ring and \mathfrak{m} a maximal ideal of R . Denote by $\lambda_{\mathfrak{m}}$ the residue of λ in $\Lambda_{\mathfrak{m}}$ and by $h_{S_{\mathfrak{m}}}$ the $S_{\mathfrak{m}}$ -algebra isomorphism: $S_{\mathfrak{m}} \otimes_{R_{\mathfrak{m}}} \Lambda_{\mathfrak{m}} \cong \text{Hom}_{S_{\mathfrak{m}}}(P_{\mathfrak{m}}^{(S)}, P_{\mathfrak{m}}^{(S)})$ induced by h_S . Further let $[\text{Pc}_{P^{(S)}}(h_S(\lambda): X)]_{\mathfrak{m}}$ be the residue of $\text{Pc}_{P^{(S)}}(h_S(\lambda): X)$ in $S_{\mathfrak{m}}[X]$. Then we see $[\text{Pc}_{P^{(S)}}(h_S(\lambda): X)]_{\mathfrak{m}} = \text{Pc}_{P_{\mathfrak{m}}^{(S)}}(h_{S_{\mathfrak{m}}}(\lambda_{\mathfrak{m}}): X)$. Since, by the preceding argument for a local ring, $\text{Pc}_{P_{\mathfrak{m}}^{(S)}}(h_{S_{\mathfrak{m}}}(\lambda_{\mathfrak{m}}): X) \in R_{\mathfrak{m}}[X]$, we have also $[\text{Pc}_{P^{(S)}}(h_S(\lambda): X)]_{\mathfrak{m}} \in R_{\mathfrak{m}}[X]$. Consequently we obtain $\text{Pc}_{P^{(S)}}(h_S(\lambda): X) \in R[X]$. It is also evident in this case that $\text{Pc}_{P^{(S)}}(h_S(\lambda): X)$ does not depend on S , $P^{(S)}$ and h_S .

Now we denote $\text{Pc}_{P^{(S)}}(h_S(\lambda): X)$ by $\text{Pcrd}_{\Lambda/R}(\lambda: X)$ and we call it the reduced characteristic polynomial of λ . Furthermore, if we put $\text{Trd}_{\Lambda/R}(\lambda) = T_{P^{(S)}}(h_S(\lambda))$ and $\text{Nrd}_{\Lambda/R}(\lambda) = N_{P^{(S)}}(h_S(\lambda))$, then they are elements of R which do not depend on S , $P^{(S)}$ and h_S and we call them the reduced trace and norm of λ , respectively.

From our definitions it follows immediately

Proposition 3.2. *For any $\lambda, \lambda_1, \lambda_2 \in \Lambda$ and any $r \in R$, we have*

$$\begin{aligned} \text{Trd}_{\Lambda/R}(\lambda_1 + \lambda_2) &= \text{Trd}_{\Lambda/R}(\lambda_1) + \text{Trd}_{\Lambda/R}(\lambda_2), \\ \text{Trd}_{\Lambda/R}(r\lambda) &= r \text{Trd}_{\Lambda/R}(\lambda), \\ \text{Trd}_{\Lambda/R}(\lambda_1 \lambda_2) &= \text{Trd}_{\Lambda/R}(\lambda_2 \lambda_1), \\ \text{Nrd}_{\Lambda/R}(\lambda_1 \lambda_2) &= \text{Nrd}_{\Lambda/R}(\lambda_1) \text{Nrd}_{\Lambda/R}(\lambda_2) \end{aligned}$$

Epecially, if Λ has rank n^2 over R , then we have

$$\text{Nrd}_{\Lambda/R}(r\lambda) = r^n \text{Nrd}_{\Lambda/R}(\lambda)$$

From this proposition, it follows that $\text{Trd}_{\Lambda/R}$ is an R -homomorphism of Λ into R and $\text{Nrd}_{\Lambda/R}$ is a semi-group homomorphism of Λ into R as the multiplicative semi-groups.

For any maximal ideal \mathfrak{m} of R , let $\overline{[\text{Prd}_{\Lambda/R}(\lambda: X)]_{\mathfrak{m}}}$ be the residue of $\text{Prd}_{\Lambda/R}(\lambda: X)$ in $(R/\mathfrak{m})[X]$ and denote by $\bar{\lambda}_{\mathfrak{m}}$ the residue of λ in $\Lambda/\mathfrak{m}\Lambda$. Now we can show $\overline{[\text{Prd}_{\Lambda/R}(\lambda: X)]_{\mathfrak{m}}} = \text{Prd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}(\bar{\lambda}_{\mathfrak{m}}: X)$. In fact, it suffices to prove this in case R is a Henselian local ring with a maximal ideal \mathfrak{m} . However, in this case, there is, by (2.1), a proper splitting local ring S of Λ such that $\mathfrak{m}S$ is a maximal ideal of S and S is a finitely generated free R -module. Then $S/\mathfrak{m}S$ becomes the splitting field of the classical central separable R/\mathfrak{m} -algebra $\Lambda/\mathfrak{m}\Lambda$, from which our result follows immediately. Accordingly, $\text{Trd}_{\Lambda/R}$ and $\text{Nrd}_{\Lambda/R}$ induce, naturally, $\text{Trd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}$ and $\text{Nrd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}$, respectively, which coincide with those in the classical sense. By summarizing these, we obtain

Proposition 3.3. *For any maximal ideal \mathfrak{m} of R , the residue of $\text{Prd}_{\Lambda/R}$ in $(R/\mathfrak{m})[X]$ coincides with $\text{Prd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}$. Especially, the residues of $\text{Trd}_{\Lambda/R}$ and $\text{Nrd}_{\Lambda/R}$ in R/\mathfrak{m} coincide with $\text{Trd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}$ and $\text{Nrd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}$, respectively.*

3. Here we shall determine the relations between the trace (norm) and reduced trace (reduced norm) of a central separable algebra, which are given in the same form as in the classical one (cf. [4]).

Assume that Λ is a projective R -module of the constant rank m . Then we may suppose $S \otimes_R \Lambda \cong M_n(S)$, where $m = n^2$. From our definitions, it follows directly that $\text{Trd}_{\Lambda/R}(1) = n$, $T_{\Lambda/R}(\lambda) = n \text{Trd}_{\Lambda/R}(\lambda)$ and $N_{\Lambda/R}(\lambda) = [\text{Nrd}_{\Lambda/R}(\lambda)]^n$. In the general case, let $R = R_1 \oplus \cdots \oplus R_t$ be the unique decomposition of R such that $R_i \otimes_R \Lambda$ has rank m_i over R_i where $m_1 < m_2 < \cdots < m_t$. Then we can put $m_i = n_i^2$ for any i . Let e_i be a unit element of R_i and λ_i the i -th component of λ . Then we obtain

Proposition 3.4. $\text{Trd}_{R_i \otimes \Lambda/R_i}(e_i) = n_i e_i$ for each i ,

$$\begin{aligned} T_{\Lambda/R}(\lambda) &= \text{Trd}_{\Lambda/R}(1) \text{Trd}_{\Lambda/R}(\lambda) = \sum_{i=1}^t n_i \text{Trd}_{R_i \otimes \Lambda/R_i}(\lambda_i) \\ N_{\Lambda/R}(\lambda) &= \sum_{i=1}^t [\text{Nrd}_{R_i \otimes \Lambda/R_i}(\lambda_i)]^{n_i} \end{aligned}$$

The following result will be used in § 4.

Proposition 3.5. $\text{Trd}_{\Lambda/R}$ is an R -epimorphism of Λ onto R .

Proof. By the remark after (3.2), it suffices to prove that $\text{Trd}_{\Lambda/R}$ is an epimorphism. By virtue of the classical result, for any maximal ideal \mathfrak{m} of R , $\text{Trd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}$ is an epimorphism of $\Lambda/\mathfrak{m}\Lambda$ onto R/\mathfrak{m} . According to (2.3), then, $\text{Trd}_{\Lambda/R}$ must be an epimorphism of Λ onto R .

Corollary 3.6. *The complete image $T_{\Lambda/R}(\Lambda)$ of $T_{\Lambda/R}$ is a principal ideal of R generated by $\text{Trd}_{\Lambda/R}(1)$. Especially, Λ is strongly separable if and only if $\text{Trd}_{\Lambda/R}(1)$ is a unit of R .*

Proof. This is an immediate consequence of (3.4) and (3.5).

4. As is remarked in § 2, we could not succeed in proving the existence of a proper splitting ring for a central separable algebra in the general case. Hence we can not define the reduced characteristic polynomial for a central separable algebra in the case where we can not show the existence of a proper splitting ring. However we can define, by using (1.2), the reduced trace for any central separable R -algebra Λ . In fact, by virtue of (1.2), there exist a Noetherian subring R' of R and a central separable R' -algebra Λ' such that $\Lambda = R \otimes_{R'} \Lambda'$. Since Λ' has a proper splitting ring by (2.2), there exists, according to 2, the reduced trace $\text{Trd}_{\Lambda'/R'}: \Lambda' \rightarrow R'$. Now we define the reduced trace $\text{Trd}_{\Lambda/R}: \Lambda \rightarrow R$, by putting $\text{Trd}_{\Lambda/R}(r \otimes \lambda') = r \text{Trd}_{\Lambda'/R'}(\lambda')$ for any $r \in R$ and for any $\lambda' \in \Lambda'$. It can be easily shown that, for any maximal ideal \mathfrak{m} of R , the $R_{\mathfrak{m}}$ -homomorphism $(\text{Trd}_{\Lambda/R})_{\mathfrak{m}}: \Lambda_{\mathfrak{m}} \rightarrow R_{\mathfrak{m}}$, which is induced on $\Lambda_{\mathfrak{m}}$ by $\text{Trd}_{\Lambda/R}$, coincides with the reduced trace $\text{Trd}_{\Lambda_{\mathfrak{m}}/R_{\mathfrak{m}}}$ of $\Lambda_{\mathfrak{m}}$ defined by using the proper splitting ring of $\Lambda_{\mathfrak{m}}$. Especially, if Λ has a proper splitting ring, $\text{Trd}_{\Lambda/R}$ coincides with that defined in 2. Furthermore we can also prove (3.2)~(3.6) in this case.

4. The symmetricity of a separable algebra

Let Λ be an R -algebra, which is a finitely generated, faithful, projective R -module. We shall consider $\Lambda^* = \text{Hom}_R(\Lambda, R)$ as a left Λ^e -module through the operations $(\lambda \cdot f)(\mu) = f(\mu\lambda)$, $(f \cdot \lambda)(\mu) = f(\lambda\mu)$ where $f \in \Lambda^*$, $\lambda, \mu \in \Lambda$. Following [6], we call Λ a *Frobenius R -algebra* if Λ^* is isomorphic to Λ as left (or equivalently right) Λ -modules, and, furthermore, is called a *symmetric R -algebra* if Λ^* is Λ^e -isomorphic to Λ . From our definitions it follows that any symmetric R -algebra is Frobenius.

We begin with

Lemma 4.1. *Let S be a symmetric, commutative R -algebra and Λ a symmetric S -algebra. Then Λ is a symmetric R -algebra.*

Proof. By our assumptions we have $\Lambda \cong \text{Hom}_S(\Lambda, S)$ as two-sided Λ -modules and $S \cong \text{Hom}_R(S, R)$ as S -modules. So we obtain $\text{Hom}_S(\Lambda, S) \cong \text{Hom}_S(\Lambda, \text{Hom}_R(S, R)) \cong \text{Hom}_R(\Lambda \otimes_S S, R) \cong \text{Hom}_R(\Lambda, R)$ as two-sided Λ -modules. This shows that Λ is a symmetric R -algebra.

It is well known, in the classical theory, that a semi-simple algebra over a field is symmetric. However, for any commutative ring R , it is an open question whether a semi-simple R -algebra is symmetric or not,

Now we give, as a partial answer to this question,

Theorem 4.2. *A separable R -algebra Λ , which is a finitely generated, faithful, projective R -module, is a symmetric R -algebra.*

Proof. Let C be the center of Λ . According to [2] (2.1), Λ is a finitely generated projective C -module. By our assumption, Λ is R -finitely generated projective, and so C is also a finitely generated projective R -module, as C is a C -direct summand of Λ . Since, by [2], A.4, a commutative separable R -algebra, which is a finitely generated, faithful, projective R -module, is symmetric, C must be a symmetric R -algebra. Therefore, by (4.1), it suffices to prove our theorem in case $R=C$.

Let Λ be a central separable R -algebra and denote by $\text{Trd}_{\Lambda/R}$ the reduced trace of Λ , defined in § 3. Then $\text{Trd}_{\Lambda/R}$ is a symmetric R -homomorphism of Λ into R : i.e., we have $\text{Trd}_{\Lambda/R}(\lambda\mu) = \text{Trd}_{\Lambda/R}(\mu\lambda)$ for any $\lambda, \mu \in \Lambda$. Hence, putting $\Phi(\lambda)(\mu) = \text{Trd}_{\Lambda/R}(\lambda\mu)$ for any $\lambda, \mu \in \Lambda$, Φ is a Λ^e -homomorphism of Λ into Λ^* . By (3.3), for any maximal ideal \mathfrak{m} of R , $\text{Trd}_{\Lambda/R}$ induces naturally the reduced trace $\text{Trd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}$ in the classical sense on $\Lambda/\mathfrak{m}\Lambda$, and therefore Φ induces, naturally, the $\Lambda^e/\mathfrak{m}\Lambda^e$ -homomorphism $\bar{\Phi}_{\mathfrak{m}}: \Lambda/\mathfrak{m}\Lambda \rightarrow \Lambda^*/\mathfrak{m}\Lambda^* \cong (\Lambda/\mathfrak{m}\Lambda)^*$ such that $\bar{\Phi}_{\mathfrak{m}}(\bar{\lambda})(\bar{\mu}) = \text{Trd}_{\Lambda/\mathfrak{m}\Lambda/R/\mathfrak{m}}(\bar{\lambda}\bar{\mu})$ for any $\bar{\lambda}, \bar{\mu} \in \Lambda/\mathfrak{m}\Lambda$. From the classical result it follows that $\bar{\Phi}_{\mathfrak{m}}$ is a $\Lambda^e/\mathfrak{m}\Lambda^e$ -isomorphism. As both Λ and Λ^* are finitely generated projective R -modules, we can easily see from this that Φ itself is an isomorphism of Λ onto Λ^* . This completes our proof.

We remark that (4.2) was known in some special cases (cf. [2], [5] and [10]).

Finally we give, as an additional remark,

Proposition 4.3. *Let Λ be a central R -algebra which is a finitely generated projective R -module. Then the following statements are equivalent:*

- (1) Λ is a separable R -algebra.
- (2) The R -module $\Lambda/[\Lambda, \Lambda]$ is isomorphic to R , and, for any maximal ideal \mathfrak{m} of R , $\Lambda/\mathfrak{m}\Lambda$ is a semi-simple R/\mathfrak{m} -algebra.

Here we denote by $[\Lambda, \Lambda]$ the R -module generated by all elements of Λ in the form $\lambda\mu - \mu\lambda$, $\lambda, \mu \in \Lambda$.

Proof. (1) \Rightarrow (2): Suppose that Λ is a separable R -algebra. Then the second assertion of (2) follows from [2], (1.6) and so it suffices to prove $\Lambda/[\Lambda, \Lambda] \cong R$. Let $\text{Trd}_{\Lambda/R}$ be the reduced trace of Λ . Then $\text{Trd}_{\Lambda/R}$ is a symmetric R -epimorphism of Λ onto R , and therefore, putting $\text{Ker } \text{Trd}_{\Lambda/R} = K$, we have an R -exact sequence:

$$0 \rightarrow K \rightarrow \Lambda \xrightarrow{\text{Trd}_{\Lambda/R}} R \rightarrow 0$$

and $K \supseteq [\Lambda, \Lambda]$. Hence we have only to show $K = [\Lambda, \Lambda]$. As is shown in § 3,

$\text{Trd}_{\Lambda/R}$ induces naturally the reduced trace $\text{Trd}_{\Lambda/m\Lambda/R/m}$ of $\Lambda/m\Lambda$ for any maximal ideal m of R , and we have $\text{Ker Trd}_{\Lambda/m\Lambda/R/m} = K/mK$. However, it is well known, in the classical theory, that the kernel of the reduced trace of a central separable R/m -algebra $\Lambda/m\Lambda$ coincides with $[\Lambda/m\Lambda, \Lambda/m\Lambda]$. Consequently we must have $K/mK = [\Lambda/m\Lambda, \Lambda/m\Lambda]$ for any maximal ideal m of R . From this we easily see $K = [\Lambda, \Lambda]$, as K is R -finitely generated. Thus the implication (1) \Rightarrow (2) is proved. (2) \Rightarrow (1). Conversely suppose (2). By (1.1) it suffices to prove that $\Lambda/m\Lambda$ has R/m as its center. By our assumption we have an R -exact sequence:

$$0 \rightarrow [\Lambda, \Lambda] \rightarrow \Lambda \xrightarrow{\alpha} R \rightarrow 0.$$

This induces an R/m -exact sequence:

$$0 \rightarrow [\Lambda, \Lambda]/m[\Lambda, \Lambda] \rightarrow \Lambda/m\Lambda \xrightarrow{\bar{\alpha}} R/mR \rightarrow 0.$$

and so we have $[\Lambda, \Lambda]/m[\Lambda, \Lambda] \cong [\Lambda/m\Lambda, \Lambda/m\Lambda]$. Therefore we have $\Lambda/m\Lambda \cong [\Lambda/m\Lambda, \Lambda/m\Lambda] \oplus R/m$. On the other hand, since $\Lambda/m\Lambda$ is R/m -semi-simple, $\Lambda/m\Lambda$ is separable over its center \bar{C} , and then we have $\Lambda/m\Lambda \cong [\Lambda/m\Lambda, \Lambda/m\Lambda] \oplus \bar{C}$. As $\bar{C} \supseteq R/m$, we see from these that \bar{C} coincides with R/m . This completes our proof.

TOKYO UNIVERSITY OF EDUCATION

References

- [1] M. Auslander and D.A. Buchsbaum: *On ramification theory in Noetherian rings*, Amer. J. Math. **81** (1959), 749–765.
- [2] M. Auslander and O. Goldman: *The Brauer group of a commutative ring*, Trans. Amer. Math. Soc. **97** (1960), 367–409.
- [3] G. Azumaya: *On maximally central algebras*, Nagoya Math. J. **2** (1951), 119–150.
- [4] N. Bourbaki: *Algèbre*, Chap. 8, Hermann, Paris, 1958.
- [5] F.R. DeMeyer: *The trace map and separable algebras*, Osaka J. Math. **3** (1966), 7–11.
- [6] S. Eilenberg and T. Nakayama: *On the dimension of modules and algebras*, II, Nagoya Math. J. **9** (1955), 1–16.
- [7] O. Goldman: *Determinants in projective modules*, Nagoya Math. J. **18** (1961), 27–36.
- [8] A. Hattori: *Semisimple algebras over a commutative ring*, J. Math. Soc. Japan **15** (1963), 404–419.
- [9] T. Kanzaki: *Special type of separable algebras over a commutative ring*, Proc. Japan Acad. **40** (1964), 781–786.
- [10] B. Müller: *Quasi-Frobenius-Erweiterungen*, II, Math. Z. **88** (1965), 380–409.
- [11] M. Nagata: *Local Rings*, Interscience, 1962.