

Title	Stochastic calculus related to nonsymmetric Dirichlet forms
Author(s)	Kim, Jai Heui
Citation	Osaka Journal of Mathematics. 1987, 24(2), p. 331-371
Version Type	VoR
URL	<a href="https://doi.org/10.18910/11837">https://doi.org/10.18910/11837</a>
rights	
Note	

***Osaka University Knowledge Archive : OUKA***

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## STOCHASTIC CALCULUS RELATED TO NON-SYMMETRIC DIRICHLET FORMS

JAI HEUI KIM

(Received April 8, 1986)

### 0. Introduction

The theory of symmetric Dirichlet spaces and the probabilistic potential theory built on Hunt processes were unified by M. Fukushima [8], M.L. Silverstein [17] and others (see References in [8]). In particular, analysis based on additive functionals (AF's) and stochastic calculus related to symmetric Dirichlet spaces were developed by M. Fukushima [8], M. Fukushima and M. Takeda [9], S. Nakao [15] and M. Takeda [20]. On the other hand, the theory of non-symmetric Dirichlet spaces was studied by J. Bliedtner [3, 4], H. Kunita [10] etc.. Furthermore S. Carrillo Menendez [5] constructed the Hunt process associated with a non-symmetric Dirichlet space. Then many results in the symmetric case have been extended to the non-symmetric case by Y. Le Jan [11, 12], M.L. Silverstein [18], S. Carrillo Menendez [6] etc.. The purpose of this paper is to extend those results in [8], [9] and [20] to the non-symmetric case and thereby enlarge the range of applications of Dirichlet space theory.

### 1. Summary of the results

We first give a precise definition of non-symmetric Dirichlet form. Let  $X$  be a locally compact Hausdorff space with countable base and  $m$  a non-negative Radon measure on  $X$  such that  $\text{supp}[m]=X$ .  $L^2(X, m)$  denotes the real  $L^2$ -space with inner product

$$(u, v)_{L^2} = \int_X u(x) v(x) m(dx), \quad u, v \in L^2(X, m).$$

Let  $\mathbf{H}$  be a dense linear subspace of  $L^2(X, m)$  which forms a Hilbert space with a norm  $\|\cdot\|_{\mathbf{H}}$  such that for some  $K>0$ ,  $\|u\|_{\mathbf{H}} \geq K\|u\|_{L^2}$  for any  $u \in \mathbf{H}$ . Moreover we assume that if  $u \in \mathbf{H}$ , then  $|u|, u \wedge 1 \in \mathbf{H}$ . In this article we consider a bilinear form  $\mathbf{a}$  on  $\mathbf{H} \times \mathbf{H}$  which satisfies the following conditions;

(a.1)  $\mathbf{a}_\alpha$  is coercive for any  $\alpha>0$ , i.e., there exists a constant  $K_1=K_1(\alpha)>0$  such that  $\mathbf{a}_\alpha(u, u) \geq K_1\|u\|_{\mathbf{H}}^2$  for every  $u \in \mathbf{H}$ ,

(a.2)  $\mathbf{a}$  is continuous in the following sense; there exists a constant  $K_2 > 0$  such that  $|\mathbf{a}(u, v)| \leq K_2 \|u\|_H \|v\|_H$  for every  $u, v \in H$ ,

(a.3)  $\mathbf{a}(T_1 u, u - T_1 u) \geq 0$ ,

(a.4)  $\hat{\mathbf{a}}(T_1 u, u - T_1 u) \geq 0$ .

Here  $\mathbf{a}_s(u, v) = \mathbf{a}(u, v) + \alpha(u, v)_L$ ,  $\hat{\mathbf{a}}(u, v) = \mathbf{a}(v, u)$  and  $T_1 u = u^+ \wedge 1$  ( $u^+ = u \vee 0$ ). A bilinear form  $\mathbf{a}$  which fulfills (a.1)~(a.4) is called a *Dirichlet form* on  $H \times H$ , and  $(\mathbf{a}, H)$  a *Dirichlet space* on  $L^2(X, m)$ . Let us note that, under the conditions (a.1) and (a.2), the conditions (a.3) and (a.4) are equivalent to the next ones respectively (see [10]);

(a.3)'  $\mathbf{a}(u + T_1 u, u - T_1 u) \geq 0$

and

(a.4)'  $\hat{\mathbf{a}}(u + T_1 u, u - T_1 u) \geq 0$ ,

In particular, if  $\mathbf{a}$  is symmetric, (a.3)' reduces to the usual sub-Markov condition for a symmetric form that  $\mathbf{a}(u, u) \geq \mathbf{a}(T_1 u, T_1 u)$ . Let  $C_0(X)$  be the set of all bounded continuous functions on  $X$  with compact support. From now on we assume that  $H$  is *regular*, i.e.,  $C_0(X) \cap H$  is dense in  $H$  with  $\|\cdot\|_H$  and dense in  $C_0(X)$  with uniform norm.

Let us now summarize the results and methods in the present paper.

In Section 2 we give some basic notions and the Beurling-Deny formula for  $\mathbf{a}$ ; for  $u, v \in C_0(X) \cap H$ ,

$$(1.1) \quad \begin{aligned} \frac{1}{2} \mathbf{a}(u, v) + \frac{1}{2} \hat{\mathbf{a}}(u, v) &= N(u, v) \\ &+ \frac{1}{2} \int_{X \times X - \Delta} (u(x) - u(y)) (v(x) - v(y)) \sigma(dx, dy) \\ &+ \frac{1}{2} \int_X u(x) v(x) \chi(dx) + \frac{1}{2} \int_X u(x) v(x) \hat{\chi}(dx). \end{aligned}$$

Here  $N$  is a local symmetric form on  $C_0(X) \cap H$ ;  $N(u, v) = 0$  if  $v$  is constant on a neighbourhood of  $\text{supp}[u]$  (=support of  $u$ ).  $\sigma$  is a positive Radon measure on  $X \times X - \Delta$  ( $\Delta$  is diagonal) satisfying

$$\int_{X \times X - \Delta} u(x) v(y) \sigma(dx, dy) = -\mathbf{a}(u, v)$$

for  $u, v \in C_0(X) \cap H$  such that  $\text{supp}[u] \cap \text{supp}[v] = \emptyset$ .  $\chi$  (resp.  $\hat{\chi}$ ) is a positive Radon measure on  $X$  satisfying

$$\begin{aligned} \int_X u(x) \chi(dx) &= \mathbf{a}(v, u) - \int_{X \times X - \Delta} u(y) (1 - v(x)) \sigma(dx, dy) \\ (\text{resp. } \int_X u(x) \hat{\chi}(dx) &= \hat{\mathbf{a}}(v, u) - \int_{X \times X - \Delta} u(x) (1 - v(y)) \sigma(dx, dy)) \end{aligned}$$

for  $u, v \in C_0(X) \cap H$  such that  $v = 1$  on a neighbourhood of  $\text{supp}[u]$  (cf. [4]).

Moreover, in Corollary 2.16, we show that the representation (1.1) of  $\mathbf{a}$  can be extended to  $u, v \in \mathbf{H}$  (cf. [12] and [18]).

Let  $\mathbf{M}=(P_x, X_t)$  and  $\hat{\mathbf{M}}=(\hat{P}_x, \hat{X}_t)$  be the Hunt processes associated with  $\mathbf{a}$  and  $\hat{\mathbf{a}}$ , respectively (see [5]). Let  $\mathbf{S}$  be the set of all smooth measures of  $\mathbf{M}$  and  $\mathbf{A}_c^+$  the set of all positive continuous AF's of  $\mathbf{M}$ . S. Carrillo Menendez [6] showed that  $\mathbf{S}$  and  $\mathbf{A}_c^+$  are in one to one correspondence which is characterized by the following relation; for  $A \in \mathbf{A}_c^+$  and  $\mu \in \mathbf{S}$ ,

$$(1.2) \quad E_{h, \mathbf{m}} \left[ \int_0^t f(X_s) dA_s \right] = \int_0^t \langle f \mu, \hat{p}_s h \rangle ds$$

for any non-negative Borel functions  $f$  and  $h$  on  $X$  and  $t > 0$  (see [8] for the symmetric case). Let the couple  $(H, N(y, dx))$  be a Levy system of  $\mathbf{M}$  (see A. Benveniste and J. Jacod [2]) and  $\nu$  the smooth measure corresponding to  $H \in \mathbf{A}_c^+$ . In the same way we can consider  $\hat{N}, \hat{\nu}$ . In Section 3 we show that  $\sigma(dx, dy) = N(y, dx) \nu(dy)$ ,  $\chi(dx) = N(y, \delta) \nu(dy)$  and  $\hat{\chi}(dy) = \hat{N}(y, \delta) \hat{\nu}(dy)$  (see Theorem 3.8 and Remark 3.9).

In Section 4, for any AF  $A$ , we define

$$(1.3) \quad e(A) = \lim_{\alpha \rightarrow \infty} \frac{\alpha^2}{2} E_{\mathbf{m}} \left[ \int_0^\infty e^{-\alpha t} A_t^2 dt \right]$$

if the limit exists.  $e(A)$  is called the energy of  $A$ . We define the mutual energy of AF's  $A$  and  $B$  by  $e(A, B) = 1/2 (e(A+B) - e(A) - e(B))$ . Then we show that  $(\mathring{\mathcal{M}}, e)$  is a real Hilbert space, where  $\mathring{\mathcal{M}}$  is the set of all martingale AF's of finite energy, and that the AF

$$A_t^{[u]} = \tilde{u}(X_t) - \tilde{u}(X_0) \quad (\tilde{u} \text{ is a q.c. version of } u \in \mathbf{H})$$

has a unique decomposition

$$(1.4) \quad A^{[u]} = M^{[u]} + N^{[u]}, \quad M^{[u]} \in \mathring{\mathcal{M}}, \quad N^{[u]} \in \mathcal{N}_e,$$

where  $\mathcal{N}_e$  is the set of all continuous AF's of zero energy. M. Fukushima [8] proved the above results in the symmetric case. Moreover we show that

$$(1.5) \quad e(A^{[u]}) = \mathbf{a}(u, u) - \frac{1}{2} \langle \hat{u}^2, \hat{\chi} \rangle.$$

M. Fukushima [8] defined the energy of AF  $A$  by

$$(1.6) \quad \bar{e}(A) = \lim_{t \downarrow 0} \frac{1}{2t} E_{\mathbf{m}} [A_t^2]$$

In the symmetric case it holds that

$$(1.7) \quad \lim_{t \downarrow 0} \frac{1}{t} (u - p_t u, u)_{L^2} = \mathbf{a}(u, u)$$

and hence

$$(1.8) \quad \bar{e}(A^{[u]}) = \mathbf{a}(u, u) - \frac{1}{2} \langle \tilde{u}^2, \hat{\chi} \rangle$$

(see Remark 4.6). But, in the non-symmetric case, we do not know the validity of (1.7), and hence we can not prove (1.8). It is known that  $M^{[u]}$  has a decomposition;

$$M^{[u]} = \overset{c}{M}^{[u]} + \overset{d}{M}^{[u]} = \overset{c}{M}^{[u]} + \overset{j}{M}^{[u]} + \overset{k}{M}^{[u]}$$

where  $\overset{c}{M}^{[u]}$  and  $\overset{d}{M}^{[u]}$  are the continuous and purely discontinuous part of  $M^{[u]}$  respectively, and  $\overset{j}{M}^{[u]}$  and  $\overset{k}{M}^{[u]}$  are defined by

$$\overset{k}{M}^{[u]} = -\widehat{\tilde{u}(X_{\zeta-}) I_{\{\zeta \leq t\}}}, \quad \overset{j}{M}^{[u]} = \overset{d}{M}^{[u]} - \overset{k}{M}^{[u]}.$$

Here,  $\zeta$  is the life time of  $\mathbf{M}$  and for AF  $A$ ,  $\widehat{A}$  denotes  $A - A^p$  with  $A^p$  being the dual predictable projection of  $A$  (cf. [7] and [9]). The smooth measure corresponding to  $A \in \mathcal{A}_c^+$  is denoted by  $\mu_A$ . If  $A_t^{(1)} = A_t^{(1)}(\omega) - A_t^{(2)}(\omega)$ ,  $A^{(1)}, A^{(2)} \in \mathcal{A}_c^+$  and if  $\mu_{A^{(i)}}$ ,  $i=1, 2$ , are bounded measures, then the measure  $\mu_A$  corresponding to  $A$  is given by  $\mu_{A^{(1)}} - \mu_{A^{(2)}}$ . In particular, for  $M^{[u]}, M^{[v]} \in \dot{\mathcal{M}}$  ( $u, v \in \mathbf{H}$ ), we use the abbreviations  $\mu_{\langle u, v \rangle}$  and  $\overset{\alpha}{\mu}_{\langle u, v \rangle}$  for  $\mu_{\langle M^{[u]}, M^{[v]} \rangle}$  and  $\mu_{\langle \overset{\alpha}{M}^{[u]}, \overset{\alpha}{M}^{[v]} \rangle}$ ,  $\alpha=c, d, j, k$ , respectively. The symbol  $\langle M, N \rangle$  is the quadratic variation of  $M, N \in \dot{\mathcal{M}}$ . In Theorem 4.9 and its collorary, we prove that for  $u, v \in \mathbf{H}$ ,

$$(1.9) \quad \overset{j}{\mu}_{\langle u, v \rangle}(dy) = \int_x (\tilde{u}(x) - \tilde{u}(y)) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy),$$

$$(1.10) \quad \overset{k}{\mu}_{\langle u, v \rangle}(dy) = \tilde{u}(y) \tilde{v}(y) \chi(dy)$$

and

$$(1.11) \quad \frac{1}{2} \overset{c}{\mu}_{\langle u, v \rangle}(X) = N(u, v).$$

In Section 5 we prove the derivation property of  $\overset{c}{\mu}_{\langle u, v \rangle}$ ; for  $u, v \in \mathbf{H}_b$  (=the set of the essentially bounded functions in  $\mathbf{H}$ ) and  $w \in \mathbf{H}$ ,

$$(1.12) \quad d\overset{c}{\mu}_{\langle uv, w \rangle}(x) = \tilde{u}(x) d\overset{c}{\mu}_{\langle v, w \rangle}(x) + \tilde{v}(x) d\overset{c}{\mu}_{\langle u, w \rangle}(x) \quad (x \in X).$$

This formula was already proved by Y. Le Jan [12] (also cf. [8]), but here we give an alternative proof based on the martingale theory. As an application of the equality (1.12), we prove a stronger local property (see Corollary 5.10) of the symmetric form  $N$ .

In the beginning of Section 6 we introduce the notion of general Dirichlet form due to H. Kunita [10], for the condition (a.1) is too restrictive to give

interesting examples. This notion is a little more general than the preceding definition of Dirichlet form in the sense that the coercivity condition (a.1) for  $\mathbf{a}_\alpha$  is only assumed for  $\alpha$  large enough. The basic analytical results on the Dirichlet form are known to hold for the general Dirichlet form with minor modifications ([4], [12]). It is also known that any regular general Dirichlet space admits an associated Hunt process ([5]). Further the arguments in the present paper only involve  $\mathbf{a}_\alpha$  for sufficiently large  $\alpha$ . Therefore all of the present assertions for the Dirichlet form persist to hold for the general Dirichlet form. We will give several examples of general Dirichlet forms which illustrate the usefulness of the theory of non-symmetric Dirichlet space.

As the first example we exhibit a class of bilinear forms given by

$$(1.13) \quad \mathbf{a}^0(u, v) = \sum_{i,j=1}^d \int_D a_{ij} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx - \sum_{i=1}^d \int_D b_i v \frac{\partial u}{\partial x_i} dx + \int_D c u v dx$$

for  $u, v \in H^1(D)$ . Here  $D$  is an open set of  $R^d$ ;  $H^1(D)$  is the Sobolev space of order 1; the coefficients  $a_{ij}$ ,  $b_i$ ,  $i, j=1, 2, \dots, d$ , and  $c$  are bounded measurable functions on  $D$  satisfying certain additional conditions. Moreover we show that this class contains the form corresponding to the differential operator  $L$ , treated by H. Osada [16], given formally by

$$L = L^0 + \sum_{i \neq j}^d \delta(x_i - x_j) \frac{\partial}{\partial x_i},$$

where  $L^0$  is the generator corresponding to  $\mathbf{a}^0$ .

Next, we consider the case with boundary conditions in an orthant. Let  $D = \{x = (\xi, x_d) \in R^d: \xi \in R^{d-1}, x_d > 0\}$ ,  $\partial D = \{x \in R^d: x_d = 0\}$  and for  $u, v \in H^1(D)$ ,

$$(1.14) \quad \mathbf{a}(u, v) = \mathbf{a}^0(u, v) - \sum_{i=1}^{d-1} \int_{\partial D} \beta_i(\xi) v(\xi, 0) \frac{u(\xi, 0)}{\partial \xi_i} d\xi,$$

where  $\beta_i$ ,  $i=1, 2, \dots, d-1$ , are functions on  $\partial D$  with bounded derivatives of first order and satisfying certain additional conditions of the same type as for  $b_i$ . Then, in Theorem 6.2, we show that  $\mathbf{a}$  is a general Dirichlet form on  $\bar{D} = D \cup \partial D$ . If the generator corresponding to  $\mathbf{a}^0$  is the Laplacian, then the process associated with  $\mathbf{a}$  given by (1.14) is a Brownian motion with oblique reflection on  $\bar{D}$ .

Finally we give an example of non-local form which is a slight generalization of Example IV. 3.2 in [5].

## 2. Basic notions and the Beurling-Deny formula

Let  $X$ ,  $m$  and  $(\mathbf{a}, H)$  be given as in Section 1.

DEFINITION 2.1. The generator  $(L, \mathcal{D}(L))$  of  $\mathbf{a}$  is the operator from  $\mathcal{D}(L)$

$= \{u \in \mathbf{H}; \text{ there exists } w \in L^2(X, m) \text{ such that } \mathbf{a}(u, v) = (-w, v)_{L^2} \text{ for all } v \in \mathbf{H}\}$   
to  $L^2(X, m)$  and defined by

$$Lu = w \quad \text{for } u \in \mathcal{D}(L).$$

We see that  $(L, \mathcal{D}(L))$  is the infinitesimal generator of a strongly continuous contraction semigroup  $(T_t)_{t>0}$  of operators on  $L^2(X, m)$ . This is contained in the following theorem.

**Theorem 2.2** (H. Kunita [10]). *For each  $\alpha > 0$ , there exists a uniquely determined bounded operator  $G_\alpha$  from  $L^2(X, m)$  to  $\mathbf{H}$  such that*

- (i)  $\mathbf{a}_\alpha(G_\alpha u, v) = (u, v)_{L^2} \quad (u \in L^2(X, m), v \in \mathbf{H})$ ,
- (ii)  $G_\alpha - G_\beta + (\alpha - \beta) G_\alpha G_\beta = 0$  for any  $\alpha, \beta > 0$ ,
- (iii)  $\|\alpha G_\alpha u\|_{L^2} \leq \|u\|_{L^2} \quad (u \in L^2(X, m))$ ,
- (iv)  $\lim_{\alpha \rightarrow \infty} \alpha G_\alpha u = u$  in  $\mathbf{H}$  as in  $L^2(X, m)$ ,
- (v)  $(\alpha I - L)^{-1} = G_\alpha$
- (vi)  $\alpha G_\alpha$  is sub-Markovian, i.e.,  $0 \leq \alpha G_\alpha u \leq 1$   $m$ -a.e. whenever  $u \in L^2(X, m)$ ,  $0 \leq u \leq 1$   $m$ -a.e..

The family  $(G_\alpha)_{\alpha>0}$  is called the *resolvent* of  $\mathbf{a}$ . By the same method we can define the *co-resolvent*  $(\hat{G}_\alpha)_{\alpha>0}$  of  $\hat{\mathbf{a}}$  such that

$$\hat{\mathbf{a}}_\alpha(\hat{G}_\alpha v, u) = \mathbf{a}_\alpha(u, \hat{G}_\alpha v) = (u, v)_{L^2}.$$

REMARK 2.3. Let

$$T_t u = \lim_{\alpha \rightarrow \infty} e^{-\alpha t} \sum_{n=0}^{\infty} \frac{(\alpha t)^n}{n!} (\alpha G_\alpha)^n u, \quad u \in L^2(X, m).$$

Then  $(T_t)_{t>0}$  determines a strongly continuous contraction sub-Markovian semigroup of  $\mathbf{a}$ .

**DEFINITION 2.4.** Let  $\Gamma$  be a non-empty convex closed subset of  $\mathbf{H}$ .  $u$  is called the  $\mathbf{a}_\alpha$ -*projection* of  $w \in \mathbf{H}$ , written by  $u = \Pi_\Gamma^{\mathbf{a}_\alpha}(w)$ , if  $u \in \Gamma$  and  $\mathbf{a}_\alpha(u - w, v - u) \geq 0$  for any  $v \in \Gamma$  ( $\alpha > 0$ ).

**DEFINITION 2.5.** A positive Radon measure  $\mu$  on  $X$  is said to be of *finite energy integral* if

$$\int_X |v(x)| \mu(dx) \leq c \|v\|_{\mathbf{H}} \quad (v \in C_0(X) \cap \mathbf{H})$$

for some constant  $c > 0$ .

We denote by  $\mathbf{S}_0$  the family of all positive Radon measures of finite energy integral.  $\mu \in \mathbf{S}_0$  if and only if there exists for each  $\alpha > 0$  a unique function

$U_\alpha \mu \in \mathbf{H}$  such that

$$\mathbf{a}_\alpha(U_\alpha \mu, v) = \int_X v(x) \mu(dx) \quad (v \in C_0(X) \cap \mathbf{H}).$$

We call  $U_\alpha \mu$  an  $\mathbf{a}_\alpha$ -potential. Similarly we can define an  $\hat{\mathbf{a}}_\alpha$ -potential  $\hat{U}_\alpha \mu$ .

REMARK 2.6. By Stampacchia's representation theorem (see [3]), there exist the  $\mathbf{a}_\alpha$ -projection and the  $\mathbf{a}_\alpha$ -potential.

Let  $A$  be an open set of  $X$  and define a closed convex subset  $\mathbf{H}_A$  of  $\mathbf{H}$  by

$$\mathbf{H}_A = \{v \in \mathbf{H}: v \geq 1 \text{ m-a.e. on } A\}.$$

If  $\mathbf{H}_A \neq \phi$ , then there exists  $u_A^\alpha = \Pi_{\mathbf{H}_A}^{\mathbf{a}_\alpha}(0) \in \mathbf{H}_A$  and  $u_A^\alpha$  is an  $\mathbf{a}_\alpha$ -potential ( $\alpha > 0$ ).

DEFINITION 2.7. We define the *capacity* of an open set  $A \subset X$  as the number

$$\text{cap}(A) = \begin{cases} \infty & \text{if } \mathbf{H}_A = \phi \\ \mathbf{a}_1(u_A^1, u_A^1) & \text{if } \mathbf{H}_A \neq \phi. \end{cases}$$

$u_A^1$  is called the *capacitary potential* of  $A$ . And for any subset  $E \subset X$  we define

$$\text{cap}(E) = \begin{cases} \infty & \text{if } \mathbf{H}_E = \phi \\ \mathbf{a}_1(u_E^1, u_E^1) & \text{if } \mathbf{H}_E \neq \phi, \end{cases}$$

where

$$\mathbf{H}_E = \overline{\bigcup_{\substack{A \subset E \\ A \text{ is open}}} \mathbf{H}_A} \quad \text{and} \quad u_E^1 = \Pi_{\mathbf{H}_E}^{\mathbf{a}_1}(0).$$

Let  $\mathcal{O}(E)$  be the set of open sets containing  $E$ . S. Carrillo Menendez [5] has proved that if there exists  $A \in \mathcal{O}(E)$  such that  $\mathbf{H}_A \neq \phi$ , then  $\text{cap}(E) = \lim_{A \in \mathcal{O}(E)} \text{cap}(A)$ . It is also known that if  $A$  and  $B$  are open sets and  $A \subset B$ , then  $\text{cap}(A) \leq \text{cap}(B)$  (see [3]). Then it follows that if  $E \subset F$ , then  $\text{cap}(E) \leq \text{cap}(F)$  and hence  $\text{cap}(E) = \inf_{\substack{A \subset E \\ A \text{ is open}}} \text{cap}(A)$ .

The *co-capacity*  $\hat{\text{cap}}(E)$  of  $E \subset X$  is defined by using  $\hat{\mathbf{a}}$  instead of  $\mathbf{a}$ .

Set

$$\mathbf{S}_0 = \{\mu \in \mathbf{S}_0: \mu(X) = 1, \|U_1 \mu\|_\infty < \infty\},$$

and

$$\hat{\mathbf{S}}_0 = \{\mu \in \mathbf{S}_0: \mu(X) = 1, \|\hat{U}_1 \mu\|_\infty < \infty\}.$$

The following lemma can be proved by the same method as Theorem 3.3.2 in [8].



**Lemma 2.8.** *The following conditions are equivalent for a Borel set  $A \subset X$ ;*

- (i)  $\text{cap}(A) = 0$ ,
- (ii)  $\text{c}\hat{\text{a}}\text{p}(A) = 0$ ,
- (iii)  $\mu(A) = 0$  for any  $\mu \in \mathbf{S}_0$ ,
- (iv)  $\mu(A) = 0$  for any  $\mu \in \mathbf{S}_{00}$ ,
- (v)  $\mu(A) = 0$  for any  $\mu \in \hat{\mathbf{S}}_{00}$ .

Let  $A$  be a subset of  $X$ . A statement depending on  $x \in X$  is said to hold q.e. on  $A$  if there exists a set  $N \subset A$  of zero capacity such that the statement is true for every  $x \in X - N$ . "q.e." is an abbreviation of "quasi-everywhere".

**DEFINITION 2.9.** An extended real valued function  $f$  on  $X$  is called *quasi-continuous* (q.c.) if for any  $\varepsilon > 0$  there exists an open set  $A$  such that

- (i)  $\text{cap}(A) < \varepsilon$
- (ii)  $f|_{X-A}$  is continuous on  $X - A$ .

**REMARK 2.10.** In the same way as in the symmetric case [8] it is shown that any element of  $\mathbf{H}$  has a q.c. version.

Let

$$\mathbf{a}^{(\alpha)}(u, v) = \alpha(u - \alpha G_\alpha u, v)_{L^2}$$

for any  $\alpha > 0$  and  $u, v \in L^2(X, m)$ .

**Lemma 2.11** (H. Kunita [10]). (i) *Let  $u \in L^2(X, m)$ . Then  $u \in \mathbf{H}$  if and only if  $\sup_{\alpha > 0} \mathbf{a}^{(\alpha)}(u, u) < \infty$ .*

- (ii) *For any  $u, v \in \mathbf{H}$ ,  $\lim_{\alpha \rightarrow \infty} \mathbf{a}^{(\alpha)}(u, v) = \mathbf{a}(u, v)$ .*

Let

$$\bar{\mathbf{a}}(u, v) = \frac{1}{2} (\mathbf{a}(u, v) + \hat{\mathbf{a}}(u, v)) \quad (u, v \in \mathbf{H}).$$

Clearly  $(\bar{\mathbf{a}}, \mathbf{H})$  is a symmetric Dirichlet space and it holds that  $\bar{\mathbf{a}}(u, u) = \mathbf{a}(u, u) = \hat{\mathbf{a}}(u, u)$ . Hence we have the following lemma (see [8]).

**Lemma 2.12.** (i) *If  $u \in \mathbf{H}$ , then  $Tu \in \mathbf{H}$  and  $\mathbf{a}(u, u) \geq \mathbf{a}(Tu, Tu)$  for any normal contraction  $T$ .*

- (ii) *If  $u, v$  are bounded functions in  $\mathbf{H}$ , then  $uv \in \mathbf{H}$  and  $\mathbf{a}(uv, uv)^{1/2} \leq \|u\|_\infty \mathbf{a}(v, v)^{1/2} + \|v\|_\infty \mathbf{a}(u, u)^{1/2}$ .*

(iii) *If  $u \in \mathbf{H}$  and  $u_n = (-n \vee u) \wedge n$ , then  $u_n \in \mathbf{H}$  and  $u_n \rightarrow u$  as  $n \rightarrow \infty$  with respect to  $\|\cdot\|_{\mathbf{H}}$ .*

- (iv) *If  $u \in \mathbf{H}$  and  $u^{(\varepsilon)} = u - ((-\varepsilon) \vee u) \wedge \varepsilon$  ( $\varepsilon > 0$ ), then  $u^{(\varepsilon)} \in \mathbf{H}$  and  $u^{(\varepsilon)} \rightarrow u$  as  $\varepsilon \rightarrow 0$  with respect to  $\|\cdot\|_{\mathbf{H}}$ .*

- (v) *Let  $D$  be an open set. Then  $C_0(D) \cap \mathbf{H}$  is dense in  $\mathbf{H}^D$  with  $\|\cdot\|_{\mathbf{H}}$ , where*

$$\mathbf{H}^D = \{v \in \mathbf{H}: v=0 \text{ q.e. on } D^c\}.$$

**Lemma 2.13** ([8]). *If the resolvent  $(G_\alpha)_{\alpha>0}$  is positive, then there exists a unique measure  $\sigma_\alpha$  on  $X \times X$  such that*

$$(2.1) \quad \alpha^2 \int_X G_\alpha u(x) v(x) m(dx) = \int_{X \times X} u(x) v(y) \sigma_\alpha(dx, dy)$$

for every Borel functions  $u, v \in L^2(X, m)$ .

Using Lemma 2.11, we can define a unique Radon measure  $\sigma$  on  $X \times X - \Delta$  such that  $\sigma_\alpha \rightarrow \sigma$  vaguely on  $X \times X - \Delta$  as  $\alpha \rightarrow \infty$  and

$$(2.2) \quad \int_{X \times X - \Delta} u(x) v(y) \sigma(dx, dy) = -\mathbf{a}(u, v)$$

for any  $u, v \in C_0(X) \cap \mathbf{H}$  such that  $\text{supp}[u] \cap \text{supp}[v] = \emptyset$ . By the same method, we can define the measures  $\hat{\sigma}_\alpha$  and  $\hat{\sigma}$  if we replace  $G_\alpha$  and  $\mathbf{a}$  by  $\hat{G}_\alpha$  and  $\hat{\mathbf{a}}$  in (2.1) and (2.2). It is easy to see that  $\sigma(dx, dy) = \hat{\sigma}(dy, dx)$ .

**Lemma 2.14.** (i) *For any relatively compact open set  $D \subset X$ , there exists a unique Radon measure  $\chi_D$  on  $X$ , supported in  $D$ , such that for every  $u \in C_0(D)$*

$$(2.3) \quad \int_X u(x) \chi_D(dx) = \lim_{\alpha \rightarrow \infty} \alpha \int_X u(x) (1 - \alpha G_\alpha I_D(x)) m(dx),$$

and it holds that if  $D'$  is a relatively compact open set such that  $D \subset D'$ , then  $\chi_D \leq \chi_{D'}$  on  $D$ .

(ii) *There exists a unique Radon measure  $\chi$  on  $X$  such that for any  $u \in C_0(X)$ ,*

$$(2.4) \quad \int_X u(x) \chi(dx) = \lim_{D \uparrow X} \int_X u(x) \chi_D(dx).$$

Moreover, if  $u, v \in C_0(X) \cap \mathbf{H}$  and  $v=1$  on a neighbourhood of  $\text{supp}[u]$ , then the measure  $\chi$  satisfies the equality

$$(2.5) \quad \int_X u(x) \chi(dx) = \mathbf{a}(v, u) - \int_{X \times X - \Delta} u(y) (1 - v(x)) \sigma(dx, dy).$$

Proof. Let  $u \in C_0(X) \cap \mathbf{H}$  and  $v=1$  on a neighbourhood of  $\text{supp}[u]$ . Then, by Lemma 2.13, we have

$$(2.6) \quad \begin{aligned} \mathbf{a}^{(\alpha)}(u, v) &= \int_{D \times D - \Delta} u(y) (1 - v(x)) \sigma_\alpha(dx, dy) \\ &\quad + \alpha \int_X u(x) (1 - \alpha G_\alpha I_D(x)) m(dx) \end{aligned}$$

which implies that the measure  $\alpha(1 - \alpha G_\alpha I_D(x)) m(dx)$  is uniformly bounded

on  $D$ . Hence there exists a subsequence  $\{\alpha_n\}$  of  $\{\alpha\}$  and a positive Radon measure  $\chi_D$  on  $D$  such that

$$\alpha_n(1-\alpha_n G_{\alpha_n} I_D) m \rightarrow \chi_D \quad \text{vaguely on } D \text{ as } \alpha_n \rightarrow \infty.$$

Thus we have

$$\begin{aligned} \int_X u(x) \chi_D(dx) &= \lim_{\alpha_n \rightarrow \infty} [\mathbf{a}^{(\alpha_n)}(v, u) - \int_{D \times D - \Delta} u(y) (1-v(x)) \sigma_{\alpha_n}(dx, dy)] \\ &= \mathbf{a}(v, u) - \int_{D \times D - \Delta} u(y) (1-v(x)) \sigma(dx, dy). \end{aligned}$$

This shows that  $\chi_D$  is uniquely determined independently of the choice of  $\{\alpha_n\}$  and satisfies the equality (2.3). Let  $D'$  be a relatively compact open set such that  $D \subset D'$ . Then it is clear that  $\chi_D \geq \chi_{D'}$  on  $D$ . It is easy to see that (ii) follows from (i). The proof is complete.

By the same method, we can define  $\hat{\chi}_D$  and  $\hat{\chi}$  if we replace  $G_{\alpha}$  by  $\hat{G}_{\alpha}$  in the above lemma. Then  $\hat{\chi}$  satisfies for the above  $u$  and  $v$ ,

$$(2.7) \quad \int_X u(x) \hat{\chi}(dx) = \mathbf{a}(u, v) - \int_{X \times X - \Delta} u(x) (1-v(y)) \sigma(dx, dy).$$

**Theorem 2.15** (Beurling-Deny formula). *The regular Dirichlet form  $\mathbf{a}$  on  $L^2(X, m)$  can be represented for  $u, v \in C_0(X) \cap \mathbf{H}$  as follows;*

$$\begin{aligned} (2.8) \quad \frac{1}{2} \mathbf{a}(u, v) + \frac{1}{2} \hat{\mathbf{a}}(u, v) &= N(u, v) \\ &+ \frac{1}{2} \int_{X \times X - \Delta} (u(x) - u(y)) (v(x) - v(y)) \sigma(dx, dy) \\ &+ \frac{1}{2} \int_X u(x) v(x) \chi(dx) + \frac{1}{2} \int_X u(x) v(x) \hat{\chi}(dx). \end{aligned}$$

Here  $N$  is a symmetric form with domain  $C_0(X) \cap \mathbf{H}$  and satisfies the following condition;

(2.9)  $N(u, v) = 0$  for  $u, v \in C_0(X) \cap \mathbf{H}$  such that  $v$  is constant on a neighbourhood of  $\text{supp}[u]$ .  $\sigma$  is a positive Radon measure on  $X \times X - \Delta$  given by (2.2).  $\chi$  (resp.  $\hat{\chi}$ ) is a positive Radon measure on  $X$  given by (2.5) (resp. (2.7)).

Proof. Since  $(\bar{\mathbf{a}}, \mathbf{H})$  is a regular symmetric Dirichlet space on  $L^2(X, m)$ , we get for  $u, v \in C_0(X) \cap \mathbf{H}$ ,

$$\begin{aligned} (2.10) \quad \bar{\mathbf{a}}(u, v) &= \bar{N}(u, v) \\ &+ \frac{1}{2} \int_{X \times X - \Delta} (u(x) - u(y)) (v(x) - v(y)) \sigma(dx, dy) \end{aligned}$$

$$+\int_X u(x) v(x) \mathcal{X}(dx),$$

where  $\bar{N}$  is a symmetric form satisfying (2.9),  $\bar{\sigma}$  is a symmetric positive Radon measure on  $X \times X - \Delta$  such that

$$(2.11) \quad \int_{X \times X - \Delta} u(x) v(y) \bar{\sigma}(dx, dy) = -\bar{\mathbf{a}}(u, v)$$

for any  $u, v \in C_0(X) \cap \mathbf{H}$  with disjoint supports, and  $\mathcal{X}$  is a positive Radon measure on  $X$  such that

$$(2.12) \quad \int_X u(x) \mathcal{X}(dx) = \bar{\mathbf{a}}(u, v) - \frac{1}{2} \int_{X \times X - \Delta} (u(x) - u(y)) (v(x) - v(y)) \bar{\sigma}(dx, dy)$$

for any  $u, v \in C_0(X) \cap \mathbf{H}$  such that  $v=1$  on a neighbourhood of  $\text{supp}[u]$ . But by the definition of  $\sigma$  and  $\hat{\sigma}$ , we can see that  $\bar{\sigma} = \frac{1}{2}(\sigma + \hat{\sigma})$ . And (2.12) implies that

$$\begin{aligned} \int_X u(x) \mathcal{X}(dx) &= \bar{\mathbf{a}}(u, v) - \int_{X \times X - \Delta} u(y) (v(y) - v(x)) \bar{\sigma}(dx, dy) \\ &= \frac{1}{2} \mathbf{a}(u, v) - \frac{1}{2} \int_{X \times X - \Delta} u(y) (1 - v(y)) \sigma(dx, dy) \\ &\quad + \frac{1}{2} \hat{\mathbf{a}}(u, v) - \frac{1}{2} \int_{X \times X - \Delta} u(y) (1 - v(y)) \hat{\sigma}(dx, dy) \\ &= \frac{1}{2} \int_X u(x) \hat{\mathcal{X}}(dx) + \frac{1}{2} \int_X u(x) \mathcal{X}(dx). \end{aligned}$$

Hence  $\mathcal{X} = \frac{1}{2}(\mathcal{X} + \hat{\mathcal{X}})$ . Now let  $N = \bar{N}$ , then  $N$  satisfies the condition (2.9) and

$$\begin{aligned} (2.13) \quad \frac{1}{2} \mathbf{a}(u, v) + \frac{1}{2} \hat{\mathbf{a}}(u, v) &= \bar{\mathbf{a}}(u, v) \\ &= N(u, v) \\ &\quad + \frac{1}{4} \int_{X \times X - \Delta} (u(x) - u(y)) (v(x) - v(y)) \sigma(dx, dy) \\ &\quad + \frac{1}{4} \int_{X \times X - \Delta} (u(x) - u(y)) (v(x) - v(y)) \hat{\sigma}(dx, dy) \\ &\quad + \frac{1}{2} \int_X u(x) v(x) \mathcal{X}(dx) + \frac{1}{2} \int_X u(x) v(x) \hat{\mathcal{X}}(dx). \end{aligned}$$

Since the second and third terms of the right hand side of (2.13) are the same, (2.13) implies (2.8). The proof is complete.

Let  $D$  be any relatively compact open subset of  $X$ . For non-negative

$u \in C_0(X) \cap \mathbf{H}$  with  $\text{supp}[u] \subset X - \bar{D}$ , we define the positive measures  $\sigma_u$ ,  $\delta_u$  and  $\bar{\sigma}_u$  on  $X$  by

$$\begin{aligned}\sigma_u(dy) &= I_D(y) \int_X u(x) \sigma(dx, dy), \\ \delta_u(dy) &= I_D(y) \int_X u(x) \delta(dx, dy)\end{aligned}$$

and

$$\bar{\sigma}_u(dy) = I_D(y) \int_X u(x) \bar{\sigma}(dx, dy).$$

We use the notation  $\mathbf{H}^D = \{v \in \mathbf{H} : v = 0 \text{ q.e. on } D^c\}$ . Since  $\bar{\sigma}_u \in \mathbf{S}_0$  with respect to the restriction  $\mathbf{a}^D = \mathbf{a}|_{\mathbf{H}^D \times \mathbf{H}^D}$  and  $\bar{\sigma} = \frac{1}{2}(\sigma + \delta)$ , so do  $\sigma_u$  and  $\delta_u$ . The equality (2.10) implies that for any  $v \in C_0(X) \cap \mathbf{H}$ ,

$$\int_X v(x)^2 \chi(dx) \leq \bar{\mathbf{a}}(v, v).$$

This implies that  $I_K \cdot \chi \in \mathbf{S}_0$  and hence  $I_K \cdot \chi$ ,  $I_K \cdot \hat{\chi} \in \mathbf{S}_0$  for any compact set  $K \subset X$ . Thus, using the same method as Lemma 4.5.4 in [8], we have the following corollary.

**Corollary 2.16.** *The representation (2.8) of Dirichlet form  $\mathbf{a}$  extends to  $u, v \in \mathbf{H}$ ;*

$$\begin{aligned}(2.14) \quad \frac{1}{2} \mathbf{a}(u, v) + \frac{1}{2} \hat{\mathbf{a}}(u, v) &= N(u, v) \\ &+ \frac{1}{2} \int_{X \times X - \Delta} (\tilde{u}(x) - \tilde{u}(y)) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy) \\ &+ \frac{1}{2} \int_X \tilde{u}(x) \tilde{v}(x) \chi(dx) + \frac{1}{2} \int_X \tilde{u}(x) \tilde{v}(x) \hat{\chi}(dx),\end{aligned}$$

where  $\tilde{u}$  and  $\tilde{v}$  denote q.e. versions and  $N$  is a symmetric form on  $\mathbf{H} \times \mathbf{H}$  satisfying, for  $u, v \in \mathbf{H}$  with compact support,  $N(u, v) = 0$  if  $v$  is constant on a neighbourhood of  $\text{supp}[u]$ .

### 3. Jumping and killing measures

Let  $\mathbf{M} = (\Omega, P_x, \zeta, X_t)$  and  $\hat{\mathbf{M}} = (\hat{\Omega}, \hat{P}_x, \hat{\xi}, \hat{X}_t)$  be the Hunt processes associated with  $\mathbf{a}$  and  $\hat{\mathbf{a}}$ , respectively, which were constructed by S. Carrillo Menendez [5]. We denote by  $\mathbf{S}$  the family of all smooth measures (see [8]). It is known that  $\mu \in \mathbf{S}$  if and only if there exists a nest  $\{F_n\}$  of  $\mu$  such that  $I_{F_n} \cdot \mu \in \mathbf{S}_0$  for each  $n$  (see [8]; Theorem 3.2.3) and that a set  $N \subset X$  is exceptional if and only if  $\text{cap}(N) = 0$  ([11]).

The transition function and resolvent of  $\mathbf{M}$  is denoted by  $(p_t)_{t>0}$  and  $(R_\alpha)_{\alpha>0}$  respectively. Then for any  $u \in L^2(X, m)$ ,  $p_t u$  and  $R_\alpha u$  are q.c. versions of  $T_t u$  and  $G_\alpha u$  ( $t, \alpha > 0$ ) respectively (see [5]). The restriction  $\mathbf{M}^D$  of  $\mathbf{M}$  by an open set  $D$  is the Hunt process associated with  $(\mathbf{a}^D, \mathbf{H}^D)$ , where  $\mathbf{H}^D = \{u \in \mathbf{H} : u = 0 \text{ q.e. on } D^c\}$  and  $\mathbf{a}^D = \mathbf{a}|_{\mathbf{H}^D \times \mathbf{H}^D}$ . Let  $(R_\alpha^D)_{\alpha>0}$  be the resolvent of  $\mathbf{M}^D$ . We define

$$H_\alpha^M(x, E) = E_x[\exp(-\alpha \sigma_M I_E(X_{\sigma_M}))] \quad (\alpha > 0, x \in X, E \in \mathcal{B}(X)),$$

where  $M = D^c$  and  $\sigma_M = \inf\{t > 0 : X_t \in M\}$ . Then we have the following relations;

$$(3.1) \quad H_\alpha^M - H_\beta^M = (\beta - \alpha) R_\alpha^D H_\beta^M$$

and

$$(3.2) \quad R_\alpha = R_\alpha^D + H_\alpha^M R_\alpha$$

Similarly we can define the kernels  $\hat{R}_\alpha$ ,  $\hat{R}_\alpha^D$ ,  $\hat{p}_t$  and  $\hat{H}_\alpha^M$  related to  $\hat{\mathbf{M}}$ .

**Lemma 3.1.** *There exists a unique positive Radon measure  $k$  on  $X$  charging no exceptional set such that for any  $u \in C_0(X) \cup \mathbf{H}_0$ ,*

$$(3.3) \quad \begin{aligned} \langle \tilde{u}, k \rangle &= \lim_{\alpha \rightarrow \infty} \alpha \int_X u(x) (1 - \alpha R_\alpha 1(x)) m(dx) \\ &= \lim_{t \downarrow 0} \frac{1}{t} \int_X u(x) (1 - p_t 1(x)) m(dx), \end{aligned}$$

where  $\mathbf{H}_0$  is a set of all elements of  $\mathbf{H}$  with compact support in  $X$  and  $\tilde{u}$  is any q.c. version of  $u$ .

Similarly  $\hat{k}$  is given by replacing  $\hat{R}_\alpha$  and  $\hat{p}_t$  for  $R_\alpha$  and  $p_t$  in (3.3).

Proof. Let  $D$ ,  $D_1$  and  $D_2$  be relatively compact open subsets of  $X$  such that  $\bar{D} \subset D_1$  and  $\bar{D}_1 \subset D_2$ . If  $u \in C_0(X) \cap \mathbf{H}^{D_1}$  and  $v \in C_0(X) \cap \mathbf{H}^{D_2}$ , then

$$\begin{aligned} \mathbf{a}^{(\omega)}(v, u) &= \int_{D_2 \times D_2} u(y) (v(y) - v(x)) \sigma_\omega(dx, dy) \\ &\quad + \alpha \int_{D_1} u(x) v(x) (1 - \alpha R_\omega I_{D_2}(x)) m(dx). \end{aligned}$$

Fix  $v$  satisfying  $v = 1$  on  $D_1$ . Then

$$\begin{aligned} \mathbf{a}^{(\omega)}(v, u) &= \int_{D_2 \times D_2} u(y) (1 - v(x)) \sigma_\omega(dx, dy) \\ &\quad + \alpha \int_{D_1} u(x) (1 - \alpha R_\omega I_{D_2}(x)) m(dx). \end{aligned}$$

This implies that the family of maps

$$(3.4) \quad F_{\alpha}^D: u \mapsto \alpha \int_X u(x) (1 - \alpha R_{\alpha} I_{D_2}(x)) m(dx)$$

is equicontinuous on  $\mathbf{H}^{D_1}$ . In fact, for any  $u \in C_0(X) \cap \mathbf{H}^{D_1}$ ,

$$(3.5) \quad |F_{\alpha}^D(u)| \leq F_{\alpha}^D(|u|) \leq \mathbf{a}^{(\omega)}(v, |u|) \\ \leq C_1 \{ \mathbf{a}^{(\omega)}(v, v) + (v, v)_{L^2} \}^{1/2} \mathbf{a}_1(|u|, |u|)^{1/2} \text{ by Lemma 3.1 of [10]} \\ \leq C \mathbf{a}_1(u, u)^{1/2} \text{ by Lemma 2.11,}$$

with some constants  $C_1$  and  $C$ . It is trivial that  $\{F_{\alpha}^D\}$  is equicontinuous on  $C_0(D)$ . Let

$$F_{\alpha}(u) = \alpha \int_X u(x) (1 - \alpha R_{\alpha} 1(x)) m(dx)$$

Then, since  $F_{\alpha}(u) \leq F_{\alpha}^D(u)$  for any positive Borel function  $u$ ,  $\{F_{\alpha}\}$  is equicontinuous both on  $\mathbf{H}^D$  and on  $C_0(D)$ . Consider the case that  $u = \hat{R}_{\beta}^D f$ ,  $\beta > 0$ ,  $f \in C_0(D)$ . Then, by (3.2), we have

$$(3.6) \quad F_{\alpha}(u) = \alpha \int_X \hat{R}_{\beta}^D f(x) (1 - \alpha R_{\alpha}^D 1(x) - \alpha H_{\alpha}^M R_{\alpha} 1(x)) m(dx) \\ = \alpha \int_X f(x) (R_{\beta}^D 1(x) - \alpha R_{\beta}^D R_{\alpha}^D 1(x)) m(dx) \\ - \alpha^2 \int_X f(x) R_{\beta}^D H_{\alpha}^M R_{\alpha} 1(x) m(dx).$$

Using the resolvent equation and (3.1), the right hand side of (3.6) is equal to

$$\alpha \int_X f(x) (R_{\alpha}^D 1(x) - \beta R_{\beta}^D R_{\alpha}^D 1(x)) m(dx) - \alpha^2 \int_X f(x) R_{\alpha}^D H_{\beta}^D R_{\alpha} 1(x) m(dx)$$

and this converges to

$$\int_X f(x) (1 - \beta R_{\beta}^D 1(x)) m(dx) - \int_X f(x) H_{\beta}^M 1(x) m(dx)$$

as  $\alpha \rightarrow \infty$ . Since the range of  $\hat{R}_{\beta}^D$ ,  $\beta > 0$ , is dense in  $\mathbf{H}^D$  and  $\{F_{\alpha}\}$  is equicontinuous on  $\mathbf{H}^D$ ,  $F_{\alpha}(u)$  converges to some limit as  $\alpha \rightarrow \infty$  for any  $u \in \mathbf{H}^D$ . By the regularity of  $(\mathbf{a}^D, \mathbf{H}^D)$ ,  $\mathbf{H}^D \cap C_0(D)$  is dense in  $C_0(D)$ . Again by the equicontinuity of  $\{F_{\alpha}\}$  on  $C_0(D)$ ,  $F_{\alpha}(u)$  converges as  $\alpha \rightarrow \infty$  for any  $u \in C_0(D)$ . Since  $D$  is arbitrary, there exists a unique positive Radon measure  $k$  such that for any  $u \in C_0(X)$ ,  $\langle u, k \rangle = \lim_{\alpha \rightarrow \infty} F_{\alpha}(u)$ . And for each  $D$ ,

$$|\langle u, k \rangle| = |\lim_{\alpha \rightarrow \infty} F_{\alpha}(u)| \leq C \mathbf{a}_1(u, u)^{1/2} \quad (u \in C_0(D) \cap \mathbf{H}^D).$$

This shows that  $k$  charges no set of zero capacity in  $D$  and  $\langle \tilde{u}, k \rangle = \lim_{\alpha \rightarrow \infty} F_{\alpha}(u)$  for any  $u \in \mathbf{H}^D$ . Since  $D$  is arbitrary,  $k$  charges no exceptional set and  $\langle \tilde{u}, k \rangle =$

$\lim_{\alpha \rightarrow \infty} F_\alpha(u)$  for any  $u \in H_0$ . Hence the measure  $k$  satisfies the first equality of (3.3). The second equality of (3.3) holds by Karamata's theorem (see D.V. Widder [22]; Theorem 5.4.3). The proof is complete.

REMARK 3.2. In the proof of the above lemma, we have seen that the family of maps  $F_\alpha^D$  in (3.4) is equicontinuous both on  $H^D$  and  $C_0(D)$ . Hence, by Lemma 2.14 and the argument similar to that for the family  $\{F_\alpha\}$ , it follows that

$$(3.7) \quad \langle \tilde{u}, \chi_{D_2} \rangle = \lim_{\alpha \rightarrow \infty} \alpha \int_X u(x) (1 - \alpha R_\alpha I_{D_2}(x)) m(dx) \quad \text{for any } u \in H^D.$$

**Lemma 3.3** (cf. [8]; Lemma 4.5.2). *The measure  $k$  in Lemma 3.1 satisfies the following conditions;*

(i) *For any  $u \in H$ ,*

$$(3.8) \quad \begin{aligned} \langle \tilde{u}^2, k \rangle &= \lim_{\alpha \rightarrow \infty} \alpha \int_X u(x)^2 (1 - \alpha R_\alpha 1(x)) m(dx) \\ &= \lim_{t \downarrow 0} \frac{1}{t} \int_X u(x)^2 (1 - p_t 1(x)) m(dx), \end{aligned}$$

(ii) *For any  $f, h \in B^+(X)$  (=the set of all positive Borel functions) and  $t > 0$ ,*

$$(3.9) \quad E_{h,m}[f(X_{\zeta-}); \zeta \leq t] = \int_0^t \langle fk, \hat{p}_s h \rangle ds,$$

(iii)  *$E_x[e^{-\alpha \zeta} f(X_{\zeta-})]$  is a q.c. version of  $U_\alpha(fk)$  for  $\alpha > 0$ ,  $f \in C_0^+(X)$ .*

Proof. Since  $\alpha \hat{R}_\alpha$  is sub-Markovian, it holds that

$$(3.10) \quad \alpha \int_X u(x)^2 (1 - \alpha R_\alpha 1(x)) m(dx) \leq 2\alpha \int_X u(x) (u(x) - \alpha R_\alpha u(x)) m(dx)$$

for all  $u \in H$  (see the proof of Proposition 1.3.3 in [12]). By Lemma 2.11 and Lemma 3.1, we have

$$\langle u^2, k \rangle \leq 2a(u, u) \quad \text{for any } u \in C_0(X) \cap H.$$

By Fatou's lemma, we have

$$(3.11) \quad \langle \tilde{u}^2, k \rangle \leq 2a(u, u) \quad \text{for any } u \in H.$$

From Lemma 3.1 and (3.11), the statements (i), (ii) and (iii) follow by the same way as Lemma 4.5.2 in [8]. The proof is complete.

REMARK 3.4. The statement (3.9) is equivalent to the following;

$$(3.12) \quad E_{h,m}[e^{-\alpha \zeta} f(X_{\zeta-})] = \langle fk, \hat{R}_\alpha h \rangle, f, h \in B^+(X), \alpha > 0.$$



We can give a direct proof of (3.12). In fact, let  $D$  be any relatively compact open set of  $X$  and  $f \in C_0^+(X)$  and  $\text{supp } [f] \subset D$ . Then  $f\hat{R}_\alpha h \in \mathbf{H}^D$  ( $h \in C_0^+(X)$ ) and it holds that

$$\begin{aligned} \langle fk, \hat{R}_\alpha h \rangle &= \langle k, f\hat{R}_\alpha h \rangle \\ &= \lim_{\beta \rightarrow \infty} \beta \int_X f \hat{R}_\alpha h (1 - \beta R_\beta 1) \, dm \\ &= \lim_{\beta \rightarrow \infty} \beta \int_X h R_\alpha [f(1 - \beta R_\beta 1)] \, dm. \end{aligned}$$

But we have

$$\begin{aligned} \beta R_\alpha [f(1 - \beta R_\alpha 1)](x) &= \beta E_x \left[ \int_0^\infty e^{-\alpha t} f(X_t) E_{x_t} [e^{-\beta \zeta}] \, dt \right] \\ &= \beta E_x \left[ \int_0^\infty e^{-\alpha t} f(X_t) e^{-\beta(\zeta - t)} \, dt \right] \end{aligned}$$

and this converges to  $E_x[e^{-\alpha \zeta} f(X_{\zeta-})]$  as  $\beta \rightarrow \infty$ .

**Theorem 3.5.**  $k = \mathcal{X}$  and  $\hat{k} = \hat{\mathcal{X}}$ .

Proof. By the definition of  $\mathcal{X}$  and  $k$ , it is clear that  $k \leq \mathcal{X}$ . Hence we prove that  $k \geq \mathcal{X}$ . Take  $D$  and  $D_2$  as in the proof of Lemma 3.1. For any  $f \in C_0^+(X) \cap \mathbf{H}$  such that  $\text{supp } [f] \subset D$  and any  $h \in C_0^+(X)$ ,

$$\begin{aligned} \langle f\mathcal{X}, \hat{R}_\alpha^D h \rangle &= \langle \mathcal{X}, f\hat{R}_\alpha^D h \rangle \leq \langle \mathcal{X}_{D_2}, f\hat{R}_\alpha^D h \rangle \\ &= \lim_{\alpha \rightarrow \infty} \alpha \int_X f \hat{R}_\alpha^D h (1 - \alpha R_\alpha I_{D_2}) \, dm \quad \text{by (3.7)} \\ &= \lim_{\alpha \rightarrow \infty} \alpha \int_X h R_\alpha^D [f(1 - \alpha R_\alpha I_{D_2})] \, dm \\ &\leq \lim_{\alpha \rightarrow \infty} \alpha \int_X h R_\alpha^D [f(1 - \alpha R_\alpha^D I_D)] \, dm \\ &= E_{h \cdot m} [e^{-(\zeta \wedge \tau_D)} f(X_{(\zeta \wedge \tau_D)-})] \end{aligned}$$

where  $\tau_D = \sigma_{X-D}$ . The last equality follows from the same calculus as Remark 3.4. Letting  $D \uparrow X$ , we have

$$\langle f\mathcal{X}, \hat{R}_\alpha h \rangle \leq E_{h \cdot m} [e^{-\zeta} f(X_{\zeta-})] = \langle fk, \hat{R}_\alpha h \rangle.$$

This shows that  $\mathcal{X} \leq k$ . Similarly we can see  $\hat{\mathcal{X}} = \hat{k}$ . The proof is complete.

In the present paper, the definitions of *additive functionals* and related concepts (*continuous*, *square integrable* etc.) are taken in the sense of Fukushima [8]; Chap. 5. Let  $\mathcal{A}_c^+$  be the set of all positive continuous AF's. Henceforth we use the notations;

$$U_A^\alpha f(x) = E_x \left[ \int_0^\infty e^{-\alpha t} f(X_t) dA_t \right] \quad \text{and} \quad (fA)_t = \int_0^t f(X_s) dA_s,$$

for  $A \in \mathbf{A}_c^+$  and  $f \in \mathbf{B}^+(X)$ . The set of all *smooth measures* on  $X$  is denoted by  $\mathbf{S}$ . The following theorem is proved in [6].

**Theorem 3.6.** *For  $\mu \in \mathbf{S}$  and  $A \in \mathbf{A}_c^+$ , the following conditions are equivalent to each other;*

- (i)  $U_A^\alpha f$  is a q.c. version of  $U_\alpha(f\mu)$  for any  $f \in \mathbf{B}^+(X)$  such that  $f\mu \in \mathbf{S}_0$ .
- (ii)  $(h, U_A^\alpha f)_{L_2} = \langle f\mu, \hat{R}_\alpha h \rangle$ ,  $\alpha > 0$ ,  $f, h \in \mathbf{B}^+(X)$ .
- (iii)  $E_{h \cdot m}[(fA)_t] = \int_0^t \langle f\mu, \hat{p}_s h \rangle ds$ ,  $t > 0$ ,  $f, h \in \mathbf{B}^+(X)$ .
- (iv)  $\lim_{\alpha \rightarrow \infty} \alpha (h, U_A^{\alpha+\gamma} f) = \langle f\mu, h \rangle$  for any  $\gamma$ -co-excessive function  $h$  and  $f \in \mathbf{B}^+(X)$ .
- (v)  $\lim_{t \downarrow 0} \frac{1}{t} E_{h \cdot m}[(fA)_t] = \langle f\mu, h \rangle$  for any  $\gamma$ -co-excessive function  $h$  and  $f \in \mathbf{B}^+(X)$ .

Moreover,  $\mathbf{S}$  and  $\mathbf{A}_c^+$  are in one to one correspondence which is characterized by one of above statements.

Let  $D \subset X$  be an open set,  $\mathbf{A}_c^+(D)$  the set of all positive continuous AF's of  $\mathbf{M}^D$  and  $\mathbf{S}^D$  the set of all smooth measures related to  $\mathbf{a}^D$ . Theorem 3.6 holds on  $D$ , i.e.,  $\mathbf{S}^D$  and  $\mathbf{A}_c^+(D)$  correspond to each other in a unique manner. For  $A \in \mathbf{A}_c^+$ , let  $A_t^D = A_{\sigma_{\mathbf{M}} \wedge t} \in \mathbf{A}_c^+(D)$ .

**Lemma 3.7.** *If  $\mu$  is the smooth measure corresponding to  $A$ , then  $\mu|_D$  is the measure corresponding to  $A^D$ .*

*Proof.* In the same way as in the proof of Lemma 5.1.5 in [8], we can show that Theorem 3.6 (iv) is valid on  $D$ , and this completes the proof.

By A. Benveniste and J. Jacod [2], there exists a couple  $(H, N(y, dx))$ , so called Levy system of  $\mathbf{M}$ , of  $H \in \mathbf{A}_c^+$  and a kernel  $N$  such that

$$E_x \left[ \sum_{0 < s \leq t} f(X_s, X_{s-}) I_{\{X_s \neq X_{s-}\}} \right] = E_x \left[ \int_0^t dH_s \int_{X \cup \{\delta\}} N(X_s, dy) f(X_s, y) \right]$$

for any  $x \in X$  and any positive Borel function  $f$  on  $(X \cup \{\delta\}) \times (X \cup \{\delta\})$  ( $\delta$  is an extra point).

Now we give the main theorem of this section.

**Theorem 3.8.** *Let  $\nu$  be the smooth measure corresponding to  $H \in \mathbf{A}_c^+$ . Then we have*

- (i)  $\sigma(dx, dy) = N(y, dx) \nu(dy)$  on  $X \times X - \Delta$ ,
- (ii)  $\chi(dy) = N(y, \delta) \nu(dy)$  on  $X$ .

*Proof.* (i) Let  $u, v \in C_0(X) \cap \mathbf{H}$  and  $D \subset X$  a relatively compact open

set satisfying  $\text{supp}[v] \subset D \subset \bar{D}$  and  $\text{supp}[u] \subset \mathring{M}$  ( $M = D^c$ ). We already remarked that the measure  $\sigma_u(dy) = I_D(y) \int_X u(x) \sigma(dx, dy)$  is of finite energy integral with respect to  $\mathbf{a}^D$ . Then

$$\begin{aligned} \mathbf{a}_\alpha^D(\alpha R_\alpha^D u + H_\alpha^M u - u, v) &= \alpha(u, v)_{L^2} + \mathbf{a}_\alpha^D(H_\alpha^M u - u, v) \\ &= \alpha(u, v)_{L^2} + \mathbf{a}_\alpha(H_\alpha^M u - u, v) \quad (\because H_\alpha^M u - u \in H^D) \\ &= \alpha(u, v)_{L^2} - \mathbf{a}_\alpha(u, v) \quad (\because \mathbf{a}_\alpha(H_\alpha^M u, v) = 0 \text{ (see [11])}) \\ &= -\mathbf{a}(u, v) = \int u(x) v(y) \sigma(dx, dy) = \int v(y) \sigma_u(dy) \\ &= \langle \sigma_u, v \rangle = \mathbf{a}_\alpha^D(U_\alpha^D(\sigma_u), v), \end{aligned}$$

and hence, for  $x \in D$ ,

$$\begin{aligned} U_\alpha^D(\sigma_u)(x) &= \alpha R_\alpha^D u(x) + H_\alpha^M u(x) - u(x) \\ &= H_\alpha^M u(x) \quad (\because \text{supp}[u] \subset \mathring{M}) \\ &= E_x[\exp(-\alpha \sigma_M) u(X_{\sigma_M})] \\ &= E_x\left[\sum_{0 < s \leq \sigma_M} \exp(-\alpha s) u(X_s)\right] \\ &= E_x\left[\int_0^{\sigma_M} \exp(-\alpha s) Nu(X_s) dH_s\right], \end{aligned}$$

where

$$Nu(y) = \int_X u(x) N(y, dx).$$

Hence  $\sigma_u$  is the smooth measure corresponding to  $((Nu)H)^D$ . On the other hand, by Lemma 3.7,  $((Nu)\nu)|_D$  is also the measure corresponding to  $((Nu)H)^D$ . Therefore  $\sigma_u = ((Nu)\nu)|_D$ . Thus

$$\begin{aligned} \int_{X \times -\Delta} u(x) v(y) \sigma(dx, dy) &= \int_X v(y) \sigma_u(dy) \\ &= \int_{X \times X - \Delta} u(x) v(y) N(y, dx) \nu(dy) \end{aligned}$$

and this implies that  $\sigma(dx, dy) = N(y, dx) \nu(dy)$ .

Now we prove (ii). For any  $f \in C_0^+(X)$ , we have, by Lemma 3.3 and Theorem 3.6,

$$\begin{aligned} \int_0^t \langle f k, \hat{p}_s, 1 \rangle ds &= E_m[f(X_{t-}) I_{\{\zeta \leq t\}}] \\ &= E_m\left[\int_0^t f(X_s) N(X_s, \delta) dH_s\right] \\ &= \int_0^t \langle f N(\cdot, \delta) \nu, \hat{p}_s, 1 \rangle ds \end{aligned}$$

which implies that  $N(\cdot, \delta) \nu = k = \chi$ . The proof is complete.

In this section, we have proved that

$$\sigma(dx, dy) = N(y, dx) \nu(dy) \text{ on } X \times X - \Delta$$

and

$$\chi(dy) = k(dy) = N(y, \delta) \nu(dy) \text{ on } X$$

which are called the *jumping* and *killing measure* of  $\mathbf{M}$  respectively.

REMARK 3.9. Let  $(\hat{H}, \hat{N}(y, dx))$  be a Lévy system of the Hunt process  $\hat{\mathbf{M}}$  and  $\hat{\nu}$  the smooth measure corresponding to  $\hat{H} \in \hat{\mathbf{A}}_c^+$ .

Then it holds that

$$\delta(dx, dy) = \hat{N}(y, dx) \hat{\nu}(dy) \text{ on } X \times X - \Delta$$

and

$$\hat{\chi}(dy) = \hat{k}(dy) = \hat{N}(y, \delta) \hat{\nu}(dy) \text{ on } X.$$

#### 4. The energy of AF's and a decomposition of AF's of finite energy

In this section, using the notion of the energy of AF's, we give a decomposition of AF's generated by functions in  $\mathbf{H}$ .

DEFINITION 4.1. For any  $A \in \mathbf{A}$  (=the set of all AF's) we define

$$(4.1) \quad e(A) = \lim_{\alpha \rightarrow \infty} \frac{\alpha^2}{2} E_m \left[ \int_0^\infty e^{-\alpha t} A_t^2 dt \right],$$

whenever the finite limit exists.  $e(A)$  is called the energy of  $A$ . And we define the mutual energy of AF's  $A$  and  $B$  by  $e(A, B) = \frac{1}{2} [e(A+B) - e(A) - e(B)]$ .

REMARK 4.2. M. Fukushima [8] defined the energy of AF  $A$  by

$$(4.2) \quad \bar{e}(A) = \lim_{t \downarrow 0} \frac{1}{2t} E_m[A_t^2].$$

If  $\bar{e}(A)$  exists, then  $e(A)$  is well defined and  $e(A) = \bar{e}(A)$ .

For any  $A \in \mathbf{A}_c^+$ , the smooth measure corresponding to  $A$  is denoted by  $\mu_A$ . We consider the families

$$\mathcal{M} = \{M \in \mathbf{A}: \text{ for q.e. } x \in X, E_x[M_t^2] < \infty \text{ and } E_x[M_t] = 0 (t > 0)\}$$

and

$$\overset{\circ}{\mathcal{M}} = \{M \in \mathcal{M} : e(M) < \infty\}.$$

For any  $M \in \overset{\circ}{\mathcal{M}}$ , it holds that

$$(4.3) \quad e(M) = \bar{e}(M) = \sup_{t>0} \frac{1}{2t} E_m[M_t^2] = \frac{1}{2} \mu_{\langle M \rangle}(X)$$

(see [8]). Here the symbol  $\langle M \rangle$  is the quadratic variation of  $M \in \overset{\circ}{\mathcal{M}}$  (see [7]).

**Lemma 4.3.** *Let  $A \in A_c^+$ . Then*

- (i)  $\langle \nu, U_A^\alpha f \rangle = \langle f \mu_A, \hat{U}_\alpha \nu \rangle$  for  $\alpha > 0$ ,  $\nu \in S_0$  and  $f \in B^+(X)$ .
- (ii)  $E_\nu[A_t] \leq e^t \|\hat{U}_1 \nu\|_\infty \mu_A(X)$  for  $t > 0$  and  $\nu \in \hat{S}_{00}$ .

Proof. (i) For any  $f \in B^+(X)$  such that  $f \mu_A \in S_0$ , it holds that

$$\begin{aligned} \langle \nu, U_A^\alpha f \rangle &= \langle \nu, U_\alpha(f \mu_A) \rangle \text{ by Theorem 3.6} \\ &= a_\alpha(U_\alpha(f \mu_A), \hat{U}_\alpha \nu) \\ &= \langle f \mu_A, U_\alpha \nu \rangle. \end{aligned}$$

Let  $f \in B^+(X)$  and  $f_n = I_{F_n}(f \wedge n)$ ,  $n = 1, 2, \dots$ , where  $\{F_n\}$  is a nest of  $\mu_A$ . Then

$$\begin{aligned} \langle \nu, U_A^\alpha f \rangle &= \lim_{n \rightarrow \infty} \langle \nu, U_A^\alpha f_n \rangle \\ &= \lim_{n \rightarrow \infty} \langle f_n \mu_A, \hat{U}_\alpha \nu \rangle \\ &= \langle f \mu_A, \hat{U}_\alpha \nu \rangle. \end{aligned}$$

$$\begin{aligned} (ii) \quad E_\nu[A_t] &\leq e^t \langle \nu, E_x[\int_0^\infty e^{-s} dA_s] \rangle \\ &= e^t \langle \nu, U_A^1 1 \rangle \\ &= e^t \langle \mu_A, \hat{U}_1 \nu \rangle \text{ by (i)} \\ &= e^t \|\hat{U}_1 \nu\|_\infty \mu_A(X). \end{aligned}$$

Using (4.3) and Lemma 4.3 (ii) instead of (5.2.17) in [8], the following theorem can be proved in the same way as Theorem 5.2.1 in [8].

**Theorem 4.4.**  $\overset{\circ}{\mathcal{M}}$  is a real Hilbert space with inner product  $e$ . Moreover, for any  $e$ -Cauchy sequence  $M^n \in \overset{\circ}{\mathcal{M}}$ , there exist a unique  $M \in \overset{\circ}{\mathcal{M}}$  and a subsequence  $n_k$  such that  $\lim_{n \rightarrow \infty} e(M^n - M) = 0$  and for q.e.  $x \in X$ ,  $P_x(\lim_{n_k \rightarrow \infty} M_t^{n_k} = M_t)$  uniformly on any finite interval of  $t = 1$ .

Let

$$\begin{aligned} \mathcal{N}_c &= \{N \in \mathcal{A} : N \text{ is continuous and for q.e. } x \in X, \\ &\quad E_x[|N_t|] < \infty \text{ (} t > 0 \text{) and } e(N) = 0\}. \end{aligned}$$

We define

$$(4.4) \quad A_t^{[u]} = \tilde{u}(X_t) - \tilde{u}(X_0) \quad (t > 0)$$

where  $\tilde{u}$  is a q.c. version of  $u \in H$ .

**Lemma 4.5.**  $A^{[u]}$  is of finite energy and for  $u, v \in H$

$$(4.5) \quad e(A^{[u]}, A^{[v]}) = \frac{1}{2} \hat{\mathbf{a}}(u, v) + \frac{1}{2} \mathbf{a}(u, v) - \frac{1}{2} \langle \tilde{u} \tilde{v}, \hat{\chi} \rangle.$$

Proof. By the definition of  $e$ ,

$$\begin{aligned} e(A^{[u]}) &= \lim_{\alpha \rightarrow \infty} \frac{\alpha^2}{2} E_m \left[ \int_0^\infty e^{-\alpha t} (\tilde{u}(X_t) - \tilde{u}(X_0))^2 dt \right] \\ &= \lim_{\alpha \rightarrow \infty} \frac{\alpha^2}{2} \int_X (R_\alpha u(x)^2 - 2u(x) R_\alpha u(x) + \frac{1}{\alpha} u(x)^2) m(dx) \\ &= \lim_{\alpha \rightarrow \infty} \frac{\alpha^2}{2} \left[ \int_X (R_\alpha u(x)^2 - \frac{1}{\alpha} u(x)^2) m(dx) + 2 \int_X \left( \frac{1}{\alpha} u(x)^2 - u(x) R_\alpha u(x) \right) m(dx) \right] \\ &= -\frac{1}{2} \lim_{\alpha \rightarrow \infty} \alpha \int_X u(x)^2 (1 - \alpha \hat{R}_\alpha 1(x)) m(dx) + \lim_{\alpha \rightarrow \infty} \alpha (u - \alpha R_\alpha u, u)_{L^2} \\ &= -\frac{1}{2} \int_X \tilde{u}(x)^2 \hat{\chi}(dx) + \mathbf{a}(u, u) \quad \text{by Lemma 2.12 and Lemma 3.3} \\ &= -\frac{1}{2} \int_X \tilde{u}(x)^2 \hat{\chi}(dx) + \mathbf{a}(u, u) \quad \text{by Theorem 3.5.} \end{aligned}$$

The equality (4.5) follows from the definition of the mutual energy. The proof is complete.

REMARK 4.6. Similarly, by the definition of  $\bar{e}$ , we have

$$(4.6) \quad \bar{e}(A^{[u]}) = \lim_{t \downarrow 0} \frac{1}{t} (u - p_t u, u)_{L^2} - \frac{1}{2} \lim_{t \downarrow 0} \frac{1}{t} \int_X \tilde{u}(x)^2 (1 - \hat{p}_t 1(x)) m(dx).$$

In the symmetric case, it holds that

$$(4.7) \quad \lim_{t \downarrow 0} \frac{1}{t} (u - p_t u, u)_{L^2} = \mathbf{a}(u, u)$$

and hence

$$(4.8) \quad \bar{e}(A^{[u]}) = \mathbf{a}(u, u) - \frac{1}{2} \int_X \tilde{u}(x)^2 \hat{\chi}(dx).$$

But, in the non-symmetric case, we do not know the validity of (4.7), and hence we can not prove (4.8).

**Lemma 4.7** (cf. [8]; Lemma 5.1.1). *For any  $u \in H$ ,  $v \in S_0$ ,  $0 < T < \infty$  and  $\varepsilon > 0$ , it holds that*

$$(4.9) \quad P_v \left( \sup_{0 \leq t \leq T} |\tilde{u}(X_t)| > \varepsilon \right) \leq \frac{C e^T}{\varepsilon} a_1(U_1 v, U_1 v)^{1/2} a_1(u, u)^{1/2}$$

for some constant  $C > 0$ .

Proof. Let  $E = \{x \in X : |\tilde{u}(x)| > \varepsilon\}$ . Then the left hand side of (4.9) is dominated by

$$\begin{aligned} (4.10) \quad e^T \int_X p(x) v(dx) \quad (p(x) = E_x[e^{-\sigma_H}]) \\ = e^T a_1(p, U_1 v) \\ \leq C_1 e^T a_1(p, p)^{1/2} a_1(U_1 v, U_1 v)^{1/2} \quad \text{by (a.1) and (a.2)} \\ = C_1 e^T a_1(u_E^1, u_E^1)^{1/2} a_1(U_1 v, U_1 v)^{1/2}, \end{aligned}$$

since  $p$  is a q.c. version of the capacitary potential  $u_E^1$  of  $E$ . By the definition of  $E$  and  $u_E^1$ , we have

$$a_1(u_E^1, u_E^1)^{1/2} \leq \frac{C_2}{\varepsilon} a_1(|u|, |u|)^{1/2} \leq \frac{C_2}{\varepsilon} a_1(u, u)^{1/2},$$

and this completes the proof of Lemma 4.7.

For the AF  $A^{[u]}$  generated by  $u \in H$ , we have the following theorem (see [8]; Theorem 5.2.2 for the symmetric case).

**Theorem 4.8.** *For any  $u \in H$ ,  $A^{[u]}$  admits a unique decomposition;*

$$(4.11) \quad A^{[u]} = M^{[u]} + N^{[u]}, \quad M^{[u]} \in \overset{\circ}{\mathcal{M}}, \quad N^{[u]} \in \mathcal{N}_c,$$

Moreover, it holds that for  $u, v \in H$ ,

$$(4.12) \quad e(M^{[u]}, M^{[v]}) = \frac{1}{2} a(u, v) + \frac{1}{2} \hat{a}(u, v) - \frac{1}{2} \langle \tilde{u} \tilde{v}, \hat{\chi} \rangle$$

Proof. The uniqueness is trivial. The equality (4.12) follows from (4.11) and Lemma 4.5. We show the existence of such  $M^{[u]}$  and  $N^{[u]}$ . First we consider the case of  $u = R_1 f$ ,  $f \in C_0^+(X)$ . Then  $Lu = u - f$ . Let

$$N_t^{[u]} = \int_0^t Lu(X_s) ds$$

and

$$M_t^{[u]} = A_t^{[u]} - \int_0^t Lu(X_s) ds.$$

Then q.e.  $x \in X$ ,

$$E_x[M_t^{[u]}] = p_t u(x) - u(x) - \int_0^t p_s(u-f) ds = 0.$$

Since

$$(M_t^{[u]})^2 \leq 3\tilde{u}(X_t)^2 + 3\tilde{u}(X_0)^2 + 3(N_t^{[u]})^2,$$

we have, for all  $t > 0$ ,

$$E_\nu[(M_t^{[u]})^2] < \infty \quad \text{for any } \nu \in \hat{S}_0.$$

By Lemma 2.8, for all  $t > 0$ ,

$$E_x[(M_t^{[u]})^2] < \infty \quad \text{q.e. } x \in X.$$

Since  $M_t^{[u]}$  is a martingale, it holds that

$$\text{q.e. } x \in X, E_x[(M_t^{[u]})^2] < \infty \quad \text{for all } t > 0.$$

Thus  $M_t^{[u]} \in \mathring{\mathcal{M}}$ . Let  $Lu = g$ . Then

$$\begin{aligned} E_m[(N_t^{[u]})^2] &= 2E_m\left[\int_0^t g(X_s) \int_s^t g(X_v) dv ds\right] \\ &= 2 \int_0^t \int_0^{t-s} (\hat{p}_s 1, g \hat{p}_v g)_{L^2} dv ds \\ &= 2 \int_0^t (\hat{p}_{t-s} 1, g \int_0^s \hat{p}_v g dv)_{L^2} ds \\ &\leq \|g\|_{L^2}^2 \cdot t^2. \end{aligned}$$

This implies that  $e(N_t^{[u]}) = 0$  and hence  $N_t^{[u]} \in \mathcal{N}_e$ . Next, for any  $u \in H$ , there exists  $u_n = R_1 f_n$  ( $f_n \in C_0(X)$ ), which converges to  $u$  in  $\|\cdot\|_H$  as  $n \rightarrow \infty$ . By the uniqueness of the decomposition (4.11) for  $u_n$ 's, we have

$$\begin{aligned} e(M^{[u_n]} - M^{[u_m]}) &= e(M^{[u_n - u_m]}) \\ &= a(u_n - u_m, u_n - u_m) - \frac{1}{2} \langle (u_n - u_m)^2, \hat{\chi} \rangle \\ &\leq a(u_n - u_m, u_n - u_m) \\ &\rightarrow 0 \quad \text{as } n, m \rightarrow \infty. \end{aligned}$$

Hence  $\{M^{[u_n]}\}$  is a Cauchy sequence in  $(\mathring{\mathcal{M}}, e)$ . Then, by Theorem 4.4, there exist a unique  $M^{[u]} \in \mathring{\mathcal{M}}$  and a subsequence  $n_k$  such that

$$M^{[u_{n_k}]} \rightarrow M^{[u]} \quad \text{in } (\mathring{\mathcal{M}}, e) \quad \text{as } n \rightarrow \infty$$

and q.e.  $x \in X$ ,



$$(4.13) \quad P_x(\lim_{n \rightarrow \infty} M_t^{[u_n]} = M_t^{[u]} \text{ uniformly on any finite interval of } t) = 1.$$

Let  $N^{[u]} = A^{[u]} - M^{[u]}$ . By virtue of Lemma 4.7 and the definition of  $A^{[u]}$ , the statement (4.13) for  $A^{[u]}$  holds. Hence the statement (4.13) for  $N^{[u]}$  holds. This implies that  $N^{[u]}$  is a continuous AF. It is clear that q.e.  $x \in X$ ,  $E_x[|N_t^{[u]}|] < \infty$  for all  $t > 0$ . Now we prove that  $e(N^{[u]}) = 0$ . Since

$$N^{[u]} = A^{[u-u_n]} - (M^{[u]} - M^{[u_n]}) + N^{[u_n]},$$

$$\begin{aligned} & \lim_{\alpha \rightarrow \infty} \frac{\alpha^2}{2} E_m \left[ \int_0^\infty e^{-\alpha t} (N_t^{[u]})^2 dt \right] \\ & \leq 3e(A^{[u-u_n]}) + 3e(M^{[u]} - M^{[u_n]}) + 3e(N^{[u_n]}) \end{aligned}$$

$\rightarrow 0$  as  $n \rightarrow \infty$ , by the same calculus as above. Thus  $e(N^{[u]}) = 0$ . The proof is complete.

Next we give a decomposition of  $M^{[u]}$  ( $u \in H$ ). Let

$$\mathcal{M}_c = \{M \in \mathcal{M} : P_x(M_t(\omega) \text{ is continuous in } t) = 1 \text{ q.e. } x \in X\}$$

and

$$\mathcal{M}_d = \{M \in \mathcal{M} : \langle M, L \rangle = 0 \text{ for any } L \in \mathcal{M}_c\}.$$

P.A. Meyer [7] showed that the martingale AF  $M^{[u]}$  has the following decomposition;

$$(4.14) \quad M^{[u]} = \overset{c}{M}^{[u]} + \overset{d}{M}^{[u]}, \quad \overset{c}{M}^{[u]} \in \mathcal{M}_c, \quad \overset{d}{M}^{[u]} \in \mathcal{M}_d$$

where

$$(4.15) \quad \overset{d}{M}_t^{[u]} = \sum_{0 < s \leq t} \Delta M_s^{[u]} I_{\{\Delta M_s^{[u]} \neq 0\}} (= \sum_{0 < s \leq t} (\tilde{u}(X_s) - \tilde{u}(X_{s-})) I_{\{X_s \neq X_{s-}\}})$$

and  $\overset{c}{M}^{[u]} = M^{[u]} - \overset{d}{M}^{[u]}$ . Here  $\Delta M_s^{[u]} = M_s^{[u]} - M_{s-}^{[u]}$  and the right hand side of (4.15) is defined by the following way. Consider the sequence  $T_n^k$ ,  $n=1, 2, \dots$ , of the stopping times defined by

$$\begin{aligned} T_1^k &= \inf \{s : \frac{1}{k} < |\tilde{u}(X_s) - \tilde{u}(X_{s-})| \leq \frac{1}{k-1}\}, \\ T_{n+1}^k &= \inf \{s > T_n^k : \frac{1}{k} < |\tilde{u}(X_s) - \tilde{u}(X_{s-})| \leq \frac{1}{k-1}\}, \quad n = 1, 2, \dots \end{aligned}$$

Let

$$A^k = \sum_{n=1}^{\infty} (\tilde{u}(X_{T_n^k}) - \tilde{u}(X_{T_n^k-})) I_{\{T_n^k \leq t\}}.$$

Since

$$E_x \left[ \sum_{n=1}^{\infty} |\tilde{u}(X_{T_n^k}) - \tilde{u}(X_{T_n^k-})| I_{\{T_n^k \leq t\}} \right] < \infty \quad \text{q.e. } x \in X,$$

there exists a dual predictable projection  $(A^k)^\flat$  of  $A^k$ . Let  $\widehat{A^k} = A^k - (A^k)^\flat$ . Then  $\sum_{k=1}^n \widehat{A^k}$  is a Cauchy sequence with respect to the family  $\{\eta_{x,t}: \eta_{x,t}(M) = E_x[M_t^2] \text{ for q.e. } x \in X \text{ and all } t > 0\}$  of the semi-norms in  $\mathcal{M}$ . Then we define the right hand side of (4.15) as  $\lim_{n \rightarrow \infty} \sum_{k=1}^n \widehat{A^k}$ .

Moreover  $\overset{d}{M}^{[u]}$  can be decomposed in the following way;

$$(4.16) \quad \overset{d}{M}^{[u]} = \overset{j}{M}^{[u]} + \overset{k}{M}^{[u]},$$

where

$$(4.17) \quad \overset{j}{M}^{[u]} = \sum_{0 < s \leq t} \overbrace{(\tilde{u}(X_s) - \tilde{u}(X_{s-})) I_{\{\xi > t\}}}^{\quad}$$

and

$$\overset{k}{M}^{[u]} = - \overbrace{\tilde{u}(X_{\xi-}) I_{\{\xi \leq t\}}}^{\quad}.$$

Here the right hand side of (4.17) is defined by the same method as the definition of  $\overset{c}{M}^{[u]}$ .

Recall that  $\mu_A$  denotes the smooth measure corresponding to  $A \in \mathbf{A}_c^+$ . If  $A_t(\omega) = A_t^{(1)}(\omega) - A_t^{(2)}(\omega)$ ,  $A_t^{(1)}, A_t^{(2)} \in \mathbf{A}_c^+$  and if  $\mu_{A^{(i)}}$ ,  $i=1, 2$ , are bounded measures, then the measure corresponding to  $A$  is defined by  $\mu_A = \mu_{A^{(1)}} - \mu_{A^{(2)}}$ . In particular, for  $M^{[u]}, M^{[v]} \in \overset{\circ}{\mathcal{M}}(u, v \in H)$ , we use the abbreviations  $\mu_{\langle u \rangle}^\alpha$ ,  $\mu_{\langle u, v \rangle}^\alpha$  and  $\mu_{\langle u, v \rangle}^\alpha$ ,  $\alpha = c, d, j, k$ , for  $\mu_{\langle M^{[u]} \rangle}^\alpha$ ,  $\mu_{\langle M^{[u]} \rangle}^\alpha$ ,  $\mu_{\langle M^{[u]}, M^{[v]} \rangle}^\alpha$  and  $\mu_{\langle M^{[u]}, M^{[v]} \rangle}^\alpha$ ,  $\alpha = c, d, j, k$ .

**Theorem 4.9.** For  $u, v \in H$ ,

$$(4.19) \quad \overset{j}{\mu}_{\langle u, v \rangle}(dy) = \int_X (\tilde{u}(x) - \tilde{u}(y)) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy)$$

and

$$(4.20) \quad \overset{k}{\mu}_{\langle u, v \rangle}(dy) = \tilde{u}(y) \tilde{v}(y) \chi(dy).$$

Proof. By Theorem 3.6, for any  $f \in \mathbf{B}^+(X)$

$$(4.21) \quad \begin{aligned} \int_X f(y) \overset{j}{\mu}_{\langle u, v \rangle}(dy) &= \lim_{t \downarrow 0} \frac{1}{t} E_m[(f \langle \overset{j}{M}^{[u]}, \overset{j}{M}^{[v]} \rangle)_t] \\ &= \lim_{t \downarrow 0} \frac{1}{t} E_m[(f([\overset{j}{M}^{[u]}, \overset{j}{M}^{[v]}])^\flat)_t] \end{aligned}$$

where  $[\overset{j}{M}^{[u]}, \overset{j}{M}^{[v]}]_t = \sum_{0 < s \leq t < \zeta} \Delta \overset{j}{M}_s^{[u]} \Delta \overset{j}{M}_s^{[v]}$ . Hence the right hand side of (4.21) is equal to

$$\begin{aligned} & \lim_{t \downarrow 0} \frac{1}{t} E_m \left[ \sum_{0 < s \leq t < \zeta} f(X_s) (\tilde{u}(X_s) - \tilde{u}(X_{s-})) (\tilde{v}(X_s) - \tilde{v}(X_{s-})) I_{\{X_s \neq X_{s-}\}} \right] \\ &= \lim_{t \downarrow 0} \frac{1}{t} E_m \left[ \int_0^t dH_s \int_X f(X_s) (\tilde{u}(x) - \tilde{u}(X_s)) (\tilde{v}(x) - \tilde{v}(X_s)) N(X_s, dx) \right] \\ &= \langle \int_X f(\cdot) (\tilde{u}(x) - \tilde{u}(\cdot)) (\tilde{v}(x) - \tilde{v}(\cdot)) N(\cdot, dx) \nu, 1 \rangle \end{aligned}$$

by Theorem 3.6 (v). This and Theorem 3.8 (i) imply (4.91). (4.20) can be proved by the same calculus as above. The proof is complete.

REMARK 4.10. Since  $\overset{d}{\mu}_{\langle u, v \rangle} = \overset{j}{\mu}_{\langle u \rangle} + \overset{k}{\mu}_{\langle v \rangle}$  ( $u, v \in \mathbf{H}$ ), it follows that

$$(4.22) \quad \overset{d}{\mu}_{\langle u, v \rangle}(dy) = \int_{X \cup \{\delta\}} (\tilde{u}(x) - \tilde{u}(y)) (\tilde{v}(x) - \tilde{v}(y)) N(y, dx) \nu(dy).$$

**Corollary 4.11.**  $\frac{1}{2} \overset{c}{\mu}_{\langle u, v \rangle}(X) = N(u, v) \quad (u, v \in \mathbf{H}).$

Proof. By (4.12) and Theorem 4.9, we have for any  $u, v \in H$ ,

$$\begin{aligned} & \frac{1}{2} \mathbf{a}(u, v) + \frac{1}{2} \hat{\mathbf{a}}(u, v) = e(M^{[u]}, M^{[v]}) + \frac{1}{2} \int_X \tilde{u}(x) \tilde{v}(x) \hat{\chi}(dx) \\ &= \frac{1}{2} \overset{c}{\mu}_{\langle u, v \rangle}(X) + \frac{1}{2} \int_X \tilde{u}(x) \tilde{v}(x) \hat{\chi}(dx) \quad \text{by (4.3)} \\ &= \frac{1}{2} \overset{c}{\mu}_{\langle u, v \rangle}(X) + \frac{1}{2} \overset{j}{\mu}_{\langle u \rangle}(X) + \frac{1}{2} \overset{k}{\mu}_{\langle v \rangle}(X) \\ &\quad + \frac{1}{2} \int_X \tilde{u}(x) \tilde{v}(x) \hat{\chi}(dx) \\ &= \frac{1}{2} \overset{c}{\mu}_{\langle u \rangle}(X) + \frac{1}{2} \int_{X \times X - \Delta} (\tilde{u}(x) - \tilde{u}(y)) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy) \\ &\quad + \frac{1}{2} \int_X \tilde{u}(x) \tilde{v}(x) \chi(dx) + \frac{1}{2} \int_X \tilde{u}(x) \tilde{v}(x) \hat{\chi}(dx). \end{aligned}$$

Comparing this with (2.14), we get the desired equality. The proof is complete.

## 5. Derivation property of $\overset{c}{\mu}_{\langle u \rangle}$ and its applications

**Lemma 5.1.** For  $f, u \in \mathbf{H}_b$  (=the set of bounded functions of  $\mathbf{H}$ ),

$$(5.1) \quad \int_X \tilde{f}(x) d\mu_{\langle u \rangle} = 2\mathbf{a}(u, uf) - \mathbf{a}(u^2, f).$$

Proof. It is sufficient to show that (5.1) holds for all  $f \in \mathbf{H}_b^+$ . By Theorem 3.6 (ii), we have

$$(f, U_{\langle M^{[u]} \rangle}^\alpha 1)_{L^2} = \langle \mu_{\langle u \rangle}, \hat{R}_\alpha f \rangle.$$

Since there exists a subsequence, also written by  $\alpha$ , such that  $\alpha R_\alpha f$  converges to  $\tilde{f}$  q.e. as  $\alpha \rightarrow \infty$ ,

$$\begin{aligned} (5.2) \quad \langle \mu_{\langle u \rangle}, \tilde{f} \rangle &= \lim_{\alpha \rightarrow \infty} \alpha (f, U_{\langle M^{[u]} \rangle}^\alpha 1)_{L^2} \\ &= \lim_{\alpha \rightarrow \infty} \alpha E_{f,m} \left[ \int_0^\infty e^{-\alpha t} d\langle M^{[u]} \rangle_t \right] \\ &= \lim_{\alpha \rightarrow \infty} \alpha E_{f,m} \left[ e^{-\alpha t} \langle M^{[u]} \rangle_t \right]_{t=0}^\infty \\ &\quad + \lim_{\alpha \rightarrow \infty} \alpha^2 E_{f,m} \left[ \int_0^\infty e^{-\alpha t} \langle M^{[u]} \rangle_t dt \right] \\ &= \lim_{\alpha \rightarrow \infty} \alpha^2 E_{f,m} \left[ \int_0^\infty e^{-\alpha t} (M_t^{[u]})^2 dt \right], \end{aligned}$$

since  $E_m[\langle M^{[u]} \rangle_t] = E_m[(M_t^{[u]})^2] \leq ct$  for some constant  $c \geq 0$ . Since

$$\lim_{\alpha \rightarrow \infty} \alpha^2 E_{f,m} \left[ \int_0^\infty e^{-\alpha t} (M_t^{[u]})^2 dt \right] = 0,$$

the right hand side of (5.2) is equal to

$$\begin{aligned} &\lim_{\alpha \rightarrow \infty} \alpha^2 E_{f,m} \left[ \int_0^\infty e^{-\alpha t} (\tilde{u}(X_t) - \tilde{u}(X_0))^2 dt \right] \\ &= 2\mathbf{a}(u, uf) - \mathbf{a}(u^2, f) \end{aligned}$$

by the same calculus as the proof of Lemma 4.5. The proof is complete.

Using the fact  $\mu_{\langle u+v \rangle} = 1/2(\mu_{\langle u+v \rangle} - \mu_{\langle u \rangle} - \mu_{\langle v \rangle})$ , Lemma 5.1 implies the following corollary.

**Corollary 5.2.** For  $f, u, v \in \mathbf{H}_b$ ,

$$\int_X \tilde{f} d\mu_{\langle u,v \rangle} = \mathbf{a}(u, fv) + \mathbf{a}(v, fu) - \mathbf{a}(uv, f).$$

**Lemma 5.3.** Let  $u_n \in \mathbf{H}$  converge to  $u \in \mathbf{H}$  with  $\| \cdot \|_{L^2}$  as  $n \rightarrow \infty$  and let  $\mathbf{a}(u_n, u_n)$  be uniformly bounded. Then

- (1)  $\mathbf{a}(u_n - u, v) \rightarrow 0$  as  $n \rightarrow \infty$  for all  $v \in \mathbf{H}$ .
- (2)  $\mathbf{a}(v, u_n - u) \rightarrow 0$  as  $n \rightarrow \infty$  for all  $v \in \mathbf{H}$ .

Proof. We only prove (1); The proof of (2) is similar. In the case  $v = \hat{R}_1 f$  ( $f \in L^2(X, m)$ ), we have

$$\mathbf{a}(u_n - u, \hat{R}_1 f) = (u_n - u, f)_{L^2} - (u_n - u, \hat{R}_1 f)_{L^2} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

For general  $v \in H$  and any  $\varepsilon > 0$ , there exists  $w \in \text{Range}(R_1)$  such that  $\|v - w\|_H < \varepsilon$ . Then

$$\limsup_{n \rightarrow \infty} |\mathbf{a}(u_n - u, v)| \leq \limsup_{n \rightarrow \infty} |\mathbf{a}(u_n - u, v - w)| + \limsup_{n \rightarrow \infty} |\mathbf{a}(u_n - u, w)| \leq C \cdot \varepsilon.$$

The proof is complete.

**Theorem 5.4** (derivation property of  $\mu_{\langle u \rangle}^c$ ). *It holds that*

$$(5.3) \quad d\mu_{\langle uv, w \rangle}^c = \tilde{u} d\mu_{\langle v, w \rangle}^c + \tilde{v} d\mu_{\langle u, w \rangle}^c \quad \text{for } u, v, w \in H_b.$$

*Proof.* This can be proved by the same method as Y. Le Jan [17] (also cf. [8], [9]), but we give an alternative proof based on the martingale theory. It is sufficient to show that for  $u, v \in H_b$ ,

$$(5.4) \quad d\mu_{\langle u^2, v \rangle}^c = 2\tilde{u} d\mu_{\langle u, v \rangle}^c.$$

Let  $\alpha_n$  be a sequence such that  $u_n = \alpha_n R_{\alpha_n} u$  converges to  $\tilde{u}$  q.e. as  $\alpha_n \rightarrow \infty$ . By Theorem 4.8 and its proof, we have the expressions;

$$\begin{aligned} u_n(X_t) - u_n(X_0) &= M_t^{[u_n]} + N_t^{[u_n]}, \\ u_n^2(X_t) - u_n^2(X_0) &= M_t^{[u_n^2]} + N_t^{[u_n^2]}, \end{aligned}$$

where  $M^{[u_n]}, M^{[u_n^2]} \in \mathring{\mathcal{M}}$  and  $N^{[u_n]}, N^{[u_n^2]} \in \mathcal{N}_c$ . Moreover  $N^{[u_n]}$  is of bounded variation. By Ito's formula,

$$u_n^2(X_t) - u_n^2(X_0) = 2 \int_0^t u_n(X_{s-}) dM_s^{[u_n]} + A_t,$$

where  $A$  is an AF of bounded variation. It then follows that

$$(5.5) \quad \langle \dot{M}^{[u_n^2]}, \dot{M}^{[v]} \rangle_t + \langle N^{[u_n^2]}, \dot{M}^{[v]} \rangle_t = 2 \int_0^t u_n(X_s) d\langle \dot{M}^{[u_n]}, \dot{M}^{[v]} \rangle_s.$$

We now prove that

$$(5.6) \quad \langle N^{[u_n^2]}, \dot{M}^{[v]} \rangle = 0 \quad P_m\text{-a.e.},$$

which implies that  $\langle N^{[u_n^2]}, \dot{M}^{[v]} \rangle = 0$   $P_x$ -a.e., q.e.  $x$ . To see this it suffices to show that

$$(5.7) \quad \langle N^{[u_n^2]} \rangle = 0 \quad P_m\text{-a.e.}$$

Write  $N$  for  $N^{[u_n^2]}$ . Since

$$E_x[\langle N \rangle_t] = \lim_{k \rightarrow \infty} \sum_{0 \leq i < 2^k} E_x(N_{(i+1)/2^k} - N_{i/2^k})^2 \text{ q.e. } x$$

and since

$$\begin{aligned} \sum_{0 \leq i < 2^k} E_m (N_{(i+1)t/2^k} - N_{it/2^k})^2 &= \sum_{0 \leq i < 2^k} E_m [E_{X_{it/2^k}} [N_{it/2^k}^2]] \\ &\leq 2^k E_m [N_{t/2^k}^2], \end{aligned}$$

we have

$$E_m [\langle N \rangle_t] \leq \liminf_{k \rightarrow \infty} 2^k E_m [N_{t/2^k}^2].$$

Hence

$$\begin{aligned} \int_0^\infty e^{-t} E_m [\langle N \rangle_t] dt &\leq \liminf_{k \rightarrow \infty} 2^k E_m \left[ \int_0^\infty e^{-t} N_{t/2^k}^2 dt \right] \\ &= \liminf_{k \rightarrow \infty} (2^k)^2 E_m \left[ \int_0^\infty e^{-2^k t} N_t^2 dt \right] = 0, \end{aligned}$$

which proves (5.7). By (5.5), (5.6) and Theorem 3.6, we have

$$(5.8) \quad d\mu_{\langle u_n^2, v \rangle}^c = 2u_n d\mu_{\langle u_n, v \rangle}^c.$$

Now we prove that for  $f \in C_0(X) \cap H$ ,

$$(5.9) \quad \lim_{n \rightarrow \infty} \int_X f d\mu_{\langle u_n^2, v \rangle}^c = \int_X f d\mu_{\langle u^2, v \rangle}^c.$$

In fact, for  $f \in C_0(X) \cap H$ ,

$$\begin{aligned} & \left| \int_X f d\mu_{\langle u_n^2, v \rangle}^c - \int_X f d\mu_{\langle u^2, v \rangle}^c \right| \\ & \leq |a(u_n^2 - u^2, fv)| + |a(v, fu_n^2 - fu^2)| + |a(u_n^2 v - u^2 v, f)| \end{aligned}$$

by Corollary 5.2, and the right hand side converges to 0 as  $n \rightarrow \infty$  by virtue of Lemma 5.3. Thus we have

$$(5.10) \quad \lim_{n \rightarrow \infty} \int_X f d\mu_{\langle u_n^2, v \rangle}^c = \int_X f d\mu_{\langle u^2, v \rangle}^c.$$

On the other hand, by Remark 4.10,

$$\begin{aligned} (5.11) \quad \int_X f d\mu_{\langle u_n^2, v \rangle}^d &= \int_{X \times X - \Delta} f(y) (u_n(x)^2 - u_n(y)^2) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy) \\ &\quad + \int_X f(y) u_n(y)^2 \tilde{v}(y) \chi(dy). \end{aligned}$$

Here

$$\int_{X \times X - \Delta} f(y) (u_n(x)^2 - u_n(y)^2) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy)$$

$$-\int_{X \times X - \Delta} f(y) (\tilde{u}(x)^2 - \tilde{u}(y)^2) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy)$$

= I + II,

where

$$\begin{aligned} \text{I} &= \int_{X \times X - \Delta} f(y) (u_n(x) + u_n(y)) [(u_n(x) - \tilde{u}(x)) (u_n(y) - \tilde{u}(y))] \\ &\quad \times (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy) \end{aligned}$$

and

$$\begin{aligned} \text{II} &= \int_{X \times X - \Delta} f(y) (u_n(x) + u_n(y) - \tilde{u}(x) - \tilde{u}(y)) (\tilde{u}(x) - \tilde{u}(y)) \\ &\quad \times (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy). \end{aligned}$$

Firstly we have

$$\begin{aligned} |\text{I}| &\leq 2 \|f\|_\infty \|u\|_\infty \left[ \int_{X \times X - \Delta} ((\tilde{u} - u_n)(x) - (\tilde{u} - u_n)(y))^2 \sigma(dx, dy) \right]^{1/2} \\ &\quad \times \left[ \int_{X \times X - \Delta} (\tilde{v}(x) - \tilde{v}(y))^2 \sigma(dx, dy) \right]^{1/2} \\ &\leq 2 \|f\|_\infty \|u\|_\infty \mathbf{a}(u - u_n, u - u_n)^{1/2} \mathbf{a}(v, v)^{1/2} \\ &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Secondly, since

$$\int_{X \times X - \Delta} |\tilde{u}(x) - \tilde{u}(y)| |\tilde{v}(x) - \tilde{v}(y)| \sigma(dx, dy) < \infty,$$

the measure  $\mu(dx, dy) = |\tilde{u}(x) - \tilde{u}(y)| |\tilde{v}(x) - \tilde{v}(y)| \sigma(dx, dy)$  on  $X \times X - \Delta$  is bounded. Let  $N$  be a properly exceptional set such that  $u_n(x) \rightarrow \tilde{u}(x)$  as  $n \rightarrow \infty$  for all  $x \in X - N$ . Then, since  $\sigma(dz, dy) = N(y, dx) \nu(dy) = \hat{N}(x, dy) \hat{\nu}(dx) = \hat{\sigma}(dy, dx)$ , we have  $\mu(X \times N) = \mu(N \times X) = 0$ . Hence

$$(u_n(x) - u_n(y) - \tilde{u}(x) + \tilde{u}(y)) \rightarrow 0 \text{ as } n \rightarrow \infty \text{ } \mu\text{-a.e.}$$

This implies that  $|\text{II}| \rightarrow 0$  as  $n \rightarrow \infty$ . Using the same method, the second term of the right hand side of (5.11) converges to

$$\int_X f(y) \tilde{u}(y)^2 \tilde{v}(y) \chi(dy)$$

as  $n \rightarrow \infty$ . Thus

$$\begin{aligned} (5.12) \quad \lim_{n \rightarrow \infty} \int_X f d\mu_{\langle u_n^2, v \rangle}^d &= \int_{X \times X - \Delta} f(y) (\tilde{u}(x)^2 - \tilde{u}(y)^2) (\tilde{v}(x) - \tilde{v}(y)) \sigma(dx, dy) \\ &\quad + \int_X f(y) \tilde{u}(y)^2 \tilde{v}(y) \chi(dy) \end{aligned}$$

$$= \int_X f d\mu_{\langle u^2, v \rangle}^d.$$

Then (5.9) holds by (5.10) and (5.12). From the fact that

$$(5.13) \quad |\mu_{\langle u_n, v \rangle}^c - \mu_{\langle u, v \rangle}^c|(X) \leq \mu_{\langle u_n - u \rangle}^c(X)^{1/2} \mu_{\langle v \rangle}^c(X)^{1/2},$$

it is easy to show that

$$\lim_{n \rightarrow \infty} \int_X f u_n d\mu_{\langle u_n, v \rangle}^c = \int_X f u d\mu_{\langle u, v \rangle}^c.$$

Combining this with (5.8) and (5.9), we have (5.4). The proof is complete.

**Corollary 5.5.** *Let  $v, w \in \mathbf{H}$  and  $v = r = \text{constant}$   $m$ -a.e. on an open set  $D \subset X$ . Then*

$$\mu_{\langle v, w \rangle}^c = 0 \text{ on } D.$$

*Proof.* Let  $v \in \mathbf{H}_b$ . Then for any  $u \in C_0(X) \cap \mathbf{H}$  such that

$$\begin{aligned} r d\mu_{\langle u, w \rangle}^c &= d\mu_{\langle ru, w \rangle}^c \text{ by Theorem 5.4,} \\ &= d\mu_{\langle vu, w \rangle}^c \\ &= \tilde{v} d\mu_{\langle u, w \rangle}^c + u d\mu_{\langle v, w \rangle}^c \\ &= r d\mu_{\langle u, w \rangle}^c + u d\mu_{\langle v, w \rangle}^c \text{ on } D. \end{aligned}$$

Hence

$$u d\mu_{\langle v, w \rangle}^c = 0 \text{ on } D.$$

Since  $u$  is arbitrary, we have

$$\mu_{\langle v, w \rangle}^c = 0 \text{ on } D.$$

Let  $v \in \mathbf{H}$  and  $v_n = ((-n) \vee v) \wedge n$ . Then since  $v_n \in \mathbf{H}_b$  and it converges to  $v$  with  $\|\cdot\|_H$ , it follows by (5.13) that

$$\mu_{\langle v_n, w \rangle}^c \rightarrow \mu_{\langle v, w \rangle}^c \text{ vaguely as } n \rightarrow \infty,$$

which completes the proof.

**REMARK 5.6.** Let  $u_1, u_2 \in \mathbf{H}$  satisfy that  $u_1 - u_2 = r = \text{constant}$   $m$ -a.e. on an open set  $D \subset X$ . Then, by Corollary 5.5,  $I_D \mu_{\langle u_1 - u_2 \rangle}^c = 0$ . Since  $I_D \mu_{\langle u_1 - u_2 \rangle}^c$  is the smooth measure corresponding to  $\langle M^{[u_1 - u_2]} \rangle_{\sigma_M \wedge I} \in A_c^+(D)$  ( $M = D^c$ ), we have

$$(5.14) \quad \bar{M}_t^{[u_1]} - \bar{M}_t^{[u_2]} = \bar{M}_t^{[u_1 - u_2]} = 0 \text{ for all } t < \sigma_{X-D}.$$

**DEFINITION 5.7.** For AF  $A$ , we define the support of  $A$  by



$$\text{supp}[A] = \{x \in X - N : P_x(R = 0) = 1\},$$

where  $N$  is the exceptional set of  $A$  and  $R = \inf \{t : A_t \neq 0\}$ .

**Theorem 5.8.** *It holds that for  $u \in H$ ,*

$$\text{supp}[\dot{M}^{[u]}] \subset \overline{\text{supp}[u]} \text{ q.e. ,}$$

where  $\overline{\text{supp}[u]} = \{x \in X : \text{for some neighbourhood } U \text{ of } x, u \text{ is constant } m\text{-a.e. on } U\}^c$ .

*Proof.* Let  $\{O_p\}_{p=1}^\infty$  be a sequence of open sets of  $X$  satisfying the following conditions;

$$(i) \quad \bigcup_{p=0}^\infty O_p = X - \overline{\text{supp}[u]},$$

$$(ii) \quad u|_{O_p} = \text{constant } m\text{-a.e. on } O_p.$$

By Remark 5.6, we have

$$\dot{M}_t^{[u]} = \dot{M}_t^{[0]} \quad (t < \sigma_{X-O_p}),$$

where 0 is the zero valued function. It is trivial that  $\dot{M}_t^{[0]} = 0$  for any  $t < \sigma_{X-O_p}$ . Hence

$$O_p \subset (\text{supp}[\dot{M}^{[u]}])^c \text{ q.e. for any } p.$$

Since  $p$  is arbitrary, this completes the proof.

**Theorem 5.9.** *It holds that for  $u, v \in H$*

$$\text{supp}[\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle] \subset \overline{\text{supp}[u]} \cap \overline{\text{supp}[v]} \text{ q.e. .}$$

*Proof.* Since

$$|\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle_t| \leq (\langle \dot{M}^{[u]} \rangle_t)^{1/2} (\langle \dot{M}^{[v]} \rangle_t)^{1/2},$$

we have

$$\begin{aligned} \text{supp}[\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle] &\subset \text{supp}[\langle \dot{M}^{[u]} \rangle] \cap \text{supp}[\langle \dot{M}^{[v]} \rangle] \\ &= \text{supp}[\dot{M}^{[u]}] \cap \text{supp}[\dot{M}^{[v]}] \\ &\subset \overline{\text{supp}[u]} \cap \overline{\text{supp}[v]}, \end{aligned}$$

by Lemma 5.8. The proof is complete.

**Corollary 5.10** (stronger local property of the form  $N$ ). *The symmetric form  $N$  satisfies the following condition;*

$$N(u, v) = 0 \quad \text{for all } u, v \in H \text{ such that } \overline{\text{supp}[u]} \cap \overline{\text{supp}[v]} = \emptyset.$$

*Proof.* Since  $\overline{\text{supp}[u]} \cap \overline{\text{supp}[v]} = \emptyset$ , by Theorem 5.9, it holds that  $\text{supp}$

$[\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle] = \phi$ . Hence there exists a properly exceptional set  $N$  such that  $\text{supp}[\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle] \subset N$ . Let  $R(\omega) = \inf \{t: \langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle_t(\omega) \neq 0\}$ . Then  $P_x(R(\omega) > 0) = 1$  and  $P_x[\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle_R = 0] = 1$ . This implies that  $E_x[\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle_t] = 0$  for all  $t > 0$  and all  $x \in X - N$  (see P.A. Meyer [13]), i.e.,  $\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle_t = 0$  for all  $t > 0$ . Hence  $N(u, v) = \frac{1}{2} \mu_{\langle u, v \rangle}(X) = e(\langle \dot{M}^{[u]}, \dot{M}^{[v]} \rangle) = 0$ . The proof is complete.

## 6. Examples

As was stated in Section 1, we extend slightly the notion of Dirichlet form according to H. Kunita [10]. A form  $\mathbf{a}$  is said to be a *general Dirichlet form* if it satisfies (a.2), (a.3), (a.4) and the following, weaker than (a.1), condition; (a.1)' There exists a constant  $\alpha_0 \geq 0$  such that  $\mathbf{a}_\alpha$  is coercive for any  $\alpha > \alpha_0$ . All the results in the first half of Section 2 concerning the Dirichlet form  $\mathbf{a}$  hold for the general Dirichlet form  $\mathbf{a}$  if  $\alpha > \alpha_0$ ,  $\mathbf{a}_1$ ,  $u_A^1$  and  $U_1\mu$  there are replaced by  $\alpha > \alpha_0$ ,  $\mathbf{a}_{\alpha_0+1}$ ,  $u_A^{\alpha_0+1}$  and  $U_{\alpha_0+1}\mu$  ([4], [12]). It is easy to see that the Beurling-Deny formula (2.8) also holds for the general Dirichlet form. Since any regular general Dirichlet space also admits an associated Hunt process (see [7]) and our arguments in preceding sections only involve  $\mathbf{a}_\alpha$  for  $\alpha$  large enough, all of the previous assertions for the Dirichlet form persist to hold for the general Dirichlet form. In this section we will give several examples of regular general Dirichlet spaces.

[I]. Let  $D$  be a relatively compact open subset of  $R^d$  and  $H^1(D)$  the Sobolev space of order 1, i.e.,

$$H^1(D) = \{u \in L^2(D): \frac{\partial u}{\partial x_i} \in L^2(D), 1 \leq i \leq d\},$$

where the derivatives  $\frac{\partial u}{\partial x_i}$  are taken in the sense of Schwartz distributions. We also consider the space  $H_0^1(D)$  the closure of  $C_0^\infty(D)$  in  $H^1(D)$ . Let  $dx$  be the Lebesgue measure on  $D$  and  $L^2 = L^2(D, dx)$ . We define the norm on  $H^1(D)$  by

$$\|u\|_{H^1} = \|u\|_{L^2} + \|u_x\|_{L^2} \quad \text{for } u \in H^1(D),$$

where

$$\|u_x\|_{L^2} = \left( \sum_{i=1}^d \left\| \frac{\partial u}{\partial x_i} \right\|_{L^2}^2 \right)^{1/2}.$$

Consider the following formal generator

$$(6.1) \quad L^0 u = \sum_{i,j=1}^d \frac{\partial}{\partial x_j} \left( a_{ij} \frac{\partial u}{\partial x_i} \right) + \sum_{i=1}^d b_i \frac{\partial u}{\partial x_i} - cu,$$

where  $a_{ij}$ ,  $b_i$ ,  $i, j=1, 2, \dots, d$ , and  $c$  are bounded measurable functions on  $D$ . The bilinear form  $\mathbf{a}^0$  corresponding to  $L^0$  is given by

$$(6.2) \quad \mathbf{a}^0(u, v) = \sum_{i,j=1}^d \int_D a_{ij} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx - \sum_{i=1}^d \int_D b_i v \frac{\partial u}{\partial x_i} dx + \int_D c u v dx$$

for  $u, v \in H^1(D)$ . It is clear that  $\mathbf{a}^0$  satisfies the condition (a.2). Moreover we assume that  $L^0$  is uniformly elliptic, i.e., there exists a constant  $\nu > 0$  such that

$$(6.3) \quad \sum_{i,j=1}^d a_{ij}(x) y_i y_j \geq \nu |y|^2 \quad \text{for all } x \in D$$

(( $a_{ij}$ ) is not necessarily symmetric). Then  $\mathbf{a}^0$  satisfies the condition (a.1)'. In fact, for any  $\alpha > 0$ ,

$$(6.4) \quad \mathbf{a}^0(u, u) + \alpha(u, u)_{L^2} \geq \nu \|u_x\|_{L^2}^2 - \sqrt{d} \|b\|_{\infty} \|u_x\|_{L^2} \|u\|_{L^2} + (\alpha - \|c\|_{\infty}) \|u\|_{L^2}^2$$

Since

$$(\nu + \sqrt{d} \|b\|_{\infty}) \|u_x\|_{L^2} \|u\|_{L^2} \leq \frac{\nu}{2} \|u_x\|_{L^2}^2 + \frac{1}{2\nu} (\nu + \sqrt{d} \|b\|_{\infty})^2 \|u\|_{L^2}^2,$$

the right hand side of (6.4) is not smaller than

$$(6.5) \quad \frac{\nu}{2} \|u_x\|_{L^2}^2 + (\alpha - \|c\|_{\infty} - \frac{1}{2\nu} (\nu + \sqrt{d} \|b\|_{\infty})^2) \|u\|_{L^2}^2 + \nu \|u_x\|_{L^2} \|u\|_{L^2}.$$

Choose  $\alpha_0 = \frac{\nu}{2} + \frac{1}{2\nu} (\nu + \sqrt{d} \|b\|_{\infty})^2 + \|c\|_{\infty}$ . Then

$$\mathbf{a}^0(u, u) + \alpha(u, u) \geq \frac{\nu}{2} (\|u_x\|_{L^2}^2 + \|u\|_{L^2}^2) = \frac{\nu}{2} \|u\|_{H^1}^2$$

for any  $\alpha > \alpha_0$ , and hence  $\mathbf{a}_\alpha^0$  is coercive for any  $\alpha > \alpha_0$ . Now we assume that

$$(6.6) \quad c \geq 0.$$

Since

$$(6.7) \quad \mathbf{a}^0(T_1 u, u - T_1 u) = \int_{u \geq 1} c(u-1) dx \geq 0,$$

$\mathbf{a}^0$  satisfies the condition (a.3). Now we assume that

$$(6.8) \quad c + \sum_{i=1}^d (b_i)_{x_i} \geq 0 \quad (\text{in the sense of distributions}).$$

Then

$$\mathbf{a}^0(u - T_1 u, T_1 u) = - \sum_{i=1}^d \int_{u \geq 1} b_i \frac{\partial(u-1)}{\partial x_i} dx + \int_{u \geq 1} c(u-1) dx,$$

i.e.,  $\mathbf{a}^0$  satisfies the condition (a.4). If  $D = R^d$ , then  $H^1(D)$  is regular.

Summarizing above we have the following theorem.

**Theorem 6.1.** *Let  $\mathbf{a}^0$  be the bilinear form given by (6.2). If the conditions (6.3), (6.6) and (6.8) are satisfied, then  $(\mathbf{a}^0, H^1(D))$  and  $(\mathbf{a}^0, H_0^1(D))$  are general Dirichlet spaces on  $L^2(D, dx)$ . Moreover the latter one is regular.*

REMARK ([19]). Let

$$L^\infty + L^n = \{b: b = b^1 + b^2, b^1 \in L^\infty, b^2 \in L^n\}.$$

Theorem 6.1 still holds if the boundedness assumption for  $b_i$  and  $c$  is replaced by the weaker condition that  $b_i \in L^\infty + L^d$ ,  $i=1, 2, \dots, d$  and  $c \in L^\infty + L^{d/2}$ .

Let  $D = R^d$  and consider the following differential operator given formally by

$$(6.9) \quad L = L^0 + \sum_{i \neq j}^d \delta(x_i - x_j) \frac{\partial}{\partial x_i}.$$

Then (6.9) can be rewritten as

$$(6.10) \quad L = L^0 + \sum_{i,j=1}^d \frac{\partial}{\partial x_j} H(x_j - x_i) \frac{\partial}{\partial x_i},$$

where  $H(y) = 1/2(I_{\{y>0\}} - I_{\{y<0\}})$ . Now we have for any  $u, v \in C_0^\infty(R^d)$ ,

$$\begin{aligned} \left( \frac{\partial}{\partial x_j} H(x_j - x_i) \frac{\partial u}{\partial x_i}, v \right)_{L^2} &= - \left( H(x_j - x_i) \frac{\partial^2 u}{\partial x_j \partial x_i}, v \right)_{L^2} \\ &\quad - \left( H(x_j - x_i) \frac{\partial u}{\partial x_i}, \frac{\partial v}{\partial x_j} \right)_{L^2}, \end{aligned}$$

and the first term of the right hand side is antisymmetric in  $i, j$  because  $H(y) = -H(-y)$ . Hence the bilinear form  $\mathbf{a}$  corresponding to  $L$  is given by, for  $u, v \in H^1(R^d)$ ,

$$(6.11) \quad \mathbf{a}(u, v) = \mathbf{a}^0(u, v) + \sum_{i,j=1}^d \int_{R^d} H(x_j - x_i) \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx,$$

and accordingly

$$(6.12) \quad \mathbf{a}(u, v) = \sum_{i,j=1}^d \int_{R^d} \tilde{a}_{ij} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx - \sum_{i=1}^d \int_{R^d} b_i v \frac{\partial u}{\partial x_i} dx + \int_{R^d} c u v dx,$$

where  $\tilde{a}_{ij} = a_{ij} + H(x_j - x_i)$ ,  $i, j=1, 2, \dots, d$ . These  $\tilde{a}_{ij}$ ,  $i, j=1, 2, \dots, d$ , are

bounded measurable functions on  $R^d$  satisfying the condition (6.3). Therefore, by Theorem 6.1, the bilinear form  $\mathbf{a}$  given by (6.11) is a general Dirichlet form which is regular and consequently generates a diffusion process on  $R^d$  by virtue of S. Carrillo Menendez [5]. The present argument can be applied to the form corresponding to formal generator

$$L = L^0 + \sum_{i,j=1}^d \frac{\partial}{\partial x_j} d_{ij} \frac{\partial}{\partial x_i},$$

if the condition

$$\sum_{i,j=1}^d d_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} = 0 \quad (u \in C_0^\infty(D))$$

holds. Under a more general condition that

$$\sum_{i,j=1}^d \int d_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} = 0,$$

H. Osada [16] has given a specific construction of the diffusion by means of the associated transition density function.

[II]. Next we consider the case with boundary conditions.

Let  $D = \{x = (\xi, x_d) \in R^d: \xi \in R^{d-1}, x_d \in R^1 \text{ and } x_d > 0\}$ ,  $\partial D = \{x \in R^d: x_d = 0\}$  and  $\bar{D} = D \cup \partial D$ , and we consider the formal generator  $L^0$  with the following boundary condition

$$(6.13) \quad \sum_{i=1}^d a_{id} \frac{\partial u}{\partial x_i} + \sum_{i=1}^{d-1} \beta_i \frac{\partial u}{\partial x_i} = 0,$$

where  $\beta_i$ ,  $i=1, 2, \dots, d-1$  are bounded measurable functions on  $\partial D$  with bounded derivatives of first order. Then the bilinear form corresponding to  $L^0$  with boundary condition (6.13) is given by, for  $u, v \in H^1(D)$ ,

$$(6.14) \quad \mathbf{a}(u, v) = \mathbf{a}^0(u, v) - \sum_{i=1}^{d-1} \int_{\partial D} \beta_i(\xi) v(\xi, 0) \frac{\partial u(\xi, 0)}{\partial \xi_i} d\xi.$$

The term  $\int_{\partial D} \beta_i(\xi) v(\xi, 0) \frac{\partial u(\xi, 0)}{\partial \xi_i} d\xi$  in (6.14) makes sense by the following argument. Let  $\gamma$  be a trace operator of  $\partial D$  (see S. Mizohara [14]). Then for  $u \in H^1(D)$ ,  $u(\xi, 0) = \gamma u(\xi) \in H^{1/2}(\partial D, d\xi)$ , where  $H^{1/2}(\partial D, d\xi) = \{f \in L^2(\partial D): (1 + |z|)^{1/2} \hat{f}(z) \in L^2(\partial D), \hat{f}(z) \text{ is the Fourier transform of } f(\xi)\}$ , and  $\frac{\partial u(\xi, 0)}{\partial \xi_i} = \frac{\partial \gamma u(\xi)}{\partial \xi_i} \in H^{-1/2}(\partial D, d\xi)$ , where  $H^{-1/2}(\partial D, d\xi)$  is the dual space of  $H^{1/2}(\partial D, d\xi)$ .

Then  $\int_{\partial D} \beta_i(\xi) v(\xi, 0) \frac{\partial u(\xi, 0)}{\partial \xi_i} d\xi$  is understood as the pairing  $\langle \frac{\partial u(\xi, 0)}{\partial \xi_i}, \beta_i(\xi) v(\xi, 0) \rangle$ .

$v(\xi, 0) \rangle$  of  $\frac{\partial u(\xi, 0)}{\partial \xi_i}$  and  $\beta_i(\xi) v(\xi, 0) \in H^{1/2}(\partial D, d\xi)$ .

**Theorem 6.2.** *Let  $\mathbf{a}^0$  be the form given in Theorem 6.1 and  $\beta_i$  satisfy the condition stated after (6.13). Moreover we assume that*

$$(6.15) \quad \sum_{i=1}^{d-1} (\beta_i)_{\xi_i} \geq 0 \quad (\text{in the sense of distributions}).$$

*Then the bilinear form  $\mathbf{a}$  given by (6.14) is a regular general Dirichlet form on  $\bar{D}$ .*

*Proof.* For any  $u \in C_0^\infty(\bar{D})$ , by integration by parts, we have

$$2 \int_{\partial D} \beta_i(\xi) u(\xi, 0) \frac{\partial u(\xi, 0)}{\partial \xi_i} d\xi = - \int_{\partial D} \frac{\partial \beta_i(\xi)}{\partial \xi_i} u(\xi, 0)^2 d\xi.$$

Thus

$$(6.16) \quad \left| \sum_{i=1}^{d-1} \int_{\partial D} \beta_i(\xi) u(\xi, 0) \frac{\partial u(\xi, 0)}{\partial \xi_i} d\xi \right| \leq C_1 \|u(\xi, 0)\|_{L^2(\partial D)}^2,$$

for some constant  $C_1 > 0$ . By Theorem 3.16 in [14], it holds that for any  $\varepsilon > 0$ ,

$$\|u(\xi, 0)\|_{L^2(\partial D)}^2 \leq \varepsilon \|u_x\|_{L^2(D)}^2 + C(\varepsilon) \|u\|_{L^2(D)}^2,$$

where  $C(\varepsilon)$  is a constant depending on  $\varepsilon > 0$ . Hence for any  $\alpha \geq 0$ ,

$$\begin{aligned} \mathbf{a}(u, u) + \alpha(u, u)_{L^2} &\geq (\nu - C_1 \varepsilon) \|u_x\|_{L^2(D)}^2 - \sqrt{d} \|b\|_\infty \|u_x\|_{L^2} \|u\|_{L^2} \\ &\quad + (\alpha - \|c\|_\infty - C_1 C(\varepsilon)) \|u\|_{L^2}^2 \end{aligned}$$

which implies that  $\mathbf{a}_\alpha$  is coercive on  $C_0^\infty(\bar{D})$  for  $\alpha$  large enough. Similarly  $\mathbf{a}$  satisfies (a.2) for any  $u, v \in C_0^\infty(\bar{D})$ . Since  $C_0^\infty(\bar{D})$  is dense in  $H^1(D)$  with respect to  $\|\cdot\|_{H^1}$ ,  $\mathbf{a}$  satisfies the conditions (a.1)' and (a.2) for  $u, v \in H^1(D)$ . Since  $\mathbf{a}(T_1 u, u - T_1 u) = \mathbf{a}^0(T_1 u, u - T_1 u)$ ,  $\mathbf{a}$  satisfies (a.3). It holds that

$$\mathbf{a}(u - T_1 u, T_1 u) = \mathbf{a}^0(u - T_1 u, T_1 u) - \sum_{i=1}^{d-1} \int_{u \geq 1} \beta_i(\xi) \frac{\partial(u(\xi, 0) - 1)}{\partial \xi_i} d\xi \geq 0,$$

by Theorem 6.1 and the condition (6.15). This completes the proof.

Now we give an alternative proof of (a.1)' and (a.2) for the form  $\mathbf{a}$  of Theorem 6.2, following the idea of S. Agmon [1]. By integration by parts, we have

$$\begin{aligned} &\int_{\partial D} \beta_i(\xi) \frac{\partial u(\xi, 0)}{\partial \xi_i} v(\xi, 0) d\xi \\ &= \int_D \frac{\partial}{\partial x_d} (\beta_i(\xi) v(x)) \frac{\partial u(x)}{\partial x_i} dx + \int_D \beta_i(\xi) v(x) \frac{\partial^2 u(x)}{\partial x_d \partial x_i} dx \end{aligned}$$

$$\begin{aligned}
&= \int_D \beta_i(\xi) \frac{\partial u(x)}{\partial x_i} \frac{\partial v(x)}{\partial x_d} dx - \int_D \frac{\partial}{\partial x_i} (\beta_i(\xi) v(x)) \frac{\partial u(x)}{\partial x_d} dx \\
&= \int_D \beta_i(\xi) \frac{\partial u(x)}{\partial x_i} \frac{\partial v(x)}{\partial x_d} dx + \int_D (-\beta_i(\xi)) \frac{\partial u(x)}{\partial x_d} \frac{\partial v(x)}{\partial x_i} dx \\
&\quad + \int_D \frac{\partial}{\partial x_i} (-\beta_i(\xi)) \frac{\partial u(x)}{\partial x_d} v(x) dx.
\end{aligned}$$

Hence the bilinear form  $\mathbf{a}$  given by (6.14) is represented by

$$\mathbf{a}(u, v) = \sum_{i,j=1}^d \int_D \tilde{a}_{ij} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx - \sum_{i=1}^d \int_D \tilde{b}_i \frac{\partial u}{\partial x_i} v dx + \int_D c u v dx,$$

where

$$\begin{aligned}
\tilde{a}_{id}(x) &= a_{id}(x) - \beta_i(\xi), \quad i = 1, 2, \dots, d-1, \\
\tilde{a}_{di}(x) &= a_{di}(x) + \beta_i(\xi), \quad i = 1, 2, \dots, d-1, \\
\tilde{a}_{ij}(x) &= a_{ij}(x), \text{ otherwise}
\end{aligned}$$

and

$$\begin{aligned}
\tilde{b}_i(x) &= b_i(x) \quad i = 1, 2, \dots, d-1, \\
\tilde{b}_d(x) &= b_d(x) + \sum_{i=1}^{d-1} \frac{\partial}{\partial x_i} \beta_i(\xi).
\end{aligned}$$

The coefficients  $\tilde{a}_{ij}$ ,  $\tilde{b}_i$ ,  $i, j=1, 2, \dots, d$ , are bounded measurable and  $\tilde{a}_{ij}$  satisfy the condition (6.3). Thus, by the proof of Theorem 6.1, the bilinear form  $\mathbf{a}$  given by (6.14) satisfies (a.1)' and (a.2).

REMARK. Let  $L^0 = \sum_{i=1}^d \frac{\partial^2}{\partial x_i^2}$ . Then the process  $X=(X^1, X^2, \dots, X^d)$  associated with the general Dirichlet form  $\mathbf{a}$  given by (6.14) is expressed by

$$\begin{aligned}
X_t^i &= x^i + B_t^i + \int_{0,t} \beta_i(\tilde{X}_s) d\phi_s, \quad i = 1, 2, \dots, d-1, \\
X_t^d &= x^d + B_t^d + \phi_t
\end{aligned}$$

where  $x^i = X_0^i$ ,  $i=1, 2, \dots, d$ ,  $B=(B^1, B^2, \dots, B^d)$  is  $d$ -dimensional Brownian motion,  $X_t=(\tilde{X}_t, X_t^d)$  and  $\phi_t$  is the local time of  $B_t^d$  (see M. Tsuchiya [21]).

[III]. Finally we consider an example with non-local form which is a slight generalization of Example IV. 3.2 in [5]. Let  $\mathbf{H}=H^1(R^d)$  and  $a_{ij}$ ,  $i, j=1, 2, \dots, d$ , be as in [I]. Let  $\phi$  be a positive measurable function on  $R^d \times R^d$  such that

$$\int_{R^d \times R^d} \phi(x, y)^2 dy dx < \infty$$

and we define the functions  $c_1$  and  $c_2$  on  $R^d$  by

$$c_1(x) = \int_{R^d} \phi(x, y) dy \quad \text{and} \quad c_2(x) = \int_{R^d} \phi(y, x) dy.$$

Assume that  $c_1$  and  $c_2$  are bounded. Let  $c$  be some bounded measurable function satisfying  $c \geq c_1 \vee c_2$ . Consider the operator

$$(6.17) \quad Lu(x) = \sum_{i,j=1}^d \frac{\partial}{\partial x_j} \left( a_{ij} \frac{\partial u(x)}{\partial x_i} \right) + \int_{R^d} \phi(x, y) u(y) dy - c(x) u(x).$$

Then the associated bilinear form  $\mathbf{a}$  is defined by

$$(6.18) \quad \begin{aligned} \mathbf{a}(u, v) = & \sum_{i,j=1}^d \int_{R^d} a_{ij}(x) \frac{\partial u(x)}{\partial x_i} \frac{\partial v(x)}{\partial x_j} dx \\ & - \int_{R^d \times R^d} \phi(x, y) u(y) v(x) dy dx + \int_{R^d} c(x) u(x) v(x) dx. \end{aligned}$$

Then it is easy to see that the form  $\mathbf{a}$  satisfies (a.1)' and (a.2). Furthermore,  $\mathbf{a}$  has another expressions;

$$(6.19) \quad \begin{aligned} \mathbf{a}(u, v) = & \sum_{i,j=1}^d \int_{R^d} a_{ij}(x) \frac{\partial u(x)}{\partial x_i} \frac{\partial v(x)}{\partial x_j} dx \\ & + \int_{R^d \times R^d} u(y) (v(y) - v(x)) \phi(x, y) dx dy \\ & + \int_{R^d} u(x) v(x) (c(x) - c_2(x)) dx \end{aligned}$$

and

$$(6.20) \quad \begin{aligned} \mathbf{a}(v, u) = & \sum_{i,j=1}^d \int_{R^d} a_{ij}(x) \frac{\partial v(x)}{\partial x_i} \frac{\partial u(x)}{\partial x_j} dx \\ & + \int_{R^d \times R^d} u(x) (v(x) - v(y)) \phi(x, y) dy dx \\ & + \int_{R^d} u(x) v(x) (c(x) - c_1(x)) dx. \end{aligned}$$

By (6.20), we have

$$(6.21) \quad \begin{aligned} \mathbf{a}(T_1 u, u - T_1 u) = & \int_{\{x \in R^d : u(x) < 0\} \times R^d} -u(x) T_1 u(y) \phi(x, y) dy dx \\ & + \int_{\{x \in R^d : u(x) > 1\} \times R^d} (u(x) - 1) (1 - T_1 u(y)) \phi(x, y) dy dx \\ & + \int_{\{x \in R^d : u(x) > 1\}} (u(x) - 1) (c(x) - c_1(x)) dx. \end{aligned}$$

Since  $0 \leq T_1 u(y) \leq 1$  for all  $y \in R^d$  and  $c \geq c_1$  and  $\phi$  is positive on  $R^d \times R^d$ , (6.21)



implies that  $\mathbf{a}(T_1 u, u - T_1 u) \geq 0$ . By the same method, (6.19) implies that  $\mathbf{a}(u - T_1 u, T_1 u) \geq 0$ . Hence  $\mathbf{a}$  given by (6.18) is a general Dirichlet form on  $H \times H$ . Moreover, the equalities (6.19) and (6.20) show that the measure  $\sigma(dx, dy) = \phi(x, y) dx dy$  is the jumping measure of  $\mathbf{a}$  and  $\chi(dx) = (c(x) - c_1(x)) dx$  (resp.  $\hat{\chi}(dx) = (c(x) - c_2(x)) dx$ ) is the killing measure of  $\mathbf{a}$  (resp.  $\hat{\mathbf{a}}$ ). And

$$N(u, v) = \sum_{i,j=1}^d \int_{R^d} \frac{1}{2} (a_{ij} + a_{ji}) \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_j} dx$$

is a symmetric form satisfying the stronger local property.

Acknowledgement. The author would like to express the deepest appreciation to Professor T. Watanabe for his valuable suggestions and encouragement. The author also wants to thank Professors M. Fukushima, N. Ikeda, S. Nakao and M. Takeda for helpful advice during the writing of this paper.

### References

- [1] S. Agmon: Lectures on elliptic boundary value problems, D. Van Nostrand Company, Princeton, 1965.
- [2] A. Benveniste and J. Jacod: *Systemes de Lévy des processus de Markov*, Invent. Math. **21** (1973), 183–198.
- [3] J. Bliedtner: *Functional spaces and their exceptional sets*, Seminar on Potential Theory, Lecture Notes in Math. 226 Springer-Verlag, 1970, 1–13.
- [4] J. Bliedtner: *Dirichlet forms on regular functional spaces*, Seminar on Potential Theory, Lecture Notes in Math. 226 Springer-Verlag, 1970, 15–61.
- [5] S. Carrillo Menendez: *Processus de Markov associé à une forme de Dirichlet non symétrique*, Z. Wahrsch. Verw. Gebiete **33** (1975), 139–154.
- [6] S. Carrillo Menendez: *Fonctionnelles additives à un ensemble polaire près*, to appear in Stochastics.
- [7] C. Dellacherie and P.A. Meyer: Probabilités et potentiel; Theories des martingales, Paris, Hermann, 1980.
- [8] M. Fukushima: Dirichlet forms and Markov processes, Amsterdam-Oxford-New York, North-Holland, Tokyo, Kodansha, 1980.
- [9] M. Fukushima and M. Takeda: *A transformation of a symmetric Markov process and the Donsker-Varadhan theory*, Osaka J. Math. **21** (1984), 331–326.
- [10] H. Kunita: *Sub-Markov semigroups in Banach lattices*, Proc. Int. Conf. Functional Analysis and Related Topics, Tokyo, 1969.
- [11] Y. LeJan: *Balayage et formes de Dirichlet*, Z. Wahrsch. Verw. Gebiete **37** (1977), 297–319.
- [12] Y. LeJan: *Mesures associées à une forme de Dirichlet. Applications*, Bull. Soc. Math. France **106** (1978), 61–112.
- [13] P.A. Meyer: *Martingales locales fonctionnelles additives I*. Séminaire de probabilités XII, Lecture Notes in Math. 649, Springer-Verlag, 1978, 775–785.
- [14] S. Mizohata: The theory of partial differential equations, Cambridge Univ.

- Press, London, 1973.
- [15] S. Nakao: *Stochastic calculus for continuous additive functionals of zero energy*, Z. Wahrsch. Verw. Gebiete **68** (1985), 557–578.
  - [16] H. Osada: *Diffusion processes associated with generalized divergence form*, preprint.
  - [17] M.L. Silverstein: *Symmetric Markov processes*, Lecture Notes in Math. 426, Springer-Verlag, 1974.
  - [18] M.L. Silverstein: *Application of the sector condition to the classification of sub-Markovian semigroups*, Trans. Amer. Math. Soc. **244** (1978), 103–146.
  - [19] G. Stampacchia: *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*, Ann. Inst. Fourier **15.1** (1965), 189–256.
  - [20] M. Takeda: *Stochastic calculus related to Dirichlet spaces and applications*, Master Thesis, Osaka University.
  - [21] M. Tsuchiya: *On the stochastic differential equation for a two dimensional Brownian motion with boundary condition*, J. Math. Soc. Japan **32** (1980), 427–436.
  - [22] D.V. Widder: *The Laplace transform*, Princeton Univ. Press, 1946.

Department of Mathematics  
College of Natural Science  
Pusan National University  
Jangjeon-Don, Dongne-Ku  
Pusan 607  
Korea

