

Title	The Y-triangle move does not preserve intrinsic knottedness
Author(s)	Flapan, Erica; Naimi, Ramin
Citation	Osaka Journal of Mathematics. 2008, 45(1), p. 107-111
Version Type	VoR
URL	https://doi.org/10.18910/11893
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

THE Y-TRIANGLE MOVE DOES NOT PRESERVE INTRINSIC KNOTTEDNESS

ERICA FLAPAN and RAMIN NAIMI

(Received March 7, 2006, revised January 31, 2007)

Abstract

We answer the question “Does the Y-triangle move preserve intrinsic knottedness?” in the negative by giving an example of a graph that is obtained from the intrinsically knotted graph K_7 by triangle-Y and Y-triangle moves but is not intrinsically knotted.

1. Introduction

A graph is said to be *intrinsically knotted* (IK) if every embedding of it in \mathbb{R}^3 contains a cycle that is a nontrivial knot. Similarly, a graph is said to be *intrinsically linked* (IL) if every embedding of it in \mathbb{R}^3 contains a nontrivial link. Sachs [5] and Conway and Gordon [1] showed that K_6 , the complete graph on six vertices, is IL. Conway and Gordon [1] also showed that K_7 is IK.


A ∇Y move on an abstract graph consists of removing the edges of a 3-cycle abc in the graph, and then adding a new vertex v and connecting it to each of the vertices a , b , and c , as shown in Fig. 1. The reverse of this move is called a $Y\nabla$ move. Note that in a $Y\nabla$ move, the vertex v cannot have degree greater than three.

Sachs [5] noticed that additional IL graphs can be obtained from K_6 by doing finite sequences of ∇Y and $Y\nabla$ moves on it. Motwani, Raghunathan, and Saran [3] showed that performing a ∇Y move on any IK or IL graph produces a graph with the same property. Robertson, Seymour, and Thomas [4] (Lemmas 1.2 and 5.1 (iii)) proved that a $Y\nabla$ move on any IL graph produces an IL graph again.

It has been an open question whether a $Y\nabla$ move on an IK graph always produces an IK graph again. We prove that the answer is negative, by giving a knotless embedding of a graph G_7 that is obtained from K_7 by ∇Y and $Y\nabla$ moves.

A graph H is a *minor* of another graph G if H can be obtained from G by a finite sequence of edge deletions and contractions and vertex deletions [2]. A graph is said to be *minor minimal* with respect to a property if the graph has that property but no minor of it has the property.

We work with connected, finite, simple graphs, i.e., graphs with no loops (an edge whose endpoints are the same) and no double-edges (two edges with the same pair

Fig. 1. ∇Y and $Y\nabla$ moves.

of endpoints). This is because loops and double-edges do not affect whether or not a graph is IK or IL: they can always be embedded such that they bound small disks with interiors disjoint from the rest of the graph. Thus, in edge contractions and $Y\nabla$ moves on an abstract graph, whenever a double-edge is introduced, one of the two edges is deleted.

2. Description of the graph G_7

We label the seven vertices of the abstract graph K_7 with the letters a through g. We perform the following five ∇Y and two $Y\nabla$ moves on $G_0 = K_7$ to obtain the graph G_7 .

1. $G_0 \rightarrow G_1$ by ∇Y on abc, with new vertex h as center.
2. $G_1 \rightarrow G_2$ by ∇Y on ade, with new vertex i as center.
3. $G_2 \rightarrow G_3$ by ∇Y on afg, with new vertex j as center.
4. $G_3 \rightarrow G_4$ by ∇Y on bdf, with new vertex k as center.
5. $G_4 \rightarrow G_5$ by ∇Y on beg, with new vertex l as center.
6. $G_5 \rightarrow G_6$ by $Y\nabla$ on hij, deleting vertex a.
7. $G_6 \rightarrow G_7$ by $Y\nabla$ on hkl, deleting vertex b.

3. G_7 is not IK

Theorem 1. *The $Y\nabla$ move does not preserve intrinsic knottedness.*

Proof. Recall that K_7 is IK, ∇Y moves preserve IKness, and G_7 is obtained from K_7 by ∇Y and $Y\nabla$ moves. Thus it suffices to prove that the embedding of G_7 shown in Fig. 2 has no nontrivial knots.

Fig. 2 contains seven crossings, numbered 1–7. Note that rotating this diagram by 180° about a horizontal line through its center leaves the embedded graph invariant, swaps crossing 1 with 2, and 6 with 7, and leaves crossings 3, 4, 5 fixed. And rotating the diagram by 180° about a vertical line through the center also leaves the embedded graph invariant, but swaps crossing 1 with 7, 2 with 6, and 3 with 5.

Suppose towards contradiction that this embedded graph contains a nontrivial knot K . The proof consists of the following three steps.

STEP 2. K must contain 1, 4, and 7.

Proof. Suppose, towards contradiction, that K does not contain $g1$. Then it contains at most three crossings, 3, 5, and 7; but ec , cf , and fe form a cycle, and therefore only links contain all three crossings 3, 5, and 7. Hence K contains $g1$. By a symmetric argument, K contains dk . Thus K contains crossings 1, 4, and 7.

STEP 3. K contains exactly one of 3 and 5.

Proof. If it contains both, it will be a link. If it contains neither, it will be trivial, since 1 and 4 do not alternate.

So, by symmetry, we can assume that K contains 1, 3, 4, and 7, and no other crossings. As K does not contain ij , this implies that K contains di . But $hidk$ is isotopic, with fixed endpoints, to hk . Thus K is isotopic to a knot that contains only crossing 1, and therefore is trivial. \square

ACKNOWLEDGMENTS. The second author thanks Caltech for its hospitality while he worked on this paper during his sabbatical leave.

References

- [1] J.H. Conway and C.McA. Gordon: *Knots and links in spatial graphs*, J. Graph Theory **7** (1983), 445–453.
- [2] R. Diestel: *Graph Theory*, Graduate Texts in Mathematics **173**, Springer-Verlag, New York, 1997.
- [3] R. Motwani, A. Raghunathan and H. Saran: *Constructive results from graph minors: Linkless embeddings*, 29th Annual Symposium on Foundations of Computer Science, IEEE, 1988, 398–409.
- [4] N. Robertson, P. Seymour and R. Thomas: *Sachs' linkless embedding conjecture*, J. Combin. Theory Ser. B **64** (1995), 185–227.
- [5] H. Sachs: *On spatial representations of finite graphs*; in *Finite and Infinite sets*, Vol. I, II (Eger, 1981), Colloq. Math. Soc. János Bolyai **37**, North-Holland, Amsterdam, 1984, 649–662.

Erica Flapan
Mathematics Department
Pomona College
Claremont, CA 91711
USA
e-mail: eflapan@pomona.edu

Ramin Naimi
Mathematics Department
Occidental College
Los Angeles, CA 90041
USA
e-mail: rnaimi@oxy.edu