



Title	On generalized decomposition numbers and Fong's reductions
Author(s)	Watanabe, Atumi
Citation	Osaka Journal of Mathematics. 1985, 22(2), p. 393-400
Version Type	VoR
URL	<a href="https://doi.org/10.18910/11982">https://doi.org/10.18910/11982</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## ON GENERALIZED DECOMPOSITION NUMBERS AND FONG'S REDUCTIONS

Dedicated to Professor Hiroshi Nagao on his 60th birthday

ATUMI WATANABE

(Received August 7, 1984)

### Introduction

In this paper we investigate how generalized decomposition numbers behave under Fong's reductions.

Let  $G$  be a finite group and  $p$  be a fixed prime number. If  $\pi$  is a  $p$ -element of  $G$  and  $B$  is a  $p$ -block of  $G$ , then for an ordinary irreducible character  $\chi$  in  $B$  and for each  $p$ -regular element  $\rho$  of the centralizer  $C_G(\pi)$  of  $\pi$ , we have

$$\chi(\pi\rho) = \sum_{\phi} d(\chi, \pi, \phi)\phi(\rho).$$

Here  $\phi$  ranges over the irreducible Brauer characters in the  $p$ -blocks of  $C_G(\pi)$  associated with  $B$ . We have the following theorem related to the Fong's first reduction.

**Theorem 1.** *Let  $H$  be a subgroup of  $G$ , and let  $B$  and  $\tilde{B}$  be  $p$ -blocks of  $G$  and  $H$ , respectively. We assume that  $\tilde{\chi} \rightarrow \tilde{\chi}^G$  is a 1-1 correspondence between the ordinary irreducible characters in  $\tilde{B}$  and those in  $B$ , where  $\tilde{\chi}^G$  is the character of  $G$  induced from  $\tilde{\chi}$ . Then the following holds.*

- (i)  *$B$  and  $\tilde{B}$  have a common defect group  $D$ .*
- (ii) *Let  $\tilde{b}$  be a root of  $\tilde{B}$  in  $C_H(D)D$ . Then  $\tilde{b}^{C_G(D)D}$  is defined in the sense of Brauer [2]. We put  $b = \tilde{b}^{C_G(D)D}$ . Then  $b$  is a root of  $B$  in  $C_G(D)D$  and  $T(b) = T(\tilde{b})C_G(D)$  where  $T(b)$  is the inertial group of  $b$  in  $N_G(D)$  and  $T(\tilde{b})$  is the inertial group of  $\tilde{b}$  in  $N_H(D)$ . In particular  $T(b)/C_G(D)D \cong T(\tilde{b})/C_H(D)D$ .*
- (iii) *Let  $\{(\pi_i, \tilde{b}_i), i=1, 2, \dots, n\}$  be a set of representatives for the conjugacy classes of subsections associated with  $\tilde{B}$ . Then  $\tilde{b}_i^{C_G(\pi_i)}$  is defined and  $\tilde{\phi} \rightarrow \tilde{\phi}^{C_G(\pi_i)}$  is a 1-1 correspondence between the irreducible Brauer characters in  $\tilde{b}_i$  and those in  $\tilde{b}_i^{C_G(\pi_i)}$ . Furthermore  $\{(\pi_i, \tilde{b}_i^{C_G(\pi_i)}), i=1, 2, \dots, n\}$  is a set of representatives for the conjugacy classes of subsections associated with  $B$ .*
- (iv) *Let  $\tilde{\chi}$  be an ordinary irreducible character in  $\tilde{B}$  and  $\tilde{\phi}$  be an irreducible Brauer character in  $\tilde{b}_i$ . Then*

$$d(\tilde{\chi}^G, \pi_i, \tilde{\phi}^{C_G(\pi_i)}) = d(\tilde{\chi}, \pi_i, \tilde{\phi}).$$

Let  $\zeta$  be an irreducible character of a normal  $p'$ -subgroup  $N$  of  $G$  and suppose that  $\zeta$  is extendible to a character  $\tilde{\zeta}$  of  $G$ . Let  $\bar{B}$  be a  $p$ -block of the factor group  $\bar{G}$  of  $G$  by  $N$  and  $\bar{\chi}_0$  be an ordinary irreducible character in  $\bar{B}$ .  $\bar{\chi}_0$  can be viewed as a character of  $G$ . We denote by  $\tilde{\zeta}\bar{B}$  the  $p$ -block of  $G$  which contains  $\tilde{\zeta}\bar{\chi}_0$ . The ordinary irreducible characters in  $\tilde{\zeta}\bar{B}$  are  $\tilde{\zeta}\bar{\chi}$ 's, where  $\bar{\chi}$  runs over the ordinary irreducible characters in  $\bar{B}$  and the irreducible Brauer characters in  $\tilde{\zeta}\bar{B}$  are  $\tilde{\zeta}\bar{\phi}$ 's, where  $\bar{\phi}$  runs over the irreducible Brauer characters in  $\bar{B}$ . If  $\bar{B}_1$  and  $\bar{B}_2$  are different  $p$ -blocks of  $\bar{G}$ , then  $\tilde{\zeta}\bar{B}_1 \neq \tilde{\zeta}\bar{B}_2$ . For an element  $x$  of  $G$ , we put  $\bar{x} = xN (\in \bar{G})$  and for a subgroup  $Q$  of  $G$ , we put  $\bar{Q} = QN/N$ . If  $Q$  is a  $p$ -subgroup, then  $C_{\bar{G}}(\bar{Q}) = \overline{C_G(Q)}$  and  $N_{\bar{G}}(\bar{Q}) = \overline{N_G(Q)}$ . We have the following theorem related to the Fong's second reduction.

**Theorem 2.** *Let  $\zeta$  be an irreducible character of a normal  $p'$ -subgroup  $N$  of  $G$  and  $\tilde{\zeta}$  be an extension of  $\zeta$  to  $G$  such that  $(o(\det \tilde{\zeta}), p) = 1$ . If  $B$  is a  $p$ -block of  $G$  and  $B = \tilde{\zeta}\bar{B}$  for some  $p$ -block  $\bar{B}$  of the factor group  $\bar{G}$ , then the following holds.*

- (i) *If  $D$  is a defect group of  $B$ , then  $\bar{D}$  is a defect group of  $\bar{B}$ .*
- (ii) *Let  $\bar{b}$  be a root of  $\bar{B}$  in  $C_{\bar{G}}(\bar{D})\bar{D}$  and let  $b$  be a  $p$ -block of  $C_G(D)D$  such that  $b^{N_{C_G(D)D}} = \tilde{\zeta}\bar{b}$ . Then  $b$  is a root of  $B$  in  $C_G(D)D$  and  $T(\bar{b}) = \overline{T(b)}$ . In particular  $T(\bar{b})/C_{\bar{G}}(\bar{D})\bar{D} \cong T(b)/C_G(D)D$ .*
- (iii) *Let  $\pi$  be a  $p$ -element of  $G$  and  $\bar{b}_1, \bar{b}_2, \dots, \bar{b}_s$  be the  $p$ -blocks of  $C_{\bar{G}}(\bar{\pi})$  associated with  $\bar{B}$ . If  $b_i$  is a  $p$ -block of  $C_G(\pi)$  such that  $b_i^{N_{C_G(\pi)}} = \tilde{\zeta}\bar{b}_i$ , then  $b_1, b_2, \dots, b_s$  are the  $p$ -blocks of  $C_G(\pi)$  associated with  $B$ . Furthermore  $b_i = \theta_{\pi}\bar{b}_i$  ( $i = 1, 2, \dots, s$ ) when  $\bar{b}_i$  is viewed as a  $p$ -block of  $C_G(\pi)/C_N(\pi)$ , where  $\theta_{\pi}$  is an ordinary irreducible character of  $C_G(\pi)$  such that  $\theta_{\pi|_{C_N(\pi)}}$  is irreducible.*
- (iv) *For each ordinary irreducible character  $\bar{\chi}$  in  $\bar{B}$ , for the above  $p$ -element  $\pi$  and for each irreducible Brauer character  $\bar{\phi}$  in  $\bar{b}_i$ , there exists a sign  $\varepsilon_{\pi} = \pm 1$  such that*

$$d(\tilde{\zeta}\bar{\chi}, \pi, \theta_{\pi}\bar{\phi}) = \varepsilon_{\pi} d(\bar{\chi}, \bar{\pi}, \bar{\phi}).$$

We remark that (ii) and (iii) in the above theorems are stated by Puig [8, Theorems 1 and 2] without proofs.

Let  $K$  be the algebraic closure of the  $p$ -adic number field  $Q_p$  and  $R$  be the ring of local integers in  $K$ . Let  $P$  denote the maximal ideal of  $R$  and  $F$  denote the residue class field  $R/P$ . For a  $p$ -block  $B$  of  $G$ , we denote the block idempotent of  $FG$  corresponding to  $B$  by  $E_B$  and for an ordinary irreducible character  $\chi$  of  $G$ , we denote the centrally primitive idempotent of  $KG$  corresponding to  $\chi$  by  $e_{\chi}$ . The number of ordinary irreducible characters in  $B$  and the number of irreducible Brauer characters in  $B$  are denoted by  $k(B)$  and  $l(B)$ , respectively.

### 1. Proof of Theorem 1

**Lemma 1.** *Let  $H$  be a subgroup of  $G$  and  $x_1, x_2, \dots, x_h$  be a set of representatives for the right cosets of  $H$  in  $G$ . For a  $p$ -block  $\tilde{B}$  of  $H$ , we assume that*

$$E_{\tilde{B}}x^{-1}E_{\tilde{B}}x = 0 \quad \text{for all } x \in G - H.$$

*Then  $\sum_{i=1}^h x_i^{-1}E_{\tilde{B}}x_i$  is a block idempotent of  $FG$ . If we put  $\sum_{i=1}^h x_i^{-1}E_{\tilde{B}}x_i = E_B$ , where  $B$  is a  $p$ -block of  $G$ , then  $\tilde{\phi} \rightarrow \tilde{\phi}^G$  is a 1-1 correspondence between the irreducible Brauer characters in  $\tilde{B}$  and those in  $B$ .*

**REMARK.** By Iizuka, Ohmori and Watanabe [6, Theorem 2], the following (i) and (ii) are equivalent.

(i)  $\tilde{\phi} \rightarrow \tilde{\phi}^G$  is a 1-1 correspondence between the irreducible Brauer characters in  $\tilde{B}$  and those in  $B$ .

(ii)  $\tilde{\chi} \rightarrow \tilde{\chi}^G$  is a 1-1 correspondence between the ordinary irreducible characters in  $\tilde{B}$  and those in  $B$ .

**Proof.** We put  $E = \sum_{i=1}^h E_{\tilde{B}}x_i$ , where  $E_{\tilde{B}}x_i = x_i^{-1}E_{\tilde{B}}x_i$ . Then  $E$  is a central idempotent of  $FG$ . By the assumption we can show  $\mathfrak{L} \rightarrow \mathfrak{L}^G$  defines a 1-1 correspondence between the isomorphism classes of (right)  $FH$ -modules  $\mathfrak{L}$  with  $\mathfrak{L}E_{\tilde{B}} = \mathfrak{L}$  and the isomorphism classes of  $FG$ -modules  $\mathfrak{M}$  with  $\mathfrak{M}E = \mathfrak{M}$ , where  $\mathfrak{L}^G$  is the induced  $FG$ -module. Furthermore if  $\mathfrak{L}$  is an irreducible or a principal indecomposable  $FH$ -module, then  $\mathfrak{L}^G$  is an irreducible or a principal indecomposable  $FG$ -module. Hence by the indecomposability of Cartan matrices,  $E$  is a block idempotent. This completes the proof.

**Proof of Theorem 1.** (i) is well known. It is also well known that if  $E_{\tilde{B}}'$  is the block idempotent of  $RH$  which corresponds to  $\tilde{B}$ , then  $E_{\tilde{B}}' = \sum_{\tilde{\chi}} e_{\tilde{\chi}}$ ,  $\tilde{\chi}$  ranges over the ordinary irreducible characters in  $\tilde{B}$ . Let  $x_1, x_2, \dots, x_h$  be a set of representatives for the cosets of  $H$  in  $G$ , where  $x_1 = 1$ . We can show that  $e_{\tilde{\chi}^G} = \sum_{i=1}^h e_{\tilde{\chi}}x_i$ , so we have  $E_B = \sum_{i=1}^h E_{\tilde{B}}x_i$ . By the assumption,  $E_{\tilde{B}}FGE_B = E_{\tilde{B}}FG$  and hence  $E_{\tilde{B}}E_B = E_{\tilde{B}}$  and

$$(1) \quad E_{\tilde{B}} \sum_{i=2}^h E_{\tilde{B}}x_i = 0.$$

By the proof of Watanabe [10, Theorem 2] and the fact  $\dim_F(E_BFG) = |G:H|^2 \dim_F(E_{\tilde{B}}FH)$ , we have

$$(2) \quad E_BFG = \sum_{i,j=1}^h \oplus x_i^{-1}E_{\tilde{B}}FHx_j.$$

From (2), we obtain  $E_{\tilde{B}}E_{\tilde{B}}x = 0$  for all  $x \in G - H$ .

Let  $Q$  be a  $p$ -subgroup of  $H$ ,  $\tilde{b}$  be a  $p$ -block of  $C_H(Q)Q$  with  $\tilde{b}^H = \tilde{B}$  and  $\text{Br}_Q$  be the Brauer morphism from  $(FG)^Q$  onto  $FC_G(Q)$ , where  $(FG)^Q = \{a \in FG \mid ya = ay \text{ for all } y \in Q\}$  (see Alperin and Broué [1]). Then we have

$$\begin{aligned}\text{Br}_Q(E_{\tilde{B}}) \text{Br}_Q(E_{\tilde{B}})^* &= 0 \quad (x \in C_G(Q)Q - C_H(Q)Q), \\ \text{Br}_Q(E_{\tilde{B}})E_{\tilde{b}} &= E_{\tilde{b}}.\end{aligned}$$

So  $E_{\tilde{b}}E_{\tilde{b}}^x = 0$  for all  $x \in C_G(Q)Q - C_H(Q)Q$ . By Reynolds [9, Theorem 2] and Lemma 1,  $\tilde{b}^{C_G(Q)Q}$  is defined and  $\tilde{\phi} \rightarrow \tilde{\phi}^{C_G(Q)Q}$  is a 1-1 correspondence between the irreducible Brauer characters in  $\tilde{b}$  and those in  $\tilde{b}^{C_G(Q)Q}$ .

Let  $z_1, z_2, \dots, z_t$  be a set of representatives for the cosets of  $T(\tilde{b})$  in  $N_H(D)$ , where  $z_1 = 1$ . Then

$$(3) \quad \text{Br}_D(E_{\tilde{B}}) = \sum_{j=1}^t E_{\tilde{b}}^{z_j}, \quad E_{\tilde{b}}E_{\tilde{b}}^{z_j} = 0$$

for  $j \geq 2$ . We assume that  $C_G(D)D = \bigcup_{i=1}^t C_H(D)Dx_i$  and  $N_G(D) = \bigcup_{i=1}^t N_H(D)x_i$ . So  $N_G(D) = \bigcup_{i=1}^t \bigcup_{j=1}^t T(\tilde{b})z_jx_i$ . For  $(i, j) \neq (1, 1)$  we have

$$(4) \quad E_{\tilde{b}}E_{\tilde{b}}^{z_j z_i} = E_{\tilde{b}}\text{Br}_D(E_{\tilde{B}})\text{Br}_D(E_{\tilde{B}})^*E_{\tilde{b}}^{z_j z_i} = 0$$

from (3). By the above argument  $\tilde{b}^{C_G(D)D}$  is defined. We put  $b = \tilde{b}^{C_G(D)D}$ . Then  $b^G = \tilde{b}^G = \tilde{B}^G = B$ . Hence  $b$  is a root of  $B$  in  $C_G(D)D$  and  $E_b = \sum_{i=1}^t E_{\tilde{b}}^{z_i}$ . If  $y \in T(b)$ , then

$$E_b = E_b E_b^y = \sum_{i,j=1}^t E_{\tilde{b}}^{z_i} E_{\tilde{b}}^{z_j y}.$$

From (4), there exist  $i$  and  $j$ ,  $1 \leq i, j \leq t$ , such that  $x_j y x_i^{-1} \in T(\tilde{b})$ , hence  $y \in T(\tilde{b})C_G(D)$ . Conversely if  $w \in T(\tilde{b})$ , then

$$E_b^w = \sum_{i=1}^t E_{\tilde{b}}^{z_i w} = \sum_{i=1}^t E_{\tilde{b}}^{w^{-1} z_i w} = E_b.$$

Therefore  $T(b) = T(\tilde{b})C_G(D)$ . This completes the proof of (ii).

Next we prove (iii) and (iv).  $\tilde{b}_i^{C_G(\pi_i)}$  is defined and  $\tilde{\phi} \rightarrow \tilde{\phi}^{C_G(\pi_i)}$  is a 1-1 correspondence between the irreducible Brauer characters in  $\tilde{b}_i$  and those in  $\tilde{b}_i^{C_G(\pi_i)}$ . Let  $\pi$  be a  $p$ -element of  $G$ . We assume that exactly  $m$  elements  $\pi_1, \pi_2, \dots, \pi_m$  are conjugate to  $\pi$  in  $G$ . We put  $\pi_i^{\#} = \pi$  ( $a_i \in G$ ,  $i = 1, 2, \dots, m$ ). Since  $\tilde{\chi} = \sum_{i=1}^m \tilde{\chi}^{(\pi_i, \tilde{b}_i)}$ ,  $\tilde{\chi}^G = \sum_{i=1}^m (\tilde{\chi}^{(\pi_i, \tilde{b}_i)})^G$ . Here  $\tilde{\chi}^{(\pi_i, \tilde{b}_i)}(\pi_i \rho) = \sum_{\tilde{\phi}} d(\tilde{\chi}, \pi_i, \tilde{\phi}) \tilde{\phi}(\rho)$  for all  $p$ -regular elements  $\rho$  of  $C_H(\pi_i)$  with  $\tilde{\phi}$  ranging over the irreducible Brauer characters in  $\tilde{b}_i$  (see Brauer [2, §1]). So we can show

$$\begin{aligned}\tilde{\chi}^G(\pi\rho) &= \sum_{i=1}^m \sum_{\tilde{\phi}} d(\tilde{\chi}, \pi_i, \tilde{\phi}) \tilde{\phi}^{C_G(\pi_i)}(\rho^{a_i^{-1}}) \\ &= \sum_{i=1}^m \sum_{\tilde{\phi}} d(\tilde{\chi}, \pi_i, \tilde{\phi}) (\tilde{\phi}^{C_G(\pi_i)})^a(\rho)\end{aligned}$$

for all  $p$ -regular elements  $\rho$  of  $C_G(\pi)$ . Hence a subsection associated with  $B$  is conjugate to some subsection  $(\pi_i, \tilde{b}_i^{C_G(\pi_i)})$  ( $i=1, 2, \dots, n$ ). In particular  $k(B) \leq \sum_{i=1}^n l(\tilde{b}_i)$ . On the other hand,  $k(B) = k(\tilde{B}) = \sum_{i=1}^n l(\tilde{b}_i)$ . Therefore, if  $i \neq j$ , then  $(\pi_i, \tilde{b}_i^{C_G(\pi_i)})$  and  $(\pi_j, \tilde{b}_j^{C_G(\pi_j)})$  are not conjugate and  $d(\tilde{\chi}^G, \pi_i, \tilde{\phi}^{C_G(\pi_i)}) = d(\tilde{\chi}, \pi_i, \tilde{\phi})$ . This completes the proof of Theorem 1.

## 2. Proof of Theorem 2

We denote the set of  $p$ -regular elements of  $G$  by  $G_{p'}$ .

If  $\chi$  is a character of  $G$  and  $T$  is a matrix representation of  $G$  affording  $\chi$ , then  $x \mapsto \det T(x)$  is a linear character of  $G$ . The linear character is denoted by  $\det \chi$  and  $o(\det \chi)$  means the order. The following lemma is a special case of Glauberman's theorem [5, Theorem 3].

**Lemma 2.** *Let  $\pi$  be a  $p$ -element of  $G$  and  $N$  be a  $p'$ -subgroup of  $G$  such that  $N^\pi = N$ . Suppose that  $\zeta$  is an irreducible character of  $N$  and  $\tilde{\zeta}$  is an extension of  $\zeta$  to  $N\langle\pi\rangle$  with  $(o(\det \tilde{\zeta}), p) = 1$ . Then there exist a unique sign  $\varepsilon = \pm 1$  and a unique irreducible character  $\beta$  of  $C_N(\pi)$  with the property that*

$$\tilde{\zeta}(\pi\rho) = \varepsilon\beta(\rho), \quad \rho \in C_N(\pi).$$

**Lemma 3.** *Let  $\zeta$  be an irreducible character of a normal  $p'$ -subgroup  $N$  of  $G$  and  $\tilde{\zeta}$  be an extension of  $\zeta$  to  $G$  such that  $(o(\det \tilde{\zeta}), p) = 1$ . For a  $p$ -element  $\pi$  of  $G$ , there exist a sign  $\varepsilon_\pi = \pm 1$  and an irreducible character  $\theta_\pi$  of  $C_G(\pi)$  with the property that  $\theta_{\pi|C_N(\pi)}$  is irreducible and*

$$\tilde{\zeta}(\pi\rho) = \varepsilon_\pi \theta_\pi(\rho) \quad \rho \in (C_G(\pi))_{p'}.$$

*In particular  $\theta_\pi$  is irreducible as a Brauer character.*

Proof. We fix a  $p$ -element  $\pi$ . By Lemma 2, there exist a unique sign  $\varepsilon = \pm 1$  and a unique irreducible character  $\beta$  of  $C_N(\pi)$  with the property that  $\tilde{\zeta}(\pi\rho) = \varepsilon\beta(\rho)$  for all  $\rho \in C_N(\pi)$ . First of all we show that  $\beta$  is extendible to  $C_G(\pi)$ . Since  $\tilde{\zeta}(\pi\rho) = \tilde{\zeta}(\pi\rho^c)$  for all  $c \in C_G(\pi)$  and all  $\rho \in C_N(\pi)$ ,  $\beta$  is  $C_G(\pi)$ -invariant. Let  $L$  be a subgroup of  $C_G(\pi)$  such that  $L/C_N(\pi)$  is a  $p$ -group. Then by Isaacs [7, (8.16)],  $\beta$  is extendible to  $L$ . Let  $M$  be a subgroup of  $C_G(\pi)$  such that  $M/C_N(\pi)$  is a  $p'$ -group. Then  $(NM)^\pi = NM$ . By Lemma 2, there exist a sign  $\varepsilon_M = \pm 1$  and an irreducible character  $\beta_M$  of  $C_{NM}(\pi)$  with the property that

$$\tilde{\zeta}(\pi\rho) = \varepsilon_M \beta_M(\rho), \quad \rho \in C_{NM}(\pi).$$

Since  $C_N(\pi) \subset C_{NM}(\pi) = M$ , we have  $\varepsilon_M = \varepsilon$  and  $\beta_{M|C_N(\pi)} = \beta$  by the uniqueness of  $\varepsilon$  and  $\beta$ . Hence by [7, (11.31)],  $\beta$  is extendible to  $C_G(\pi)$ .

Let  $\theta_0$  be an extension of  $\beta$ . For a  $p'$ -subgroup  $M$  of  $C_G(\pi)$  with  $M \supset C_N(\pi)$ , there exists a unique linear character  $\lambda_M$  of  $M/C_N(\pi)$  which satisfies

$$\theta_{0|M} \lambda_M = \beta_M.$$

Furthermore for  $p'$ -subgroups  $M, M'$  of  $C_G(\pi)$  with  $M, M' \supset C_N(\pi)$ , if  $M \supset M'$  then  $\lambda_{M'} = (\lambda_M)_{|M'}$  and if  $M' = M^x$  for some  $x \in C_G(\pi)$  then  $\lambda_{M'} = \lambda_M^x$ . Here we define a class function  $\lambda$  of  $C_G(\pi)/C_N(\pi)$  as follows. For an element  $c$  of  $C_G(\pi)/C_N(\pi)$

$$\lambda(c) = \lambda_M(c_{p'}),$$

where  $c_{p'}$  is the  $p'$ -part of  $c$  and  $M$  satisfies that  $M/C_N(\pi) = \langle c_{p'} \rangle$ . Then  $\lambda$  is a generalized character of  $C_G(\pi)/C_N(\pi)$  by Brauer's theorem on generalized characters. Since the inner product  $(\lambda, \lambda)$  and  $\lambda(1)$  are equal to 1,  $\lambda$  is a linear character. If we put  $\theta = \theta_0 \lambda$ , then  $\xi(\pi\rho) = \varepsilon\theta(\rho)$  for all  $\rho \in (C_G(\pi))_{p'}$ . This completes the proof.

Proof of Theorem 2. (i) is well known. Let  $Q$  be a  $p$ -subgroup of  $G$  and  $\bar{b}$  be a  $p$ -block of  $C_{\bar{G}}(\bar{Q})\bar{Q}$  associated with  $\bar{B}$ . We show  $(\xi\bar{b})^G = B$ . Let  $C$  be an arbitrary conjugacy class of  $G$  and  $\bar{\chi}$  and  $\bar{\psi}$  be ordinary irreducible characters in  $\bar{B}$  and  $\bar{b}$ , respectively. Since  $\bar{b}^G = \bar{B}$ ,

$$\bar{\chi}\left(\sum_{x \in C} \bar{x}\right)/\bar{\chi}(1) \equiv \bar{\psi}\left(\sum_{x \in C \cap NC_G(Q)Q} \bar{x}\right)/\bar{\psi}(1) \pmod{P}.$$

If  $x_0$  is an element of  $C$ , then

$$\begin{aligned} (\xi\bar{\chi})\left(\sum_{x \in C} \bar{x}\right)/\xi\bar{\chi}(1) &= \xi(x_0)\bar{\chi}\left(\sum_{x \in C} \bar{x}\right)/\xi\bar{\chi}(1), \\ (\xi\bar{\psi})\left(\sum_{x \in C \cap NC_G(Q)Q} \bar{x}\right)/\xi\bar{\psi}(1) &= \xi(x_0)\bar{\psi}\left(\sum_{x \in C \cap NC_G(Q)Q} \bar{x}\right)/\xi\bar{\psi}(1). \end{aligned}$$

Hence  $(\xi\bar{b})^G = \xi\bar{B} = B$ . Since a defect group of  $\bar{b}$  is  $\bar{D}$ ,  $D$  is a defect group of  $\xi\bar{b}$ . Let  $b$  be a root of  $\xi\bar{b}$  in  $C_G(D)D$ .  $b$  is a root of  $B$  in  $C_G(D)D$  and is determined uniquely, because  $N_G(D) \cap NC_G(D)D = C_G(D)D$ . If  $x \in T(b)$ , then

$$\xi\bar{b} = (b^x)^{NC_G(D)D} = (b^{NC_G(D)D})^x = (\xi\bar{b})^x = \xi\bar{b}^x.$$

Hence  $\bar{b} = \bar{b}^x$ , so  $\bar{x} \in T(\bar{b})$ . If  $y \in N_G(D)$  and  $\bar{y} \in T(\bar{b})$ , then

$$\xi\bar{b} = (\xi\bar{b})^y = (b^y)^{NC_G(D)D}.$$

By the uniqueness of a root  $b$  of  $\xi\bar{b}$ ,  $b = b^y$  and hence  $y \in T(b)$ . So we have  $T(\bar{b}) = \overline{T(b)}$ .

Next we prove (iii) and (iv). By Lemma 3, there exist a sign  $\varepsilon_\pi = \pm 1$  and an ordinary irreducible character  $\theta_\pi$  of  $C_G(\pi)$  such that  $\theta_\pi$  is irreducible as a

Bauer character and  $\xi(\pi\rho) = \varepsilon_\pi \theta_\pi(\rho)$  for all  $\rho \in (C_G(\pi))_{p'}$ .  $\bar{b}_i$  can be viewed as a  $p$ -block of  $C_G(\pi)/C_N(\pi)$ . We put  $b_i = \theta_\pi \bar{b}_i$ . Since

$$\bar{\chi}(\bar{\pi}\bar{\rho}) = \sum_{i=1}^s \sum_{\bar{\phi}} d(\bar{\chi}, \bar{\pi}, \bar{\phi}) \bar{\phi}(\bar{\rho}) \quad \rho \in (C_G(\pi))_{p'}.$$

we have

$$(5) \quad (\xi\bar{\chi})(\pi\rho) = \sum_{i=1}^s \sum_{\bar{\phi}} \varepsilon_\pi d(\bar{\chi}, \bar{\pi}, \bar{\phi}) (\theta_\pi \bar{\phi})(\rho) \quad \rho \in (C_G(\pi))_{p'}.$$

Here  $\bar{\phi}$  ranges over the irreducible Brauer characters in  $\bar{b}_i$ . By the second main theorem on  $p$ -blocks,  $b_1, b_2, \dots, b_s$  are the  $p$ -blocks of  $C_G(\pi)$  associated with  $B$ . In particular we see that  $b_i$  is a unique  $p$ -block of  $C_G(\pi)$  such that  $b_i^{N_{C_G(\pi)}} = \xi \bar{b}_i$ . From (5),  $d(\xi\bar{\chi}, \pi, \theta_\pi \bar{\phi}) = \varepsilon_\pi d(\bar{\chi}, \bar{\pi}, \bar{\phi})$ . This completes the proof of Theorem 2.

We have the following as a corollary of Theorem 1 and 2.

**Corollary.** *Suppose that  $G$  is a  $p$ -solvable group. Let  $B$  be a  $p$ -block of  $G$  with an abelian defect group  $D$  and  $b$  be a root of  $B$  in  $C_G(D)$ . We assume that  $T(b)/C_G(D)$  is cyclic and any element of  $T(b)/C_G(D) - \{\bar{1}\}$  does not fix any element of  $D - \{1\}$ . Let  $\pi_1, \pi_2, \dots, \pi_t$  be a set of representatives for the  $T(b)$ -conjugacy classes of  $D - \{1\}$  and  $\Lambda$  be a set of representatives for the  $T(b)$ -conjugacy classes of non-trivial linear characters of  $D$ , where  $t = (p^d - 1)/e$ ,  $e = |T(b) : C_G(D)|$  and  $p^d = |D|$ . Then the following holds.*

(i)  *$B$  contains exactly  $e$  irreducible Brauer characters  $\phi_1, \phi_2, \dots, \phi_e$  and exactly  $e + (p^d - 1)/e$  ordinary irreducible characters  $\chi_1, \chi_2, \dots, \chi_e, \chi_\lambda$  ( $\lambda \in \Lambda$ ).*

(ii) *For  $i$ ,  $1 \leq i \leq e$ , and  $\lambda, \lambda \in \Lambda$ ,*

$$\begin{aligned} \chi_i &= \phi_i \quad \text{on } G_{p'}, \\ \chi_\lambda &= \phi_1 + \dots + \phi_e \quad \text{on } G_{p'}. \end{aligned}$$

(iii)  *$(1, B), (\pi_j, b^{C_G(\pi_j)})$  ( $j = 1, 2, \dots, t$ ) form a set of representatives for the conjugacy classes of subsections associated with  $B$ .  $b^{C_G(\pi_j)}$  contains a unique irreducible Brauer character  $\phi^{(j)}$ .*

(iv) *There exist  $t$  signs  $\varepsilon_j = \pm 1$  such that*

$$\begin{aligned} d(\chi_i, \pi_j, \phi^{(j)}) &= \varepsilon_j, \\ d(\chi_\lambda, \pi_j, \phi^{(j)}) &= (\varepsilon_j / |C_G(D)|) \sum_{x \in T(b)} \lambda^x(\pi_j). \end{aligned}$$

for  $i$ ,  $1 \leq i \leq e$ ,  $\lambda, \lambda \in \Lambda$  and  $j$ ,  $1 \leq j \leq t$ .

**Proof.** If  $\pi \in D - \{1\}$ , then  $C_G(\pi) \cap T(b) = C_G(D)$ . Hence  $b^{C_G(\pi)}$  contains a unique irreducible Brauer character by [2, (7A)] and Brauer [3, (6C)]. Hence (iii) follows from [3, (6C)]. By Fong's reductions (see Feit [4, Chapter X, Lemma 1.1]) and Theorems 1 and 2, we may assume that  $D$  is a normal sub-

group of  $G$  and  $T(b)=G$ . Then  $B$  is a unique  $p$ -block of  $G$  which covers  $b$ . Let  $\Lambda_0$  be the set of all linear characters of  $D$ . By [9, Theorem 3],  $b$  contains a unique irreducible Brauer character  $\phi$  and exactly  $p^d$  ordinary irreducible characters  $\tilde{\chi}_\mu$ ,  $\mu \in \Lambda_0$ , where if  $\pi \in D$  and  $\rho \in (C_G(D))_{p'}$ , then  $\tilde{\chi}_\mu(\pi\rho)=\mu(\pi)\phi(\rho)$ . Since  $\phi$  is  $G$ -invariant and  $G/C_G(D)$  is cyclic,  $B$  contains exactly  $e$  irreducible Brauer characters  $\phi_1, \phi_2, \dots, \phi_e$ . Since  $\tilde{\chi}_1$  is also  $G$ -invariant,  $B$  contains exactly  $e$  ordinary irreducible characters  $\chi_1, \chi_2, \dots, \chi_e$  such that  $\chi_i|_{C_G(D)}=\tilde{\chi}_1$ . We may assume  $\chi_i=\phi_i$  on  $G_{p'}$ . For an element  $\pi \in D - \{1\}$ ,  $\chi_i(\pi\rho)=\phi(\rho)$  ( $\rho \in (C_G(D))_{p'}$ ). Here we note  $C_G(D)=C_G(\pi)$ . By the assumption, if  $\mu \neq 1$ , then the stabilizer of  $\tilde{\chi}_\mu$  in  $G$  is equal to  $C_G(D)$ . Hence  $\tilde{\chi}_\mu^G$  is irreducible and

$$\begin{aligned}\tilde{\chi}_\mu^G &= \phi_1 + \dots + \phi_e \quad \text{on } G_{p'}, \\ \tilde{\chi}_\mu^G(\pi\rho) &= (1/|C_G(D)|) \sum_{\pi \in G} \mu^*(\pi) \phi(\rho) \quad \rho \in (C_G(D))_{p'}.\end{aligned}$$

This completes the proof.

---

### References

- [1] J. Alperin and M. Broué: *Local methods in block theory*, Ann. of Math. **110** (1979), 143–157.
- [2] R. Brauer: *On blocks and sections in finite groups* II, Amer. J. Math. **90** (1968), 895–925.
- [3] ———: *On the structure of blocks of characters of finite groups*, Lecture Note in Math. 372, Springer, Berlin, 103–130.
- [4] W. Feit: *The representation theory of finite groups*, North-Holland, Amsterdam, 1982.
- [5] G. Glauberman: *Correspondences of characters for relatively prime operator groups*, Canad. J. Math. **20** (1968), 1465–1488.
- [6] K. Iizuka, F. Ohmori and A. Watanabe: *A remark on the representations of finite groups* VI, Mem. Fac. Gen. Ed. Kumamoto Univ. **18** (1983), 1–8 (in Japanese).
- [7] I.M. Isaacs: *Character theory of finite groups*, Academic Press, New York, 1976.
- [8] L. Puig: *Local block theory in  $p$ -solvable groups*, Proc. Sympos Pure Math. **37** (1980), 385–388.
- [9] W.F. Reynolds: *Blocks and normal subgroups of finite groups*, Nagoya Math. J. **22** (1963), 15–32.
- [10] A. Watanabe: *Relations between blocks of a finite group and its subgroup*, J. Algebra **78** (1982), 282–291.

Department of Mathematics  
 Faculty of General Education  
 Kumamoto University  
 Kurokami 2-40-1  
 Kumamoto 860, Japan