



Title	Smooth actions of special unitary groups on cohomology complex projective spaces
Author(s)	Uchida, Fuichi
Citation	Osaka Journal of Mathematics. 1975, 12(2), p. 375-400
Version Type	VoR
URL	https://doi.org/10.18910/12022
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

SMOOTH ACTIONS OF SPECIAL UNITARY GROUPS ON COHOMOLOGY COMPLEX PROJECTIVE SPACES

FUICHI UCHIDA

(Received July 24, 1974)

0. Introduction

The purpose of this paper is to study smooth $SU(n)$ -actions on a compact orientable $2m$ -manifold whose rational cohomology ring is isomorphic to $H^*(P_m(\mathbb{C}); \mathbb{Q})$. First we show the following result.

Theorem 2.1. *Let $n \geq 7$ and $0 \leq k < n-4$. Let M be a compact orientable smooth $2(n+k)$ -manifold with*

$$H^*(M; \mathbb{Q}) = H^*(P_{n+k}(\mathbb{C}); \mathbb{Q}).$$

Then for any non-trivial smooth $SU(n)$ -action on M , the stationary point set $F = F(SU(n), M)$ is an orientable $2k$ -manifold with

$$H^*(F; \mathbb{Q}) = H^*(P_k(\mathbb{C}); \mathbb{Q})$$

and there is an equivariant diffeomorphism

$$M = \partial(D^{2n} \times X)/S^1.$$

Here X is a compact connected orientable $(2k+2)$ -manifold which is acyclic over rationals, X admits a smooth S^1 -action which is free on dX , the $SU(n)$ -action is standard on D^{2n} and trivial on X , and

$$\pi_1(X) = \pi_1(M).$$

Furthermore, if

$$H^*(M; \mathbb{Z}) = H^*(P_{n+k}(\mathbb{C}); \mathbb{Z}),$$

then X is acyclic over integers, the S^1 -action on X is semi-free, and

$$H^*(F; \mathbb{Z}) = H^*(P_k(\mathbb{C}); \mathbb{Z}).$$

Corollary 2.2. *Let $n \geq 7$ and $0 \leq k < n-4$. Let M be a compact connected smooth $2(n+k)$ -manifold which is homotopy equivalent to $P_{n+k}(\mathbb{C})$. If M admits a non-trivial smooth $SU(n)$ -action, then M is diffeomorphic to $P_{n+k}(\mathbb{C})$.*

Examples of $SU(n)$ -actions on cohomology complex projective spaces are constructed in section 3. And we have the following results.

Theorem 3.1. *Let $n \geq 2$, $k \geq 1$ and $p \geq 1$. Then there is a compact orientable $2(n+k)$ -manifold M such that*

$$\pi_1(M) = \mathbf{Z}/p\mathbf{Z} \text{ and } H^*(M; \mathbf{Q}) = H^*(P_{n+k}(\mathbf{C}); \mathbf{Q})$$

and M admits a smooth $SU(n)$ -action with

$$F(SU(n), M) = P_k(\mathbf{C}).$$

Theorem 3.2. *Let $n \geq 2$ and $k \geq 3$. Let G be a finitely presentable group with $H_1(G; \mathbf{Z}) = H_2(G; \mathbf{Z}) = 0$. Then*

(a) *there is a compact orientable $2(n+k)$ -manifold M such that*

$$\pi_1(M) = G \text{ and } H^*(M; \mathbf{Z}) = H^*(P_{n+k}(\mathbf{C}); \mathbf{Z})$$

and M admits a smooth $SU(n)$ -action with

$$F(SU(n), M) = P_k(\mathbf{C}),$$

(b) *there is a smooth $SU(n)$ -action on $P_{n+k}(\mathbf{C})$ such that*

$$\pi_1(F) = G \text{ and } H^*(F; \mathbf{Z}) = H^*(P_k(\mathbf{C}); \mathbf{Z}),$$

where $F = F(SU(n), P_{n+k}(\mathbf{C}))$.

Next, in section 4, we study a signature of closed orientable manifold which admits a smooth G -action with isotropy groups of uniform dimension, and we have a result which is a generalization of the fact that $\text{Sign}(M) = 0$ if M admits a smooth circle action without stationary points.

Next we study smooth $SU(3)$ -actions on orientable 8-manifolds in section 5, and as an application we show a similar result as Theorem 2.1 for non-trivial smooth $SU(3)$ -action on a cohomology complex projective 4-space. We construct examples of stationary point free $SU(3)$ -actions on orientable 8-manifolds with non-zero signature in section 6.

As a concluding remark, classification of smooth $SU(n)$ -actions on orientable $2n$ -manifolds is done in the final section.

1. $SU(n)$ -actions with certain isotropy types

Let E be a manifold with smooth $SU(n)$ -action ($n \geq 3$). Assume that the identity component of each isotropy group is conjugate to $SU(n-1)$ or $NSU(n-1)$, the normalizer of $SU(n-1)$ in $SU(n)$. Then $S^1 = NSU(n-1)/SU(n-1)$ acts naturally on

$$X = F(SU(n-1), E),$$

the stationary point set of $SU(n-1)$. It is easily seen that

$$(1.1) \quad SU(n)/SU(n-1) \times X \rightarrow E, \quad [gSU(n-1), x] \rightarrow gx$$

is an equivariant diffeomorphism as $SU(n)$ -manifolds, since $g \in SU(n)$ and $g^{-1}SU(n-1)g \subset NSU(n-1)$ imply $g \in NSU(n-1)$.

Lemma 1.2. *Let V be a real vector space with linear $SU(n)$ -action ($n \geq 3$). Assume that the identity component of each isotropy group on the invariant unit sphere $S(V)$ is conjugate to $SU(n-1)$ or $NSU(n-1)$. Then $S(V) = SU(n)/SU(n-1)$ as $SU(n)$ -spaces.*

Proof. By (1.1), there is an equivariant diffeomorphism

$$S(V) = SU(n)/SU(n-1) \times F(SU(n-1), S(V)),$$

where $F(SU(n-1), S(V))$ is a sphere. Then it is easily seen that

$$F(SU(n-1), S(V)) = S^1$$

by the homotopy exact sequence of the fibre bundle

$$F(SU(n-1), S(V)) \rightarrow S(V) \rightarrow P_{n-1}(\mathbb{C}).$$

Considering S^1 -actions on S^1 , we have

$$S(V) = SU(n)/SU(n-1)$$

as $SU(n)$ -spaces.

q.e.d.

Lemma 1.3. *Let V be a real vector space with linear $SU(n)$ -action such that $S(V) = SU(n)/SU(n-1)$ as $SU(n)$ -spaces ($n \geq 3$). Then the $SU(n)$ -action on $V = \mathbb{R}^{2n}$ is equivalent to the standard action.*

Proof. This is a known result (see [8], Theorem I), but we give an elementary proof for the completeness. It is well-known that a real irreducible $SU(n)$ -vector space \mathbb{R}^{2n} with an invariant complex structure is equivalent to \mathbb{R}^{2n} with the standard $SU(n)$ -action. So we prove the existence of an invariant complex structure on V . Denote by Z_n , the center of $SU(n)$. Then Z_n is a cyclic group of order n , and the Z_n -action on $S(V)$ is free, since

$$Z_n \cap SU(n-1) = \{1\}.$$

Consider a direct sum decomposition

$$V = V_1 \oplus \cdots \oplus V_k$$

as \mathbf{Z}_n -vector space, where $V_i (i=1, \dots, k)$ are irreducible. Leaving a non-zero vector $v_1 \in V_1$ fixed, we have an element $g_i \in SU(n)$ such that

$$v_i = g_i v_1 \in V_i \quad (i=1, \dots, k)$$

by the transitivity of the $SU(n)$ -action on $S(V)$. Then

$$V_i = g_i V_1 \quad (i=1, \dots, k).$$

Since the \mathbf{Z}_n -action on $S(V_1)$ is free, there is a complex structure \mathbf{J}_1 on V_1 such that

$$\sigma \mathbf{J}_1 = \mathbf{J}_1 \sigma, \quad \sigma v_1 = av_1 + b\mathbf{J}_1 v_1$$

for some $a, b \in \mathbf{R}$, $b \neq 0$, where σ is a generator of \mathbf{Z}_n , moreover the real vector space V_1 is spanned by $\{v_1, \mathbf{J}_1 v_1\}$. Therefore there is a complex structure J on V such that

$$\mathbf{J}v_1 - \mathbf{J}_1 v_1, \quad \mathbf{J}g_i v_1 - g_i \mathbf{J}_1 v_1 \quad \text{and} \quad \sigma v = av + b\mathbf{J}v$$

for each $v \in V$. Then

$$g\sigma v = agv + bg\mathbf{J}v,$$

$$\sigma g v = agv + b\mathbf{J}g v$$

for any $g \in SU(n)$. Therefore the complex structure \mathbf{J} is $SU(n)$ -invariant, since $g\sigma = \sigma g$ and $b \neq 0$. q.e.d.

Let M be a closed connected manifold with smooth $SU(n)$ -action ($n \geq 3$). Assume that the identity component of each isotropy group is conjugate to one of the following

$$SU(n), \quad SU(n-1) \quad \text{and} \quad NSU(n-1).$$

Assume that the stationary point set $F = F(SU(n), M)$ is non-empty. Let U be an invariant closed tubular neighborhood of F in M . Then there is an equivariant decomposition

$$M = U \cup (SU(n)/SU(n-1) \times_{S^1} X) \sqcup (S^{2n-1} \times_{S^1} X),$$

where $X = F(SU(n-1), M - \text{int } U)$ with the natural S^1 -action. Since

$$dU = SU(n)/SU(n-1) \times_{S^1} \partial X = S^{2n-1} \times_{S^1} \partial X$$

as $SU(n)$ -manifolds, the S^1 -action on ∂X is free, $F = \partial X/S^1$, and the disk bundle $U \rightarrow F$ with $SU(n)$ -action is equivariantly isomorphic to the disk bundle

$$D^{2n} \times_{S^1} \partial X \rightarrow \partial X/S^1,$$

where the $SU(n)$ -action on D^{2n} is standard by Lemma 1.2 and Lemma 1.3.

Therefore the codimension of F in M is $2n$, X is connected, and there is an equivariant diffeomorphism

$$(1.4) \quad M = \partial(D^{2n} \times X)/S^1 \cong \underset{S^1}{D^{2n}} \times \partial X \cup \underset{S^1}{S^{2n-1}} \times X$$

as $SU(n)$ -manifolds.

Lemma 1.5. *Let G be a closed connected proper subgroup of $SU(n)$, ($n \geq 7$).*

If

$$\dim G > n^2 - 4n + 7 = \dim N(SU(n-2), SU(n)),$$

then G is conjugate to $SU(n-1)$ or $NSU(n-1)$ in $SU(n)$.

Proof. The inclusion $\rho: G \subset SU(n)$ gives an n -dimensional complex representation of G . First we show that the representation ρ is reducible. Suppose that ρ is irreducible. Then G is semi-simple from the Shur's lemma. If G is not simple, then there are integers $p \geq q \geq 2$ with $n = pq$, such that G is conjugate to a subgroup of the tensor product

$$SU(p) \otimes SU(q)$$

in $SU(pq)$, by considering the induced representation of the universal covering group of G . Therefore

$$\dim G \leq p^2 + q^2 - 2 \leq \left(\frac{n}{2}\right)^2 + 2 \leq \frac{n(n+1)}{2}.$$

If G is simple but not one of the type

$$A_k, D_{2k+1} \text{ and } E_6,$$

then G is conjugate to a subgroup of $SO(n)$ or $Sp(n/2)$, (see [6], p. 336, Theorem 0.20). But

$$\dim SO(n) = \frac{n(n-1)}{2}, \quad \dim Sp\left(\frac{n}{2}\right) = \frac{n(n+1)}{2}$$

and hence

$$\dim G \leq \frac{n(n+1)}{2}.$$

If G is of type D_{2k+1} ($k \geq 2$), then the lowest dimensional non-trivial irreducible complex representation is $(4k+2)$ -dimensional (see [6], p. 378, Table 30). Therefore $4k+2 \leq n$ and hence

$$\dim G = \dim SO(4k+2) = (2k+1)(4k+1) \leq \frac{n(n-1)}{2}.$$

If G is of type E_6 , then $n \geq 27$ (see [6], p. 378, Table 30). Therefore

$$\dim G = 78 \leq 3n \leq \frac{n(n+1)}{2}.$$

Finally, if G is of type A_{k-1} ($k < n$), then

$$\frac{k(k-1)}{2} \leq n,$$

by the Weyl's formula (see [14], Theorem 7.5). Therefore

$$\dim G = \dim SU(k) = k^2 - 1 \leq 3n - 2 \leq \frac{n(n+1)}{2}.$$

Consequently

$$\dim G \leq \frac{n(n+1)}{2},$$

if p : $G \subset SU(n)$ is irreducible ($n \geq 4$). Therefore p is reducible, if

$$\dim G > n^2 - 4n + 7 \quad \text{and} \quad n \geq 7.$$

Since p is reducible, G is conjugate to a subgroup of

$$N(SU(n-p), SU(n)), \left(1 \leq p \leq \frac{n}{2}\right)$$

the normalizer of $SU(n-p)$ in $SU(n)$. But

$$\dim N(SU(n-p), SU(n)) \leq n^2 - 4n + 7$$

for $2 \leq p \leq \frac{n}{2}$. Therefore G is conjugate to a subgroup G' of $NSU(n-1)$. If $G' \neq NSU(n-1)$, then

$$\dim G' \leq \dim G'' + 1$$

where $G'' = G' \cap SU(n-1)$, by the isomorphism

$$NSU(n-1)/SU(n-1) \cong S^1.$$

If $G'' = SU(n-1)$ then $G' = G'' = SU(n-1)$. If $G'' \neq SU(n-1)$, then

$$\dim G'' \leq (n-2)^2 = \dim N(SU(n-2), SU(n-1)),$$

by making use of the first part of the proof of this lemma for $SU(n-1)$ instead of $SU(n)$, and hence

$$\dim G' \leq (n-2)^2 + 1 < n^2 - 4n + 7.$$

Consequently we see that G is conjugate to $SU(n-1)$ or $NSU(n-1)$ in $SU(n)$.
q.e.d.

Lemma 1.6. *Let M be a manifold with smooth $SU(n)$ -action. If $\dim M < 4n-8$, then*

$$\dim SU(n)_x > n^2 - 4n + 7$$

for each $x \in M$.

Proof. Since $SU(n)/SU(n)_x$ is equivariantly embedded in M ,

$$\dim SU(n) - \dim SU(n)_x \leq \dim M < 4n - 8.$$

Hence $\dim SU(n)_x > \dim SU(n) - (4n - 8) = n^2 - 4n + 7$. q.e.d.

2. $SU(n)$ -actions on cohomology complex projective spaces

In this section we prove the following results.

Theorem 2.1. *Let $n \geq 7$ and $0 \leq k < n-4$. Let M be a compact connected orientable smooth $2(n+k)$ -manifold with*

$$H^*(M; \mathbb{Q}) = H^*(P_{n+k}(\mathbb{C}); \mathbb{Q}).$$

Then for any non-trivial smooth $SU(n)$ -action on M , the stationary point set $F = F(SU(n), M)$ is an orientable $2k$ -manifold with

$$H^*(F; \mathbb{Q}) = H^*(P_k(\mathbb{C}); \mathbb{Q})$$

and there is an equivariant diffeomorphism

$$M = \partial(D^{2n} \times X)/S^1.$$

Here X is a compact connected orientable $(2k+2)$ -manifold which is acyclic over rationals, X admits a smooth S^1 -action which is free on ∂X , the $SU(n)$ -action is standard on D^{2n} and trivial on X , and

$$\pi_1(X) = \pi_1(M).$$

Furthermore, if

$$H^*(M; \mathbb{Z}) = H^*(P_{n+k}(\mathbb{C}); \mathbb{Z}),$$

then X is acyclic over integers, the S^1 -action on X is semi-free, and

$$H^*(F; \mathbb{Z}) = H^*(P_k(\mathbb{C}); \mathbb{Z}).$$

Corollary 2.2. *Let $n \geq 7$ and $0 \leq k < n-4$. Let M be a compact connected smooth $2(n+k)$ -manifold which is homotopy equivalent to $P_{n+k}(\mathbb{C})$. If M admits a non-trivial smooth $SU(n)$ -action, then M is diffeomorphic to $P_{n+k}(\mathbb{C})$.*

Proof of Theorem 2.1. By Lemma 1.5, Lemma 1.6 and the assumption $n \geq 7$ and $0 \leq k < n-4$, the identity component of each isotropy group of the

given $SU(n)$ -action on M is conjugate to one of the following

$$SU(n), SU(n-1) \quad \text{and} \quad NSU(n-1).$$

(i) First we show that the stationary point set $F = F(SU(n), M)$ is non-empty. Assume $F = \emptyset$, then by (1.1) there is a smooth fibre bundle

$$F(SU(n-1), M) \rightarrow M \rightarrow P_{n-1}(C).$$

Thus

$$\chi(M) = \chi(P_{n-1}(C)) \cdot \chi(F(SU(n-1), M))$$

and hence

$$k+1 \equiv 0 \pmod{n}.$$

This is impossible by the assumption $0 \leq k < n-4$. Thus $F \neq \emptyset$. Then by (1.4) there is an equivariant diffeomorphism

$$M = \partial(D^{2n} \times_{S^1} X) / S^1 = D^{2n} \times_{S^1} \partial X \cup S^{2n-1} \times_{S^1} X$$

as $SU(n)$ -manifolds. Here X is a compact connected orientable $(2k+2)$ -manifold with smooth S^1 -action which is free on ∂X .

(ii) Next we show that X is acyclic over rationals. Since

$$D^{2n} \times_{S^1} \partial X \rightarrow \partial X / S^1 = F$$

is a $2n$ -disk bundle, there is an isomorphism

$$H^i(M, S^{2n-1} \times_{S^1} X; \mathbb{Q}) = H^{i-2n}(F; \mathbb{Q}).$$

Thus

$$(2.3) \quad H^i(M; \mathbb{Q}) = H^i(S^{2n-1} \times_{S^1} X; \mathbb{Q}) \quad \text{for } i \leq 2n-2.$$

Now we show that the euler class $e(p)$ of the principal S^1 -bundle

$$p: \partial(D^{2n} \times X) \rightarrow M$$

is non-zero in $H^2(M; \mathbb{Q})$. Assume $e(p) = 0$, then the euler class of the principal S^1 -bundle

$$S^{2n-1} \times X \rightarrow S^{2n-1} \times_{S^1} X$$

is zero in $H^2(S^{2n-1} \times_{S^1} X; \mathbb{Q})$, and hence there is an isomorphism

$$H^*(S^{2n-1}; \mathbb{Q}) \otimes H^*(X; \mathbb{Q}) = H^*(S^1; \mathbb{Q}) \otimes H^*(S^{2n-1} \times_{S^1} X; \mathbb{Q}).$$

Therefore

$$H^i(X; \mathbb{Q}) = \mathbb{Q} \quad \text{for } 0 \leq i \leq 2n-2$$

by (2.3) and the assumption

$$H^*(M; \mathbf{Q}) = H^*(P_{n+k}(C); \mathbf{Q}).$$

But

$$\dim X = 2k+2 \leq 2n-2.$$

Thus $H^{2k+2}(X; \mathbf{Q}) = \mathbf{Q}$ and this is a contradiction, since the connected manifold X has a non-empty boundary. Therefore $e(p) \neq 0$ and hence

$$(2.4) \quad H^*(\partial(D^{2n} \times X); \mathbf{Q}) = H^*(S^{2n+2k+1}; \mathbf{Q}).$$

There is an isomorphism

$$H^i(D^{2n} \times X; \mathbf{Q}) = H_{2n+2k+2-i}(D^{2n} \times X, \partial(D^{2n} \times X); \mathbf{Q})$$

by the Poincaré-Lefschetz duality, and the homomorphism

$$H_{2n+2k+2-i}(D^{2n} \times X; \mathbf{Q}) \rightarrow H_{2n+2k+2-i}(D^{2n} \times X, \partial(D^{2n} \times X); \mathbf{Q})$$

is onto for $0 < i < 2n+2k+2$ by (2.4). Since X is a connected $(2k+2)$ -manifold with a non-empty boundary,

$$H_{2n+2k+2-i}(D^{2n} \times X; \mathbf{Q}) = 0 \quad \text{for } i \leq 2n,$$

and hence

$$H^i(X; \mathbf{Q}) = 0 \quad \text{for } 0 < i \leq 2n.$$

Therefore X is acyclic over rationals. Then

$$H^*(\partial X; \mathbf{Q}) = H^*(S^{2k+1}; \mathbf{Q}),$$

by the Poincaré-Lefschetz duality, and hence

$$H^*(F; \mathbf{Q}) = H^*(P_k(C); \mathbf{Q}).$$

Furthermore $F(S^1, X)$ consists just one point by the P.A. Smith theory (see [2], chapter IV) from the fact that X is acyclic over rationals and the S^1 -action is free on ∂X .

(iii) Next we show $\pi_1(X) = \pi_1(M)$. Since $F(S^1, X) = \{x_0\}$, there is an S^1 -map

$$s: \tilde{M} \rightarrow \partial(D^{2n} \times X)$$

given by $s(y) = (y, x_0)$. Then we have an isomorphism

$$\pi_1(M) = \pi_1(\partial(D^{2n} \times X))$$

from the following commutative diagram:

$$\begin{array}{ccc}
 \pi_1(S^1) & \longrightarrow & \pi_1(S^{2n-1}) \\
 \downarrow id & & \downarrow s_* \\
 \pi_1(S^1) & \longrightarrow & \pi_1(\partial(D^{2n} \times X)) \xrightarrow{p_*} \pi_1(M).
 \end{array}$$

Applying the van Kampen theorem (see [5], p. 63) to the decomposition

$$\partial(D^{2n} \times X) = D^{2n} \times \partial X \cup S^{2n-1} \times X,$$

we have

$$\pi_1(X) = \pi_1(\partial(D^{2n} \times X)),$$

and hence

$$\pi_1(X) = \pi_1(M).$$

(iv) Finally we show that the assumption

$$H^*(M; Z) = H^*(P_{n+k}(C); Z)$$

implies $H^*(X, x_0; Z) = 0$. There is a commutative diagram:

$$\begin{array}{ccc}
 S^{2n-1} & \xrightarrow{s} & \partial(D^{2n} \times X) \\
 \downarrow p_0 & & \downarrow p \\
 P_{n-1}(C) & \xrightarrow{t} & M.
 \end{array}$$

Since $t^*e(p) = e(p_0)$ is a generator of $H^*(P_{n-1}(C); Z)$, $e(p)$ is a generator of $H^*(M; Z)$. Therefore

$$H^*(\partial(D^{2n} \times X); Z) = H^*(S^{2n+2k+1}; Z)$$

by the Gysin sequence for the principal S^1 -bundle

$$p: \partial(D^{2n} \times X) \rightarrow M,$$

and hence X is acyclic over integers and

$$H^*(F; Z) = H^*(P_k(C); Z)$$

by the same argument as in (ii). Then the S^1 -action on X is semi-free by the P.A. Smith theory from the fact that X is acyclic over integers and the S^1 -action is free on ∂X . This completes the proof of Theorem 2.1.

Proof of Corollary 2.2. If M admits a non-trivial smooth $SU(n)$ -action, then by Theorem 2.1, there is an equivariant diffeomorphism

$$M = \partial(D^{2n} \times X)/S^1$$

as $SU(n)$ -manifolds. Here X is a compact contractible $(2k+2)$ -manifold with smooth semi-free S^1 -action with just one stationary point x_0 . Therefore the

S^1 -action on $D^{2n} \times X$ is semi-free and its stationary point is only $(0, x_0)$. Let U be an invariant closed disk around the point $(0, x_0)$. One may assume that the S^1 -action on U is linear. Put

$$W = (D^{2n} \times X - \text{int } U)/S^1.$$

Then

$$\partial W = dU/S^1 \cup \partial(D^{2n} \times X)/S^1 = P_{n+k}(\mathbf{C}) \cup M.$$

Since

$$\begin{aligned}\pi_1(M) &= \pi_1(W) = 0, \\ H_*(W, \mathbf{Z}) &= 0\end{aligned}$$

and

$$\dim W = 2n + 2k + 1 \geq 6,$$

we have

$$M = P_{n+k}(\mathbf{C})$$

by applying the h -cobordism theorem (see [10], Theorem 9.1) to the triad $(W; M, P_{n+k}(\mathbf{C}))$. This completes the proof of Corollary 2.2.

3. Construction of $SU(n)$ -actions

In this section we construct $SU(n)$ -actions on cohomology complex projective spaces, and we have the following results.

Theorem 3.1. *Let $n \geq 2$, $k \geq 1$ and $p \geq 1$. Then there is a compact orientable $2(n+k)$ -manifold M such that*

$$\pi_1(M) = \mathbf{Z}/p\mathbf{Z} \quad \text{and} \quad H^*(M; \mathbf{Q}) = H^*(P_{n+k}(\mathbf{C}); \mathbf{Q})$$

and M admits a smooth $SU(n)$ -action with

$$F(SU(n), M) = P_k(\mathbf{C}).$$

Theorem 3.2. *Let $n \geq 2$ and $k \geq 3$. Let G be a finitely presentable group with $H_1(G; \mathbf{Z}) = H_2(G; \mathbf{Z}) = 0$. Then*

(a) *there is a compact orientable $2(n+k)$ -manifold M such that*

$$\pi_1(M) = G \quad \text{and} \quad H^*(M; \mathbf{Z}) = H^*(P_{n+k}(\mathbf{C}); \mathbf{Z})$$

and M admits a smooth $SU(n)$ -action with

$$F(SU(n), M) = P_k(\mathbf{C}),$$

(b) *there is a smooth $SU(n)$ -action on $P_{n+k}(\mathbf{C})$ such that*

$$\pi_1(F) = G \quad \text{and} \quad H^*(F; \mathbf{Z}) = H^*(P_k(\mathbf{C}); \mathbf{Z}),$$

where $F = F(SU(n), P_{n+k}(\mathbf{C}))$.

First we prepare the following lemma. It is proved by a similar argument as in the proof of Theorem 2.1 and Corollary 2.2, so we omit the proof.

Lemma 3.3. *Let X be a compact orientable $(2k+2)$ -manifold which is acyclic over Z (resp. Q). Assume that X admits a smooth S^1 -action which is free on ∂X . If $n \geq 2$, then*

(a) $M = \partial(D^{2n} \times X)/S^1$ is a cohomology $P_{n+k}(\mathbf{C})$ over Z (resp. Q),

(b) $\pi_1(M) = \pi_1(X)$.

Moreover if $n+k \geq 3$ and X is contractible, then $M = P_{n+k}(\mathbf{C})$.

Now we construct an acyclic S^1 -manifold. Let W be a closed orientable smooth homology $(2k+1)$ -sphere over Z (resp. Q) and let

$$(3.4) \quad Y = P_k(\mathbf{C}) \times [0, 1] \# W, \quad (k \geq 1).$$

Then F is a compact connected orientable smooth $(2k+1)$ -manifold with boundary

$$\partial Y = P_k(\mathbf{C}) \times 0 \cup P_k(\mathbf{C}) \times 1.$$

It is easily seen that

$$(3.5) \quad \pi_1(Y) = \pi_1(W),$$

$$(3.6) \quad H^i(Y; \mathbf{Z}) = H^i(P_k(\mathbf{C}); \mathbf{Z}) \oplus H^i(W; \mathbf{Z}), \quad (0 < i \leq 2k).$$

Furthermore there is a smooth principal S^1 -bundle

$$p: E \rightarrow Y$$

such that $\partial_i E \rightarrow P_k(\mathbf{C}) \times i$, ($i=0, 1$) is equivalent to the Hopf bundle $S^{2k+1} \rightarrow P_k(\mathbf{C})$, where $\partial_i E = p^{-1}(P_k(\mathbf{C}) \times i)$. Then

$$(3.7) \quad \pi_1(E) = \pi_1(Y),$$

$$(3.8) \quad H^*(E, \partial_i E; A) = 0$$

where $A = \mathbf{Z}$ (resp. Q), by (3.6) and the Gysin sequence for S^1 -bundles. Furthermore

$$X = E \cup_{\partial_1 E} D^{2k+2}$$

is a compact orientable manifold with a semi-free smooth S^1 -action which is linear and free on $\partial X = \partial_0 E = S^{2k+1}$. It is easily seen that

$$(3.9) \quad \pi_1(X) = \pi_1(W), \quad \text{by (3.5) and (3.7),}$$

$$(3.10) \quad X \text{ is acyclic over } Z \text{ (resp. } Q), \quad \text{by (3.8).}$$

Proof of Theorem 3.1. Put $W = S^{2k+1}/\mathbf{Z}_p$ lens space, in (3.4). Then there is a compact orientable $(2k+2)$ -manifold X with a semi-free smooth S^1 -action which is linear and free on $\partial X = S^{2k+1}$, such that $\pi_1(X) = \mathbf{Z}_p$ and X is acyclic over \mathbf{Q} . Then by Lemma 3.3, the $SU(n)$ -manifold

$$M = \partial(D^{2n} \times X)/S^1$$

is a compact orientable $2(n+k)$ -manifold such that

$$\pi_1(M) = \mathbf{Z}_p, H^*(M; \mathbf{Q}) = H^*(P_{n+k}(\mathbf{C}); \mathbf{Q})$$

and

$$F(SU(n), M) = \partial X/S^1 = P_k(\mathbf{C}). \quad \text{q.e.d.}$$

REMARK 3.11. It is known that if G is a finitely presentable group with $H_1(G; \mathbf{Z}) = H_2(G; \mathbf{Z}) = 0$, then for each $m \geq 7$, there is a compact contractible smooth n -manifold P such that

$$\pi_1(\partial P) = G \quad (\text{see [12]}).$$

It is known that there are infinitely many groups satisfying the above condition.

Proof of Theorem 3.2 (a). Let $k \geq 3$. Put $W = \partial P$, a smooth homology $(2k+1)$ -sphere over \mathbf{Z} with $\pi_1(\partial P) = G$, in (3.4). Then there is a compact orientable $(2k+2)$ -manifold X with a semi-free smooth S^1 -action which is linear and free on $\partial X = S^{2k+1}$, such that $\pi_1(X) = G$ and X is acyclic over \mathbf{Z} . Then by Lemma 3.3, the $SU(n)$ -manifold

$$M = \partial(D^{2n} \times X)/S^1$$

is a compact orientable $2(n+k)$ -manifold such that

$$\pi_1(M) = G, H^*(M; \mathbf{Z}) = H^*(P_{n+k}(\mathbf{C}); \mathbf{Z})$$

and

$$F(SU(n), M) = P_k(\mathbf{C}). \quad \text{q.e.d.}$$

Proof of Theorem 3.2 (b). Let $k \geq 3$. For a given group G satisfying the hypothesis, there is a compact contractible smooth $(2k+1)$ -manifold P such that

$$\pi_1(\partial P) = G$$

by Remark 3.11. Let

$$Y = P_k(\mathbf{C}) \times [0, 1] \# P,$$

a boundary connected sum with boundary

$$\partial Y = P_k(\mathbf{C}) \# \partial P \cup P_k(\mathbf{C}) \times 1.$$

Then $P_k(\mathbf{C}) \times 1$ is a deformation retract of Y , and hence there is a smooth principal S^1 -bundle

$$p : E \rightarrow Y,$$

such that $\partial_1 E \rightarrow P_k(\mathbf{C}) \times 1$ is equivalent to the Hopf bundle $S^{2k+1} \rightarrow P_k(\mathbf{C})$, where $\partial_1 E = p^{-1}(P_k(\mathbf{C}) \times 1)$. Then

$$X = E \cup_{\partial_1 E} D^{2k+2}$$

is a compact contractible $(2k+2)$ -manifold with a semi-free smooth S^1 -action. Then by Lemma 3.3, the $SU(n)$ -manifold

$$M = \partial(D^{2n} \times X)/S^1$$

is diffeomorphic to $P_{n+k}(\mathbf{C})$ for $n \geq 2$, and

$$F(SU(n), M) = \partial X/S^1 = P_k(\mathbf{C}) \# \partial P.$$

Therefore there is a smooth $SU(n)$ -action on $P_{n+k}(\mathbf{C})$ such that

$$\pi_1(F) = G \quad \text{and} \quad H^*(F; \mathbf{Z}) = H^*(P_k(\mathbf{C}); \mathbf{Z}),$$

where $F = F(SU(n), P_{n+k}(\mathbf{C}))$.

q.e.d.

4. Signature of certain smooth G -manifolds

The purpose of this section is to study a signature of closed orientable manifold which admits a smooth G -action with isotropy groups of uniform dimension. We have the following result.

Theorem 4.1. *Let G be a compact Lie group and H a closed connected subgroup. Let M be a compact orientable manifold without boundary. Assume that M admits a smooth G -action such that the identity component of an isotropy group G_x is conjugate to H in G for each point x of M . Then $F(H, M)$, the stationary point set with respect to the H -action, is orientable, and*

- (a) *if $\dim N(H) \neq \dim H$, then $\text{Sign}(M) = 0$,*
- (b) *if $\dim N(H) = \dim H$, then*

$$|N(H)/H| \text{Sign}(M) = \text{Sign}(G/H) \text{Sign}(F(H, M)).$$

Here $N(H)$ is the *normalizer* of H in G , $|N(H)/H|$ is the order of the finite group $N(H)/H$.

The result is a generalization of the fact that $\text{Sign}(M) = 0$ if M admits a smooth circle action without stationary points.

Lemma 4.2. *Let G be a compact Lie group and H a closed connected subgroup. Let M be a smooth G -manifold such that the identity component of G_x is*

conjugate to H in G for each point x of M . Then

- (a) the $W(H)$ -action on $F(H, M)$ is almost free (i.e. all isotropy groups are discrete), where $W(H) = N(H)/H$,
 (b) there is an equivariant diffeomorphism

$$M = G \times_{N(H)} F(H, M) = G/H \times_{W(H)} F(H, M),$$

- (c) if M is orientable, then $F(H, M)$ is orientable.

Proof. By the assumption, the identity component of G_x is equal to H for each point x of $F(H, M)$, and the mapping

$$/: G \times F(H, M) \rightarrow M$$

given by $f(g, x) = gx$ is surjective. Moreover $f(g, x)$ is in $F(H, M)$ if and only if $g \in N(H)$, thus $W(H)$ acts on $F(H, M)$ naturally and (b) is proved. Next, if an isotropy group $W(H)_x$ is not discrete for a point x of $F(H, M)$, then

$$\dim G_x \neq \dim H.$$

This contradicts our assumption, and (a) is proved. By (b), the product manifold $G/H \times F(H, M)$ is a total space of a principal $W(H)$ -bundle over M . Therefore $G/H \times F(H, M)$ is orientable, if M is orientable, and hence $F(H, M)$ is orientable. q.e.d.

Lemma 4.3. Let G be a compact Lie group which is not discrete. Let M be a compact orientable smooth manifold without boundary. Then, $\text{Sign}(M) = 0$ if M admits an almost free smooth G -action.

Proof. G contains a circle subgroup and the circle action on M has no stationary points. Therefore $\text{Sign}(M) = 0$. q.e.d.

Proof of Theorem 4.1. Denote by $W(H)^0$, the identity component of $W(H)$. Then

$$G/H \times_{W(H)^0} F(H, M)$$

is a total space of a principal $W(H)/W(H)^0$ -bundle over M by Lemma 4.2. (b). Therefore

$$|W(H)/W(H)^0| \cdot \text{Sign}(M) = \text{Sign}(G/H \times_{W(H)^0} F(H, M)).$$

Next, $G/H \times_{W(H)^0} F(H, M)$ is a total space of a smooth fibre bundle over an orientable manifold $(G/H)/W(H)$ with a fibre $F(H, M)$ and a structure group $W(H)^0$ which is connected. Therefore

$$\text{Sign}(G/H \times_{W(H)^0} F(H, M)) = \text{Sign}((G/H)/W(H)^0) \cdot \text{Sign}(F(H, M))$$

for a certain orientation of $F(H, M)$ by [4]. By the above equations,

$$I W(H)/W(H)^0 \text{Sign}(M) = \text{Sign}((G/H)/W(H)^0) \text{Sign}(F(H, M)).$$

Now, if $\dim W(H) \neq 0$ then $\text{Sign}(F(H, M)) = 0$ by Lemma 4.2 (a) and Lemma 4.3. If $\dim W(H) = 0$, then

$$I W(H) \text{Sign}(M) = \text{Sign}(G/H) \text{Sign}(F(H, M)).$$

This completes the proof.

REMARK 4.4. Let G be an arbitrary compact connected Lie group and T be a maximal torus. Then $\text{Sign}(G/T) = 0$, since G/T is stably parallelizable (see [3], section 5.4).

REMARK 4.5. Let G be a compact connected Lie group and H a closed connected subgroup. Then $\text{Sign}(G/H) = 0$ if

$$\text{rank } G \neq \text{rank } H \quad (\text{see [7]}).$$

Because the left translation on G/H of a maximal torus of G has no stationary points.

5. $SU(3)$ -actions on orientable 8-manifolds

The purpose of this section is to prove the following result.

Theorem 5.1. *Let M be a closed connected orientable 8-manifold. Assume that M admits a non-trivial smooth $SU(3)$ -action with a principal isotropy type (H) . Then*

- (a) $H^4(M; \mathbb{Q}) = 0$, if $\dim H = 0$,
- (b) $\text{Sign}(M) = 0$, if $\dim H = 1$ and M has not isotropy types $(NSU(2))$ and $(T_{(2)})$,
- (c) $\text{Sign}(M) = 0$, if $\dim H = 2$,
- (d) $H^4(M; \mathbb{Q}) = 0$, if $\dim H = 3$ and M has not an isotropy type $(NSU(2))$,
- (e) $M = P_2(C) \times F(NSU(2), M)$, if $\dim H = 4$.

Here $NSU(2)$ is the normalizer of $SU(2)$ in $SU(3)$, the identity component of $T_{(2)}$ is a maximal torus of $SU(3)$ and $T_{(2)}$ has 2-components.

First we recall an additivity property of the signature due to S.P. Novikov (see [1], p. 588). Suppose that Y is a compact oriented $4n$ -manifold with boundary dY . Let $\hat{H}^{2n}(Y; \mathbb{Q})$ denote the image of the natural homomorphism

$$j^*: H^{2n}(Y, \partial Y; \mathbb{Q}) \rightarrow H^{2n}(Y; \mathbb{Q}).$$

Then the bilinear form B on $\hat{H}^{2n}(Y \setminus \partial Y; \mathbb{Q})$ defined by

$$B(j^*(a), j^*(b)) = ab[Y]$$

is symmetric and non-degenerate by Poincare-Lefschetz duality. We can now define $\text{Sign}(Y)$ as the signature of B . Suppose now that Y' is another compact oriented $4n$ -manifold with boundary $\partial Y' = -\partial Y$. Then $X = Y \cup_{\partial Y} Y'$ is a closed oriented $4n$ -manifold and

$$(5.2) \quad \text{Sign}(X) = \text{Sign}(Y) + \text{Sign}(Y').$$

REMARK 5.3. Let ξ be an orientable k -plane bundle over a closed orientable manifold X . Denote by $t(\xi)$, $e(\xi)$ and $D(\xi)$, the Thom class, the Euler class and the disk bundle of ξ , respectively. Then $D(\xi)$ is a compact orientable manifold and there is a commutative diagram:

$$\begin{array}{ccc} H^*(D(\xi), \partial D(\xi)) & \xrightarrow{j^*} & H^*(D(\xi)) \\ \cong \uparrow \psi & & \cong \uparrow \pi^* \\ H^*(X) & \xrightarrow{\cdot e(\xi)} & H^*(X). \end{array}$$

Here ψ is the Thom isomorphism defined by

$$\psi(a) = \pi^*(a) \cdot t(\xi).$$

There is an equation

$$\psi(a) \cdot \psi(b) = (-1)^{k^2} \psi(ab \cdot e(\xi)) \quad \text{for } b \in H^p(X).$$

Therefore we can calculate $\text{Sign}(D(\xi))$ from the information about the cohomology ring $H^*(X)$ and the Euler class $e(\xi)$.

Now we prepare the following results.

Lemma 5.4.

- (a) $H^*(SU(3); \mathbf{Z}) = \wedge_{\mathbf{Z}}(x_3, x_5)$, $\deg x_i = i$, $(i=3, 5)$.
- (b) $H^*(SU(3)/SU(2); \mathbf{Z}) = H^*(S^5; \mathbf{Z})$ and the right translation of $NSU(2)/SU(2) = S^1$ induces a trivial action on $H^*(SU(3)/SU(2); \mathbf{Z})$.
- (c) $H^*(SU(3)/SO(3); \mathbf{Q}) = H^*(S^5; \mathbf{Q})$, and the right translation of $NSO(3)/SO(3) = \mathbf{Z}_3$ induces a trivial action on $H^*(SU(3)/SO(3); \mathbf{Q})$.
- (d) $H^*(SU(3)/T; \mathbf{Z}) = \mathbf{Z}[u_1, u_2, u_3]/(s_1, s_2, s_3)$, where T is a maximal torus of $SU(3)$ consists of all diagonal matrices, s_k is the k -th elementary symmetric polynomials, and $\deg u_i = 2$, $(i=1, 2, 3)$. Furthermore the induced action of $N(T)/T = S_3$, the symmetric group on 3-elements, is given by

$$a^*(u_i) = u_{a(i)}, \quad a \in S_3.$$

- (e) $H^*(SU(3)/D(m, n); \mathbf{Q}) = \wedge_{\mathbf{Q}}(x_2, x_5)$, $\deg x_i = i$, $(i=2, 5)$. Here $D(m, n)$ is a closed one-dimensional subgroup defined by

$$D(m, n) = \left\{ \begin{pmatrix} z^m & & \\ & z^n & \\ & & z^{-(m+n)} \end{pmatrix} ; z \in \mathbb{C}, |z|=1 \right\}$$

for any pair of integers $(m, n) \neq (0, 0)$.

Proof. Since $SU(3)/SU(2) = S^5$ (b) is true. (a) is proved by making use of the Gysin sequence for

$$SU(2) \rightarrow SU(3) \rightarrow S^5.$$

(c) is proved from

$$\pi_1(SU(3)/SO(3)) = 0 \text{ and } \pi_2(SU(3)/SO(3)) = \mathbb{Z}_2.$$

(d) is a classical result (see [9]). In fact $u_i = p_i^*(u)$, where u is a generator of $H^2(P_2(\mathbb{C}); \mathbb{Z})$ and $p_i: SU(3)/T \rightarrow P_2(\mathbb{C})$ is defined by

$$p_i((x_{ab}) \cdot T) = (x_{1i}: x_{2i}: x_{3i}).$$

Finally (e) is proved from the fact that the Euler class of principal S^1 -bundle $\pi: SU(3)/D(m, n) \rightarrow SU(3)/T$ is

$$e(\pi) = nu_1 + mu_2,$$

and hence the homomorphism

$$H^2(SU(3)/T; \mathbb{Q}) \xrightarrow{\cdot e(\pi)} H^4(SU(3)/T; \mathbb{Q})$$

is an isomorphism.

q.e.d.

Lemma 5.5.

(a) Let φ be an 8-dimensional non-trivial real representation of $SU(3)$. Let (H_φ) be the principal isotropy type of the linear action given by φ . Then there are only the following cases:

- (i) $\varphi = Ad_{SU(3)}, H_\varphi = T$: a maximal torus of $SU(3)$,
- (ii) $\varphi = \rho_3 + \text{trivial summand}, H_\varphi = SU(2)$,

where $\rho_3: SU(3) \rightarrow O(6)$ is the standard representation.

(b) Let ψ be a 4-dimensional non-trivial real representation of $NSU(2)$. Let (H_ψ) be the principal isotropy type of the linear action given by ψ . Then there are only the following cases:

- (i) $\psi = Ad_{NSU(2)}, H_\psi = T$: a maximal torus of $NSU(2)$,
- (ii) $\psi = \sigma_k, H_\psi = D(k-1, -k), (k \in \mathbb{Z})$,

where the representation $\sigma_k: NSU(2) \rightarrow U(2) \subset O(4)$ is given by

$$\sigma_k \begin{pmatrix} x_{11} & x_{12} & 0 \\ x_{21} & x_{22} & 0 \\ 0 & 0 & y \end{pmatrix} = \begin{pmatrix} y^k x_{11} & y^k x_{12} \\ y^k x_{21} & y^k x_{22} \end{pmatrix}.$$

(iii) ψ is induced from a non-trivial real representation of S^1 , via the natural projection $NSU(2) \rightarrow NSU(2)/SU(2) \cong S^1$, and $H_\psi^0 = SU(2)$, where H_ψ^0 is the identity component of H_ψ .

We omit the proof (see [8], Theorem 1).

From now on we assume that M is a closed connected orientable smooth 8-manifold and M admits a non-trivial smooth $SU(3)$ -action with a principal isotropy type (H) . Then $SU(3)/H$ is orientable by the differentiable slice theorem (see [11], Lemma 3.1).

We will prove Theorem 5.1 by the following many propositions.

Proposition 5.6. Assume that $SU(3)_x^0$ is conjugate to H^0 in $SU(3)$ for each $x \in M$. Here G^0 is the identity component of G and $SU(3)_x$ is the isotropy group at x . Then,

- (a) $\text{Sign}(M) = 0$, if $\dim H = 1$ or 2 ,
- (b) $H^*(M; \mathbb{Q}) = 0$, if $\dim H = 0$ or 3 ,
- (c) $M = P_2(\mathbb{C}) \times F(NSU(2), M)$, if $\dim H = 4$.

Proof. If $\dim H = 1$ or 2 , then $\text{Sign}(M) = 0$ by Theorem 4.1 and Remarks 4.4, 4.5. If $\dim H = 0$, then $M = SU(3)/H$ and hence $H^*(M; \mathbb{Q}) = 0$ by Lemma 5.4 (a). By Lemma 4.2, there is an equivariant diffeomorphism

$$M = SU(3)/H^0 \times_{\kappa} F\mathbb{K} = N(H^0)/H^0, F = F(H^0, M).$$

If $\dim H = 4$, then H^0 is conjugate to $NSU(2)$ in $SU(3)$ and $N(NSU(2)) = NSU(2)$. Therefore

$$M = P_2(\mathbb{C}) \times F(NSU(2), M).$$

Finally if $\dim H = 3$, then H^0 is conjugate to $SO(3)$ or $SU(2)$ in $SU(3)$. If $H^0 = SO(3)$, then $\dim F = 3$ and

$$H^*(M; \mathbb{Q}) = H^*(SU(3)/SO(3) \times F\mathbb{Q}) = 0$$

by Lemma 5.4 (c). Next if $H^0 = SU(2)$, then $\dim F = 4$, F admits a smooth S^1 -action without stationary points and there is an equivariant diffeomorphism

$$M = S^5 \times_{S^1} F.$$

There is a sufficiently large integer n such that the S^1/\mathbb{Z}_n -action on the orbit space F/\mathbb{Z}_n is free. Then there is an isomorphism

$$H^*(M; \mathbf{Q}) = H^*(M'; \mathbf{Q}),$$

where

$$M' = (S^5/\mathbf{Z}_n \times F/\mathbf{Z}_n)/(S^1/\mathbf{Z}_n),$$

and there is a fibre bundle

$$S^5/\mathbf{Z}_n \rightarrow M' \rightarrow F/S^1$$

with a structure group S^1/\mathbf{Z}_n . Here $F/S^1 = (F/\mathbf{Z}_n)/(S^1/\mathbf{Z}_n)$ is a 3-dimensional rational cohomology manifold. Therefore

$$H^*(M; \mathbf{Q}) = H^*(M'; \mathbf{Q}) = 0. \quad \text{q.e.d.}$$

REMARK 5.7. Now Theorem 5.1 is proved for $\dim H=0$ or 4. Moreover, Theorem 5.1 is proved for the case $H^0=SO(3)$, since $SO(3)$ is not conjugate to any subgroup of $NSU(2)$ in $SU(3)$ and H with $H^0=SO(3)$ is not a principal isotropy group of any 8-dimensional real representation of $SU(3)$ by Lemma 5.5.

Proposition 5.8. Suppose $\dim H=1$. Then $\text{Sign}(M)=0$, if M has not isotropy types $(NSU(2))$ and $(T_{(2)})$.

Proof. By Proposition 5.6 (a), one may assume that there is an isotropy type (K_1) with $\dim K_1 > 1$. Then by making use of the differentiable slice theorem, there is an isotropy type (K_2) and there is an equivariant decomposition

$$M = D(\nu_1) \cup D(\nu_2),$$

where $D(\nu_i)$ is an equivariant normal disk bundle of an embedding $SU(3)/K_i$ in CM , and

$$\partial D(\nu_1) = -\partial D(\nu_2) = SU(3)/H.$$

Thus

$$\text{Sign}(M) = \text{Sign}(D(\nu_1)) + \text{Sign}(D(\nu_2)).$$

Since

$$H^*(SU(3)/K; \mathbf{Q}) = 0 \quad \text{for } \dim K \neq 2, 4,$$

by Lemma 5.4, $\text{Sign}(D(\nu_i))=0$ for $\dim K_i \neq 2, 4$. Let K be a 2-dimensional closed subgroup of $SU(3)$. Then K is conjugate to one of the following

$$T, T_{(2)}, T_{(3)} \quad \text{and} \quad N(T) = T_{(6)}.$$

Here $T_{(i)}^0 = T$ and $T_{(i)}$ has i -components. By Lemma 5.4 (d),

$$H^*(SU(3)/T; \mathbf{Q}) = \mathbf{Q} \oplus \mathbf{Q},$$

$$H^*(SU(3)/T_{(2)}; \mathbf{Q}) = \mathbf{Q},$$

$$H^*(SU(3)/T_{(3)}; \mathbf{Q}) = H^*(SU(3)/N(T); \mathbf{Q}) = 0.$$

Thus $\text{Sign}(D(v_i))=0$, if $K_i=T_{(3)}$ or $N(T)$. If $K_i=T$, then $\text{Sign}(D(v_i))=0$ from Lemma 5.4 (d) and Remark 5.3. q.e.d.

REMARK 5.9. If $\dim H=2$ in Theorem 5.1, then $H=T$ or $T_{(3)}$, since $SU(3)/T_{(2)}$ and $SU(3)/N(T)$ are non-orientable by Lemma 5.4(d). Theorem 5.1 is proved for $H=T_{(3)}$ by Proposition 5.6, since $T_{(3)}$ is not conjugate to any subgroup of $NSU(2)$ in $SU(3)$ and $T_{(3)}$ is not a principal isotropy group of any 8-dimensional real representation of $SU(3)$ by Lemma 5.5. Therefore, it remains to prove Theorem 5.1 for the cases $H=T$ and $H^0=SU(2)$.

Proposition 5.10. *Suppose $H=T$. Then $\text{Sign}(M)=0$.*

Proof. If $F(NSU(2), M)$ is empty, then $\text{Sign}(M)=0$ by Proposition 5.6. Now we assume that $F(NSU(2), M)$ is not empty. Then

$$\dim F(NSU(2), M) = 1$$

by Lemma 5.5, and any stationary point (if exists) of $SU(3)$ is isolated by Lemma 5.5 (a). Let

$$F(SU(3), M) = \{x_1, \dots, x_k\}, \quad (k \geq 0)$$

and let D_i be an invariant closed disk around x_i , such that

$$D_i \cap D_j = \emptyset \quad \text{for } i \neq j.$$

Let $D = D_1 \cup \dots \cup D_k$ and $E = M - \text{int } D$. Then

$$D_i \cap F(NSU(2), E) \neq \emptyset, \quad (i = 1, \dots, k)$$

by Lemma 5.5 (a). Let

$$E_0 = \{x \in E \mid (SU(3)_x) = (NSU(2))\},$$

let U_0 be an invariant closed tubular neighborhood of E_0 in E , and let $U = U_0 \cup D$. Then $M - \text{int } U$ is connected and

$$(SU(3)_x^0) = (T), \quad \text{for } x \in M - \text{int } U.$$

Therefore, there is an equivariant diffeomorphism

$$M - \text{int } U \cong SU(3)/T \times_{N(T)/T} F, \quad F = F(T, M - \text{int } U)$$

by Lemma 4.2, and there is a commutative diagram:

$$\begin{array}{ccc} H^*(M - \text{int } U; \mathbb{Q}) & \xrightarrow{i^*} & H^*(\partial(M - \text{int } U); \mathbb{Q}) \\ \cong \downarrow p^* & & \cong \downarrow p^* \\ H^*(SU(3)/T \times F; \mathbb{Q})^{N(T)/T} & \xrightarrow{i_0^*} & H^*(SU(3)/T \times \partial F; \mathbb{Q})^{N(T)/T} \end{array}$$

Here i_0^* is injective, since $H^{odd}(SU(3)/T\mathbf{Q})=0$ by Lemma 5.4 (d), $\dim F=2$, and each connected component of F has non-empty boundary from the connectedness of $M - \text{int } U$. Thus

$$\hat{H}^*(M - \text{int } U; \mathbf{Q}) = 0 ,$$

and hence $\text{Sign}(M - \text{int } U)=0$. Next, let U_1, \dots, U_n be connected components of U . Then we can prove that

$$\begin{aligned} \hat{H}^*(U_i; \mathbf{Q}) &= 0 , & \text{if } U \cap D = \emptyset , \\ H^*(U_i; \mathbf{Q}) &= 0 , & \text{if } U_i \cap D \neq \emptyset , \end{aligned}$$

and hence

$$\text{Sign}(U) = \text{Sign}(U_1) + \dots + \text{Sign}(U_n) = 0 .$$

Therefore

$$\text{Sign}(M) = \text{Sign}(M - \text{int } U) + \text{Sign}(U) = 0 . \quad \text{q.e.d.}$$

We recall the following result which is essentially proved in the proof of Proposition 5.6 (b).

Lemma 5.11. *Let X be a compact connected orientable smooth n -manifold (∂X is empty or not). Let $n=7$ or 8 . Assume that X admits a smooth $SU(3)$ -action with*

$$(SU(3)_x^0) = (SU(2)) \quad \text{for } x \in X.$$

Then

$$H^{n-*}(X; \mathbf{Q}) = 0 .$$

Proposition 5.12. *Assume that $H^0 = SU(2)$ and M has not an isotropy type $(NSU(2))$. Then $H^*(M; \mathbf{Q})=0$.*

Proof. If $F(SU(3), M) = \emptyset$, then $H^*(M; \mathbf{Q})=0$ by Lemma 5.11. Next if $F(SU(3), M) \neq \emptyset$, then $\dim F(SU(3), M)=2$ by Lemma 5.5 (a). Let U be an invariant closed tubular neighborhood of $F(SU(3), M)$ in M . Then there is an exact sequence:

$$H^3(\partial U; \mathbf{Q}) \rightarrow H^*(M; \mathbf{Q}) \rightarrow H^*(U; \mathbf{Q}) \oplus H^*(M - \text{int } U; \mathbf{Q}) .$$

Here

$$H^3(\partial U; \mathbf{Q}) = H^*(M - \text{int } U; \mathbf{Q}) = 0$$

by Lemma 5.11, and

$$H^*(U; \mathbf{Q}) = H^*(F(SU(3), M); \mathbf{Q}) = 0 .$$

Therefore

$$H^*(M; \mathbf{Q}) = 0 . \quad \text{q.e.d.}$$

This completes the proof of Theorem 5.1.

6. SZ7(3)-actions on cohomology $P_4(\mathbb{C})$

In the previous paper [13] we have considered smooth $SU(3)$ -actions on homotopy $P_3(\mathbb{C})$. In this section, first we prove the following result as an application of Theorem 5.1.

Theorem 6.1. *Let M be a compact connected orientable 8-manifold such that*

$$H^*(M; \mathbb{Q}) = H^*(P_4(\mathbb{C}); \mathbb{Q}).$$

Then for any non-trivial smooth $SU(3)$ -action on M , the stationary point set is a 2-sphere and the principal isotropy type is $(SU(2))$. Furthermore there is an equivariant diffeomorphism

$$M = \partial(D^6 \times X)/S^1.$$

Here X is a compact connected orientable 4-manifold which is acyclic over rationals, X admits a smooth S^1 -action which is free on ∂X , the $SU(3)$ -action is standard on D^6 and trivial on X .

Proof. Denote by (H) , the principal isotropy type of the given $SU(3)$ -action on M . Since $\text{Sign}(M) \neq 0$, the following are the only possible cases from Theorem 5.1,

- (a) $\dim H = 1$ and M has an isotropy type $(NSU(2))$ or $(T_{(2)})$,
- (b) $H = SU(2)$ and M has an isotropy type $(NSU(2))$,
- (c) $H = NSU(2)$ and $M = P_2(\mathbb{C}) \times F(NSU(2), M)$.

If $H = NSU(2)$, then $\chi(M) = 5$ is divisible by $\chi(P_2(\mathbb{C})) = 3$, and this is a contradiction. Next if $\dim H = 1$, then there is a decomposition

$$M = D(\nu_1) \cup D(\nu_2)$$

as in the proof of Proposition 5.8, where $D(\nu_i)$ is a normal disk bundle over $SU(3)/K_i$. One may assume $K_1 = NSU(2)$ or $T_{(2)}$, and hence

$$\chi(SU(3)/K_1) = 3$$

by Lemma 5.4. On the other hand,

$$5 = \chi(M) = \chi(SU(3)/K_1) + \chi(SU(3)/K_2)$$

Thus $\chi(SU(3)/K_2) = 2$, and hence $K_2 = T_{(3)}$ by Lemma 5.4. Since $H^2(SU(3)/T_{(3)}; \mathbb{Q}) = 0$, there is a contradiction in the following exact sequence of rational cohomology groups:

$$\begin{aligned} H^1(\partial D(\nu_1)) \rightarrow H^2(M) &\rightarrow H^2(SU(3)/K_1) \oplus H^2(SU(3)/K_2) \\ &\rightarrow H^2(\partial D(\nu_1)) \rightarrow H^3(M). \end{aligned}$$

Therefore we obtain $H^0 = SU(2)$. If $F(SU(3), M) = \emptyset$, then there is a fibre bundle

$$F(SU(2), M) \rightarrow M \rightarrow P_2(\mathbb{C}).$$

Thus $\chi(M) = 5$ is divisible by $\chi(P_2(\mathbb{C})) = 3$, and this is a contradiction. Hence $F(SU(3), M) \neq \emptyset$ and this implies $H = SU(2)$ by Lemma 5.5 (a). Let U be an invariant tubular neighborhood of $F(SU(3), M)$ in M . Then

$$X = F(SU(2), M - \text{int } U)$$

is a compact connected orientable 4-manifold with the natural action of $NSU(2)/SU(2) = S^1$ which is free on ∂X . Furthermore there is an equivariant diffeomorphism

$$M = \partial(D^6 \times X)/S^1,$$

and X is acyclic over rationals by the same argument as in the proof of Theorem 2.1. Finally,

$$F(SU(3), M) = \partial X/S^1 = S^2. \quad \text{q.e.d.}$$

Next, as a complementary part of Theorem 5.1, we give examples of certain $SU(3)$ -actions on 8-manifolds with non-zero signature.

Let $\psi: NSU(2) \rightarrow U(3)$ be a unitary representation of $NSU(2)$. Then ψ induces a smooth $NSU(2)$ -action ψ_* on $P_2(\mathbb{C})$. Denote by $M(\psi)$, the orbit manifold of the free smooth action of $NSU(2)$ on $SU(3) \times P_2(\mathbb{C})$ given by

$$h \cdot (g, x) = (gh^{-1}, \psi_*(h, x)), \quad g \in SU(3), \quad h \in NSU(2), \quad x \in P_2(\mathbb{C}).$$

Then the compact connected orientable 8-manifold $M(\psi)$ admits a natural smooth $SU(3)$ -action without stationary points and

$$\text{Sign}(M(\psi)) = 1.$$

EXAMPLE 6.2. Let $\alpha_k: NSU(2) \rightarrow U(3)$ be a unitary representation given by

$$\alpha_k \left(\begin{array}{ccc|ccc} * & * & 0 & 1 & 0 & 0 \\ * & * & 0 & 0 & 1 & 0 \\ 0 & 0 & y & 0 & 0 & y^k \end{array} \right).$$

Then $M(\alpha_k)$ has just two isotropy types

$$(SU(2)_{(k)}) \quad \text{and} \quad (NSU(2)),$$

where $SU(2)_{(k)}$ has k -components and its identity component is $SU(2)$. (see Theorem 5.1 (d))

EXAMPLE 6.3. Let $\beta_k: NSU(2) \rightarrow U(3)$ be a unitary representation given by

$$\beta_k \begin{pmatrix} x_{11} & x_{12} & 0 \\ x_{21} & x_{22} & 0 \\ 0 & 0 & y \end{pmatrix} = \begin{pmatrix} x_{11} & x_{12} & 0 \\ x_{21} & x_{22} & 0 \\ 0 & 0 & y^k \end{pmatrix}.$$

Then $M(\beta_k)$ has just three isotropy types

$$(D(k, -k-1)), (T) \quad \text{and} \quad (NSU(2)),$$

where $D(k, -k-1)$ is a closed one-dimensional subgroup defined in Lemma 5.4. (see Theorem 5.1 (b))

EXAMPLE 6.4. Let $\gamma: NSU(2) \rightarrow U(3)$ be a unitary representation given by

$$\gamma \begin{pmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & * \end{pmatrix} = \begin{pmatrix} a^2 & \sqrt{2}ab & b^2 \\ \sqrt{2}ac & ad+bc & \sqrt{2}bd \\ c^2 & \sqrt{2}cd & d^2 \end{pmatrix}.$$

Then $M(\gamma)$ has just three isotropy types

$$(D(1, 1)_{(2)}), (T) \quad \text{and} \quad (T_{(2)}),$$

where $G_{(2)}$ is a subgroup of $SU(3)$ such that $G_{(2)}$ has 2-components and its identity component is G . (see Theorem 5.1 (b))

7. Classification of smooth $SU(n)$ -actions on orientable $2n$ -manifolds

Let M be a compact connected $2n$ -manifold with non-trivial smooth $SU(n)$ -action, then the identity component of each isotropy group is conjugate to one of the following

$$SU(n), SU(n-1) \quad \text{and} \quad NSU(n-1),$$

for $n \geq 5$. This is proved similarly as Lemma 1.5. Therefore there is an equivariant diffeomorphism

$$M = \partial(D^{2n} \times X)/S^1$$

as $SU(n)$ -manifolds by (1.1) and (1.4). Here X is a compact connected 2-dimensional S^1 -manifold and the S^1 -action on dX is free if dX is non-empty. Furthermore if M is orientable, then X is also orientable. Next we remark that for orientable 2-dimensional S^1 -manifold X , if the isotropy group $S_x^1 \neq S^1$ for $x \in X$, then S_x^1 is a principal isotropy group by the differentiable slice theorem, and hence the S^1 -space $X - F(S^1, X)$ has just one isotropy type.

(i) If X has just one isotropy type (S^1), then $\partial X = \emptyset$ and

$$M = P_{n-1}(C) \times X.$$

(ii) If X has just one isotropy type (\mathbf{Z}_k) , then

$$\begin{aligned} M &= S^{2n} && \text{if } \partial X \neq \emptyset, \\ M &= L^{2n-1}(k) \times S^1 && \text{if } \partial X = \emptyset. \end{aligned}$$

Here $L^{2n-1}(k) = S^{2n-1}/\mathbf{Z}_k$ is a standard lens space.

(iii) If X has just two isotropy types (\mathbf{Z}_k) and (S^1) , then

$$\begin{aligned} M &= P_n(\mathbf{C}) && \text{if } \partial X \neq \emptyset, \\ M &= S^{2n-1} \times_{S^1} S^2_{(k)} && \text{if } \partial X = \emptyset. \end{aligned}$$

Here $S^2_{(k)}$ is a 2-sphere with the S^1 -action given by

$$e^{i\theta} (x_0, x_1, x_2) = (x_0, x_1 \cos k\theta + x_2 \sin k\theta, -x_1 \sin k\theta + x_2 \cos k\theta).$$

This completes the classification.

OSAKA UNIVERSITY

References

- [1] M.F. Atiyah and I.M. Singer: *The index of elliptic operators* III, Ann. of Math. 87 (1968), 546-604.
- [2] A. Borel et al.: *Seminar on Transformation Groups*, Ann. of Math. Studies, 46, Princeton Univ. Press, 1960.
- [3] A. Borel and F. Hirzebruch: *Characteristic classes and homogeneous spaces* III, Amer. J. Math. 82 (1960), 491-504.
- [4] S.S. Chern, F. Hirzebruch and J.P. Serre: *On the index of a fibered manifolds*, Proc. Amer. Math. Soc. 8 (1957), 587-596.
- [5] R. Crowell and R. Fox: *Introduction to Knot Theory*, Ginn and Co., 1963.
- [6] E.B. Dynkin: *The maximal subgroups of the classical groups*, Amer. Math. Soc. Transl. 6 (1957), 245-378.
- [7] A. Hattori: *The index of coset spaces of compact Lie groups*, J. Math. Soc. Japan 14 (1962), 26-36.
- [8] W.Y. Hsiang: *On the principal orbit type and P.A. Smith theory of $SU(p)$ actions*, Topology 6 (1967), 125-135.
- [9] J. Leray: *Sur l'anneau d'homologie de l'espace homogène*, C. R. Acad. Sci. Paris, 223 (1946), 412-415.
- [10] J. Milnor: *Lectures on the h -cobordism Theorem*, Princeton Math. Notes, 1965.
- [11] D. Montgomery, H. Samelson and C.T. Yang: *Exceptional orbits of highest dimension*, Ann. of Math. 64 (1956), 131-141.
- [12] I. Tamura: *Variety of varieties* (Japanese), Sugaku 21 (1969), 275-285.
- [13] F. Uchida: *Linear $SU(n)$ -actions on complex projective spaces*, Osaka J. Math. 11 (1974), 473-481.
- [14] H. Weyl: *The Classical Groups*, 2nd ed. Princeton Univ. Press, 1946.