

Title	Growth properties of $p$ -th means of biharmonic Green potentials in the unit ball
Author(s)	Futamura, Toshihide; Mizuta, Yoshihiro
Citation	Osaka Journal of Mathematics. 42(1) p.85-p.99
Issue Date	2005-03
oaire:version	VoR
URL	<a href="https://doi.org/10.18910/12070">https://doi.org/10.18910/12070</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## GROWTH PROPERTIES OF $p$ -TH MEANS OF BIHARMONIC GREEN POTENTIALS IN THE UNIT BALL

Dedicated to Professor Masakazu Shiba on the occasion of his sixtieth birthday

TOSHIHIDE FUTAMURA and YOSHIHIRO MIZUTA

(Received July 22, 2003)

### Abstract

Let  $u$  be a biharmonic Green potential on the unit ball  $\mathbf{B}$  of  $\mathbf{R}^n$ . We show that

$$\lim_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p} \mathcal{M}_p(u, r) = 0$$

for  $p$  such that  $1 \leq p < (n-1)/(n-4)$  in case  $n \geq 5$  and  $1 \leq p < \infty$  in case  $n \leq 4$ . Further, if  $n \geq 5$  and  $(n-1)/(n-4) \leq p < (n-1)/(n-5)$ , then it is shown that

$$\liminf_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p} \mathcal{M}_p(u, r) = 0.$$

Finally we show that these limits characterize biharmonic Green potentials among super-biharmonic functions on  $\mathbf{B}$ .

### 1. Introduction and statement of results

A function  $u$  on an open set  $\Omega \subset \mathbf{R}^n$  ( $n \geq 2$ ) is called biharmonic if  $u \in C^4(\Omega)$  and  $\Delta^2 u = 0$  on  $\Omega$ , where  $\Delta$  denotes the Laplacian and  $\Delta^2 u = \Delta(\Delta u)$ . We say that a lower semicontinuous and locally integrable function  $u$  on  $\Omega$  is super-biharmonic in  $\Omega$  if every point of  $\Omega$  is a Lebesgue point of  $u$  and  $\Delta^2 u$  is a nonnegative measure on  $\Omega$  in the weak sense, that is,

$$\int_{\Omega} u(x) \Delta^2 \varphi(x) dx \geq 0 \quad \text{for all nonnegative } \varphi \in C_0^\infty(\Omega).$$

The open ball and the sphere centered at  $x$  with radius  $r$  are denoted by  $B(x, r)$  and  $S(x, r)$ . We write  $B(r) = B(0, r)$  and  $S(r) = S(0, r)$ . We also denote by  $\mathbf{B}$  and  $\mathbf{S}$  the unit ball  $B(1)$  and the unit sphere  $S(1)$ . We write  $x^* = |x|^{-2}x$ , so that  $x^*$  is the inverse point of  $x$  relative to the unit sphere  $\mathbf{S}$ .

Let  $G_2(x, y)$  denote the biharmonic Green function in the unit ball  $\mathbf{B}$  (cf. [8]), that

is,

$$G_2(x, y) = \begin{cases} \alpha_n \left( |x-y|^{4-n} - (|y||x-y^*|)^{4-n} - \frac{n-4}{2}(1-|x|^2)(1-|y|^2)(|y||x-y^*|)^{2-n} \right) & \text{in case } n \neq 2, 4, \\ \alpha_n \left( |x-y|^{4-n} \log \left( \frac{|x-y|}{|y||x-y^*|} \right)^2 + (1-|x|^2)(1-|y|^2)(|y||x-y^*|)^{2-n} \right) & \text{in case } n = 2, 4, \end{cases}$$

where  $\alpha_n^{-1} = 2(4-n)(2-n)\sigma_n$  for  $n \neq 2, 4$  and  $\alpha_n^{-1} = (-1)^{n/2+1}8\sigma_n$  for  $n = 2, 4$ . Here  $\sigma_n$  denotes the surface measure of the unit sphere  $\mathbf{S}$ . For a nonnegative measure  $\mu$  on  $\mathbf{B}$ , we define

$$G_2\mu(x) = \int_{\mathbf{B}} G_2(x, y) d\mu(y).$$

If  $\mu$  has a density  $f \in L^1_{\text{loc}}(\mathbf{B})$ , then we write  $G_2f$  instead of  $G_2\mu$ . The function  $G_2\mu$  is called a biharmonic Green potential if  $G_2\mu \neq \infty$ .

For a Borel measurable function  $u$  on  $S(r)$ , we define the average integral over  $S(r)$  by

$$\mathcal{M}(u, r) = \int_{S(r)} u dS = \frac{1}{|S(r)|} \int_{S(r)} u dS,$$

where  $|S(r)|$  denotes the surface measure of  $S(r)$ . For  $p > 0$  and a Borel measurable function  $u$  on  $S(r)$ , define  $\mathcal{M}_p(u, r) = \{\mathcal{M}(|u|^p, r)\}^{1/p}$ .

Gardiner [5] and the second author [9] studied the limiting behavior of  $\mathcal{M}_p(v, r)$  for (harmonic) Green potentials on  $\mathbf{B}$ . We also refer to Stoll [13, 14] for invariant potentials in the unit ball of  $\mathbf{C}^n$ .

In this paper we are concerned with biharmonic Green potentials on  $\mathbf{B}$ .

**Theorem 1.1.** *Let  $G_2\mu$  be a biharmonic Green potential on  $\mathbf{B}$ . If  $1 \leq p < (n-1)/(n-4)$  in case  $n \geq 5$  and  $1 \leq p < \infty$  in case  $n \leq 4$ , then*

$$\lim_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p} \mathcal{M}_p(G_2\mu, r) = 0.$$

**Theorem 1.2.** *Let  $G_2\mu$  be a biharmonic Green potential on  $\mathbf{B}$ . If  $n \geq 5$  and  $(n-1)/(n-4) \leq p < (n-1)/(n-5)$ , then*

$$\liminf_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p} \mathcal{M}_p(G_2\mu, r) = 0.$$

Finally we give a characterization for a super-biharmonic function to be a biharmonic Green potential on  $\mathbf{B}$ .

**Theorem 1.3.** *Let  $u$  be a super-biharmonic function on  $\mathbf{B}$ . If  $u$  satisfies*

$$\liminf_{r \rightarrow 1} (1-r)^{-1} \mathcal{M}_1(u, r) = 0,$$

*then it is a biharmonic Green potential on  $\mathbf{B}$ .*

## 2. $p$ -th means of biharmonic Green potentials

Throughout this paper, let  $M$  denote various constants independent of the variables in question.

We need the following fundamental estimates for the biharmonic Green function on the unit ball  $\mathbf{B}$  (cf. [1] and [7]).

**Lemma 2.1.** *There exist positive constants  $C_i$ ,  $1 \leq i \leq 4$ , satisfying the following conditions:*

(1) *If  $n \geq 5$ , then for every  $(x, y) \in \mathbf{B} \times \mathbf{B}$*

$$0 < C_1 \frac{(1-|x|^2)^2(1-|y|^2)^2}{|x-y|^{n-4}(|y||x-y^*|)^4} \leq G_2(x, y) \leq C_2 \frac{(1-|x|^2)^2(1-|y|^2)^2}{|x-y|^{n-4}(|y||x-y^*|)^4}.$$

(2) *If  $n = 4$ , then for every  $(x, y) \in \mathbf{B} \times \mathbf{B}$*

$$\begin{aligned} 0 < C_1 \frac{(1-|x|^2)^2(1-|y|^2)^2}{(|y||x-y^*|)^4} \log \left( \frac{2|y||x-y^*|}{|x-y|} \right) \\ \leq G_2(x, y) \leq C_2 \frac{(1-|x|^2)^2(1-|y|^2)^2}{(|y||x-y^*|)^4} \log \left( \frac{2|y||x-y^*|}{|x-y|} \right). \end{aligned}$$

(3) *If  $n = 2, 3$ , then for every  $(x, y) \in \mathbf{B} \times \mathbf{B}$*

$$C_1 \frac{(1-|x|^2)^2(1-|y|^2)^2}{(|y||x-y^*|)^n} \leq G_2(x, y) \leq C_2 \frac{(1-|x|^2)^2(1-|y|^2)^2}{(|y||x-y^*|)^n}.$$

*Furthe, in all cases,*

$$C_3 \frac{(1-|x|^2)^2(1-|y|^2)^2}{(|y||x-y^*|)^n} \leq G_2(x, y) \leq C_4 \frac{(1-|x|^2)^2(1-|y|^2)^2}{|x-y|^n}.$$

By Lemma 2.1, we have the following result; see [7] and [10].

**Corollary 2.2.** *Let  $\mu$  be a nonnegative measure on  $\mathbf{B}$ . Then  $G_2\mu$  is a biharmonic Green potential if and only if*

$$(2.1) \quad \int_{\mathbf{B}} (1-|y|)^2 d\mu(y) < \infty.$$

**Lemma 2.3.** *If  $(n-1)/n < p < \infty$  and  $1/2 < r < 1$ , then*

$$\mathcal{M}_p(G_2(\cdot, y), r) \leq M(1-r)^{2-n+(n-1)/p} \left( \frac{1-r}{|r-|y||} \right)^{n-(n-1)/p} (1-|y|)^2.$$

*In particular, if  $n = 2, 3$  and  $(n-1)/n < p < \infty$ , then*

$$\mathcal{M}_p(G_2(\cdot, y), r) \leq M(1-r)^{2-n+(n-1)/p} \left( \frac{1-r}{1-r|y|} \right)^{n-(n-1)/p} (1-|y|)^2.$$

This follows from Lemma 2.1 and the fact that, if  $\beta < 1-n$ , then

$$(2.2) \quad \int_{S(r)} |x-y|^\beta dS(x) \leq M(r+|y|)^{1-n} |r-|y||^{\beta+n-1},$$

where  $M$  is a positive constant depending only on  $n$  and  $\beta$ .

Set  $A(r) = \{y \in \mathbf{B} : (5r-1)/4 < |y| < (3r+1)/4\}$  for  $0 < r < 1$ .

**Lemma 2.4.** *Let  $\mu$  be a nonnegative measure on  $\mathbf{B}$  satisfying (2.1). If  $(n-1)/n < p < \infty$ , then*

$$\lim_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p} \int_{\mathbf{B} \setminus A(r)} \mathcal{M}_p(G_2(\cdot, y), r) d\mu(y) = 0.$$

*Proof.* By Lemma 2.3, we obtain

$$\begin{aligned} & (1-r)^{n-2-(n-1)/p} \int_{\mathbf{B} \setminus A(r)} \mathcal{M}_p(G_2(\cdot, y), r) d\mu(y) \\ & \leq M \int_{\mathbf{B} \setminus A(r)} \left( \frac{1-r}{|r-|y||} \right)^{n-(n-1)/p} (1-|y|)^2 d\mu(y). \end{aligned}$$

Since  $(1-r)/|r-|y|| \leq 4$  for  $y \in \mathbf{B} \setminus A(r)$ , Lebesgue's dominated convergence theorem implies that

$$\lim_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p} \int_{\mathbf{B} \setminus A(r)} \mathcal{M}_p(G_2(\cdot, y), r) d\mu(y) = 0.$$

In case  $n = 2$  and  $3$ , we can show by Lemma 2.3 that

$$\lim_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p} \int_{\mathbf{B}} \mathcal{M}_p(G_2(\cdot, y), r) d\mu(y) = 0. \quad \square$$

**Lemma 2.5.** *Let  $1/2 < r < 1$  and  $y \in A(r)$ . If  $n \geq 5$ , then*

$$\mathcal{M}_p(G_2(\cdot, y), r) \leq M(1-r)^{2-n+(n-1)/p}(1-|y|)^2 \\ \times \begin{cases} 1 & \text{if } \frac{n-1}{n} < p < \frac{n-1}{n-4}, \\ \left(\log \frac{1-r}{|r-|y||}\right)^{1/p} & \text{if } p = \frac{n-1}{n-4}, \\ \left(\frac{1-r}{|r-|y||}\right)^{n-4-(n-1)/p} & \text{if } p > \frac{n-1}{n-4}. \end{cases}$$

If  $n = 4$ , then

$$\mathcal{M}_p(G_2(\cdot, y), r) \leq M(1-r)^{2-n+(n-1)/p}(1-|y|)^2.$$

*Proof.* For  $y \in A(r)$  and  $r > 1/2$ , setting

$$I_1(y, r) = S(r) \setminus B\left(y, \frac{1-r}{2}\right) \quad \text{and} \quad I_2(y, r) = S(r) \cap B\left(y, \frac{1-r}{2}\right),$$

we write

$$\mathcal{M}_p(G_2(\cdot, y), r)^p = \frac{1}{\sigma_n r^{n-1}} \left( \int_{I_1(y, r)} G_2(x, y)^p dS(x) + \int_{I_2(y, r)} G_2(x, y)^p dS(x) \right) \\ = u_1(y) + u_2(y).$$

Since  $-np + n - 1 < 0$ , we have

$$u_1(y) \leq M(1-r)^{2p}(1-|y|)^{2p} \int_{I_1(y, r)} |x-y|^{-np} dS(x) \\ \leq M(1-r)^{2p+n-1-np}(1-|y|)^{2p}.$$

On the other hand, if  $x \in I_2(y, r)$ , then  $1-r \leq |y||x-y^*| \leq 3(1-r)$ . In case  $n \geq 5$  we see that

$$G_2(x, y) \leq M(1-r)^{-2}(1-|y|)^2|x-y|^{4-n},$$

so that

$$\begin{aligned}
u_2(y) &\leq M(1-r)^{-2p}(1-|y|)^{2p} \int_{I_2(y,r)} |x-y|^{(4-n)p} dS(x) \\
&\leq M(1-r)^{-2p}(1-|y|)^{2p} |y-r|^{(4-n)p+n-1} \\
&\quad \times \begin{cases} \left(\frac{1-r}{|r-|y||}\right)^{n-1+(4-n)p} & \text{if } \frac{n-1}{n} < p < \frac{n-1}{n-4}, \\ \log \frac{1-r}{|r-|y||} & \text{if } p = \frac{n-1}{n-4}, \\ 1 & \text{if } p > \frac{n-1}{n-4}. \end{cases}
\end{aligned}$$

Since  $(1-r)/|y-r| > 4$  on  $A(r)$ , we obtain the required inequality.

Similarly, in case  $n=4$ , we find

$$\begin{aligned}
u_2(y) &\leq M(1-r)^{(2-n)p}(1-|y|)^{2p} \int_{I_2(y,r)} \left(\log \frac{2(1-r)}{|x-y|}\right)^p dS(x) \\
&\leq M(1-r)^{(2-n)p}(1-|y|)^{2p}(1-r)^{n-1}.
\end{aligned}$$

Hence it follows that

$$\mathcal{M}_p(G_2(\cdot, y), r) \leq M(1-r)^{2-n+(n-1)/p}(1-|y|)^2. \quad \square$$

### 3. Proofs of Theorems 1.1 and 1.2

In this section, we give proofs of Theorems 1.1 and 1.2.

**Proof of Theorem 1.1.** Let  $1 \leq p < (n-1)/(n-4)$  when  $n \geq 5$  and  $1 \leq p < \infty$  when  $n \leq 4$ . By applying Minkowski's inequality for integrals, we have

$$\begin{aligned}
\mathcal{M}_p(G_2\mu, r) &\leq \int_{\mathbf{B}} \mathcal{M}_p(G_2(\cdot, y), r) d\mu(y) \\
(3.1) \quad &= \int_{\mathbf{B} \setminus A(r)} \mathcal{M}_p(G_2(\cdot, y), r) d\mu(y) + \int_{A(r)} \mathcal{M}_p(G_2(\cdot, y), r) d\mu(y).
\end{aligned}$$

Thus Theorem 1.1 follows from Lemmas 2.4 and 2.5. □

**Proof of Theorem 1.2.** First, we give a proof in case  $(n-1)/(n-4) < p < (n-1)/(n-5)$ . Set  $\beta = n-4 - (n-1)/p$  and  $dv(x) = (1-|x|^2) d\mu(x)$ . Here note that  $0 < \beta < 1$ . By Lemmas 2.3, 2.5 and (3.1), we see that

$$(3.2) \quad (1-r)^{n-2-(n-1)/p} \mathcal{M}_p(G_2\mu, r) \leq o(1) + M(1-r)^\beta \int_{A(r)} |y-r|^{-\beta} dv(y).$$

Hence it suffices to show that

$$(3.3) \quad \liminf_{r \rightarrow 1} (1-r)^\beta \int_{A(r)} ||y| - r|^{-\beta} d\nu(y) = 0.$$

For this purpose, we see that

$$\int_{1-2^{-j+1}}^{1-2^{-j}} ||y| - r|^{-\beta} dr \leq M2^{-j(1-\beta)}.$$

Hence it follows that

$$\begin{aligned} & \int_{1-2^{-j+1}}^{1-2^{-j}} \left( (1-r)^\beta \int_{A(r)} ||y| - r|^{-\beta} d\nu(y) \right) \frac{dr}{1-r} \\ & \leq 2^{-j(\beta-1)} \int_{\{y: 2^{-j-1} < 1-|y| < 2^{-j+2}\}} \left( \int_{1-2^{-j+1}}^{1-2^{-j}} ||y| - r|^{-\beta} dr \right) d\nu(y) \\ & \leq M\nu(\{y: 2^{-j-1} < 1-|y| < 2^{-j+2}\}). \end{aligned}$$

Since  $\nu(\mathbf{B}) < \infty$ , we can find a sequence  $\{r_j\}$  such that  $2^{-j} < 1-r_j < 2^{-j+1}$  and

$$\lim_{j \rightarrow \infty} (1-r_j)^\beta \int_{A(r_j)} ||y| - r_j|^{-\beta} d\nu(y) = 0,$$

which implies (3.3). Thus the case  $(n-1)/(n-4) < p < (n-1)/(n-5)$  now follows from (3.2) and (3.3).

Next, we deal with the case  $p = (n-1)/(n-4)$ . We see that

$$(1-r)^{n-2-(n-1)/p} \mathcal{M}_p(G_2\mu, r) \leq o(1) + M \int_{A(r)} \left( \log \frac{1-r}{||y| - r|} \right)^{1/p} d\nu(y).$$

In the same way as above, we have

$$\begin{aligned} & \int_{1-2^{-j+1}}^{1-2^{-j}} \left( \int_{A(r)} \left( \log \frac{1-r}{||y| - r|} \right)^{1/p} d\nu(y) \right) \frac{dr}{1-r} \\ & \leq 2^j \int_{\{y: 2^{-j-1} < 1-|y| < 2^{-j+2}\}} \left( \int_{1-2^{-j+1}}^{1-2^{-j}} \left( \log \frac{1-r}{||y| - r|} \right)^{1/p} dr \right) d\nu(y) \\ & \leq M\nu(\{y: 2^{-j-1} < 1-|y| < 2^{-j+2}\}), \end{aligned}$$

which implies that the left hand-side is zero by letting  $j \rightarrow \infty$ . Thus the theorem is established.  $\square$

REMARK 3.1. Our theorems are best possible as to the power of  $1-r$ .



To show this, for  $p \geq 1$  and  $\delta > 0$ , we give an example of biharmonic Green potential  $v$  satisfying

$$(3.4) \quad \lim_{r \rightarrow 1} (1-r)^{n-2-(n-1)/p-\delta} \mathcal{M}_p(v, r) = \infty.$$

Letting

$$f(x) = (1-|x|)^{-3+\delta/2} |x - e_1|^{1-n},$$

where  $e_1 = (1, 0, \dots, 0)$ , we consider the potential

$$v(x) = \int_{\mathbf{B}} G_2(x, y) f(y) dy.$$

Then  $\int_{\mathbf{B}} (1-|x|)^2 f(x) dx < \infty$ . Further, in case  $n \geq 5$ , we have

$$\begin{aligned} v(x) &\geq \int_{B(x, (1-|x|)/2)} G_2(x, y) f(y) dy \\ &\geq M(1-|x|)^{-3+\delta/2} |x - e_1|^{1-n} \int_{B(x, (1-|x|)/2)} |x - y|^{4-n} dy \\ &= M(1-|x|)^{1+\delta/2} |x - e_1|^{1-n}, \end{aligned}$$

so that

$$\begin{aligned} \mathcal{M}_p(v, r) &\geq M(1-r)^{1+\delta/2} \left( \int_{S(r)} |x - e_1|^{p(1-n)} dS(x) \right)^{1/p} \\ &\geq M(1-r)^{1+\delta/2+1-n+(n-1)/p} \\ &= M(1-r)^{2-n+(n-1)/p+\delta/2}. \end{aligned}$$

Thus (3.4) follows. We can show the remaining case in the same manner.

#### 4. Proof of Theorem 1.3

First we study the spherical means of super-biharmonic functions.

**Lemma 4.1** (cf. [6]). *Let  $u$  be a super-biharmonic function on  $\mathbf{B}$ . Then*

$$u(x) \geq \frac{1}{r_2^2 - r_1^2} \left( r_2^2 \int_{S(x, r_1)} u dS - r_1^2 \int_{S(x, r_2)} u dS \right)$$

whenever  $x \in \mathbf{B}$  and  $0 < r_1 < r_2 < 1 - |x|$ .

*Proof.* Denote by  $K_2$  the fundamental solution for the operator  $\Delta^2$  in  $\mathbf{R}^n$ , that is,

$$K_2(x) = \begin{cases} \alpha_n |x|^{4-n} & \text{in case } n \neq 2, 4, \\ 2\alpha_n |x|^{4-n} \log |x| & \text{in case } n = 2, 4. \end{cases}$$

For  $x \in \mathbf{B}$  and  $0 < r < 1 - |x|$ , we can find a biharmonic function  $h$  in  $B(x, r)$  such that

$$u(y) = \int_{B(x,r)} K_2(z-y) d\mu(z) + h(y)$$

for every  $y \in B(x, r)$ , where  $\mu = \Delta^2 u$ . By the Almansi expansion, we have

$$h(x) = \frac{r_2^2}{r_2^2 - r_1^2} \int_{S(x,r_1)} h(y) dS(y) - \frac{r_1^2}{r_2^2 - r_1^2} \int_{S(x,r_2)} h(y) dS(y)$$

for every  $0 < r_1 < r_2 < r$ . Hence we have only to show that

$$(4.1) \quad K_2(x) \geq \frac{r_2^2}{r_2^2 - r_1^2} \int_{S(r_1)} K_2(x-y) dS(y) - \frac{r_1^2}{r_2^2 - r_1^2} \int_{S(r_2)} K_2(x-y) dS(y)$$

for every  $x \in \mathbf{R}^n$  and  $0 < r_1 < r_2$ . We define

$$g_2(x) = K_2(x) - \frac{r_2^2}{r_2^2 - r_1^2} \int_{S(r_1)} K_2(x-y) dS(y) + \frac{r_1^2}{r_2^2 - r_1^2} \int_{S(r_2)} K_2(x-y) dS(y)$$

and

$$g_1(x) = -\Delta g_2(x).$$

We see that

$$g_1(x) = K_1(x) - \frac{r_2^2}{r_2^2 - r_1^2} \int_{S(r_1)} K_1(x-y) dS(y) + \frac{r_1^2}{r_2^2 - r_1^2} \int_{S(r_2)} K_1(x-y) dS(y)$$

where  $K_1(x) = (n-2)^{-1} \sigma_n^{-1} |x|^{2-n}$  if  $n > 2$  and  $K_1(x) = \sigma_2^{-1} \log(1/|x|)$  if  $n = 2$ . Note that  $g_i(x) = g_i(x')$  for  $|x| = |x'|$  and  $g_i \in C^{2(i-1)}(\mathbf{R}^n \setminus \{0\})$ . Further we see that  $g_i(x) = 0$  for  $|x| \geq r_2$ ,  $g_2(0) = \infty$  if  $n \geq 4$  and  $g_2(0) > 0$  if  $n = 2, 3$ . Setting  $t = K_1(x)$ , we define  $f_i(t) = g_i(x)$ . Then

$$f_1(t) = \begin{cases} 0 & \text{if } K_1(\infty) < t \leq K_1(r_2), \\ -\frac{r_1^2}{r_2^2 - r_1^2} (t - K_1(r_2)) & \text{if } K_1(r_2) < t \leq K_1(r_1), \\ t - t_0 & \text{if } t > K_1(r_1), \end{cases}$$

where  $t_0 = (r_2^2 K_1(r_1) - r_1^2 K_1(r_2)) / (r_2^2 - r_1^2)$  and  $K_1(r) = K_1(x)$  with  $|x| = r$ . Hence we see that  $f_1(t) < 0$  on  $K_1(r_2) < t < t_0$  and  $f_1(t) > 0$  on  $t > t_0$ . Since  $f_2''(t) = -c(t) f_1(t)$  with  $c(t) > 0$  and  $\lim_{t \rightarrow \infty} f_2(t) > 0$ , we obtain  $f_2(t) > 0$  for  $t > K_1(r_2)$ . Thus (4.1) follows.  $\square$

**Lemma 4.2.** *Let  $u$  be a super-biharmonic function on  $\mathbf{B}$ . Then*

$$\lim_{r \rightarrow 1} \mathcal{M}(u, r) \text{ exists in } (-\infty, \infty].$$

*In particular, if  $\liminf_{r \rightarrow 1} \mathcal{M}(u^+, r) = 0$ , then*

$$\mathcal{M}(u, r) \leq M(1 - r^2) \text{ for } r_0 < r < 1.$$

*Proof.* Let  $u$  be a super-biharmonic function on  $\mathbf{B}$  and  $\mu = \Delta^2 u$ . For  $0 < t < 1$ , there exists a biharmonic function  $h_t$  on  $B(t)$  such that

$$u(x) = \int_{\overline{B(t)}} G_2(x, y) d\mu(y) + h_t(x) \quad (x \in B(t)).$$

Set

$$v_t(x) = \begin{cases} h_{(1+t)/2}(x) + \int_{\overline{B((1+t)/2)} \setminus \overline{B(t)}} G_2(x, y) d\mu(y) & \text{if } x \in B((1+t)/2), \\ u(x) - \int_{\overline{B(t)}} G_2(x, y) d\mu(y) & \text{if } x \in \mathbf{B} \setminus \overline{B(t)}. \end{cases}$$

Then  $v_t$  is well defined. Further  $v_t$  is biharmonic on  $B(t)$ , super-biharmonic on  $\mathbf{B}$ ,  $v_t(0) < \infty$  and

$$(4.2) \quad u(x) = v_t(x) + \int_{\overline{B(t)}} G_2(x, y) d\mu(y) \quad (x \in \mathbf{B}).$$

In view of Lemma 2.3, we see that

$$(4.3) \quad \int_{S(r)} \left( \int_{\overline{B(t)}} G_2(x, y) d\mu(y) \right) dS(x) \leq M(1 - r)^2 \frac{\mu(\overline{B(t)})}{r - t}$$

for  $t < r < 1$ ,

where  $M$  is a positive constant independent of  $t$  and  $r$ . By Lemma 4.1, we have

$$\mathcal{M}(v_t, r_2) \geq \frac{r_2^2}{r_1^2} \mathcal{M}(v_t, r_1) - \frac{r_2^2 - r_1^2}{r_1^2} v_t(0)$$

for  $0 < r_1 < r_2 < 1$ , which implies that

$$(4.4) \quad \liminf_{r_2 \rightarrow 1} \mathcal{M}(v_t, r_2) \geq \frac{1}{r_1^2} \mathcal{M}(v_t, r_1) - \frac{1 - r_1^2}{r_1^2} v_t(0) > -\infty$$

for  $0 < r_1 < 1$ . Hence we have

$$\liminf_{r_2 \rightarrow 1} \mathcal{M}(v_t, r_2) \geq \limsup_{r_1 \rightarrow 1} \mathcal{M}(v_t, r_1).$$

In view of (4.4), we see that  $\lim_{r \rightarrow 1} \mathcal{M}(v_t, r)$  exists in  $(-\infty, \infty]$ , and so  $\lim_{r \rightarrow 1} \mathcal{M}(u, r)$  exists in  $(-\infty, \infty]$ .

Moreover, assume that  $\liminf_{r \rightarrow 1} \mathcal{M}(u^+, r) = 0$ . By Lemma 4.1, we have

$$(r_2^2 - r_1^2)v_t(0) \geq r_2^2 \mathcal{M}(v_t, r_1) - r_1^2 \mathcal{M}(v_t, r_2) \geq r_2^2 \mathcal{M}(v_t, r_1) - r_1^2 \mathcal{M}((v_t)^+, r_2)$$

for  $0 < r_1 < r_2 < 1$ . Since  $(v_t)^+ \leq u^+$  and  $\liminf_{r_2 \rightarrow 1} \mathcal{M}(u^+, r_2) = 0$ , we have  $\liminf_{r_2 \rightarrow 1} \mathcal{M}((v_t)^+, r_2) = 0$ . Hence we obtain

$$(4.5) \quad \mathcal{M}(v_t, r_1) \leq v_t(0)(1 - r_1^2).$$

Combining (4.3) and (4.5) we conclude that

$$\mathcal{M}(u, r) \leq M(1 - r)^2 \quad \text{for } t < r < 1,$$

where  $M$  is independent of  $r$ . The claim follows.  $\square$

**REMARK 4.3.** Let  $u$  be a super-biharmonic function on  $\mathbf{B}$  satisfying  $u(0) < \infty$  and  $\liminf_{r \rightarrow 1} \mathcal{M}(u^+, r) = 0$ . Then

$$\mathcal{M}(u, r) \leq u(0)(1 - r^2) \quad \text{for } 0 < r < 1.$$

**Corollary 4.4.** Let  $u$  be a super-biharmonic function on  $\mathbf{B}$  satisfying

$$(4.6) \quad \liminf_{r \rightarrow 1} \mathcal{M}(u^+, r) = 0.$$

Then the following are equivalent.

- (1)  $\liminf_{r \rightarrow 1} (1 - r)^{-1} \mathcal{M}_1(u, r) = 0$  (resp.  $< \infty$ ).
- (2)  $\liminf_{r \rightarrow 1} (1 - r)^{-1} \mathcal{M}(u^-, r) = 0$  (resp.  $< \infty$ ).

**Lemma 4.5.** Let  $u$  be a super-biharmonic function on  $\mathbf{B}$  and  $\mu = \Delta^2 u$ . Suppose  $u$  satisfies

$$\liminf_{r \rightarrow 1} (1 - r)^{-1} \mathcal{M}_1(u, r) < \infty.$$

Then (2.1) is satisfied and  $u$  is of the form

$$u(x) = G_2 \mu(x) + (1 - |x|^2)h(x),$$

where  $h$  is harmonic on  $\mathbf{B}$  satisfying

$$\sup_{r < 1} \mathcal{M}_1(h, r) < \infty.$$

Proof. By considering the function  $v_{1/2}$  in the proof of Lemma 4.2, we may assume that  $u(0) < \infty$ . Take a function  $v_t$  as in the proof of Lemma 4.2. First we show that

$$(4.7) \quad \inf_{0 < t < 1} v_t(0) > -\infty.$$

By Corollary 4.4, we see that  $\liminf_{r \rightarrow 1} \mathcal{M}(u^+, r) = 0$ , and so (4.5) holds. Using (4.3) and (4.5), we establish

$$\begin{aligned} v_t(0) &\geq \frac{1}{1-r^2} \mathcal{M}(v_t, r) \\ &\geq -\frac{1}{1-r^2} \left( \mathcal{M}_1(u, r) + \int_{S(r)} \left( \int_{B(t)} G_2(x, y) d\mu(y) \right) dS(x) \right) \\ &\geq -\frac{1}{1-r^2} \left( \mathcal{M}_1(u, r) + M(1-r)^2 \frac{\mu(\overline{B}(t))}{r-t} \right) \\ &= -\frac{1}{1+r} \left( (1-r)^{-1} \mathcal{M}_1(u, r) + M(1-r) \frac{\mu(\overline{B}(t))}{r-t} \right) \end{aligned}$$

for  $t < r < 1$ . Letting  $r \rightarrow 1$ , we obtain

$$v_t(0) \geq -\frac{1}{2} \liminf_{r \rightarrow 1} (1-r)^{-1} \mathcal{M}_1(u, r),$$

which implies (4.7).

By Lemma 2.1, we have

$$\begin{aligned} C_3 \int_{\overline{B}(t)} (1-|y|^2)^2 d\mu(y) &\leq \int_{\overline{B}(t)} G_2(0, y) d\mu(y) \\ &= u(0) - v_t(0) \\ &\leq u(0) - \inf_{0 < t < 1} v_t(0) < \infty \end{aligned}$$

with the positive constant  $C_3$  in Lemma 2.1. Hence (2.1) follows. In view of Corollary 2.2,  $G_2\mu$  is a biharmonic Green potential, and hence there exists a biharmonic function  $v$  on  $\mathbf{B}$  such that  $u = G_2\mu + v$  on  $\mathbf{B}$ . By Theorem 1.1, we see that

$$(4.8) \quad \liminf_{r \rightarrow 1} (1-r)^{-1} \mathcal{M}_1(v, r) < \infty.$$

We note that  $v$  has an Almansi representation (see [3] and [11]):

$$v(x) = g(x) + (1-|x|^2)h(x)$$

where  $g$  and  $h$  are harmonic on  $\mathbf{B}$ . If  $|x| < r < 1$ , then we apply the Poisson integrals of both sides and find that

$$\liminf_{r \rightarrow 1} (1-r)^{-1} |g(x) + (1-r^2)h(x)| < \infty.$$

This shows that  $g$  is identically zero in  $\mathbf{B}$ . □

The above proof also gives the following.

**Lemma 4.6.** *If  $u$  is a biharmonic function on  $\mathbf{B}$  satisfying*

$$\liminf_{r \rightarrow 1} (1-r)^{-1} \mathcal{M}_1(u, r) = 0,$$

*then  $u \equiv 0$  on  $\mathbf{B}$ .*

Proof of Theorem 1.3. By Lemma 4.5, we see that  $u$  is of the form

$$u(x) = G_2\mu(x) + v(x),$$

where  $v$  is biharmonic on  $\mathbf{B}$  and

$$\int_{\mathbf{B}} (1-|y|)^2 d\mu(y) < \infty.$$

By Theorem 1.1 we find

$$\liminf_{r \rightarrow 1} (1-r)^{-1} \mathcal{M}_1(v, r) = 0.$$

Hence we see from Lemma 4.6 that  $v \equiv 0$ . □

## 5. Remarks

Suppose  $u$  is super-biharmonic on  $\mathbf{B}$ , and set  $\mu = \Delta^2 u \geq 0$ . Further, suppose there exists a sequence  $\{r_j\}$ ,  $0 < r_1 < r_2 < \dots < s < 1$ , tending to 1 such that  $\{u(r_j \cdot) dS\}$  converges weak-star to some finite Borel measure  $\nu$  on  $\mathbf{S}$  and

$$\liminf_{r \rightarrow 1} (1-r)^{-1} \mathcal{M}((u - F[\nu])^-, r) < \infty.$$

Here  $F[\nu](x) = \int_{\mathbf{S}} F(x, y) d\nu(y)$  with  $F(x, y) = D_{\mathbf{n}(y)}(\Delta_y G_2(x, y))$ , where  $D_{\mathbf{n}(y)}$  denotes differentiation with respect to the outward unit normal. Then, as in Abkar and Hedenmalm [1],  $u$  can be represented as

$$u(x) = G_2\mu(x) + F[\nu](x) + (1-|x|^2)P[\lambda](x),$$

where  $P[\lambda]$  denotes the Poisson integral of some finite Borel measure  $\lambda$  on  $\mathbf{S}$  and

$$\int_{\mathbf{B}} (1 - |x|^2)^2 d\mu(x) < \infty.$$

---

### References

- [1] A. Abkar and H. Hedenmalm: *A Riesz representation formula for super-biharmonic functions*, Ann. Acad. Sci. Fenn. Math. **26** (2001), 305–324.
- [2] D.H. Armitage and S.J. Gardiner: *Classical Potential Theory*, Springer-Verlag London, 2001.
- [3] N. Aronszajn, T.M. Creese and L.J. Lipkin: *Polyharmonic functions*, Clarendon Press, 1983.
- [4] T. Futamura and Y. Mizuta: *Isolated singularities of super-polyharmonic functions*, Hokkaido Math. J. **33** (2004), 675–695.
- [5] S.J. Gardiner: *Growth properties of  $p$ -th means of potentials in the unit ball*, Proc. Amer. Math. Soc. **103** (1988), 861–869.
- [6] W. Haußmann and O. Kounchev: *Definiteness of the Peano kernel associated with the polyharmonic mean value property*, J. London Math. Soc. **62** (2000), 149–160.
- [7] W.K. Hayman and P.B. Kennedy: *Subharmonic functions*, Academic Press, London-New York, 1976.
- [8] W.K. Hayman and B. Korenblum: *Representation and uniqueness theorems for polyharmonic functions*, J. Anal. Math. **60** (1993), 113–133.
- [9] Y. Mizuta: *Growth properties of  $p$ -th hyperplane means of Green potentials in a half space*, J. Math. Soc. Japan, **44** (1992), 1–12.
- [10] Y. Mizuta: *Potential theory in Euclidean spaces*, Gakkōtōsyō, Tokyo, 1996.
- [11] M. Nicolesco: *Recherches sur les fonctions polyharmoniques*, Ann. Sci. École Norm. Sup. **52** (1935), 183–220.
- [12] P. Pizetti: *Sulla media deivalori che una funzione dei punti dello spazio assume alla superficie di una sfera*, Rend. Lincei **5** (1909), 309–316.
- [13] M. Stoll: *Rate of growth of  $p$ th means of invariant potentials in the unit ball of  $\mathbf{C}^n$* , J. Math. Anal. Appl. **143** (1989), 480–499.
- [14] M. Stoll: *Rate of growth of  $p$ th means of invariant potentials in the unit ball of  $\mathbf{C}^n$ , II*, J. Math. Anal. Appl. **165** (1992), 374–398.

Toshihide FUTAMURA  
Department of Mathematics  
Graduate School of Science  
Hiroshima University  
Higashi-Hiroshima 739-8526, Japan  
e-mail: toshi@mis.hiroshima-u.ac.jp

Current adress:  
Department of Mathematics  
Daido Institute of Technology  
Nagoya 457-8530, Japan  
e-mail: futamura@daido-it.ac.jp

Yoshihiro Mizuta  
The Division of Mathematical and Information Sciences  
Faculty of Integrated Arts and Sciences  
Hiroshima University  
Higashi-Hiroshima 739-8521, Japan  
e-mail: mizuta@mis.hiroshima-u.ac.jp