



Title	On complex cobordism groups of classifying spaces for dihedral groups
Author(s)	Kamata, Masayoshi
Citation	Osaka Journal of Mathematics. 1974, 11(2), p. 367-378
Version Type	VoR
URL	<a href="https://doi.org/10.18910/12117">https://doi.org/10.18910/12117</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Kamata, M.  
 Osaka J. Math.  
 11 (1974), 367-378

## ON COMPLEX COBORDISM GROUPS OF CLASSIFYING SPACES FOR DIHEDRAL GROUPS

MASAYOSHI KAMATA

(Received December 30, 1973)

### 1. Introduction

Let  $G=H\cdot\Gamma$  be a semi-direct product of a finite group  $H$  by a finite group  $\Gamma$ ,  $X$  a compact  $G$ -manifold which induces by restriction a principal  $H$ -manifold and  $Y$  a principal  $\Gamma$ -manifold. Then we have a principal  $G$ -space  $X\times Y$  with a  $G$ -action defined by  $h\gamma(x, y)=(h\gamma x, \gamma y)$ ,  $h\gamma\in H\cdot\Gamma$ . The equivariant map  $i: X\rightarrow X\times Y$  defined by  $i(x)=(x, y_0)$ , induces a homomorphism

$$i^*: U^*((X\times Y)/G) \rightarrow U^*(X/H).$$

We can define a  $\Gamma$ -action over  $U^*(X/H)$  corresponding to a  $\Gamma$ -action over the complex bordism group of unitary  $G$ -manifolds defined by (1.3) of [7]. The action is denoted by  $x^\gamma$ ,  $x\in U^*(X/H)$ ,  $\gamma\in\Gamma$ .

In this paper, we define a homomorphism

$$i_*: U^*(X/H) \rightarrow U^*((X\times Y)/G)$$

and obtain the following.

**Theorem 1.1.** *For  $x\in U^*(X/H)$ ,  $i^*i_*(x)=\sum_{\gamma\in\Gamma} x^\gamma$ .*

Let  $D_p(m, n)$  be the orbit manifold of  $S^{2m+1}\times S^n$  by the dihedral group  $D_p$  whose action is given in [7]. Making use of Theorem 1.1 and the Atiyah-Hirzebruch spectral sequence of the complex cobordism group, we have the following.

**Theorem 1.2.** *Suppose that  $p$  is an odd prime. There exists an isomorphism*

$$\tilde{U}^{2m}(D_p(2k+1, 4k+3)) \cong \tilde{U}^{2m}(L^{2k+1}(p))^{Z_2} \oplus \tilde{U}^{2m}(RP^{4k+3}) \oplus U^{2m-8k-6},$$

where  $L^i(p)=S^{2i+1}/Z_p$  is a  $(2i+1)$ -dimensional lens space,  $RP^s$  is an  $s$ -dimensional real projective space and  $U^*(\ )^{Z_2}$  is the subgroup consisting of the elements which are fixed under the  $Z_2$ -action.

Let  $BZ_p$  be a classifying space for  $Z_p$ . There exists an isomorphism  $U^{ev}(BZ_p) \cong U^*[[X]]/([p]_F(X))$ ,  $U^{ev}(\ ) = \sum U^{2i}(\ )$  [8]. Consider the  $Z_2$ -action on  $U^{ev}(BZ_p)$  defined by

$$f(X)^t = f([-1]_F(X)) ,$$

where  $t$  is a generator of  $Z_2$ . We use Milnor's short exact sequence [10] and Theorem 1.2 to compute the complex cobordism group of a classifying space for the dihedral group  $D_p$ .

**Theorem 1.3.** *Suppose that  $p$  is an odd prime. There exist isomorphisms*

$$\tilde{U}^{2m}(BD_p) \cong \tilde{U}^{2m}(BZ_p)^{Z_2} \oplus \tilde{U}^{2m}(BZ_2)$$

and

$$\tilde{U}^{2m+1}(BD_p) \cong 0 .$$

Making use of the Conner and Floyd isomorphism

$$\tilde{K}(X) \cong \tilde{U}^{ev}(X) \otimes_{U^*} Z$$

and Theorem 1.2, we can deduce the structure of the  $K$ -group of  $D_p(2k+1, 4k+3)$  which is also obtained in [5] and [6].

## 2. The homomorphism $i^*: U^*(X/H) \rightarrow U^*((X \times Y)/G)$

By a  $G$ -manifold we mean a  $C^\infty$ -manifold which can be embedded equivariantly in some Euclidean  $G$ -space [11]. Let  $M$  and  $X$  be  $G$ -manifolds. By a complex orientation of a  $G$ -map  $f: M \rightarrow X$  we mean an equivalence class of factorizations

$$Z \xrightarrow{i} E \xrightarrow{p} X$$

where  $p: E \rightarrow X$  is a complex  $G$ -vector bundle over  $X$  and where  $i$  is an equivariant  $G$ -embedding endowed with a complex structure compatible with the  $G$ -action on its normal bundle  $\nu_i$ . As Quillen [12] we can define equivariantly a cobordant relation joining such proper complex oriented  $G$ -maps for a  $G$ -manifold  $X$ . We denote by  $U_G^m(X)$  the set of cobordism classes of proper complex oriented  $G$ -maps of dimension  $-m$ . Assume that  $X$  is a principal  $G$ -manifold which is a  $G$ -manifold such that no element of the group other than the identity has a fixed point [2]. Then the complex cobordism group  $U_G^m(X)$  is isomorphic to  $U^m(X/G)$  by sending the equivariant cobordism class  $[Z \xrightarrow{i} E \xrightarrow{p} X]_G$  to  $[Z/G \xrightarrow{i'} E/G \xrightarrow{p'} X/G]$ , where  $i'$  and  $p'$  are quotient maps.

From now on, we suppose that  $G$  is a semi-direct product  $H \cdot \Gamma$  of a finite group  $H$  by a finite group  $\Gamma$  and that  $X$  is a  $G$ -manifold whose action restricted to  $H$  is free and  $Y$  is a principal  $\Gamma$ -manifold. The element  $\gamma$  of  $\Gamma$  acts on the group  $H$  by the inner automorphisms  $h^\gamma = \gamma^{-1} h \gamma$  and the group operation of  $H \cdot \Gamma$  is given by

$$(h_1 \gamma_1)(h_2 \gamma_2) = h_1 h_2^{\gamma_1^{-1}} \gamma_1 \gamma_2.$$

The map  $i: X \rightarrow X \times Y$ ,  $i(x) = (x, y_0)$ , is an equivariant map. Then, there exists a composition homomorphism

$$i^*: U^*((X \times Y)/G) \xrightarrow{r^*} U^*((X \times Y)/H) \xrightarrow{i_H^*} U^*(X/H)$$

where  $r^*$  sends an equivariant cobordism class  $[Z \rightarrow E \rightarrow X]_G$  to the class  $[Z \rightarrow E \rightarrow X]_H$  obtained by restriction of the group action and  $i_H^*$  is the quotient map of  $i$ . Suppose that  $X$  is a compact principal  $G$ -manifold,  $G = H \cdot \Gamma$ . Let  $[Z \xrightarrow{i} E \xrightarrow{p} X]_H$  be an element of  $U_H^m(X)$  represented by an  $H$ -equivariant factorization. Since  $q: X \rightarrow X/H$  is a principal bundle, a functor  $q^*$  from the category of vector bundles and homomorphisms over  $X/H$  to the category of  $H$ -vector bundles and  $H$ -homomorphisms over  $X$  is an equivalence [1]. There exists an  $H$ -complex vector bundle  $F$  over  $X$  such that  $E \oplus F = X \times C^n$  where  $H$  acts on  $X \times C^n$  by the rule  $h(x, z) = (hx, z)$ . Therefore,

$$[Z \xrightarrow{i} E \xrightarrow{p} X]_H = [Z \xrightarrow{i} X \times C^n \xrightarrow{\tilde{p}} X]_H$$

as equivariant cobordism classes, where  $i(z) = (i(z), 0)$  and  $\tilde{p}(x, z) = x$ . We form the quotient space  $G \times_H Z$ . The group  $G$  acts on  $G \times_H Z$  by  $\hat{g}(g \times_H z) = (\hat{g}g \times_H z)$ . We have then the equivariant embedding

$$\begin{aligned} i_1: G \times_H Z \times Y &\rightarrow X \times C^n \times Y \times V \\ i_1(h\gamma \times_H z, y) &= (h\gamma i(z), y, e(\gamma)) \end{aligned}$$

where  $G \times_H Z \times Y$  is a  $G$ -space by  $h\gamma(g \times_H z, y) = (h\gamma g \times_H z, \gamma y)$ ,  $V$  is a complex Euclidean  $\Gamma$ -space, for example a regular representation space of  $\Gamma$ ,  $X \times C^n \times Y \times V$  is a  $G$ -space by  $h\gamma(x, z, y, v) = (h\gamma x, z, \gamma y, \gamma v)$  and  $e: \Gamma \rightarrow V$  is a  $\Gamma$ -equivariant embedding.

**Lemma 2.1.** *If the normal bundle  $\nu$  of  $i: Z \rightarrow X \times C^n$  has a complex structure compatible with the  $H$ -action, then the normal bundle  $\nu_1$  of  $i_1: G \times_H Z \times Y \rightarrow X \times C^n \times Y \times V$  has a complex structure compatible with the  $G$ -action.*

Proof. Let  $J: \nu \rightarrow \nu$  be a complex structure compatible with  $H$ -action, that is,  $hJ = Jh$ . We may consider that  $X$  and  $Y$  are embedded in a Euclidean  $G$ -space  $V_x$  and a Euclidean  $\Gamma$ -space  $V_y$ , respectively and that each element of  $G$  operates on  $V_x \times C^n \times V_y \times V$  as an orthogonal linear transformation. The total space of the normal bundle  $\nu_1$  is described as follows:

$$E(\nu_1) = \{(i_1(h\gamma \times_H z, y), (h\gamma w, v)): w \text{ is a vector of a fiber of } \nu \text{ over } i(z) \text{ and } v \in V\}.$$

We put

$$\tilde{J}(i_1(h\gamma \times_H z, y), (w, v)) = (i_1(h\gamma \times_H z, y), (\gamma J \gamma^{-1} w, \sqrt{-1} v)).$$

The homomorphism  $\tilde{J}$  is a complex structure of the bundle  $\nu_1$  q.e.d.

From Lemma 2.1, we have a factorization

$$G \times_H Z \times Y \xrightarrow{i_1} X \times C^* \times Y \times V \xrightarrow{p_1} X \times Y,$$

$p_1(x, z, y, v) = (x, y)$ , which is a complex orientation of a map  $p_1 \circ i_1$ . We set

$$i_*[Z \xrightarrow{i} E \xrightarrow{p} X]_H = [G \times_H Z \times Y \xrightarrow{i_1} X \times C^* \times Y \times V \xrightarrow{p_1} X \times Y]_G.$$

This defines a  $U^*$ -module homomorphism

$$i_*: U^*(X/H) \rightarrow U^*((X \times Y)/G)$$

of degree 0.

We define a  $\Gamma$ -action on  $U^*(X/H)$ : We take an equivariant cobordism class  $[Z \xrightarrow{i} X \times C^* \xrightarrow{p} X]_H \in U_H^*(X) = U^*(X/H)$ , with an  $H$ -action  $\phi: H \times Z \rightarrow Z$ . Let  $Z^\gamma$  be a copy of  $Z$  whose action  $\phi^\gamma: H \times Z^\gamma \rightarrow Z^\gamma$  is given by

$$\phi^\gamma(h, z) = \phi(h^\gamma, z)$$

and  $i^\gamma: Z^\gamma \rightarrow X \times C^*$  be an equivariant  $H$ -map given by

$$i^\gamma(z) = \gamma i(z).$$

Denote by  $\nu$  the normal bundle of  $i: Z \rightarrow X \times C^*$  and  $\nu_x$  the fiber over  $x$ . The total space  $E$  of the normal bundle  $\nu^\gamma$  of  $i^\gamma: Z^\gamma \rightarrow X \times C^*$  is

$$E = \{(i^\gamma(z), \gamma v) : v \text{ is a vector in the fiber } \nu_{i(z)}\}.$$

Let  $J: \nu \rightarrow \nu$  be a complex structure compatible with the  $H$ -action. Then, a bundle map  $J^\gamma: E \rightarrow E$ ,  $J^\gamma(i^\gamma(z), w) = (i^\gamma(z), \gamma J \gamma^{-1} w)$ , is a complex structure of  $\nu^\gamma$  compatible with the  $H$ -action. We set

$$[Z \xrightarrow{i} X \times C^* \xrightarrow{p} X]_H^\gamma = [Z^\gamma \xrightarrow{i^\gamma} X \times C^* \xrightarrow{p} X]_H.$$

Proof of Theorem 1.1.

We recall that  $i_*[Z \xrightarrow{i} X \times C^* \xrightarrow{\tilde{p}} X]_H = [G \times_H Z \times Y \xrightarrow{i_1} X \times C^* \times Y \times V \xrightarrow{\tilde{p}_1} X \times Y]_G$ . Consider the map  $j: X \times C^* \times V \rightarrow X \times C^* \times Y \times V$ ,  $j(x, z, v) = (x, z, y_0, v)$ . The map  $j$  is an  $H$ -map and transversally regular on  $i_1(G \times_H Z \times Y)$ . Let  $\Gamma$  be the set consisting of  $\gamma_1, \gamma_2, \dots, \gamma_m$ . It follows that

$$j^{-1}(i_1(G \times_H Z \times Y)) = \bigcup_k Z_k$$

where  $Z_k = \{(h\gamma_k i(z), e(\gamma_k)) : h \in H, z \in Z\} \subset X \times C^n \times V$ . Clearly,  $Z_k$  is equivariantly diffeomorphic to  $Z^{\gamma_k}$  and  $[Z_k \xrightarrow{i_k} X \times C^n \times V \xrightarrow{\tilde{p}} X]_H = [Z \xrightarrow{i} X \times C^n \xrightarrow{\tilde{p}} X]_H^{\gamma_k}$ , where  $i_k$  is an inclusion. Therefore, we have  $i^* i_* [Z \xrightarrow{i} X \times C^n \xrightarrow{\tilde{p}} X]_H = \Sigma [Z \xrightarrow{i} X \times C^n \xrightarrow{\tilde{p}} X]_H^{\gamma_k}$ . q.e.d.

### 3. The structure of $\tilde{U}^{2m}(D_p(2k+1, 4k+3))$

In [7], the manifold  $D_p(l, n) = (S^{2l+1} \times S^n)/D_p$  was useful to determine the structure of complex bordism group of principal dihedral group  $D_p$ -actions. In this section, we determine the additive structure of  $\tilde{U}^{2m}(D_p(2k+1, 4k+3))$ . Consider an action of the dihedral group  $D_p = Z_p \cdot Z_2$  over  $S^{2l+1} \times S^n$  given by

$$(1) \quad (g^i t^j)(z, x) = (\rho^i c^j(z), (-1)^j x), \quad \rho = \exp 2\pi\sqrt{-1}/p$$

where  $g$  is a generator of order  $p$  and  $t$  is the generator of order 2 and  $c(z)$  is the conjugation operator. The manifold  $D_p(l, n)$  is the orbit space. This manifold is an example of manifolds described in §2. We take a  $Z_p$ -space  $S^{2l+1}$  with  $g \cdot z = \rho z$  ( $z \in S^{2l+1}$ ,  $g$  is a generator of  $Z_p$ ), a  $Z_2$ -space  $S^n$  with  $t \cdot x = (-1)x$  ( $x \in S^n$ ,  $t$  is the generator of  $Z_2$ ) and a  $D_p$ -space  $S^{2l+1} \times S^n$  with the  $D_p$ -action given by (1). Then, there are equivariant maps

$$\begin{aligned} i: S^{2l+1} &\rightarrow S^{2l+1} \times S^n & i(z) &= (z, (1, 0, \dots, 0)) \\ j: S^n &\rightarrow S^{2l+1} \times S^n & j(x) &= ((1, 0, \dots, 0), x) \end{aligned}$$

and

$$p: S^{2l+1} \times S^n \rightarrow S^n \quad p(z, x) = x$$

with respect to inclusions  $i: Z_p \rightarrow D_p$ ,  $j: Z_2 \rightarrow D_p$  and a projection  $p: D_p \rightarrow Z_2$  respectively. Denote by  $U^*(S^{2l+1}/Z_p)^{Z_2}$  the subgroup consisting of elements fixed under the  $Z_2$ -action over  $U^*(S^{2l+1}/Z_p)$  described in §2. Then we have the following.

**Proposition 3.1.** *If  $p$  is an odd prime, the homomorphism  $\Phi: \tilde{U}^{2m}(S^{2l+1}/Z_p)^{Z_2} \oplus \tilde{U}^{2m}(S^n/Z_2) \rightarrow \tilde{U}^{2m}(D_p(l, n))$  given by  $\Phi(x, y) = i_*(x) + p^*(y)$  is injective.*

**Proof.** We remark that  $\tilde{U}^{2m}(S^{2l+1}/Z_p)$  is a  $p$ -group and  $\tilde{U}^{2m}(S^n/Z_2)$  is a 2-group. Hence,  $i^* p^* = 0$ . Since  $j^* p^* = 1$  and from Theorem 1.1  $i^* i_*(x) = 2x$ ,  $\Phi$  is injective. q.e.d.

Denote by  $L'(p)$  a  $(2l+1)$ -dimensional lens space. The manifold  $D_p(l, n)$  is homeomorphic to the orbit space of  $L'(p) \times S^n$  by a  $Z_2$ -action  $t([z], x) = ([cz], -x)$ ,  $t \in Z_2$  the generator. Let  $C_i$  and  $D_j$  be the standard cells of  $L'(p)$  and  $S^n$  respectively. The images  $(C_i, D_j)$  of the  $C_i \times D_j$  by the quotient map  $L'(p) \times S^n \rightarrow D_p(l, n)$  give a cellular decomposition of  $D_p(l, n)$ . Denote by  $(c^i, d^j)$  the dual

cochain element to  $(C_i, D_j)$ . Then we have the following coboundary relations

$$\begin{aligned}\delta(c^{2i+1}, d^j) &= \{(-1)^i + (-1)^j\}(c^{2i+1}, d^{j+1}) + p(c^{2i+2}, d^j) \\ \delta(c^{2i}, d^j) &= \{(-1)^i + (-1)^{j+1}\}(c^{2i}, d^{j+1}).\end{aligned}$$

Therefore, we have the following.

**Proposition 3.2.** *The integral cohomology group  $\tilde{H}^*(D_p(l, n); \mathbb{Z})$  is a direct sum of the following groups*

- (i) *case l: even and n: even*  
a free group generated by  $(c^{2l+1}, d^n)$ , torsion groups generated by the  $(c^0, d^{2j})$  and the  $(c^{2l+1}, d^{2j-1})$  whose orders are 2 and torsion groups generated by the  $(c^{4i}, d^0)$  and the  $(c^{4i-2}, d^n)$  whose orders are  $p$ ,
- (ii) *case l: even and n: odd*  
a free group generated by  $(c^0, d^n)$ , torsion groups generated by the  $(c^0, d^{2j})$  and the  $(c^{2l+1}, d^{2j+1})$  whose orders are 2 and torsion groups generated by the  $(c^{4i}, d^0)$  and the  $(c^{4i}, d^n)$  whose orders are  $p$ ,
- (iii) *case l: odd and n: even*  
a free group generated by  $(c^{2l+1}, d^0)$ , torsion groups generated by the  $(c^0, d^{2j})$  and the  $(c^{2l+1}, d^{2j})$  whose orders are 2 and torsion groups generated by the  $(c^{4i}, d^0)$  and the  $(c^{4i-2}, d^n)$  whose orders are  $p$ ,
- (iv) *case l: odd and n: odd*  
free groups generated by  $(c^0, d^n)$ ,  $(c^{2l+1}, d^0)$  and  $(c^{2l+1}, d^n)$ , torsion groups generated by the  $(c^0, d^{2j})$  and the  $(c^{2l+1}, d^{2j})$  whose orders are 2 and torsion groups generated by the  $(c^{4i}, d^0)$  and the  $(c^{4i}, d^n)$  whose orders are  $p$ ,

where  $0 \leq 2j \leq n$  and  $0 \leq 2i \leq l$ .

Let  $Y_k$  be the  $(8k+5)$ -skeleton of  $D_p(2k+1, 4k+3)$ . Denote by  $(E_r^{s,t}(X), d_r^{s,t})$  the Atiyah-Hirzebruch spectral sequence for  $U^*(X)$ .

**Lemma 3.3.** *If  $s \neq 8k+6$  then an inclusion  $\iota: Y_k \rightarrow D_p(2k+1, 4k+3)$  induces the isomorphism for any  $r$*

$$E_r^{s,t}(Y_k) \cong E_r^{s,t}(D_p(2k+1, 4k+3)).$$

**Proof.** Using Proposition 3.2, it follows that  $\iota^*: E_2^{s,t}(D_p(2k+1, 4k+3)) \rightarrow E_2^{s,t}(Y_k)$  is isomorphic if  $s \neq 8k+6$ . We note that the images of the differentials  $d_r^{s,t}$  for any  $r$  are torsion groups [4]. By induction on  $r$  we have the lemma. q.e.d.

**Proposition 3.4.** *There exists a short exact sequence*

$$0 \rightarrow U^{2m-8k-6} \rightarrow \tilde{U}^{2m}(D_p(2k+1, 4k+3)) \rightarrow \tilde{U}^{2m}(Y_k) \rightarrow 0.$$

**Proof.** Consider the exact sequence of complex cobordism groups for a pair  $(D_p(2k+1, 4k+3), Y_k)$ :

$$\dots \rightarrow \tilde{U}^*(D_p(2k+1, 4k+3)) \rightarrow \tilde{U}^*(Y_k) \rightarrow \tilde{U}^{*+1}(D_p(2k+1, 4k+3)/Y_k) \rightarrow$$

From Lemma 3.3  $\iota^* : \tilde{U}^i(D_p(2k+1, 4k+3)) \rightarrow \tilde{U}^i(Y_k)$  is isomorphic for  $i$  odd. Since  $\tilde{H}^i(D_p(2k+1, 4k+3)/Y_k; Z) = 0$  if  $i \neq 8k+6$  and  $\tilde{H}^{8k+6}(D_p(2k+1, 4k+3)/Y_k; Z) \cong Z$ , we have that  $\tilde{U}^{2m}(D_p(2k+1, 4k+3)/Y_k) \cong U^{2m-8k-6}$ . q.e.d.

We investigate the Thom homomorphism  $\mu : U^*(X) \rightarrow H^*(X)$  which is the edge homomorphism of the spectral sequence associated with  $U^*(X)$ . Let  $X$  be an orientable manifold. We take an element  $[M \xrightarrow{i} X \xrightarrow{id} X] \in U^*(X)$  which is represented by an inclusion map  $M \xrightarrow{i} X$  with the normal bundle  $\nu$  equipped with a complex structure. Denote by  $N(\nu)$  the tubular neighborhood of  $M$ , and we have a canonical map  $j : (X, \phi) \rightarrow (X, \{\text{Int } N(\nu)\}^c)$ . Then, we can describe the Thom homomorphism as  $\mu[M \xrightarrow{i} X \xrightarrow{id} X] = j^* \tau(\nu)$ ,  $\tau(\nu)$  is the Thom class of  $\nu$ , and

$$(2) \quad \mu[M \xrightarrow{i} X \xrightarrow{id} X] = Di_* \sigma(M)$$

where  $D$  is the Poincaré duality isomorphism  $H_*(M) \cong H^*(M)$  and  $\sigma(M)$  is a fundamental class of  $M$ .

We put

$$L_{k-m} = [S^{4m+3} \xrightarrow{i} S^{4k+3} \xrightarrow{id} S^{4k+3}]_{Z_p} \in U_{Z_p}^{4(k-m)}(S^{4k+3}),$$

where  $S^{4k+3}$  and  $S^{4m+3}$  are  $Z_p$ -spaces with canonical action  $g \cdot z = \rho z$  and  $i$  is the canonical inclusion, and

$$R_{2k+1-n} = [S^{2n+1} \xrightarrow{i} S^{4k+3} \xrightarrow{id} S^{4k+3}]_{Z_2} \in U_{Z_2}^{4k+2-2m}(S^{4k+3})$$

where  $S^{2n+1}$  and  $S^{4k+3}$  are  $Z_2$ -spaces with the canonical action  $t \cdot x = (-1)x$ , and  $i$  is the canonical inclusion.

**Proposition 3.5.** *Suppose that  $p$  is an odd prime, then*

$$\mu i_*(L_{k-m} + L_{k-m}^t) = a(c^{4(k-m)}, d^0), \quad a \not\equiv 0 \pmod{p}$$

and

$$\mu p^*(R_{2k+1-n}) = (c^0, d^{4k+2-2m}).$$

**Proof.** The manifold  $D_p(2k+1, 4k+3)$  is orientable. Using Theorem 1.1 and (2), we have the proposition. q.e.d.

**Proof of Theorem 1.2.**

Proposition 3.5 shows that in the Atiyah-Hirzebruch spectral sequence for  $\tilde{U}^*(D_p(2k+1, 4k+3))$ , the  $(c^{4i}, d^0)$  and the  $(c^0, d^{2j})$  are permanent cycles. It is

easy to prove that the spectral sequence is trivial. Therefore it follows from Propositions 3.1 and 3.5 that there exists an isomorphism

$$\begin{aligned} \lambda^* + i_* + p^* : \tilde{U}^{2m}(D_p(2k+1, 4k+3)/Y_k) &\oplus \tilde{U}^{2m}(S^{4k+3}/Z_p)^{Z_2} \oplus \tilde{U}^{2m}(S^{4k+3}/Z_2) \\ &\rightarrow \tilde{U}^{2m}(D_p(2k+1, 4k+3)) \end{aligned}$$

where  $\lambda : D_p(2k+1, 4k+3) \rightarrow D_p(2k+1, 4k+3)/Y_k$  is the projection map. q.e.d.

#### 4. $\tilde{U}^*(BZ_p)$ , $p$ an odd prime

The complex cobordism group  $\tilde{U}^{ev}(L^n(p)) \cong \tilde{U}^{ev}(S^{2n+1}/Z_p)$  is a  $U^*$ -module with a generating set  $\{[S^{2k+1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_p}; Z_p\text{-equivariant cobordism classes which are represented by the canonical equivariant inclusion map } i(z_0, \dots, z_k) = (z_0, \dots, z_k, 0, \dots, 0), 0 \leq k \leq n-1\}$ .

$$\begin{aligned} \text{Lemma 4.1. } &\{i_n^*([S^{2k+1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_p})\}^t \\ &= i_n^*([S^{2k+1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_p}^t), \end{aligned}$$

where  $i_n : L^{n-1}(p) \rightarrow L^n(p)$  is the inclusion map  $i_n(z_0, \dots, z_{n-1}) = (z_0, \dots, z_{n-1}, 0)$ .

Proof. By the definition of the  $Z_2$ -action,  $[S^{2k+1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_p}^t = [(S^{2k+1})^t \xrightarrow{i^t} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_p}$  with  $i^t(z) = ci(z)$ . Let  $H_n : S^{2n-1} \times I \rightarrow S^{2n+1}$  be a map defined by

$$H_n(z_0, \dots, z_{n-1}, t) = \frac{1}{A}(tz_0, tz_1 + (1-t)z_0, \dots, tz_{n-1} + (1-t)z_{n-2}, (1-t)z_{n-1})$$

where  $A$  is the norm of  $(tz_0, tz_1 + (1-t)z_0, \dots, (1-t)z_{n-1})$ .  $H_n$  is an equivariant  $Z_p$ -map. Put

$$j_n(z) = H_n(z, 0),$$

then we have that  $j_n^* = i_n^*$ . Moreover  $j_n : S^{2n-1} \rightarrow S^{2n+1}$  is transverse regular on  $i^t(S^{2k+1})$ . Therefore, we have

$$j_n^*[(S^{2k+1})^t \xrightarrow{i^t} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_p} = [(S^{2k-1})^t \xrightarrow{i^t} S^{2n-1} \xrightarrow{id} S^{2n-1}]_{Z_p}.$$

q.e.d.

Let  $F(X, Y)$  be the formal group of the complex cobordism theory. Denote by  $[-1]_F(X)$  the element of  $U^*[[X]]$  satisfying  $F(X, [-1]_F(X)) = 0$  and by  $[k]_F(X)$  the element of  $U^*[[X]]$  defined by the following formulae

$$\begin{cases} [1]_F(X) = X \\ F(X, [k]_F(X)) = [k+1]_F(X). \end{cases}$$

We define a  $Z_2$ -action on  $U^*[[X]]$  by

$$f(X)^t = f([-1]_F(X)).$$

By the definition of the formal group law, it follows immediately that  $\{[p]_F(X)\}^t$  and  $(X^{n+1})^t$  belong to the ideal  $([p]_F(X), X^{n+1})$  generated by  $[p]_F(X)$  and  $X^{n+1}$  in  $U^*[[X]]$  and thus  $Z_2$  acts on  $U^*[[X]]/([p]_F(X), X^{n+1})$ . We can see that the element  $[S^{2n-1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_2}$  corresponds to the cobordism 1-st Chern class  $c_1(\xi_n)$  of the canonical line bundle  $\xi_n$  over  $L^n(p)$  and that  $[S^{2n-1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_2}^t$  is the cobordism 1-st Chern class  $c_1(\bar{\xi}_n)$  of the conjugate bundle  $\bar{\xi}_n$ . Therefore, we have the following.

**Lemma 4.2.**  $U^{ev}(L^n(p))^{Z_2} \cong \{U^*[[X]]/([p]_F(X), X^{n+1})\}^{Z_2}$ .

Proof. From the definition of the multiplication in  $U^{ev}(L^n(p))$  we have that for  $0 \leq k, l \leq n$

$$\begin{aligned} & [S^{2k+1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_2} [S^{2l+1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_2} \\ &= \begin{cases} [S^{2(n-k+l)+1} \xrightarrow{i} S^{2n+1} \xrightarrow{id} S^{2n+1}]_{Z_2} & \\ 0 & \text{if } n-k-l > 0. \end{cases} \end{aligned}$$

Then, it follows immediately that the  $Z_2$ -action on  $U^*(L^n(p))$  is multiplicative. There exists an isomorphism  $U^{ev}(L^n(p)) \cong U^*[[X]]/([p]_F(X), X^{n+1})$  which maps  $c_1(\xi_n)$  to  $X$  [13]. Since  $F(c_1(\xi_n), c_1(\bar{\xi}_n)) = c_1(\xi_n \otimes \bar{\xi}_n) = 0$ , the lemma follows. q.e.d.

Denote by  $j_k: D_p(2k-1, 4k-1) \rightarrow D_p(2k+1, 4k+3)$  and  $\hat{j}_k: L^{2k-1}(p) \rightarrow L^{2k+1}(p)$  respectively, the maps induced by the inclusions  $S^{4k-1} \times S^{4k-1} \subset S^{4k+3} \times S^{4k+3}$  and  $S^{4k-1} \subset S^{4k+3}$ . The following diagram is commutative

$$\begin{array}{ccc} \tilde{U}^{2m}(L^{2k+1}(p)) & \xrightarrow{i_*} & \tilde{U}^{2m}(D_p(2k+1, 4k+3)) \\ \downarrow \hat{j}_k^* & & \downarrow j_k^* \\ \tilde{U}^{2m}(L^{2k-1}(p)) & \xrightarrow{i_*} & \tilde{U}^{2m}(D_p(2k-1, 4k-1)). \end{array}$$

Since the  $Z_2$ -action on  $U^*(L^n(p))$  and  $\hat{j}_k^*$  are  $U^*$ -homomorphisms, it follows from Lemma 4.1 that  $i_*$  induces a homomorphism of inverse systems

$$i_*: \{\tilde{U}^{2m}(L^{2k+1}(p))^{Z_2}, \hat{j}_k^*\} \rightarrow \{\tilde{U}^{2m}(D_p(2k+1, 4k+3)), j_k^*\}.$$

Consider the quotient map of  $j_k$

$$\tilde{j}_k: D_p(2k-1, 4k-1)/Y_{k-1} \rightarrow D_p(2k+1, 4k+3)/Y_k,$$

where  $Y_k$  is a  $(8k+5)$ -skeleton of  $D_p(2k+1, 4k+3)$ . Maps  $\lambda: D_p(2k+1, 4k+3)$

$\rightarrow D_p(2k+1, 4k+3)/Y_k$  and  $p$ ;  $D_p(2k+1, 4k+3) \rightarrow RP^{4k+3}$  induce homomorphisms of inverse systems

$$\begin{aligned} \lambda^*: \{ \tilde{U}^{2m}(D_p(2k+1, 4k+3)/Y_k) &\cong U^{2m-8k-6}, \tilde{j}_k^* \} \\ &\rightarrow \{ \tilde{U}^{2m}(D_p(2k+1, 4k+3)), j_k^* \} \end{aligned}$$

and

$$p^*: \{ \tilde{U}^{2m}(RP^{4k+3}), \hat{j}_k^* \} \rightarrow \{ \tilde{U}^{2m}(D_p(2k+1, 4k+3)), j_k^* \},$$

where  $\hat{j}_k: RP^{4k-1} \rightarrow RP^{4k+3}$  is the inclusion map. From Theorem 1.2, we have an isomorphism

$$(4.1) \quad \begin{aligned} i_* + p^*: \lim_{\leftarrow} \tilde{U}^{2m}(L^{2k+1}(p))^{Z_2} \oplus \lim_{\leftarrow} \tilde{U}^{2m}(RP^{4k+3}) \\ \rightarrow \lim_{\leftarrow} \tilde{U}^{2m}(D_p(2k+1, 4k+3)), \end{aligned}$$

because  $\tilde{j}_k^*: \tilde{U}^{2m}(D_p(2k+1, 4k+3)/Y_k) \rightarrow \tilde{U}^{2m}(D_p(2k-1, 4k-1)/Y_{k-1})$  is a zero homomorphism.

**Lemma 4.3.**  $j_k^*: \tilde{U}^{2m+1}(D_p(2k+1, 4k+3)) \rightarrow \tilde{U}^{2m+1}(D_p(2k-1, 4k-1))$  is a zero homomorphism.

Proof. Let  $\tilde{Y}_k$  be a  $(8k+2)$ -skeleton of  $D_p(2k, 4k+2)$ . We consider the map  $j_k$  as a composition map  $j_k: D_p(2k-1, 4k-1) \rightarrow \tilde{Y}_k \rightarrow D_p(2k, 4k+2) \rightarrow D_p(2k+1, 4k+3)$ . By Proposition 3.2 case (i), it follows that  $\tilde{H}^{odd}(Y_k; Z) \cong 0$  and  $\tilde{U}^{2m+1}(\tilde{Y}_k) \cong 0$ . Therefore,  $j_k^*$  is the zero homomorphism. q.e.d.

**Lemma 4.4.**  $\lim_{\leftarrow}^1 \tilde{U}^{2m}(D_p(2k+1, 4k+3)) = 0$ .

Proof. From Proposition 3.5 and Theorem 1.2 it follows that  $\{L_{k-m} + L_{k-m}^t\}$  is a generating set for  $U^*$ -module  $\tilde{U}^{ev}(S^{4k+3}/Z_p)^{Z_2}$ . By Lemma 4.1,  $\hat{j}_k^*: \tilde{U}^{2m}(L^{4k+3}(p))^{Z_2} \rightarrow \tilde{U}^{2m}(L^{4k-1}(p))^{Z_2}$  is surjective. Therefore, it follows that an inverse system  $\{ \tilde{U}^{2m}(D_p(2k+1, 4k+3)), j_k^* \}$  satisfies the Mittag-Leffler condition and the lemma follows. q.e.d.

Proof of Theorem 1.3.

There exists Milnor's short exact sequence

$$(4.2) \quad \begin{aligned} 0 \rightarrow \lim_{\leftarrow}^1 \tilde{U}^{*-1}(D_p(2k+1, 4k+3)) &\rightarrow \tilde{U}^*(BD_p) \\ &\rightarrow \lim_{\leftarrow} \tilde{U}^*(D_p(2k+1, 4k+3)) \rightarrow 0 [10]. \end{aligned}$$

Using Lemma 4.3 and 4.4, we have  $\tilde{U}^{2m+1}(BD_p) = 0$ .

Lemma 4.3 implies that the inverse system  $\{ \tilde{U}^{2m+1}(D_p(2k+1, 4k+3)), j_k^* \}$  satisfies the Mittag-Leffler condition. Therefore we have that

$$\tilde{U}^{2m}(BD_p) \cong \lim_{\leftarrow} \tilde{U}^{2m}(D_p(2k+1, 4k+3)).$$

Using Theorem 1.2 and Lemma 4.2 we complete the proof.

### 5. The structure of $\tilde{K}(D_p(2k+1, 4k+3))$

In [3], Conner and Floyd gave the isomorphism

$$(5.1) \quad c: \tilde{K}(X) \cong \tilde{U}^{ev}(X) \otimes_{U^*} Z,$$

which maps  $\eta_n - n$  to  $c_1(\eta_n) \times 1$ . Consider a  $Z_2$ -action on  $K(L^n(p))$  defined by  $\eta^t = \bar{\eta}$ ,  $t$  a generator of  $Z_2$ . Since  $Z_2$ -action on  $U^*(L^n(p))$  is multiplicative, we have the commutative diagram

$$(5.2) \quad \begin{array}{ccc} \tilde{K}(L^n(p)) & \xrightarrow{c} & \tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z \\ \downarrow t & & \downarrow t \otimes_{U^*} id \\ \tilde{K}(L^n(p)) & \xrightarrow{c} & \tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z \end{array}$$

**Lemma 5.1.**  $\tilde{U}^{ev}((L^n(p)) \otimes_{U^*} Z)^{Z_2} = \tilde{U}^{ev}(L^n(p))^{Z_2} \otimes_{U^*} Z$ , where  $(\tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z)^{Z_2}$  is an invariant subgroup of  $\tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z$  under the  $Z_2$ -action  $\cdot^t \times_{U^*} id$ .

Proof. By the definition of  $Z_2$ -action of  $\tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z$ , it follows that  $\tilde{U}^{ev}(L^n(p))^{Z_2} \otimes_{U^*} Z \subset (\tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z)^{Z_2}$ . Suppose that  $x \otimes_{U^*} m \in \tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z$  and  $x^t \otimes_{U^*} m = x \otimes_{U^*} m$ . Since  $c$  is isomorphic, there exists an element  $\eta \in \tilde{K}(L^n(p))$  with  $c(\eta) = x \otimes_{U^*} m$ . By the commutative diagram (5.2),

$$c(\eta) = c(\eta)^t = c(\eta^t) \quad \text{and} \quad \eta = \eta^t.$$

N. Mahammed [9] proved that  $\tilde{K}(L^n(p)) = Z[\xi_n]/(\xi_n^n - 1, (\xi_n - 1)^{n+1})$ ,  $\xi_n$  is the canonical line bundle over  $L^n(p)$ . Put  $X = c_1(\xi_n)$ . Then, the element  $c_1(\eta)$  is described as a polynomial  $f(X)$  with the coefficient in  $U^*$ . We can see that  $c_1(\bar{\eta}) = f([-1]_F(X))$ . By the observation in Lemma 4.2, it follows that  $c_1(\eta) \in \tilde{U}^{ev}(L^n(p))^{Z_2}$ . Therefore, we have that if  $x \otimes_{U^*} m \in (\tilde{U}^{ev}(L^n(p)) \otimes_{U^*} Z)^{Z_2}$ , then there exists an element  $\eta \in \tilde{K}(L^n(p))$  such that

$$x \otimes_{U^*} m = c_1(\eta) \otimes_{U^*} 1, \quad c_1(\eta) \in \tilde{U}^{ev}(L^n(p))^{Z_2}.$$

q.e.d.

From the isomorphism (5.1), Lemma 5.1 and Theorem 1.2, we have the following.

**Theorem 5.2** ([5] and [6]).

$$\tilde{K}(D_p(2k+1, 4k+3)) \cong Z \oplus \tilde{K}(L^{2k+1}(p))^{Z_2} \oplus \tilde{K}(RP^{4k+3}).$$

KYUSHU UNIVERSITY

## References

- [1] M.F. Atiyah and G.B. Segal: *Equivariant K-theory*, Lecture Note, University of Warwick (1965).
- [2] P.E. Conner and E.E. Floyd: *Differentiable Periodic Maps*, *Ergebnisse der Math. und ihrer Grenzgebiete*, Springer Verlag, 1964.
- [3] P.E. Conner and E.E. Floyd: *The Relation of Cobordism to K-theories*, Springer, Berlin, 1966.
- [4] A. Dold: *Relations between ordinary and extraordinary cohomology*, Notes, Aarhus Colloquim on Algebraic Topology, Aarhus, 1962.
- [5] T. Fujino, N. Ishikawa and M. Kamata: *On the complex K-group of certain manifold*, *Math. Rep. College Ed. Kyushu Univ.* **9** (1973), 1–6.
- [6] M. Imaoka and M. Sugawara: *On the K-ring of the orbit manifold  $(S^{2m+1} \times S^l)/D_n$  by the dihedral group  $D_n$* , *Hiroshima Math. J.* **4** (1974), 53–70.
- [7] M. Kamata and H. Minami: *Bordism groups of dihedral groups*, *J. Math. Soc. Japan* **25** (1973), 334–341.
- [8] P. S. Landweber: *Coherence, flatness and cobordism of classifying spaces*, Proc. Adv Study inst., Aarhus, 1970.
- [9] N. Mahammed: *A propos de la K-theories des espaces lenticulaires*, *C.R. Acad. Sci. Paris*, **271** (1970), 639–642.
- [10] J. Milnor: *On axiomatic homology theory*, *Pacific J. Math.* **12** (1962), 337–341.
- [11] R.S. Palais: *The classification of G-spaces*, *Mem. Amer. Math. Soc.* No. **36** (1960).
- [12] D. Quillen: *Elementary proofs of some results of cobordism theory using Steenrod operations*, *Advances in Math.* **7** (1971), 29–56.
- [13] K. Shibata: *Oriented and weakly complex bordism algebra of free periodic maps*, *Trans. Amer. Math. Soc.* **177** (1973), 199–220.