

Title	Schottky groups and Bers boundary of Teichmüller space
Author(s)	Ito, Kentaro
Citation	Osaka Journal of Mathematics. 2003, 40(3), p. 639-657
Version Type	VoR
URL	https://doi.org/10.18910/12325
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

SCHOTTKY GROUPS AND BERS BOUNDARY OF TEICHMÜLLER SPACE

KENTARO ITO

(Received July 15, 1999)

1. Introduction

We will show that every Kleinian group on the Bers boundary of the Teichmüller space is an algebraic limit of a sequence of Schottky groups. This claim has already proved by Otal [26], but our argument is completely different from that of Otal. In this paper, we extend the action of the mapping class group on a Bers slice to that on a wider class of Kleinian groups. We obtain a sufficient condition for the action of the mapping class group to be continuous at a given point and show that the orbit of every maximal cusp is dense in the Bers boundary.

Here, we explain the fundamental idea of extending the action of the mapping class group. Let S be an oriented compact surface possibly with boundary ∂S , and let $T(S)$ be the Teichmüller space of complete hyperbolic structures on the interior of S with finite area. Let $R(S)$ be the space of conjugacy classes $[\rho, G]$ of representations $\rho: \pi_1(S) \rightarrow G \subset \mathrm{PSL}_2(\mathbb{C})$ of $\pi_1(S)$ which map each component of the boundary ∂S to parabolic elements. The subspace $QF(S)$ of $R(S)$ consisting of discrete faithful representations whose images are quasi-Fuchsian groups is naturally identified with a product of Teichmüller spaces $T(S) \times T(\tilde{S})$, where \tilde{S} denotes S with its orientation reversed. We denote the canonical homeomorphism by

$$Q: T(S) \times T(\tilde{S}) \rightarrow QF(S).$$

The mapping class group $\mathrm{Mod}(S)$ of S naturally acts on $T(S)$ and $T(\tilde{S})$ and hence on the Bers slice $B_X = Q(\{X\} \times T(\tilde{S}))$ by

$$Q(X, \tilde{Y}) \mapsto Q(X, \sigma\tilde{Y})$$

for $(X, \tilde{Y}) \in T(S) \times T(\tilde{S})$ and $\sigma \in \mathrm{Mod}(S)$. A crucial observation is that the representation $Q(X, \sigma\tilde{Y})$ has another description as follows;

$$Q(X, \sigma\tilde{Y}) = Q(\sigma^{-1}X, \tilde{Y}) \circ \sigma_*^{-1},$$

where σ_* is the group isomorphism of $\pi_1(S)$ induced by σ . The right-hand side of the above equation suggests us a possibility to define the action of $\mathrm{Mod}(S)$ even when \tilde{Y}

is pinched or degenerated. From this point of view, we extend the action of $\text{Mod}(S)$ on the Bers slice B_X to that on the subset C_X of $R(S)$, which is called the *extended Bers slice* for X . A representation $[\rho, G]$ is an element of C_X by definition if G is a function group with an invariant component $\Omega_0(G)$ of the region of discontinuity $\Omega(G)$ which is covering $X \in T(S)$. We remark that the extended Bers slice C_X contains the closure \bar{B}_X of the Bers slice B_X and that the restriction of the action to \bar{B}_X coincides with the action defined by Bers in [4].

The set $\partial B_X = \bar{B}_X - B_X$ is called the *Bers boundary*. Bers [4] obtained a sufficient condition for the action of $\text{Mod}(S)$ to be continuous at a give point in ∂B_X . On the other hand, it is known by that the action of $\text{Mod}(S)$ on ∂B_X is not always continuous (see Kerckhoff-Thurston [13]). In Section 3, we extend the Bers' result and obtain a sufficient condition for the action to be continuous at a given point in C_X :

Corollary 3.2. *Let $[\rho, G]$ be an element of C_X such that all components of $\Omega(G)/G$ except for $X = \Omega_0(G)/G$ have no moduli of deformation. Then $\text{Mod}(S)$ acts continuously at $[\rho, G]$.*

It is known by McMullen [24] that the set of maximal cusps is dense in ∂B_X . Moreover, one can see that the set of maximal cusps in ∂B_X decomposes into finitely many orbits under the action of $\text{Mod}(S)$. In Section 5, we will show that *each* orbit is dense in ∂B_X :

Theorem 5.5. *For any maximal cusp in ∂B_X , its orbit under the action of $\text{Mod}(S)$ is dense in ∂B_X .*

Suppose that S is a closed surface of genus $g \geq 2$. Now let S_X be the subset of C_X consisting of Schottky groups. In Section 6, we combine Corollary 3.2 (continuity at maximal cusps) and Theorem 5.5 (orbit density for maximal cusps) to obtain the following theorem:

Theorem 6.1. *Suppose that S is a closed surface of genus ≥ 2 . The set of accumulation points of S_X contains the Bers boundary ∂B_X .*

REMARK. Gallo [10] has proved the claim of Theorem 6.1 for the case of genus 2. More generally, Otal [26] has already proved the claim of Theorem 6.1 for the case of any genus ≥ 2 by using hyperbolic Dehn surgery theorem. In our proof, we will make use of Thurston's compactness theorem [28] instead of hyperbolic Dehn surgery theorem.

This paper is organized as follows: In Section 2, we give a definition of an extended Bers slice C_X on which we shall define the action of the mapping class group.

In Section 3, we obtain a sufficient condition for the action of $\text{Mod}(S)$ to be continuous at a point C_X (Corollary 3.2). In Section 5, we show that the orbit of every maximal cusp is dense in the Bers boundary ∂B_X (Theorem 5.5). In Section 6, we prove our main theorem (Theorem 6.1) as a consequence of the preceding sections. In Sections 5 and 6, one of the crucial tools is Thurston's compactness theorem [28], which will be introduced in Section 4. In Section 7, we collect some properties of the set of Schottky groups S_X which can be easily seen.

ACKNOWLEDGEMENTS. The author would like to express his gratitude to Hiroshige Shiga for his useful suggestions. He also appreciates the referee's valuable comments on the previous manuscript of this paper.

2. Preliminaries

2.1. Teichmüller spaces and Kleinian groups. Let S be a compact oriented surface of negative Euler characteristic possibly with boundary ∂S . The *Teichmüller space* $T(S)$ of S is the set of equivalence classes of pairs (f, X) ; where X is a complete hyperbolic Riemann surface of finite area and $f: \text{int}(S) \rightarrow X$ is an orientation preserving homeomorphism from the interior of S . Two pairs (f_1, X_1) and (f_2, X_2) are said to be equivalent if there is a conformal map $g: X_1 \rightarrow X_2$ such that $g \circ f_1$ is isotopic to f_2 . The equivalence class of (f, X) is simply denoted by X .

The *mapping class group* $\text{Mod}(S)$ is the group consisting of isotopy classes of orientation preserving homeomorphisms of S . The natural action of $\sigma \in \text{Mod}(S)$ on $T(S)$ is given by

$$\sigma(f, X) = (f \circ \sigma^{-1}, X).$$

We also consider the *Teichmüller space* $T(\bar{S})$ of \bar{S} , where \bar{S} denotes S with its orientation reversed. Then we have a canonical isomorphism $\iota: T(S) \rightarrow T(\bar{S})$ defined by $(f: S \rightarrow X) \mapsto (\bar{f}: \bar{S} \rightarrow \bar{X})$, where \bar{X} is the reflection of X . The action of $\sigma \in \text{Mod}(S) = \text{Mod}(\bar{S})$ on $T(\bar{S})$ is given by $\sigma(\bar{f}, \bar{X}) = (\bar{f} \circ \sigma^{-1}, \bar{X})$, so that $\sigma \circ \iota = \iota \circ \sigma$ is satisfied.

A Kleinian group G is a discrete subgroup of $\text{PSL}_2(\mathbb{C})$, which acts on the hyperbolic space \mathbf{H}^3 as isometries, and on the sphere at infinity $S_\infty^2 = \hat{\mathbb{C}}$ as conformal automorphisms. The limit set of G in $\hat{\mathbb{C}}$ is denoted by $\Lambda(G)$ and its complement $\hat{\mathbb{C}} - \Lambda(G)$, which is called the region of discontinuity of G , is denoted by $\Omega(G)$. A Kleinian group G is called a *function group* if its region of discontinuity $\Omega(G)$ has an invariant component. If a function group G has exactly two invariant components, it is called a *quasi-Fuchsian group*; otherwise, it has a unique invariant component.

2.2. Projective structures. For a given $X \in T(S)$, let Γ_X be a Fuchsian group acting on the unit disc $\Delta = \{z \in \hat{\mathbb{C}} : |z| < 1\}$ such that $X = \Delta/\Gamma_X$. A *bounded holomorphic quadratic differential* on Δ for Γ_X is a holomorphic function φ on Δ satis-

fying $\varphi \circ \gamma(\gamma')^2 = \varphi$ for any $\gamma \in \Gamma_X$ and $\|\varphi\|_\infty < \infty$, where $\|\varphi\|_\infty$ is the hyperbolic sup-norm of φ defined by

$$\|\varphi\|_\infty = \sup_{z \in \Delta} (1 - |z|^2)^2 |\varphi(z)|.$$

We let $B_2(\Gamma_X)$ denote the set of bounded holomorphic quadratic differentials on Δ for Γ_X . Then $B_2(\Gamma_X)$ is a finite dimensional complex Banach space.

The *developing map* for $\varphi \in B_2(\Gamma_X)$ is a meromorphic local homeomorphism

$$f_\varphi: \Delta \rightarrow \hat{\mathbb{C}}$$

whose Schwarzian derivative $S(f_\varphi)$ is equal to φ . We always assume that the developing map f_φ is normalized by the conditions $f_\varphi(0) = 0$, $f'_\varphi(0) = 1$ and $f''_\varphi(0) = 0$. Associated to the developing map f_φ , there is a group homomorphism

$$\rho_\varphi: \Gamma_X \rightarrow \mathrm{PSL}_2(\mathbb{C})$$

satisfying $f_\varphi \circ \gamma = \rho_\varphi(\gamma) \circ f_\varphi$ for all $\gamma \in \Gamma_X$. This group homomorphism ρ_φ is said to be the *holonomy representation* for φ . We call the pair $(f_\varphi, \rho_\varphi)$ the *(normalized) projective structure* for $\varphi \in B_2(\Gamma_X)$. Then there is a bijective correspondence between the set of normalized projective structures and $B_2(\Gamma_X)$.

We denote by $\hat{C}(\Gamma_X)$ the subset of $B_2(\Gamma_X)$ consisting of elements $\varphi \in B_2(\Gamma_X)$ whose developing maps f_φ are covering maps onto their images. For an element φ of $\hat{C}(\Gamma_X)$, the holonomy image $G = \rho_\varphi(\Gamma_X)$ of Γ_X is a function group (possibly with torsion) and $f_\varphi(\Delta)$ is an invariant component of G , which is denoted by $\Omega_0(G)$ (see [14] and [16] for more information). Moreover, we consider a subset $C(\Gamma_X)$ of $\hat{C}(\Gamma_X)$ as follows. An element φ of $\hat{C}(\Gamma_X)$ is contained in $C(\Gamma_X)$ by definition if the developing map $f_\varphi: \Delta \rightarrow f_\varphi(\Delta) = \Omega_0(G) \subset \hat{\mathbb{C}}$ descends to a conformal isomorphism $X = \Delta/\Gamma_X \rightarrow f_\varphi(\Delta)/\rho_\varphi(\Gamma_X) = \Omega_0(G)/G$ where G is the holonomy image $\rho_\varphi(\Gamma_X)$ of Γ_X . In summary,

$$\begin{aligned} \hat{C}(\Gamma_X) &= \{\varphi \in B_2(\Gamma_X) \mid f_\varphi: \Delta \rightarrow f_\varphi(\Delta) \subset \hat{\mathbb{C}} \text{ is a covering map}\}, \\ C(\Gamma_X) &= \{\varphi \in \hat{C}(\Gamma_X) \mid X = \Delta/\Gamma_X \cong f_\varphi(\Delta)/\rho_\varphi(\Gamma_X)\}. \end{aligned}$$

If Γ_X is maximal (i.e. there are no Fuchsian groups which properly contain Γ_X) $C(\Gamma_X)$ coincides with $\hat{C}(\Gamma_X)$. Note that Γ_X is maximal for almost every $X \in T(S)$. On the other hand, in [18], one can find examples of elements of $\hat{C}(\Gamma_X)$ but not of $C(\Gamma_X)$ for some Γ_X . For $\varphi \in C(\Gamma_X)$, $G = \rho_\varphi(\Gamma_X)$ may have an elliptic element whose fixed points are not contained in the invariant component $\Omega_0(G) = f_\varphi(\Delta)$.

2.3. Extended Bers slices. A sequence of representations $\rho_n: \pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbb{C})$ is said to converges *algebraically* to a representation $\rho: \pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbb{C})$ if $\rho_n(g) \rightarrow$

$\rho(g)$ in $\mathrm{PSL}_2(\mathbf{C})$ for all $g \in \pi_1(S)$. The conjugacy class of a representation $\rho: \pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbf{C})$ with $\rho(\pi_1(S)) = G$ is denoted by $[\rho, G]$ or simply by $[\rho]$. Let $R(S)$ denote the space of conjugacy classes $[\rho]$ of irreducible representations $\rho: \pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbf{C})$ such that $\rho(\gamma)$ is parabolic for every $\gamma \in \pi_1(\partial S)$. The space $R(S)$ is a complex manifold endowed with the topology of algebraic convergence.

It is known by Kra [15] that the map

$$\mathrm{hol}: B_2(\Gamma_X) \rightarrow R(S)$$

defined by $\varphi \mapsto [\rho_\varphi]$ is a holomorphic embedding, where $[\rho_\varphi]$ is the conjugacy class of the representation $\rho_\varphi: \pi_1(S) \cong \Gamma_X \rightarrow \mathrm{PSL}_2(\mathbf{C})$. (Here and hereafter, we frequently identify a representation of $\pi_1(S)$ with a representation of Γ_X .) For any $X \in T(S)$, we define subsets \hat{C}_X and C_X of $R(S)$ by

$$\begin{aligned}\hat{C}_X &= \mathrm{hol}(\hat{C}(\Gamma_X)), \\ C_X &= \mathrm{hol}(C(\Gamma_X)).\end{aligned}$$

We call C_X the *extended Bers slice*, on which we will define an action of the mapping class group.

The *Bers slice* B_X is the subset of C_X consisting of faithful representations whose images are quasi-Fuchsian groups. It is known by Bers [3] that the Bers slice B_X can be identified with the Teichmüller space $T(S)$, and that it is relatively compact in $R(S)$. The set $\partial B_X = \bar{B}_X - B_X$ is called the *Bers boundary*, where \bar{B}_X is the closure of B_X in $R(S)$. Moreover, we denote by \hat{B}_X the subset of C_X consisting of faithful representations. It is conjectured that $\bar{B}_X = \hat{B}_X$ in Bers [3].

The following are the sets which we want to consider in this paper:

$$B_X \subset \bar{B}_X \subseteq \hat{B}_X \subset C_X \subset \hat{C}_X.$$

EXAMPLE. In the case that S is a closed surface, a typical example of an element of $C_X - \hat{B}_X$ is a Schottky group. A Kleinian group G is a *Schottky group* if G is torsion-free and if its Kleinian manifold $N_G = (\mathbf{H}^3 \cup \Omega(G))/G$ is homeomorphic to a handlebody H_g of genus g . Let G be a Schottky group which uniformizes X , that is $X = \Omega(G)/G = \partial N_G$, then the representation $\rho: \pi_1(S) \cong \pi_1(\partial N_G) \rightarrow G \cong \pi_1(N_G)$ induced by the inclusion map $\partial N_G \hookrightarrow N_G$ is an element of C_X but not of \hat{B}_X .

Lemma 2.1. *For any $X \in T(S)$, C_X is a compact subset of $R(S)$.*

Proof. To show that $C_X = \mathrm{hol}(C(\Gamma_X))$ is compact, it is enough to see that $C(\Gamma_X)$ is closed and bounded subset of $B_2(\Gamma_X)$. Since it is known by Kra and Maskit [18] that $\hat{C}(\Gamma_X)$ is a closed and bounded subset of $B_2(\Gamma_X)$, we only have to show that $C(\Gamma_X)$ is closed. Let $\varphi_n \in C(\Gamma_X)$ be a sequence converging to $\varphi \in \hat{C}(\Gamma_X)$. Let f_n and

f be the developing maps corresponding to φ_n and φ , respectively. Then f_n converges to f locally uniformly on Δ . Suppose that the map $g: X \rightarrow f(\Delta)/\rho(\Gamma_X)$ induced by $f: \Delta \rightarrow f(\Delta)$ is not injective. Then there are two points $x, y \in X (x \neq y)$ such that $g(x) = g(y)$, and hence there are lifts $\tilde{x}, \tilde{y} \in \Delta$ of x and y such that $f(\tilde{x}) = f(\tilde{y})$. Since $f_n(\tilde{x}) \rightarrow f(\tilde{x})$ and $f_n(\tilde{y}) \rightarrow f(\tilde{y})$, the hyperbolic distances d_n between $f_n(\tilde{x})$ and $f_n(\tilde{y})$ in $f_n(\Delta)$ tend to 0 as $n \rightarrow \infty$. On the other hand, since $\varphi_n \in C(\Gamma_X)$, the maps $g_n: X \rightarrow f_n(\Delta)/\rho_n(\Gamma_X)$ induced by $f_n: \Delta \rightarrow f_n(\Delta)$ are conformal isomorphisms. Hence the hyperbolic distance between x and y on X is equal to the hyperbolic distance between $g_n(x)$ and $g_n(y)$ on $f_n(\Delta)/\rho_n(\Gamma_X)$ which are less or equal to d_n . This contradicts to $x \neq y$. \square

2.4. Quasiconformal deformations. For a given Kleinian group G with $\Omega(G) \neq \emptyset$, a measurable function μ on $\hat{\mathbb{C}}$ is called a *Beltrami differential* for G if

$$\mu(g(z))\overline{g'(z)} = \mu(z)g'(z)$$

holds for a.e. $z \in \hat{\mathbb{C}}$ and for all $g \in G$. The space of all Beltrami differentials μ for G whose essential sup-norm satisfying $\|\mu\|_\infty < 1$ is denoted by $\text{Belt}(G)_1$. For a G -invariant open set $U \subset \Omega(G)$, we denote by $\text{Belt}(G, U)_1$ the subset of $\text{Belt}(G)_1$ consisting of all elements with support in U . For $\mu \in \text{Belt}(G, U)_1$, there is a unique quasiconformal homeomorphism

$$w_\mu: \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$$

satisfying $(w_\mu)_z/(w_\mu)_{\bar{z}} = \mu$ (a.e.) and fixing 0, 1 and ∞ . Two elements $\mu, \nu \in \text{Belt}(G, U)_1$ are *equivalent* (denoted by $\mu \sim \nu$) if w_μ and w_ν induce the same group isomorphism; that is, $w_\mu \circ g \circ (w_\mu)^{-1} = w_\nu \circ g \circ (w_\nu)^{-1}$ for all $g \in G$.

In the rest of this subsection, we restrict our attention to the following situation: For a given $X \in T(S)$, let $[\rho, G]$ be an element of C_X and let $f: \Delta \rightarrow \Omega_0(G)$ be the developing map inducing the holonomy representation $\rho: \pi_1(S) \cong \Gamma_X \rightarrow G$.

We denote by $QC([\rho])$ the space of quasi-conformal deformations of $[\rho] = [\rho, G]$ induced by elements of $\text{Belt}(G)_1$. That is, $[\rho'] \in R(S)$ is an element of $QC([\rho])$ if $\rho'(\gamma) = w_\mu \circ \rho(\gamma) \circ (w_\mu)^{-1}$ is satisfied for all $\gamma \in \pi_1(S)$ for some $\mu \in \text{Belt}(G)_1$. This space $QC([\rho])$ is identified with the quotient space $\text{Belt}(G)_1 / \sim$. Moreover, we denote by $QC_0([\rho])$ the space of quasi-conformal deformations of $[\rho] = [\rho, G]$ induced by elements of $\text{Belt}(G, \Omega_0(G))_1$, that is $QC_0([\rho]) = \text{Belt}(G, \Omega_0(G))_1 / \sim$.

Now let consider the case that ρ is the identity representation $\text{id}: \pi_1(S) \cong \Gamma_X \rightarrow \Gamma_X$ induced by the identity map $\text{id}: \Delta \rightarrow \Delta = \Omega_0(\Gamma_X)$. Then $\Omega(\Gamma_X)$ has exactly two component $\Delta = \Omega_0(\Gamma_X)$ and $\Delta^* = \{z \in \hat{\mathbb{C}} : |z| > 1\}$. The space of quasiconformal deformations $QC([\text{id}])$ of $[\text{id}]$ is said to be the *quasi-Fuchsian space* and denoted by $QF(S)$. On the other hand, the quotient space $QC_0([\text{id}]) = \text{Belt}(\Gamma_X, \Delta)_1 / \sim$ can be identified with the Teichmüller space $T(X) \cong T(S)$. Similarly, the quotient space $\text{Belt}(\Gamma_X, \Delta^*)_1 / \sim$ can be identified with the Teichmüller space $T(\bar{X}) \cong T(\bar{S})$, where

$\bar{X} = \Delta^*/\Gamma_X$ is the reflection of X . It is the well known fact (cf. Bers [2]) that the map

$$\text{Belt}(\Gamma_X, \Delta) \times \text{Belt}(\Gamma_X, \Delta^*) \rightarrow \text{Belt}(\Gamma_X)$$

defined by $(\mu, \nu) \mapsto \mu + \nu$ descends to the canonical homeomorphism

$$Q: T(S) \times T(\bar{S}) \rightarrow QF(S).$$

We remark that the Bers slice B_X defined in Subsection 2.3 is equal to $Q(\{X\} \times T(\bar{S}))$.

Recall that the holonomy representation $\rho: \pi_1(S) \cong \Gamma_X \rightarrow G$ is induced by the developing map $f: \Delta \rightarrow \Omega_0(G)$. For $\mu \in \text{Belt}(G, \Omega_0(G))_1$, the pull-back $f^*\mu$ of μ by f is defined by

$$f^*\mu(z) = \mu \circ f(z) \frac{\overline{f'(z)}}{f'(z)}.$$

Since f descends to an isomorphism $\Delta/\Gamma_X \rightarrow \Omega_0(G)/G$, the map

$$f^*: \text{Belt}(G, \Omega_0(G))_1 \rightarrow \text{Belt}(\Gamma_X, \Delta)_1$$

defined by $\mu \mapsto f^*\mu$ is an isomorphism. This is the reason why we can define the action on C_X but cannot on \hat{C}_X . We denote the inverse $(f^*)^{-1}$ of f^* by $f_*: \text{Belt}(\Gamma_X, \Delta)_1 \rightarrow \text{Belt}(G, \Omega_0(G))_1$. Then, it was shown by Maskit [21] (see also Kra [17]) that the map $f_*: \text{Belt}(\Gamma_X, \Delta)_1 \rightarrow \text{Belt}(G, \Omega_0(G))_1$ descends to an unbranched covering map

$$\underline{f}_*: T(S) \rightarrow QC_0([\rho])$$

with $\underline{f}_*(X) = [\rho]$. We use the notation

$$\text{qc}([\rho], Y) = \underline{f}_*(Y)$$

for any $Y \in T(S)$. Then the representation $\text{qc}([\rho], Y) \in QC_0([\rho])$ can be regarded as the quasiconformal deformation of $[\rho]$ corresponding to the quasiconformal deformation Y of X in $T(S)$. Note that $\text{qc}([\rho], X) = [\rho]$ and that $\text{qc}([\rho], Y) \in C_Y$ for all $Y \in T(S)$.

2.5. The action of the mapping class group. Now we define the action of $\text{Mod}(S)$ on C_X by

$$\sigma([\rho]) = \text{qc}([\rho], \sigma^{-1}X) \circ \sigma_*^{-1}$$

for $[\rho] \in C_X$ and $\sigma \in \text{Mod}(S)$, where σ_* is the group isomorphism of $\pi_1(S)$ induced by σ . Since $[\rho'] \circ \sigma_*^{-1} \in C_{\sigma Y}$ for any $[\rho'] \in C_Y$ and $Y \in T(S)$, one can see that $\sigma([\rho])$ also contained in C_X .




Fig. 1. The action of the mapping class group

Let take $[\rho] \in C_X$ and $\sigma \in \text{Mod}(S)$ and put $[\rho'] = \sigma([\rho])$. Then $\ker \rho' = \sigma_*(\ker \rho)$ is satisfied. Therefore, if $\ker \rho \neq \sigma_*(\ker \rho)$, $\sigma([\rho])$ is not a quasiconformal deformation of $[\rho]$ and hence $\sigma([\rho])$ is not contained in the connected component of C_X containing $[\rho]$.

Here we explain that the action of $\text{Mod}(S)$ on the Bers slice $B_X (\subset C_X)$ defined above coincides with the action on the Teichmüller space $T(\bar{S})$ under the identification $B_X = T(\bar{S})$ (see Fig. 1). The mapping class group $\text{Mod}(S)$ acts on $B_X = Q(\{X\} \times T(\bar{S}))$ by $Q(X, \bar{Y}) \mapsto Q(X, \sigma\bar{Y})$ for $Q(X, \bar{Y}) \in B_X$ and $\sigma \in \text{Mod}(S)$. Now we put $[\rho] = Q(X, \bar{Y})$ and are going to show that $Q(X, \sigma\bar{Y}) = \sigma([\rho])$. By definition, $\sigma([\rho]) = qc([\rho], \sigma^{-1}X) \circ \sigma_*^{-1} = Q(\sigma^{-1}X, \bar{Y}) \circ \sigma_*^{-1}$. Since $Q(\sigma X, \sigma Y) = Q(X, Y) \circ \sigma_*^{-1}$ holds for any $(X, \bar{Y}) \in T(S) \times T(\bar{S})$, one obtain the desired equation $Q(X, \bar{Y}) = Q(\sigma^{-1}X, \bar{Y}) \circ \sigma_*^{-1}$.

3. Continuity of the action of the mapping class group

In this section, we obtain a sufficient condition for $[\rho] \in C_X$ so that the action of $\text{Mod}(S)$ at $[\rho]$ is continuous. The same result, where C_X is replaced by \hat{B}_X , was obtained by Bers [4].

We first show the continuity for the change of base points.

Proposition 3.1. *Let $[\rho, G]$ be an element of C_X such that all components of $\Omega(G)/G$ except for $X = \Omega_0(G)/G$ have no moduli of deformation. Then the following holds: If $[\rho_n] \rightarrow [\rho]$ in C_X then $qc([\rho_n], Y) \rightarrow qc([\rho], Y)$ in C_Y for all $Y \in T(S)$.*

Proof. Let φ_n and φ be elements in $C(\Gamma_X)$ such that $\text{hol}(\varphi_n) = [\rho_n]$ and $\text{hol}(\varphi) = [\rho]$, respectively. Since $C(\Gamma_X)$ is compact and the map $\text{hol}: B_2(\Gamma_X) \rightarrow R(S)$ is injective, $\varphi_n \rightarrow \varphi$ in $C(\Gamma_X)$. Let (f_n, ρ_n) and (f, ρ) be normalized projective structures for φ_n and φ , respectively. Then f_n converges to f locally uniformly on Δ . Let $\mu \in \text{Belt}(\Gamma_X, \Delta)_1$ be a representative of $Y \in T(S) = \text{Belt}(\Gamma_X, \Delta)_1 / \sim$. We may assume that μ is continuous function on Δ . Put $\hat{\mu}_n = (f_n)_* \mu \in \text{Belt}(G_n, \Omega_0(G_n))_1$ and $\hat{\mu} = f_* \mu \in \text{Belt}(G, \Omega_0(G))_1$, where $G_n = \rho_n(\pi_1(S))$ and $G = \rho(\pi_1(S))$. Since $\{w_{\hat{\mu}_n}\}$ fix 0, 1 and ∞ and their dilatations are uniformly bounded, it has a subsequence (which we denote by the same symbol) $\{w_{\hat{\mu}_n}\}$ converging uniformly to some quasiconformal homeomorphism w_∞ of $\hat{\mathbb{C}}$. Since the representatives of $\text{qc}([\rho_n], Y)$ are induced by $w_{\hat{\mu}_n} \circ f_n$, $\{\text{qc}([\rho_n], Y)\}$ converges algebraically to the conjugacy class of the representation induced by $w_\infty \circ f$. Therefore, we have only to show that w_∞ and $w_{\hat{\mu}}$ induce the same group isomorphism from G into $\text{PSL}_2(\mathbb{C})$.

Since injectivity radii (with respect to the Poincaré metric on Δ) of f_n are uniformly bounded below (see [18, Lemma 5.1]), for any $z \in \Omega_0(G)$, there exist an open neighborhood U of z and suitable branches of the inverse maps f_n^{-1} and f^{-1} on U such that f_n^{-1} converges to f^{-1} uniformly on U . Hence one can see that $\hat{\mu}_n$ converges to $\hat{\mu}$ locally uniformly on $\Omega_0(G)$. Therefore, $w_{\hat{\mu}_n} \circ (w_{\hat{\mu}})^{-1}$ converges to a conformal map $w_\infty \circ (w_{\hat{\mu}})^{-1}$ locally uniformly on $w_{\hat{\mu}}(\Omega_0(G))$, and hence, the Beltrami coefficient of w_∞ is equal to $\hat{\mu}$ almost everywhere on $\Omega_0(G)$. Since there is no essential deformation on $\Omega(G) - \Omega_0(G)$ by assumption and on $\Lambda(G)$ by Sullivan's rigidity theorem [27], w_∞ and $w_{\hat{\mu}}$ induce the same group isomorphism. \square

Corollary 3.2. *Let $[\rho, G]$ be an element of C_X such that all components of $\Omega(G)/G$ except for $X = \Omega_0(G)/G$ have no moduli of deformation. Then the action of $\text{Mod}(S)$ is continuous at $[\rho]$; that is, if $[\rho_n] \rightarrow [\rho]$ in C_X then $\sigma([\rho_n]) \rightarrow \sigma([\rho])$ for all $\sigma \in \text{Mod}(S)$.*

Proof. By Proposition 3.1, $\text{qc}([\rho_n], \sigma^{-1}X) \rightarrow \text{qc}([\rho], \sigma^{-1}X)$ for all $\sigma \in \text{Mod}(S)$. Therefore, $\sigma([\rho_n]) = \text{qc}([\rho_n], \sigma^{-1}X) \circ \sigma_*^{-1}$ converges algebraically to $\sigma([\rho]) = \text{qc}([\rho], \sigma^{-1}X) \circ \sigma_*^{-1}$. \square

REMARK. In [13], Kerckhoff and Thurston showed that there is a Bers slice B_X and a point $[\rho] \in \partial B_X$ at which the action of $\text{Mod}(S)$ is not continuous.

4. Thurston's compactness theorem

In this section, we introduce Thurston's compactness theorem [28], which will play an important role in the following sections.

Let M be a compact 3-manifold with boundary ∂M . A non-trivial closed curve γ on ∂M is said to be *compressible* if it is null homotopic in M ; otherwise it is *incompressible*. A proper map $f: (A, \partial A) \rightarrow (M, \partial M)$ of an annulus A into M is said to

be *essential* if $f_*: \pi_1(A) \rightarrow \pi_1(M)$ is an injection and f is not homotopic (as a map of pairs) to a map into ∂M .

DEFINITION. Let M be a compact 3-manifold with boundary ∂M . Let λ be a system of non-trivial, mutually disjoint, homotopically distinct simple closed curves on ∂M . Then a pair (M, λ) is *doubly incompressible* if

- (1) every compressible simple closed curve on ∂M intersects λ at least three times,
- (2) there are no essential annuli with boundary in $\partial M - \lambda$, and
- (3) every maximal abelian subgroup of $\pi_1(\partial M - \lambda)$ is mapped to a maximal abelian subgroup of $\pi_1(M)$.

Let M be as above. In addition, we assume that the interior of M admits a hyperbolic structure. We denote by $AH(M)$ the space of conjugacy classes $[\rho, G]$ of discrete faithful representations $\rho: \pi_1(M) \rightarrow G \subset \mathrm{PSL}_2(\mathbf{C})$. The space $AH(M)$ is equipped with the algebraic topology. Let γ be an incompressible closed curve on ∂M . For $[\rho, G] \in AH(M)$, $\mathrm{length}_\rho(\gamma)$ denotes the length of the geodesic representative of γ in the hyperbolic manifold \mathbf{H}^3/G if $\rho(\gamma)$ is loxodromic and $\mathrm{length}_\rho(\gamma) = 0$ if $\rho(\gamma)$ is parabolic. For a positive constant $K > 0$, we denote by $AH(M, \lambda, K)$ the set of elements $[\rho, G] \in AH(M)$ such that $\mathrm{length}_\rho(\lambda) \leq K$, where $\mathrm{length}_\rho(\lambda)$ is the total sum of the lengths of all components of λ .

Now we can state Thurston's compactness theorem:

Theorem 4.1 (Thurston [28]). *Let M be a compact 3-manifold with boundary ∂M whose interior $\mathrm{int}(M)$ admits a hyperbolic structure. If (M, λ) is doubly incompressible, then $AH(M, \lambda, K)$ is compact for all $K > 0$.*

Let S be an oriented compact surface possibly with boundary ∂S . A curve system $\lambda = \{\alpha_j\}_{j=1}^N$ on S is called *homotopically independent* if it has the following properties: (1) each α_j is a simple closed curve on S and $\alpha_i \cap \alpha_j = \emptyset$ for $i \neq j$, (2) each α_j is non-trivial and not freely homotopic to a component of ∂S , and (3) α_i is not freely homotopic to α_j if $i \neq j$. A homotopically independent curve system $\lambda = \{\alpha_j\}_{j=1}^N$ on S is *maximal* if it divides S into pairs of pants. (If S is a closed surface of genus g with n open discs removed, then $N = 3g - 3 + n$.) A pair (λ, λ') of maximal curve systems on S is said to be *binding* S if they have no curves in common and if each component of $S - (\lambda \cup \lambda')$ is a simply connected domain or an annulus containing a component of ∂S in its boundary after realizing λ and λ' by geodesics for a hyperbolic structure on S .

The following lemma is discussed in a more general setting in Ohshika [25].

Lemma 4.2. *Let S be an oriented compact surface possibly with boundary ∂S . Let (λ', λ'') be a pair of maximal curve systems which binds S . For this pair, we de-*

fine a maximal curve system λ on $\partial(S \times I)$, where I is the closed interval $[0, 1]$, as

$$\lambda = (\lambda' \times \{0\}) \cup (\lambda'' \times \{1\}) \cup \left(\partial S \times \left\{ \frac{1}{2} \right\} \right).$$

Then $(S \times I, \lambda)$ is doubly incompressible.

Proof. We only consider the case $\partial S \neq \emptyset$, since the proof of the case $\partial S = \emptyset$ is easier. If $\partial S \neq \emptyset$, then $S \times I$ is homeomorphic to a handlebody H_g of some genus g . We identify $S \times I$ with H_g via this homeomorphism. We first check the condition (1) in the definition of double incompressibility. Let γ be a compressible simple closed curve on ∂H_g . Since $S \times \{0\}$ and $S \times \{1\}$ contain no compressible curves, γ must intersect a component of $\partial S \times \{1/2\}$. If $i(\gamma, \lambda) \leq 2$ (here $i(\cdot, \cdot)$ denotes the geometric intersection number), one can easily see that there exists a component δ of $\partial S \times \{1/2\}$ and a component W of $\partial H_g - ((\lambda' \times \{0\}) \cup (\lambda'' \times \{1\}))$ homeomorphic to a four-time-punctured sphere such that $\gamma \cup \delta \subset W$ and $i(\gamma, \delta) = 2$. Let α and β be components of ∂W such that α, β and γ bound a pair of pants T . Then, after choosing suitable orientations of α, β and γ , we have $[\gamma] = [\alpha] + [\beta]$ in $H_1(\partial H_g, \mathbf{Z})$, where $[\gamma]$ is the homology class of γ and so on. Since (λ', λ'') binds S , $\langle [\alpha], [\beta] \rangle$ are rank 2 free abelian subgroup of $H_1(\partial H_g, \mathbf{Z})$ which is mapped into $H_1(H_g, \mathbf{Z})$ injectively. This contradicts the assumption that γ is null homotopic and hence null homologous.

Now we check the condition (2). Suppose that there exists an essential annulus $f: (A, \partial A) \rightarrow (H_g, \partial H_g)$ with boundary in $\partial H_g - \lambda$. Let γ and γ' be the components of the image of ∂A in ∂H_g . Since $(\gamma \cup \gamma') \cap (\partial S \times \{1/2\}) = \emptyset$, γ and γ' may be assumed to be contained in $(S \times \{0\}) \cup (S \times \{1\})$. Let $p: S \times I \rightarrow S$ be the canonical retraction. Then $p(\gamma)$ is homotopic to $p(\gamma')$ in S . Since (λ', λ'') binds S , both γ and γ' are contained in $S \times \{0\}$ or $S \times \{1\}$. Now the retraction above gives a homotopy between f and a map into ∂H_g . This is a contradiction.

Finally, we check the condition (3). Since all non-trivial abelian subgroups of $\pi_1(\partial H_g - \lambda)$ or $\pi_1(H_g)$ are isomorphic to \mathbf{Z} , we have only to show that all primitive elements of $\pi_1(\partial H_g - \lambda)$ are also primitive in $\pi_1(H_g)$. This follows from the fact that H_g is homotopically equivalent to S by the retraction. \square

5. Orbit density for maximal cusps

Let $[\rho] \in \hat{B}_X$. The *accidental parabolic locus* of $[\rho]$ is a homotopically independent curve system $\lambda = \{\alpha_j\}$ on S such that $\rho(\alpha_j)$ is (the conjugacy class of) a parabolic element of $G = \rho(\pi_1(S))$ for every j , and no simple closed curve which is not homotopic to a component of λ has this property. For $[\rho] \in \hat{B}_X$, its accidental parabolic locus is uniquely determined up to homotopy. An element $[\rho] \in \hat{B}_X$ is called a *maximal cusp* if its accidental parabolic locus is maximal. It is a well known fact that every maximal cusp is contained in ∂B_X and that, for any maximal curve system λ on S , there exists a unique maximal cusp whose accidental parabolic locus

is λ (see Abikoff [1] and Maskit [20]).

For a simple closed curve α on S , let $D_\alpha \in \text{Mod}(S)$ denote the Dehn twist once around α .

Proposition 5.1. *Let (λ', λ'') be a binding pair of maximal curve systems on S . Let $[\rho] \in \partial B_X$ be a maximal cusp whose accidental parabolic locus is λ'' . Put $\sigma = D_{\alpha_1} \circ \cdots \circ D_{\alpha_N} \in \text{Mod}(S)$, where $\lambda' = \{\alpha_j\}_{j=1}^N$. Then the sequence $\{\sigma^n([\rho])\}_{n \in \mathbb{Z}}$ converges to the maximal cusp $[\rho_\infty] \in \partial B_X$ whose accidental parabolic locus is λ' as $|n| \rightarrow \infty$.*

In the proof of Proposition 5.1, we will make use of the following two lemmas; the first one is due to Canary [6] and the second one is a well known technical lemma.

Lemma 5.2 (Canary [6]). *Given $A > 0$, there exists a constant $R > 0$ such that if G is a non-elementary, torsion-free Kleinian group such that every incompressible closed geodesic on $\Sigma = \Omega(G)/G$ has hyperbolic length at least A , then for any closed curve γ on Σ ,*

$$\text{length}_N(\gamma) \leq R \cdot \text{length}_\Sigma(\gamma),$$

where $\text{length}_N(\gamma)$ and $\text{length}_\Sigma(\gamma)$ are hyperbolic lengths of geodesic representatives of γ in $N = \mathbf{H}^3/G$ and in Σ , respectively.

Lemma 5.3. *Let F_2 be a rank 2 free group and let $\{\chi_n: F_2 \rightarrow \text{PSL}_2(\mathbb{C})\}$ be a sequence of discrete faithful representations which converges algebraically to χ_∞ . If a sequence $\{\chi'_n = \psi_n \cdot \chi_n \cdot \psi_n^{-1}\}$ also converges algebraically to χ'_∞ for a sequence $\{\psi_n\}$ in $\text{PSL}_2(\mathbb{C})$, then ψ_n converges to some element ψ_∞ in $\text{PSL}_2(\mathbb{C})$.*

Proof of Proposition 5.1. Our argument is almost parallel to that of Kerckhoff and Thurston [13] (see also [5]).

Since C_X is compact, the sequence $\{[\rho_n] = \sigma^n([\rho])\}_{n \in \mathbb{Z}}$ has a convergent subsequence. We also denote this subsequence by the same symbol. In fact, in the following argument, we can see that any convergent subsequence of $\{[\rho_n]\}_{n \in \mathbb{Z}}$ converges to a unique maximal cusp $[\rho_\infty]$, and hence $\{[\rho_n]\}_{n \in \mathbb{Z}}$ converges without passing to a subsequence.

On the other hand, we shall show that the sequence $\{[\bar{\rho}_n] = \text{qc}([\rho], \sigma^{-n}X)\}_{n \in \mathbb{Z}}$ also has a convergent subsequence. Recall that $AH(S \times I)$ is the space of conjugacy classes of discrete faithful representations $\chi: \pi_1(S \times I) \rightarrow \text{PSL}_2(\mathbb{C})$. We denote by $AH_{\partial S}(S \times I)$ the set of representations $[\chi] \in AH(S \times I)$ such that $\chi(g)$ are parabolic for all $g \in \pi_1(\partial S \times I)$. Then we can regard $AH_{\partial S}(S \times I)$ as a subset of the representation space $R(S)$. Now we have a sequence $\{[\bar{\rho}_n]\}_{n \in \mathbb{Z}}$ in $AH_{\partial S}(S \times I) \subset R(S)$. Let λ be the

maximal curve system on $\partial(S \times I)$ defined by

$$\lambda = (\lambda' \times \{0\}) \cup (\lambda'' \times \{1\}) \cup \left(\partial S \times \left\{ \frac{1}{2} \right\} \right).$$

Then $(S \times I, \lambda)$ is doubly incompressible by Lemma 4.2. We can see that the sequence $\{[\bar{\rho}_n]\}_{n \in \mathbb{Z}}$ is contained in $AH(S \times I, \lambda, K)$ for some $K > 0$ since we have

$$\text{length}_{\bar{\rho}_n}(\lambda'' \times \{1\}) = \text{length}_{\bar{\rho}_n} \left(\partial S \times \left\{ \frac{1}{2} \right\} \right) = 0$$

and

$$\text{length}_{\bar{\rho}_n}(\lambda' \times \{0\}) \leq R \cdot \text{length}_{\sigma^{-n}X}(\lambda' \times \{0\}) = R \cdot \text{length}_X(\lambda' \times \{0\})$$

from Lemma 5.2. Since $AH(S \times I, \lambda, K)$ is a compact subset in $AH(S \times I)$ by Theorem 4.1, we have a convergent subsequence of $\{[\bar{\rho}_n]\}_{n \in \mathbb{Z}}$ in $AH(S \times I)$ and hence in $AH_{\partial S}(S \times I) \subset R(S)$. Again, we denote this subsequence by the same symbol.

Take representatives ρ_n of $[\rho_n] = \sigma^n([\rho])$ so that the sequence $\{\rho_n\}$ converges to a representation ρ_∞ . Since $[\rho_n] = \sigma^n([\rho]) = \text{qc}([\rho], \sigma^{-n}X) \circ \sigma_*^{-n} = [\bar{\rho}_n] \circ \sigma_*^{-n}$, we may assume that $\bar{\rho}_n = \rho_n \circ \sigma_*^n$. In addition, there are elements $\psi_n \in \text{PSL}_2(\mathbb{C})$ such that the sequence $\{\psi_n \cdot \bar{\rho}_n \cdot \psi_n^{-1}\}$ converges to a representation $\bar{\rho}_\infty$, since $\{[\bar{\rho}_n]\}_{n \in \mathbb{Z}}$ is a convergent sequence.


Now let α be a component of λ' . We are going to show that $\rho_\infty(\alpha)$ is a parabolic element. Let T be a component of $S - \lambda'$ containing α in its boundary and $\alpha' (\neq \alpha)$ be a component of λ' or a component of ∂S contained in the boundary of T . Choose a base point x in T and regard $\pi_1(S) = \pi_1(S, x)$. By abuse of notation, α and α' also denote the elements of $\pi_1(S, x)$ freely homotopic to α and α' respectively. Moreover, we assume that (the representatives of) $\alpha, \alpha' \in \pi_1(S, x)$ contained in T . Note that $\langle \alpha_1, \alpha_2 \rangle$ is a rank 2 free subgroup of $\pi_1(S, x)$. Since $\bar{\rho}_n(\alpha) = \rho_n \circ \sigma_*^n(\alpha) = \rho_n(\alpha)$ and $\bar{\rho}_n(\alpha') = \rho_n \circ \sigma_*^n(\alpha') = \rho_n(\alpha')$, the elements $\psi_n \in \text{PSL}_2(\mathbb{C})$ may be taken to be the identity by Lemma 5.3.

One can find non-trivial elements $\gamma_1, \gamma_2 \in \pi_1(S, x)$ which satisfy the following conditions (see Fig. 2): (1) γ_j intersects α twice in the opposite direction for $j = 1, 2$, (2) γ_j does not intersect any other components of λ' for $j = 1, 2$, and (3) $\langle \gamma_1, \gamma_2 \rangle$ is a rank 2 free subgroup of $\pi_1(S, x)$.

Then we have

$$\begin{cases} \bar{\rho}_n(\gamma_1) = \rho_n(\alpha^n) \cdot \rho_n(\gamma_1) \cdot \rho_n(\alpha^{-n}), \\ \bar{\rho}_n(\gamma_2) = \rho_n(\alpha^n) \cdot \rho_n(\gamma_2) \cdot \rho_n(\alpha^{-n}). \end{cases}$$

Since both the sequences $\{\rho_n\}$ and $\{\bar{\rho}_n\}$ are convergent sequence, Lemma 5.3 again implies that $\rho_n(\alpha^n)$ converges to an element $\hat{\alpha}$ in $\text{PSL}_2(\mathbb{C})$. Since $\rho_n(\alpha)$ commutes with $\rho_n(\alpha^n)$ for all n , $\rho_\infty(\alpha)$ commutes with $\hat{\alpha}$. If the abelian subgroup $\langle \rho_\infty(\alpha), \hat{\alpha} \rangle$ were

Fig. 2. γ_1 and γ_2

isomorphic to \mathbf{Z} , then $\rho_\infty(\alpha^k) = \hat{\alpha}^l$ for some integers k and l , and thus $\rho_n(\alpha^{nl-k}) \rightarrow \text{id}$. This contradicts the fact that $[\rho_n]$ are discrete faithful representations (cf. Jørgensen [12, Lemma 2]). Therefore we conclude that $\langle \rho_\infty(\alpha), \hat{\alpha} \rangle$ is isomorphic to $\mathbf{Z} \oplus \mathbf{Z}$, and hence it is a rank 2 parabolic subgroup in $\text{PSL}_2(\mathbf{C})$. In particular, $\rho_\infty(\alpha)$ is parabolic. The same argument works well for all components of λ' . Therefore we can conclude that $[\rho_\infty]$ is a maximal cusp whose accidental parabolic locus is λ' . \square

Lemma 5.4. *For any two maximal curve systems $\lambda = \{\alpha_j\}_{j=1}^N$ and $\lambda' = \{\beta_j\}_{j=1}^N$ on S , there exists a maximal curve system $\nu = \{\gamma_j\}_{j=1}^N$ such that the pairs (λ, ν) and (ν, λ') are binding S .*

Proof. There exists a simple closed curve δ on S such that $i(\delta, \alpha_j) > 0$ for all j (see [8]). Put $\sigma = D_{\alpha_1} \circ \cdots \circ D_{\alpha_N}$. If $i(\beta_j, \lambda) = 0$ then $\beta_j = \alpha_i$ for some i and hence $i(\beta_j, \sigma^n(\delta)) > 0$ for all n . If $i(\beta_j, \lambda) > 0$ then $i(\beta_j, \alpha_i) > 0$ for some i . In this case, $i(\beta_j, \sigma^n(\delta)) > 0$ for all but finitely many n . Therefore, for sufficiently large n , $i(\beta_j, \sigma^n(\delta)) > 0$ holds for all j . Fix such n and let $\gamma_1 = \sigma^n(\delta)$. Choose simple closed curves $\gamma_2, \dots, \gamma_N$ so that $\nu = \{\gamma_j\}_{j=1}^N$ is a maximal curve system. This ν satisfies the desired condition. \square

It was shown by McMullen [24] that the set of maximal cusps is dense in ∂B_X . Since the number of ways to decompose S into pairs of pants up to the action of $\text{Mod}(S)$ is finite, the set of maximal cusps in ∂B_X decomposes into finitely many orbits under the action of $\text{Mod}(S)$. The next theorem shows that *each* orbit is dense in ∂B_X .

Theorem 5.5. *For any maximal cusp $[\rho] \in \partial B_X$, its orbit $\{\sigma([\rho])\}_{\sigma \in \text{Mod}(S)}$ under the action of $\text{Mod}(S)$ is dense in ∂B_X .*

Proof. Since the set of maximal cusps is dense in ∂B_X , we have only to show that, for arbitrary fixed two maximal cusps $[\rho]$ and $[\rho']$ in ∂B_X , the orbit

$\{\sigma([\rho])\}_{\sigma \in \text{Mod}(S)}$ of $[\rho]$ contains a sequence converging to $[\rho']$. Let λ and λ' be accidental parabolic loci for $[\rho]$ and $[\rho']$, respectively. Then we can find a maximal curve system $\nu = \{\gamma_j\}_{j=1}^N$ such that both the pairs (λ, ν) and (ν, λ') are binding S (Lemma 5.4). Put $\sigma = D_{\gamma_1} \circ \cdots \circ D_{\gamma_N}$ and $\tau = D_{\beta_1} \circ \cdots \circ D_{\beta_N}$, where $\lambda' = \{\beta_j\}_{j=1}^N$. Then $\sigma^n([\rho])$ converges to a maximal cusp $[\rho''] \in \partial B_X$ whose accidental parabolic locus is ν by Proposition 5.1. Similarly $\tau^n([\rho''])$ converges to $[\rho']$. Since the action of $\text{Mod}(S)$ is continuous at maximal cusps (Corollary 3.2), we can find a desired sequence by a diagonal argument. \square

6. Orbits of Schottky groups and Bers boundary

In this section, we assume that S is a closed surface of genus ≥ 2 . We denote by S_X the set of $[\rho, G] \in C_X$ such that G is a Schottky group. The aim of this section is to prove the following theorem.

Theorem 6.1. *Let S be a closed surface of genus ≥ 2 . For any $X \in T(S)$, the set of accumulation points of S_X contains the boundary ∂B_X of the Bers slice B_X .*

REMARK. It is known by Gallo [9] that there is an accumulation point of S_X which is not contained in ∂B_X . This fact can be seen also from a slight modification of the proof of the above theorem.

In the proof of Theorem 6.1, we need the following lemma which give a sufficient condition for an element $[\rho] \in C_X$ to be contained in ∂B_X .

Lemma 6.2. *Let S be a compact surface of hyperbolic type possibly with boundary. Let $X \in T(S)$, $[\rho] \in C_X$ and $\lambda = \{\alpha_j\}$ be a maximal curve system on S . If $\rho(\alpha_j)$ are parabolic for all j , then $[\rho]$ is the maximal cusp in ∂B_X whose accidental parabolic locus is λ .*

Proof. We have only to show that $[\rho]$ is a faithful representation. Suppose that $\rho: \pi_1(S) \rightarrow G$ is not faithful. Then the covering map $p: \Omega_0(G) \rightarrow X = \Omega_0(G)/G$ is not universal, where $\Omega_0(G)$ is the unique invariant component of G . Then, by the planarity theorem (see [22], X.A.4), there exist a non-trivial simple closed curve δ on X and a simple closed curve $\tilde{\delta}$ on $\Omega_0(G)$ such that $p|_{\tilde{\delta}}: \tilde{\delta} \rightarrow \delta$ is a finite-sheeted covering map; say k -sheeted. Let $g \in G$ be a generator for the stabilizer of $\tilde{\delta}$ in G . Since $\lambda(\subset X)$ is maximal and δ is not parallel to a component of λ , it follows that δ must intersect some component of λ , say α_1 . We may assume that the number of intersection points of δ and α_1 is equal to $i(\delta, \alpha_1)$. Let $\tilde{\alpha}_1$ be a connected component of $p^{-1}(\alpha_1)$ on $\Omega_0(G)$ which intersects $\tilde{\delta}$. Let h be a parabolic element which is conjugate to $\rho(\alpha_1)$ in G and stabilizing $\tilde{\alpha}_1$. By adjoining the fixed point of h to $\tilde{\alpha}_1$, we obtain a simple closed curve, which divides \hat{C} into two domains. Let D be one of the

two domains such that D satisfies $D \cap g(D) = \emptyset$ if $k > 1$. Let η_1 be a connected component of $D \cap \tilde{\delta}$ and β be the arc in $\tilde{\alpha}_1$ which connects the end points of η_1 . Let $\tilde{\delta}_1 = \eta_1 \cup \beta$ and let $\tilde{\delta}_2$ be the closed curve $\tilde{\delta}$ with $\eta_1, g(\eta_1), \dots, g^{k-1}(\eta_1)$ replaced by $\beta, g(\beta), \dots, g^{k-1}(\beta)$. Then, for $j = 1, 2$, $\tilde{\delta}_j$ projects to a simple closed curve δ_j on X such that $p \mid \tilde{\delta}_j: \tilde{\delta}_j \rightarrow \delta_j$ is a finite-sheeted covering map. Moreover, note that $i(\delta_j, \lambda)$ is strictly less than $i(\delta, \lambda)$ for $j = 1, 2$. Since δ is non-trivial and $\delta = \delta_1 \cdot \delta_2$, either δ_1 or δ_2 are non-trivial. After a finite number of steps as above, we obtain a non-trivial simple closed curve δ' such that $i(\delta', \lambda) = 0$ and that, for a connected component $\tilde{\delta}'$ of $p^{-1}(\delta')$, $p \mid \tilde{\delta}': \tilde{\delta}' \rightarrow \delta'$ is a finite-sheeted covering map. This is a contradiction. \square

Proof of Theorem 6.1. Let S be a closed surface of genus $g \geq 2$. Let $[\rho, G]$ be an element of S_X . We claim that there exists an element $\sigma \in \text{Mod}(S)$ such that the sequence $\{\sigma^n([\rho])\}_{n \in \mathbb{Z}}$ converges to some maximal cusp $[\rho_\infty] \in \partial B_X$ as $|n| \rightarrow \infty$. If it has been shown, the similar argument in Theorem 5.5 reveals that the claim of the theorem holds: In fact, for any element $[\rho'] \in \partial B_X$, there exists a sequence $\{\tau_n\}$ in $\text{Mod}(S)$ such that $\tau_n([\rho_\infty])$ converges to $[\rho']$ by Theorem 5.5. Since the action of $\text{Mod}(S)$ is continuous at maximal cusps (Corollary 3.2), we can find a sequence in S_X which converges to $[\rho']$ by a diagonal argument.

Now we will show that there exists an element $\sigma \in \text{Mod}(S)$ such that the sequence $\{\sigma^n([\rho])\}_{n \in \mathbb{Z}}$ converges to some maximal cusp $[\rho_\infty] \in \partial B_X$ as $|n| \rightarrow \infty$. (Most of the following argument is similar to that of the proof of Proposition 5.1.) Note that the Kleinian manifold $N_G = (\mathbf{H}^3 \cup \Omega(G))/G$ is homeomorphic to a handlebody H_g of genus g whose boundary ∂H_g is homeomorphic to S . Under the identification $G = \pi_1(H_g)$, we have a proper embedding $\Psi: AH(H_g) \rightarrow R(S)$ where Ψ maps the conjugacy class of $\chi: \pi_1(H_g) \rightarrow \text{PSL}_2(\mathbb{C})$ to the conjugacy class of $\chi \circ \rho: \pi_1(S) \rightarrow G = \pi_1(H_g) \rightarrow \text{PSL}_2(\mathbb{C})$. By identifying $AH(H_g)$ with its image $\Psi(AH(H_g))$, we regard $AH(H_g)$ as a subset of $R(S)$.

Let Σ be a compact oriented surface with boundary $\partial \Sigma$ such that $\Sigma \times I$ is homeomorphic to H_g . (For example, let Σ be a closed disk with g open disk removed.) We can find a pair (λ', λ'') of maximal curve systems on Σ which binds Σ (cf. Lemma 5.4). Using this pair, we define a maximal curve system λ on $S = \partial(\Sigma \times I) = \partial H_g$, as

$$\lambda = (\lambda' \times \{0\}) \cup (\lambda'' \times \{1\}) \cup \left(\partial \Sigma \times \left\{ \frac{1}{2} \right\} \right).$$

Then (H_g, λ) is doubly incompressible by Lemma 4.2 and hence $AH(H_g, \lambda, K)$ is compact by Theorem 4.1. Put $\sigma = D_{\alpha_1} \circ \dots \circ D_{\alpha_N} \in \text{Mod}(S)$, where $\lambda = \{\alpha_j\}_{j=1}^N$. Now we consider the sequence $\{[\bar{\rho}_n] = \text{qc}([\rho], \sigma^{-n}X)\}_{n \in \mathbb{Z}}$ in $AH(H_g) \subset R(S)$. Then we can see that the sequence $\{[\bar{\rho}_n]\}_{n \in \mathbb{Z}}$ is contained in a compact set $AH(H_g, \lambda, K) \subset R(S)$

for some $K > 0$ since we have

$$\text{length}_{\bar{\rho}_n}(\lambda) \leq R \cdot \text{length}_{\sigma^{-n}X}(\lambda) = R \cdot \text{length}_X(\lambda).$$

from Lemma 5.2 for some $R > 0$. Therefore the sequence $\{[\bar{\rho}_n]\}_{n \in \mathbb{Z}}$ has a convergent subsequence.

On the other hand, since C_X is compact, $\{[\rho_n] = \sigma^n([\rho])\}_{n \in \mathbb{Z}}$ also has a convergent subsequence. Take representatives ρ_n of $[\rho_n] = \sigma^n([\rho])$ so that the sequence $\{\rho_n\}$ converges to a representation ρ_∞ . Since $[\rho_n] = [\bar{\rho}_n] \circ \sigma_*^{-n}$, we may assume that $\bar{\rho}_n = \rho_n \circ \sigma_*^n$. In addition, there are elements $\psi_n \in \text{PSL}_2(\mathbb{C})$ such that the sequence $\{\psi_n \cdot \bar{\rho}_n \cdot \psi_n^{-1}\}$ converges to a representation $\bar{\rho}_\infty$.

For any component α of λ , we claim that $\rho_\infty(\alpha)$ is a parabolic element. But this can be seen from the same argument in the proof of Proposition 5.1. Therefore, we leave the proof to the reader.

Since $\rho_\infty(\alpha)$ are parabolic for any component α of λ , we can conclude that $[\rho_\infty]$ is a maximal cusp in ∂B_X whose accidental parabolic locus is λ by Lemma 6.2. \square

7. Some property of the set of Schottky groups

Let S be a closed surface of genus ≥ 2 . In this section, we collect some property of S_X which can be easily seen. For a representation ρ of $\pi_1(S)$ onto a Kleinian group G , we denote by N_G its Kleinian manifold $(\mathbb{H}^3 \cup \Omega(G))/G$.

Lemma 7.1. *The mapping class group $\text{Mod}(S)$ acts on S_X transitively; that is, $S_X = \{\sigma([\rho])\}_{\sigma \in \text{Mod}(S)}$ for any $[\rho] \in S_X$.*

Proof. Let $[\rho_1, G_1]$ and $[\rho_2, G_2]$ be arbitrary two elements of S_X . Then there exists a homeomorphism $N_{G_1} \rightarrow N_{G_2}$ such that the restriction of this map to the boundaries is a quasiconformal map $\Omega_0(G_1)/G_1 \rightarrow \Omega_0(G_2)/G_2$. Now one can see that $[\rho_2] = \sigma([\rho_1])$, where $\sigma \in \text{Mod}(S)$ is the isotopy class of a homeomorphism of S induced by the quasiconformal map. \square

A Kleinian group is called *geometrically finite* if it has a finite sided convex fundamental polyhedron in \mathbb{H}^3 .

Lemma 7.2 (Hejhal [11], Matsuzaki [23]). *Each element $[\rho] \in S_X$ is an isolated point in C_X . On the other hand, if a torsion-free, geometrically finite element $[\rho] \in C_X$ is isolated in C_X , then $[\rho] \in S_X$.*

Proof. The first statement is due to Hejhal [11], who showed that any element $[\rho, G] \in \hat{C}_X$ such that G is a Schottky group is isolated in \hat{C}_X . Conversely, let take

an element $[\rho] \in C_X$ which is isolated in C_X . Since the same argument of Lemma 2.1 reveals that $\hat{C}_X - C_X$ is closed, $[\rho]$ is also isolated in \hat{C}_X . It was shown by Matsuzaki ([23, Theorem 3]) that, if a torsion-free, geometrically finite element $[\rho] \in \hat{C}_X$ is isolated in \hat{C}_X , then $[\rho]$ is a Schottky group. Thus, the second statement is proved. \square

REMARK. In Matsuzaki [23], obtained is a necessary and sufficient condition for a (not necessarily torsion-free) geometrically finite element of \hat{C}_X to be isolated in \hat{C}_X .

For $[\rho] \in S_X$, the following lemma gives a characterization of the elements of $\text{Mod}(S)$ which stabilize $[\rho]$.

Lemma 7.3. *Let $[\rho, G] \in S_X$ and $\sigma \in \text{Mod}(S)$. Then the following are equivalent:*

- (1) $\sigma([\rho]) = [\rho]$,
- (2) $\sigma_*(\ker \rho) = \ker \rho$, and
- (3) σ can be extended to a homeomorphism of the Kleinian manifold N_G , where σ is regarded as a homeomorphism of $X = \partial N_G$.

Proof. (1) \Rightarrow (2) and (3) \Rightarrow (2) are trivial. (2) \Rightarrow (1) can be seen from Matsuzaki [23, Theorem 2] and Lemma 7.2. We will show that (2) \Rightarrow (3). Let (f, ρ) be the projective structure corresponding to $[\rho]$. We may assume that $\sigma: X \rightarrow X$ is a quasiconformal map. Let $\tilde{\sigma}: \Delta \rightarrow \Delta$ be a lift of $\sigma: X \rightarrow X$. If $\sigma_*(\ker \rho) = \ker \rho$, then $\tilde{\sigma}$ descends to a quasiconformal map $\hat{\sigma}: f(\Delta) \rightarrow f(\Delta)$, because the covering group $f: \Delta \rightarrow f(\Delta)$ is $\ker \rho$. Since $G = \rho(\pi_1(S))$ is geometrically finite and $\Omega(G) = f(\Delta)$, Marden's isomorphism theorem [19] implies that $\hat{\sigma}$ can be extended to a G -compatible quasiconformal automorphism of $\hat{\mathbf{C}}$. This quasiconformal map can be extended to a G -compatible homeomorphism of $\mathbf{H}^3 \cup \hat{\mathbf{C}}$, which descends to a homeomorphism of N_G (cf. [7]). \square

References

- [1] W. Abikoff: *On boundaries of Teichmüller spaces and on Kleinian groups: III*, Acta Math. **134** (1975), 211–234.
- [2] L. Bers: *Simultaneous uniformization*, Bull. Amer. Math. Soc. **66** (1960), 94–97.
- [3] L. Bers: *On boundaries of Teichmüller spaces and on Kleinian groups: I*, Ann. of Math. **91** (1970), 570–600.
- [4] L. Bers: *The action of the modular group on the complex boundary*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference, Ann. of Math. Stud. **97**, Princeton Univ. Press, Princeton, 1981, 33–52.
- [5] J.F. Brock: *Iteration of mapping classes on a Bers slice: examples of algebraic and geometric*

- limits of hyperbolic 3-manifolds*, Lipa's legacy (New York, 1995), Contemporary Math. **211**, Amer. Math. Soc., Providence, 1997, 81–106.
- [6] R.D. Canary: *The Poincaré metric and a conformal version of a theorem of Thurston*, Duke Math. J. **64** (1991), 349–359.
 - [7] A. Douady and C.J. Earle: *Conformally natural extension of homeomorphisms of the circle*, Acta Math. **157** (1986), 23–48.
 - [8] A. Fathi, F. Raudenbach and V. Poenaru: *Travaux de Thurston sur le surfaces*, Astérisque **66–67**, Société Mathématique de France, Paris, 1979.
 - [9] D.M. Gallo: *Some special limits of Schottky groups*, Proc. Amer. Math. Soc. **118** (1993), 877–883.
 - [10] D.M. Gallo: *Schottky groups and the boundary of Teichmüller space: genus 2*, The mathematical legacy of Wilhelm Magnus: groups, geometry and special functions (Brooklyn, NY, 1992), Contemporary Math. **169**, Amer. Math. Soc., Providence, 1994, 283–305.
 - [11] D.A. Hejhal: *On Schottky and Koebe-like uniformizations*, Duke Math. J. **55** (1987), 267–286.
 - [12] T. Jørgensen: *On discrete groups of Möbius transformations*, Amer. J. Math. **98** (1976), 739–749.
 - [13] S.P. Kerckhoff and W.P. Thurston: *Non-continuity of the action of the modular group at Bers' boundary of Teichmüller space*, Invent. Math. **100** (1990), 25–47.
 - [14] I. Kra: *Deformations of Fuchsian groups*, Duke Math. J. **36** (1969), 537–546.
 - [15] I. Kra: *A generalization of a theorem of Poincaré*, Proc. Amer. Math. Soc. **27** (1971), 299–302.
 - [16] I. Kra: *Deformations of Fuchsian groups*, II, Duke Math. J. **38** (1971), 499–508.
 - [17] I. Kra: *On spaces of Kleinian groups*, Comm. Math. Helv. **47** (1972), 53–69.
 - [18] I. Kra and B. Maskit: *Remarks on projective structures*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference, Ann. of Math. Stud. **97**, Princeton Univ. Press, Princeton, 1981, 343–359.
 - [19] A. Marden: *The geometry of finitely generated Kleinian groups*, Ann. of Math. **99** (1974), 384–462.
 - [20] B. Maskit: *On boundaries of Teichmüller spaces and on kleinian groups*: II, Ann. of Math. **91** (1970), 607–639.
 - [21] B. Maskit: *Self-maps of Kleinian groups*, Amer. J. Math. **93** (1971), 840–856.
 - [22] B. Maskit: *Kleinian groups*, Springer-Verlag, Berlin, 1988.
 - [23] K. Matsuzaki: *Projective structures inducing covering maps*, Duke Math. J. **78** (1995), 413–425.
 - [24] C.T. McMullen: *Cusps are dense*, Ann. of Math. **133** (1991), 217–247.
 - [25] K. Ohshika: *Geometrically finite Kleinian groups and parabolic elements*, Proc. Edinburgh Math. Soc. **41** (1998), 141–159.
 - [26] J.P. Otal: *Sur le bord du prologement de Bers de l'espace de Teichmüller*, C.R. Acad. Sci. Paris, **316** (1993), 157–160.
 - [27] D.P. Sullivan: *On the ergodic theory at infinity of an arbitrary discrete group of hyperbolic motions*, Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference, Ann. of Math. Stud. **97**, Princeton Univ. Press, Princeton, 1981, 465–496.
 - [28] W.P. Thurston: *Hyperbolic structures on 3-manifolds, III: Deformations of 3-manifolds with incompressible boundary*, preprint.

Graduate School of Mathematics
 Nagoya University
 Nagoya 464-8602, Japan
 e-mail: itoken@math.nagoya-u.ac.jp