

Title	Stonian spaces and the second conjugate spaces of AM spaces
Author(s)	Wada, Junzo
Citation	Osaka Mathematical Journal. 1957, 9(2), p. 195-200
Version Type	VoR
URL	<a href="https://doi.org/10.18910/12339">https://doi.org/10.18910/12339</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## ***Stonian Spaces and the Second Conjugate Spaces of AM Spaces***

By Junzo WADA

Let  $X$  be a compact space and let  $C(X)$  be the set of all real-valued continuous functions on  $X$ . If any non-void subset of  $C(X)$  with an upper bound has a least upper bound in  $C(X)$ , such a compact space  $X$  is called a stonian space.<sup>1)</sup> Stone [10] has shown that a compact space  $X$  is stonian if and only if it is extremally disconnected, that is to say, if for any open set  $U$  in  $X$  its closure  $\bar{U}$  is open. While, Kelley [9] has proved that if for any Banach space  $F$  containing a Banach space  $E$  there exists a projection of  $F$  on  $E$  whose norm is 1,  $E$  is isometric to  $C(X)$ , where  $X$  is stonian. Also Dixmier [4] has considered a compact space  $X$  such that  $C(X)$  is isomorphic to an  $L^\infty(R, \mu)$  as Banach algebras, where  $R$  is a locally compact space and  $\mu$  is a positive measure on  $R$ . He called such a space  $X$  a hyperstonian space. A hyperstonian space is stonian. We shall see that a compact space  $X$  is hyperstonian if and only if  $C(X)$  is lattice-isomorphic and isometric to a conjugate space of an  $AL$  space.<sup>2)</sup>

In §1 we state some general properties of stonian spaces, and in §2 we consider an  $AM$  space  $C(X)$  which is the second conjugate space of an  $AM$  space. Such a space  $X$  is hyperstonian, and if the character of  $X$  is countable, then  $X$  is the space  $\beta N_0$ , where  $N_0$  is a discrete space whose cardinal number is at most countable (cf. Theorem 3, Corollary).

### **§ 1. Stonian spaces**

For a completely regular space  $X$ , let  $\beta X$  denote the Čech compactification of  $X$ . (cf. Čech [2]). Dixmier [4] has shown that  $\beta U = X$  for any open dense set  $U$  in a stonian space  $X$ . Therefore we obtain easily the following:

---

1) See Stone [10] and Dixmier [4]. Numbers in bracket refer to the reference cited at the end of the paper.

2) See Kakutani [7] and [8].

- (i)  $X$  is stonian if and only if for any open set  $U$  in  $X$ ,  $\beta U \subset X$ .
- (ii) If  $X$  is a stonian space and if  $X$  has an open dense subspace which satisfies the 1st axiom of countability,  $X$  is a space  $\beta N$ , where  $N$  is a discrete space.
- (iii) If  $X$  is a stonian space and if a point  $x$  of  $X$  is not isolated, then  $\{x\}$  is not a  $G_\delta$  set.

We shall prove here the following theorem.

**Theorem 1.** *Let  $X$  be a stonian space. If  $X$  is a product  $R \times R$  of a compact space  $R$ , then  $X$  is finite.<sup>3)</sup>*

Proof. Suppose that a stonian  $X$  be of the form  $R \times R$ , where  $R$  is a compact space with infinite points. Then  $R$  is stonian. Let  $U$  be a dense open set in  $R$ . Then  $R \times R = \beta U \times \beta U \supset U \times U$  and  $U \times U$  is a dense open set in  $R \times R$ . Since  $R \times R$  is stonian,  $\beta(U \times U) = R \times R = \beta U \times \beta U$ .

Now if for a fully normal space<sup>4)</sup>  $S$   $\beta(S \times S) = \beta S \times \beta S$ , then  $S$  is compact. (Ishiwata [6]). Therefore it is sufficient to prove that there exists an open dense subset  $U$  in  $R$  which is not compact but fully normal. We shall construct such an open set  $U$ . Since  $R$  is an infinite set, there exists a countable family  $\{U_i\}_{i=1}^{\infty}$  of mutually disjoint non void sets which are both open and closed. Let  $V = R - \sum_{i=1}^{\infty} U_i$  and let  $U = V \cup \sum_{i=1}^{\infty} U_i$ . Since the set  $U$  is the union of a countable family of open and compact sets in  $R$  which are mutually disjoint,  $U$  is fully normal. Clearly,  $U$  is not compact and is dense in  $R$ .

REMARK. Theorem 1 shows that there exists no stonian space  $S$  with infinite points of the form  $R \times R$ . But we can find easily a totally disconnected compact space  $S$  which is a product space  $R \times R$ , where  $R$  is compact and infinite.

## § 2. Second conjugate spaces of $AM$ spaces.

Let  $X$  be a stonian space and let  $M(X)$  be the set of all measures on  $X$ . A positive measure  $\mu$  on  $X$  is called a *normal measure* if for any nowhere dense set  $A$ ,  $\mu(A) = 0$ . A real measure  $\mu$  on  $X$  is called *normal* if its positive part and its negative part are both normal. Let  $M'(X)$

3) Henrikson and Isbell announced the following theorem (Bull. Amer. Soc. Vol. 63. 1957 Abstract): if  $X$  and  $Y$  are infinite completely regular spaces such that  $\beta(X \times Y) = \beta X \times \beta Y$ , then  $X \times Y$  is pseudo-compact, that is, any continuous function on  $X \times Y$  is always bounded. If we make use of this theorem, we obtain moreover that if a stonian space  $X$  is a product  $R \times S$  of compact spaces  $R$  and  $S$ , then either  $R$  or  $S$  is finite.

4) See Tukey [11], p. 53.

denote the set of all normal measures on  $X$ . A stonian space  $X$  is called *hyperstonian* if it has positive normal measures, the union of whose carriers is dense in  $X$ . We shall see that a compact space  $X$  is hyperstonian if and only if  $C(X)$  is lattice-isomorphic and isometric to the conjugate space of an  $AL$  space. Let  $E$  be an  $AM$  space. Then the second conjugate space of  $E$  is lattice-isomorphic and isometric to  $C(X)$ , where  $X$  is hyperstonian, and the conjugate space of  $E$  is lattice-isomorphic and isometric to  $M'(X)$ .  $M'(X)$  is also lattice-isomorphic and isometric to an  $L^1(\Omega, \mu)$ , where  $\Omega$  is an open dense set in  $X$  and  $\mu$  is a suitable positive measure on  $\Omega$ . (cf. [4]). We consider now an  $AM$  space  $C(X)$  which is the second conjugate space of an  $AM$  space. Let  $E$  be a Banach space and  $E^*$ ,  $E^{**}$  denote the conjugate space of  $E$  and the second conjugate space of  $E$  respectively. For any closed linear subspace  $V$  in  $E^*$  we define its *characteristic*  $r$  by  $r = \inf_{x \in E} \sup_{f \in V \cap S} \frac{|f(x)|}{\|x\|}$ , where  $S$  is a unit sphere in  $E^*$ . A closed linear subspace  $V$  in  $E^*$  is called *minimally weakly dense* if it is weakly dense in  $E^*$  and if any other closed subspace in  $V$  is not weakly dense in  $E^*$ .

The following lemma was proved by Dixmier [3].

**Lemma.** (i) *Let  $E$  be a Banach space. Then  $E$  is a minimally weakly dense subspace in  $E^{**}$  which is characteristic one.*

(ii) *If  $V$  is a minimally weakly dense subspace in  $E^*$  which is characteristic one, then  $E^{**} = E \oplus V^+$  and  $\|x\| \leq \|x+z\|$  for  $x \in E$ ,  $z \in V^+$ , where  $V^+$  denotes the set  $\{z | z \in E^{**}, z(f) = 0 \text{ for any } f \in V\}$ .*

Let  $K$  be an open set in a hyperstonian space. Then the *character* of  $K$  is said to be countable if any family of non-void open and closed sets in  $K$  which are mutually disjoint is at most countable.

We can prove the following theorem. Hereafter  $X$  denotes a hyperstonian space.

**Theorem 2.**  *$C(X)$  is lattice-isomorphic and isometric to the second conjugate space of an  $AM$  space with a unit<sup>2)</sup> if and only if there exists a lattice-closed and (topologically) closed linear subspace  $V$  in  $C(X)$  which has constant functions such that*

- (I) *for any  $f \in C(X)$  and for any open and closed set  $K$  (in  $X$ ) whose character is countable, there exists a sequence  $\{f_n\}$  in  $V$  such that  $f_n$  pointwise converges to  $f$  on  $K$  except a nowhere dense set.*
- (II)  *$V$  is a minimal closed linear space which has the property (I): any other closed subspace in  $V$  does not satisfy (I).*

Proof. (a) Let  $C(X)$  be lattice-isomorphic and isometric to  $E^{**}$

and let  $E$  be an  $AM$  space with a unit. Then, by Lemma, there exists a minimally weakly dense subspace  $V$  in  $C(X)$  which is lattice-closed. We see here that  $E$  and  $E^*$  are lattice-isomorphic and isometric to  $V$  and  $M'(X)$  respectively. Since  $E$  has a unit, we can assume that  $V$  has constant functions in  $C(X)$ . In order to prove (I) and (II), we are only to prove the equivalence of (I) and that  $V$  is weakly dense. Now if the property (I) is satisfied, then we see easily that  $V$  is weakly dense. Conversely, if  $V$  is weakly dense, then we see easily that for any  $f \in C(X)$  and for any open and closed set  $K$  (in  $X$ ) whose character is countable, there exist  $f_n \in V$  such that

$$\int_{K \cap \Omega} |f(x) - f_n(x)| d\mu(x) < \frac{1}{n} \quad (n = 1, 2, \dots),$$

where  $\Omega$  is an open dense set in  $X$  and  $\mu$  is a suitable positive measure on  $\Omega$ . Therefore, as is well known, a subsequence  $f_{n_i}$  of  $\{f_n\}$  pointwise converges to  $f$  almost everywhere on  $K \cap \Omega$ .<sup>5)</sup> Since any set of measure null on  $K \cap \Omega$  is nowhere dense,  $f_{n_i}$  pointwise converges to  $f$  on  $K$  except a nowhere dense set.

(b) If properties (I) and (II) are satisfied, we see easily that  $V$  is a minimally weakly dense subspace in  $C(X)$ . (cf. (a)). We shall prove that  $V$  is of characteristic one. For any  $u \in M'(X)$ , let  $A$  and  $B$  be carriers of the positive part  $u^+$  of  $u$  and of the negative part  $u^-$  of  $u$  respectively and let the function  $f$  take the value 1 on  $A$  and the value  $-1$  on  $B$ . Then, by (I), there exists a sequence  $\{f_n\}$  in  $V$  such that  $f_n$  pointwise converges to  $f$  on  $A \cup B$  except a nowhere dense set. We may assume here that for any  $n$ ,  $\|f_n\|_\infty \leq 1$ , since  $V$  has constant functions.<sup>6)</sup> Since  $u(f_n)$  converges to  $u(f) = \|u\|$ ,  $V$  is of characteristic one. By Lemma,  $M(X) = M'(X) \oplus V^+$  and  $\|u\| \leq \|u+z\|$  for  $u \in M'(X)$ ,  $z \in V^+$ . Therefore if  $F$  is a linear functional on  $V$ , then there exists  $u \in M'(X)$  such that  $F(f) = u(f)$  for any  $f \in V$  and  $\|F\| = \|u\|$ , that is,  $M'(X)$  is lattice-isomorphic and isometric to  $V^*$ , and  $C(X)$  is lattice-isomorphic and isometric to  $V^{**}$ . This concludes the proof.

We consider next an  $AL$  space with an  $F$ -unit. Let  $l^1$  be the set of all sequences  $\{\xi_i\}$  of real numbers with convergent  $\sum_{i=1}^\infty |\xi_i|$ .  $l^1$  is a Banach space where the norm of  $x = \{\xi_i\} \in l^1$  is  $\sum_{i=1}^\infty |\xi_i|$ . (cf. Banach [1]).

**Theorem 3.** *If an  $AL$  space  $E$  with an  $F$ -unit<sup>2)</sup> is lattice-isomorphic*

5) See, for example, Halmos [5].

6)  $\|f\|_\infty$  denotes the uniform norm:  $\|f\|_\infty = \sup_{x \in X} |f(x)|$ .

and isometric to a conjugate space of an AM space,  $E$  is lattice-isomorphic and isometric to  $l^1$ .

Proof. Let  $E$  be of the form  $L^1(\Omega, \mu)$ , where  $\Omega$  is an open set in a hyperstonian space. Since  $E$  has an  $F$ -unit, the character of  $\Omega$  is countable. If  $L^1(\Omega, \mu)$  is the conjugate space of an AM space  $F$  and if  $F$  is of the form of  $C(Y, y_\alpha, y'_\alpha, \lambda_\alpha) = \{f \mid f \in C(Y), f(y_\alpha) = \lambda_\alpha f(y'_\alpha), 0 \leq \lambda_\alpha < 1, \alpha \in m\}$ <sup>7)</sup>, then function  $g_\alpha$  in  $L^1(\Omega, \mu)$  which correspond to  $\mu_{y'_\alpha} \in F^*$  are mutually distinct, where  $\mu_{y'_\alpha}$  is a dirac measure, that is to say  $\mu_{y'_\alpha}(f) = f(y'_\alpha)$  for any  $f \in F$ . We see easily that the carrier of function  $g_\alpha$ <sup>8)</sup> is a one-point set  $x_\alpha$ , and therefore,  $x_\alpha$  is an isolated point in  $\Omega$ . Since the character of  $\Omega$  is countable, the cardinal number of  $Z_0 = \{y'_\alpha\}_{\alpha \in m}$  is at most countable. Since  $C(Y) \supset F$ , any linear functional  $\xi$  on  $F$  can be extended to a linear functional  $\xi'$  on  $C(Y)$ .  $\xi'$  is a measure on  $Y$  and for any  $f \in F$ ,  $\xi(f) = \xi'(f) = \int_Y f(x) d\xi'(x)$ . Since the cardinal number of  $Y_0$  is countable, we can put  $Z_0 = \{z_1, z_2, \dots\}$ . For any  $n$ , let  $Y_n$  denote the set of  $y_\beta$  with  $y'_\beta = z_n$ . Then we have  $\xi(f) = \sum_{i=1}^{\infty} \int_{Y_n} f(y) d\xi'(y) = \sum_{n=1}^{\infty} \left( \int_{Y_n \ni y_\beta} \lambda_\beta d\xi'(y_\beta) \right) f(z_n)$ . If we put  $p_n = \int_{Y_n \ni y_\beta} \lambda_\beta d\xi'(y_\beta)$ , we obtain that  $\xi(f) = \sum_{n=1}^{\infty} p_n f(z_n)$ . We see easily that if  $\xi$  is positive, any  $p_n$  is non-negative and  $\|\xi\| = \sum_{n=1}^{\infty} p_n$ .

**Corollary.** *If  $C(X)$  is lattice-isomorphic and isometric to the second conjugate space of an AM space and if the character of  $X$  is countable, then  $X$  is the space  $\beta N_0$ , where  $N_0$  is a discrete space whose cardinal number is at most countable.*

(Received October 10, 1957)

### Bibliography

- [ 1 ] S. Banach: Théorie des opérations linéaires, 1932.
- [ 2 ] E. Čech: On bicomact spaces, Ann. of Math. **38** (1937).
- [ 3 ] J. Dixmier: Sur un théorème de Banach, Duke Math. J. **15** (1948).
- [ 4 ] ———: Sur certains espaces considérés par M. H. Stone, Summa Brasil. Math. **2** (1951).
- [ 5 ] P. R. Halmos: Measure theory, 1954.
- [ 6 ] T. Ishiwata: Uniform spaces with complete structures, Sûgaku Kenkyûroku **1** (1952) (Japanese).

7) See Kakutani [8], Theorem 1.

8) We may assume that  $g_\alpha$  is a continuous function on  $\Omega$ . See [4].

- [ 7 ] S. Kakutani: Concrete representation of abstract (L)-spaces and the mean ergodic theorem, *Ann. of Math.* **42** (1941)
- [ 8 ] —————: Concrete representation of abstract (M)-spaces, *Ann. of Math.* **42** (1941).
- [ 9 ] J. L. Kelly: Banach spaces with the extension property, *Trans. Amer. Math. Soc.* **72** (1952).
- [10] M. H. Stone: Boundedness properties in function-lattices, *Canad. J.* **1** (1949).
- [11] J. W. Tukey: *Convergence and uniformity in Topology*, 1940.