



Title	Strongly semiprime rings and nonsingular quasi-injective modules
Author(s)	Kutami, Mamoru; Oshiro, Kiyochi
Citation	Osaka Journal of Mathematics. 1980, 17(1), p. 41-50
Version Type	VoR
URL	https://doi.org/10.18910/12487
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

STRONGLY SEMIPRIME RINGS AND NONSINGULAR QUASI-INJECTIVE MODULES

MAMORU KUTAMI AND KIYOICHI OSHIRO

(Received February 3, 1979)

Following Handelman [8] we call a ring R is a right strongly semiprime ring provided if I is a two-sided ideal of R and is essential as a right ideal, then it contains a finite subset whose right annihilator is zero.

In this paper, we first show that a ring R is a right strongly semiprime ring if and only if

- (1) $Q(R)$ is a direct sum of simple rings, and
- (2) $eQ(R)eR = eQ(R)$ for all idempotents e in $Q(R)$ where $Q(R)$ denotes the maximal ring of right quotients of R .

Using these conditions (1) and (2), we shall investigate the following conditions:

- (a) Every nonsingular quasi-injective right R -module is injective.
- (b) Any finite direct sum of nonsingular quasi-injective right R -modules is quasi-injective.
- (c) Any direct sum of nonsingular quasi-injective right R -modules is quasi-injective.
- (d) Any direct product of nonsingular quasi-injective right R -modules is quasi-injective.

It is shown that the conditions (a), (b) and (d) are equivalent; indeed, the rings satisfying one of these conditions are determined as rings R such that $R/G(R)$ is a right strongly semiprime ring, where $G(R)$ denotes the right Goldie torsion submodule of R . A ring R satisfying the condition (c) is also characterized as a ring R such that $R/G(R)$ is a semiprime right Goldie ring.

1. Preliminaries and notations

Throughout this paper all rings considered have identity and all modules are unitary.

Let R be a ring. $Q(R)$ denotes its maximal ring of right quotients. Let M be a right R -module. By $E_R(M)$, nM , $Z(M)$ and $G(M)$ we denote its injective hull, the direct product of n -copies, its singular submodule and its Goldie torsion submodule, respectively. (Note that $Z(M/Z(M)) = G(M)/Z(M)$.) For

a given two right R -modules N and M , we adopt the symbol $N \subseteq M$ to denote the fact that N is isomorphic to a submodule of M , and use the symbol $N \subseteq_e M$ to indicate N to be an essential submodule of M .

Now, for a nonsingular right R -module M , the following statements hold:

- (1) $MG(R)=0$; so M become a right $R/G(R)$ -module by usual way,
- (2) M is also nonsingular as a right $R/G(R)$ -module, and
- (3) M is R -injective (R -quasi-injective) if and only if M is $R/G(R)$ -injective ($R/G(R)$ -quasi-injective).

Noting that $R/G(R)$ is a right nonsingular ring, we conclude from [4, Theorem 2.2] that any nonsingular injective right R -module has a unique right $Q(R/G(R))$ -module structure compatible with the R -module structure. So, for a nonsingular right R -module M , we have $M \subseteq_e MQ(R/G(R)) \subseteq_e E_R(M)$.

It is well known (e.g. [4, Theorem 3.2]) that every finitely generated nonsingular right module over a right self-injective regular ring is both projective and injective. Therefore, if M is a finitely generated nonsingular injective right R -module, then M is both $Q(R/G(R))$ -projective and $Q(R/G(R))$ -injective.

For a subset S of a ring R , $(0:S)_R^r((0:S)_R^l)$ denotes the right (left) annihilator of S in R .

Lemma 1.1. *Let R be a ring and set $\bar{R}=R/G(R)$ and $Q=Q(\bar{R})$. If M is a nonsingular right Q -module, then the following statements hold:*

- (a) *M is nonsingular as a right R -module. (Of course, M becomes a right R -module by a natural way.)*
- (b) *M is Q -quasi-injective if and only if M is R -quasi-injective.*

Proof. (a) Let x be an element in M such that $(0:x)_R^r \subseteq_e R$. Inasmuch as $G(R) \subseteq (0:x)_R^r \subseteq_e R$, we see from [4, Proposition 1.28] that $(0:x)_R^r \subseteq_e \bar{R}$. Hence it follows $(0:x)_Q^r \subseteq_e Q$, whence $x=0$.

(b) Clearly $M \subseteq_e E_{\bar{R}}(M)$ as a right Q -module. It is also easily seen that $M \subseteq_e E_Q(M)$ as a right \bar{R} -module. As a result we get $E_{\bar{R}}(M)=E_Q(M)$, whence $E_R(M)=E_Q(M)$. On the other hand we see that $End_R(E_R(M))=End_{\bar{R}}(E_{\bar{R}}(M))=End_Q(E_{\bar{R}}(M))$ and $End_R(E_Q(M))=End_{\bar{R}}(E_Q(M))=End_Q(E_Q(M))$; consequently $End_R(E_R(M))=End_Q(E_Q(M))$, where $End_*(\#)$ denotes the endomorphism ring of a right $*$ -module $\#$. The proof is now easily done by applying the well known fact that a module is quasi-injective if and only if it is a fully invariant submodule of its injective hull.

The following lemma is frequently used in this paper.

Lemma 1.2. *If M is a quasi-injective right R -module such that $R \subseteq_n nM$ for some positive integer n , then M is injective.*

Proof. By virtue of Harada [9, Proposition 2.4], nM is also quasi-injective.

Hence we can easily see from $R \subseteq_n nM$ that nM is injective, whence so is M .

2. Strongly semiprime rings

We recall some definitions introduced by Handelman and Lawrence [7] and Handelman [8]. An right ideal I of a ring R is *insulated* if there exists a finite set $\subseteq I$ whose right annihilator in R is zero. For a non-zero element a in R , a finite set $\{r_1, \dots, r_n\} \subseteq R$ is a *right insulator of a* if the right annihilator of $\{ar_1, \dots, ar_n\}$ is zero. A ring R is said to be a right strongly prime ring provided every non-zero ideal of R is insulated as a right ideal, and said to be a right strongly semiprime ring if every ideal I of R with $I \subseteq_e R$ as a right ideal is insulated as a right ideal. As is easily seen, a ring R is right strongly prime if and only if every non-zero element in R has a right insulator.

The notion ‘insulated’ coincides with ‘cofaithful’ in Beachy-Blair [1] and is connected with ‘finite intersection property on annihilator right ideals’ in Zermanowitz [14]. The class of right strongly prime rings is just that of right absolutely torsion-free rings in the sense of Rubin [11]. For details of strongly prime rings and strongly semiprime rings, the reader is referred to [1], [6], [7], [8] and [11].

DEFINITION. For an element a in a ring R , we call a finite set $\{r_1, \dots, r_n; b\} \subseteq R$ is a right semi-insulator of a when $RaR \cap RbR = 0$ and the right annihilator of $\{ar_1, \dots, ar_n\} \cup bR$ is zero.

Proposition 2.1. *If R is a ring such that every element in R has a right semi-insulator, then R is a semiprime right nonsingular ring.*

Proof. Let $a \in R$. Then there exists a finite set $\{r_1, \dots, r_n; b\} \subseteq R$ satisfying $RaR \cap RbR = 0$ and $[\bigcap_{i=1}^n (0: ar_i)_R] \cap (0: bR)_R = 0$. If $a \in Z(R)$ and $a \neq 0$, then $ar_i \in Z(R)$ for each i and $0 \neq ar \in \bigcap_{i=1}^n (0: ar_i)_R$ for some $r \in R$. But it follows from $bRar = 0$ that $ar = 0$, a contradiction. If $aRa = 0$, then $a = 0$ because $a \in [\bigcap_{i=1}^n (0: ar_i)_R] \cap (0: bR)_R = 0$. Thus R is a semiprime right nonsingular ring.

Lemma 2.2. *Let R be a semiprime ring.*

(a) *If I is an ideal of R and J is a right ideal of R such that $I \cap J = 0$, then $I \cap RJ = 0$ and moreover $Q(R)IQ(R) \cap Q(R)JQ(R) = 0$.*

(b) *For ideals I and J of R , $I \subseteq_e J$ as a right ideal if and only if $I \subseteq_e J$ as a left ideal.*

(c) *If $\{I_\lambda \mid \lambda \in \Lambda\}$ is an independent family of ideals of R , then so is $\{Q(R)I_\lambda Q(R) \mid \lambda \in \Lambda\}$.*

Proof. (b) and (c) easily follow from (a).

(a). Set $Q=Q(R)$. Since $I \cap J=0$, we see $JI=0$ and it follows $(IQJ \cap R)^2=0$. Hence $IQJ=0$, from which we have $(QIQ \cap QJQ \cap R)^2=0$ and therefore $QIQ \cap QJQ=0$.

NOTE. Let I and J be ideals of a semiprime ring R . When we use ' $I \subseteq_e J$ ' instead of ' $I \subseteq_e J$ as a right ideal' or ' $I \subseteq_e J$ as a left ideal', no confusion arises by Lemma 2.2(b).

Proposition 2.3. *The following conditions are equivalent for a semiprime right nonsingular ring R :*

- (a) $Q(R)$ is a direct sum of prime rings.
- (b) The set of all central idempotents of $Q(R)$ is a finite set.
- (c) R contains no infinite direct sums of ideals.
- (d) Every ideal of R is essentially cyclic generated, i.e., if I is an ideal of R , then there exists a in I such that $RaR \subseteq_e I$.

Proof. Set $Q=Q(R)$. (a) \Rightarrow (b) is clear.

(b) \Rightarrow (c). Suppose that R contains an infinite independent set $\{I_\lambda | \lambda \in \Lambda\}$ of non-zero ideals. Lemma 2.2 (c) says that $\{QI_\lambda Q | \lambda \in \Lambda\}$ is independent and so is $\{E_Q(QI_\lambda Q) | \lambda \in \Lambda\}$. However, inasmuch as each $E_Q(QI_\lambda Q)$ is an ideal of Q , each $E_Q(QI_\lambda Q)$ is generated by a central idempotent in Q by [5, Corollary 1.10]. This contradicts (b).

(c) \Rightarrow (d). Let I be a non-zero ideal of R . For $0 \neq a_1 \in I$, if Ra_1R is not essential in I , we can take $0 \neq a_2$ in I such that $\{Ra_1R, Ra_2R\}$ is independent by Lemma 2.2(a). Similarly when $Ra_1R \oplus Ra_2R$ is not essential in I , then there exists a_3 in I such that $\{Ra_1R, Ra_2R, Ra_3R\}$ is independent. Continuing this manner, by (c), we must reach to n such that $\{Ra_1R, \dots, Ra_nR\}$ is independent and $Ra_1R \oplus \dots \oplus Ra_nR \subseteq_e I$. Here we claim $R(a_1 + \dots + a_n)R \subseteq_e I$. From Lemma 2.2(c), $\{Qa_1Q, \dots, Qa_nQ\}$ is independent. This implies $a_1Q \oplus \dots \oplus a_nQ = (a_1 + \dots + a_n)Q$ since Q is a regular ring. Hence we see $(Ra_1R \oplus \dots \oplus Ra_nR)Q = R(a_1 + \dots + a_n)Q$, which shows $R(a_1 + \dots + a_n)R \subseteq_e Ra_1R \oplus \dots \oplus Ra_nR$. Therefore surely $R(a_1 + \dots + a_n)R \subseteq_e I$.

(d) \Rightarrow (a). It is easily seen from (d) that Q is a direct sum of indecomposable rings, say $Q=Q_1 \oplus \dots \oplus Q_n$. To show that each Q_i is prime, let X be an ideal of Q_i . Then $E_{Q_i}(X)$ is generated by a central idempotent in Q_i by again [5, Corollary 1.10]. So, $X \subseteq_e Q_i$ as a right Q_i -module from which we see that Q_i is a prime ring.

REMARK. The equivalence of (a) and (b) is due to J. Kado (see [10, Proof of Proposition 3.2]).

Lemma 2.4 ([8]). *If R is a right strongly semiprime ring, then*

- (a) R is a semiprime right nonsingular ring, and

(b) $Q(R)$ is a direct sum of simple rings.

Proof. (a). Let I be an ideal of R such that $I^2=0$. Clearly $I^2=0$ implies $(0: I)_R^l \subseteq_e R$ as a right ideal. So $(0: I)_R^l$ is insulated as a right ideal. Inasmuch as $(0: I)_R^l I=0$, it follows $I=0$. Hence R is a semiprime ring. Since R is semiprime, using Lemma 2.2(a), there exists an ideal $K \subseteq R$ such that $Z(R) \oplus K \subseteq_e R$. Since $Z(R) \oplus K$ is insulated as a right ideal, there exists a finite set $\{z_1, \dots, z_n\} \subseteq Z(R)$ and $\{k_1, \dots, k_n\} \subseteq K$ such that $\bigcap_{i=1}^n (0: z_i + k_i)_R^r = (\bigcap_{i=1}^n (0: z_i)_R^r) \cap (\bigcap_{i=1}^n (0: k_i)_R^r)$. Let $a \in Z(R)$ and suppose $a \neq 0$. Then $0 \neq ar \in \bigcap_{i=1}^n (0: z_i)_R^r$ for some r in R . But, since each $k_i ar=0$, we infer $ar=0$, a contradiction. Thus $Z(R)=0$.

(b). Inasmuch as every non-zero essential ideal of R is insulated, clearly, R contains no infinite direct sums of non-zero ideals. Hence, by Proposition 2.3, $Q(R)$ is a direct sum of prime rings, say $Q(R)=Q_1 \oplus \dots \oplus Q_n$. In order to show that each Q_i is simple, let X_i be a non-zero ideal of Q_i , $i=1, \dots, n$. Since Q_i is a prime right self-injective regular ring, we see $X_i \subseteq_e Q_i$ by [5, Proposition 1.10]. As a result, $(X_1 \oplus \dots \oplus X_n) \cap R \subseteq_e R$. So $(X_1 \oplus \dots \oplus X_n) \cap R$ is insulated as a right ideal, whence $R \subseteq k((X_1 \oplus \dots \oplus X_n) \cap R) \subseteq k(X_1 \oplus \dots \oplus X_n)$ for some positive integer k . Since $X_1 \oplus \dots \oplus X_n$ is an ideal of Q , it is Q -quasi-injective and so is by Lemma 1.1, R -quasi-injective. Therefore we see that $X_1 \oplus \dots \oplus X_n$ is R -injective, whence $Q(R)=X_1 \oplus \dots \oplus X_n$. Therefore $Q_i=X_i$, $i=1, \dots, n$.

Theorem 2.5. For a given ring, R , the following conditions are equivalent:

- (a) R is a right strongly semiprime ring.
- (b) (1) $Q(R)$ is a direct sum of simple rings, and
(2) $Q(R)eR=Q(R)eQ(R)$, or equivalently, $eQ(R)eR=eQ(R)$ for all idempotents e in $Q(R)$.
- (c) (1) R contains no infinite direct sums of ideals,
(2) every element of R has a right semi-insulator.
- (d) $Q(R)I=Q(R)$ for any essential right ideal I of R .
- (e) There exists a ring extension S of R with the same identity satisfying $SI=S$ for any essential right ideal I of R .

Proof. Set $Q=Q(R)$. (a) \Rightarrow (b). According to Lemma 2.4, Q is a direct sum of simple rings. So every ideal of Q is a direct summand. Let $e=e^2 \in Q$ and take an ideal T of Q such that $QeQ \oplus T=Q$. Since $(QeR \cap R) \oplus (T \cap R)$ is essential in R , it is insulated as a right ideal, hence there exists a positive integer k such that $R \subseteq k((QeR \cap R) \oplus (T \cap R))$ as a right R -module. Since $QeR \oplus T$ is a left ideal of Q , $QeR \oplus T$ is Q -quasi-injective and so is R -quasi-injective (Lemma 1.1). Hence Lemma 1.2 says that $QeR \oplus T$ is R -injective, whence we have $QeR=QeQ$.

(b) \Rightarrow (c). In order to show R to be semiprime, let $a \in R$ such that $aRa = 0$. Since Q is a direct sum of simple rings, clearly it is a right nonsingular ring; whence it is a regular ring. Thus $Qa = Qe$ for some $e = e^2$ in Q . Since $QaR = QeR = QeQ = QaQ$, we see $0 = QaRa = QaQa$, from which we have $a = 0$. (1) now follows from Proposition 2.3. Let us write $Q = Q_1 \oplus \cdots \oplus Q_n$, where each Q_i is simple, and let $1 = e_1 + \cdots + e_n$ in this decomposition. $\{e_1, \dots, e_n\}$ is a set of non-zero central orthogonal idempotents. Now, to show (2), let $a \in R$. Then $QaR = QaQ = \sum_{i \in I} \oplus Q_i$ for some $I \subseteq \{1, \dots, n\}$. Without loss of generality, we can assume $I = \{1, \dots, s\}$. Let us express $e_1 + \cdots + e_s$ in QeR as $e_1 + \cdots + e_s = \sum_{i=1}^t q_i ar_i$, where $q_i \in Q$ and $r_i \in R$. We can take r in R satisfying $0 \neq e_m r \in R$, $m = s+1, \dots, n$. Put $b = r(e_{s+1} + \cdots + e_n)$. Here we claim that $\{r_1, \dots, r_t; b\}$ is right semi-insulator for a . $RaR \cap RbR = 0$ is obvious. If x is in $[\bigcap_{i=1}^t (0: ar_i)_R^r] \cap (0: bR)_R^r$, then $(e_1 + \cdots + e_s)x = 0$. Further, inasmuch as $Qe_m rQ = Q_m$ for $m = s+1, \dots, n$, we infer $QbR = Q_{s+1} \oplus \cdots \oplus Q_n$; whence $(e_{s+1} + \cdots + e_n)x = 0$. Therefore $x = 0$ as required.

(c) \Rightarrow (a). Proposition 2.1 says that R is a semiprime right nonsingular ring. If I is an essential ideal of R , then there exists a in I such that $RaR \subseteq_e I$ ($\subseteq_e R$) by Proposition 2.3. Let $\{r_1, \dots, r_n; b\}$ be a right semi-insulator of a . Since $RaR \subseteq_e R$ and $RaR \cap RbR = 0$, we see $b = 0$. Consequently $\bigcap_{i=1}^n (0: ar_i) = 0$. Therefore I is insulated as a right ideal.

(b) \Rightarrow (d). If I is an essential right ideal of R , then $QI \subseteq_e Q$ as a right R -module. As is seen in the proof of (b) \Rightarrow (c), it follows from (1) that Q is regular. Therefore (2) easily implies $QI = QIQ$. As a result $QI = QIQ < \oplus Q$ and hence $QI = Q$.

(d) \Rightarrow (e) \Rightarrow (a) is obvious.

Corollary 2.6. *A ring R is a right strongly prime ring if and only if $Q(R)$ is simple and $Q(R)eR = Q(R)eQ(R)$ for all idempotents e in $Q(R)$.*

Corollary 2.7. *The following conditions are equivalent for a given ring R .*

- (a) R is a semiprime right Goldie ring.
- (b) R is a right finite dimensional right strongly semiprime ring.
- (c) $IQ(R) = Q(R)I = Q(R)$ for every essential right ideal I of R .

Proof. (a) \Rightarrow (b). Since every essential ideal of R contains a regular element, clearly R is a right strongly semiprime ring.

(b) \Rightarrow (a) follows from Lemma 2.4, and (b) \Leftrightarrow (c) follows from Theorem 2.5 and [12, Theorem 1.6].

Corollary 2.8 ([8, Corollary 16]). *A regular right strongly semiprime ring R*

is a direct sum of simple rings. Therefore R is also a left strongly semiprime ring.

Proof. Inasmuch as R contains no infinite direct sums of ideals, it is sufficient to show that R contains no proper essential ideals. Let I be an essential ideal of R . Then $QI=Q$ by Theorem 2.5, whence it follows from regularity of R that $1=qe$ for some $q \in Q$ and $e=e^2 \in I$. Then, clearly, $1=e$. So $I=R$.

3. Nonsingular quasi-injective modules

Lemma 3.1 ([1]). *If R is a right strongly prime ring, then every nonsingular quasi-injective right R -module is injective.*

Proof. Let $M (\neq 0)$ be a nonsingular quasi-injective right R -module and let $0 \neq x \in M$. Since $xQ(R)$ is $Q(R)$ -projective there exists e in $Q(R)$ and an isomorphism $\psi: xQ(R) \approx eQ(R)$ with $\psi(x)=e$. We can take r in R such that $0 \neq er \in R$. Then $R \subseteq n(erR)$ for some positive integer n , since er has a right insulator. Inasmuch as $R \subseteq n(erR) \approx n(xrR) \subseteq nM$, M is injective by Lemma 1.2.

Lemma 3.2. *Let R be a right self-injective regular ring such that every nonsingular quasi-injective right R -module is injective. Then R is a direct sum of simple rings.*

Proof. According as every ideal of R is a nonsingular quasi-injective right R -module, every ideal of R is a direct summand. Hence R contains no infinite direct sums of ideals. Hence by Proposition 2.3, R is written as a direct sum of prime rings, say $R=R_1 \oplus \cdots \oplus R_n$. Since R_i is prime and every ideal of R_i is a direct summand, R_i must be simple, $i=1, \dots, n$.

Proposition 3.3. *If R is a right nonsingular ring, then the following conditions are equivalent:*

- (a) $Q(R)$ is a direct sum of simple rings.
- (b) $E_R(M)=MQ(R)$ for all nonsingular quasi-injective right R -module M .

Proof. Set $Q=Q(R)$. (a) \Rightarrow (b). If M is a nonsingular quasi-injective right R -module, then MQ is nonsingular Q -quasi-injective. Hence, by Lemma 3.1, MQ is Q -injective; whence MQ is R -injective.

(b) \Rightarrow (a). If M is a nonsingular quasi-injective right Q -module, then M is nonsingular R -quasi-injective (Lemma 1.1). Hence $M=MQ=E_R(M)=E_Q(M)$, which shows that M is Q -injective. Thus, by Lemma 3.2, we conclude that Q is a direct sum of simple rings.

We are now in a position to show our main theorem.

Theorem 3.4. *For a given ring R , the following conditions are equivalent:*

- (a) $R/G(R)$ is a right strongly semiprime ring.

- (b) *Every nonsingular quasi-injective right R -module is injective.*
 (c) *Any finite direct sum of nonsingular quasi-injective right R -module is also quasi-injective.*
 (d) *Any direct product of nonsingular quasi-injective right R -module is quasi-injective.*

Proof. Set $\bar{R}=R/G(R)$ and $Q=Q(R/G(R))$. (b) \Rightarrow (d) \Rightarrow (c): Obvious.

(a) \Rightarrow (b). Since \bar{R} is a right strongly semiprime ring, Theorem 2.5 says that Q is a direct sum of simple rings and $eQe\bar{R}=eQ$ for all idempotents e in Q . Now, let $M(\neq 0)$ be a nonsingular quasi-injective right R -module. In order to show M is injective, we may show $M=MQ$ by Proposition 3.3. Let $0\neq x\in M$. Since xQ is Q -projective, there exists an idempotent e in Q and an isomorphism $\psi: xQ\approx eQ$ with $\psi(x)=e$. Inasmuch as xQ is Q -injective, $E_R(M)=xQ\oplus Y$ for some submodule Y . Since M is quasi-injective, this yields $M=(xQ\cap M)\oplus(Y\cap M)$. As a result, $xQ\cap M$ is quasi-injective. Put $Z=\psi(xQ\cap M)$. Inasmuch as $xR\subseteq xQ\cap M\subseteq xQ$, we infer that $E_R(xQ\cap M)=xQ$; whence $E_R(Z)=eQ$. Observing $eQ=eQeR=End_Q(eQ)eR=End_R(eQ)eR\subseteq End_R(eQ)Z=Z$, we see $eQ=Z=\psi(xQ\cap M)$. Consequently $xQ=xQ\cap M$ and it follows $xQ\subseteq M$. Therefore $MQ=M$ as desired.

(c) \Rightarrow (a). In view of Theorem 2.5, it is enough to show that $eQeR=eQ$ for all idempotents e in Q and Q is a direct sum of simple rings.

Let $e=e^2\in Q$ and set $T=eQeR\oplus(1-e)Q(1-e)R$. Then T is a nonsingular quasi-injective right R -module because both $eQeR$ and $(1-e)Q(1-e)R$ are so. Since $R\subseteq T$, it follows that T is injective; whence so is $eQeR$. Thus we get $eQeR=eQeQ=eQ$. Now, assume that Q can not be expressed as a direct sum of prime rings. Then, by Proposition 2.3, we see that there exist infinite orthogonal non-zero central idempotents $\{e_i|i=1,2,\dots\}$ in Q . Since $\sum_{i=1}^{\infty}e_iQ$ is nonsingular Q -quasi-injective, it is also nonsingular R -quasi-injective (Lemma 1.1). Putting $T=(1-e_1)Q\times(\sum_{i=1}^{\infty}e_iQ)$, T is then a nonsingular quasi-injective right R -module, since both $(1-e_1)Q$ and $\sum_{i=1}^{\infty}e_iQ$ are so. As a result, it follows from $R\subseteq T$ that T is injective and $\sum_{i=1}^{\infty}e_iQ<\oplus Q$, a contradiction. Hence Q must be written as a direct sum of prime rings, say $Q=Q_1\oplus\cdots\oplus Q_n$. Let X be a non-zero ideal of Q_i . Then X is a nonsingular quasi-injective right Q -module and hence it is nonsingular R -quasi-injective by Lemma 1.1. Take a non-zero idempotent e in X and consider $X\times(1-e)Q$. Since both X and $(1-e)Q$ are nonsingular quasi-injective right R -module, so is $X\times(1-e)Q$. Inasmuch as $R\subseteq X\times(1-e)Q$, it follows that $X\times(1-e)Q$ is injective; whence $X\oplus Q_i$. Since Q_i is a prime ring, this shows $X=Q_i$. Accordingly each Q_i is simple.

Combining Theorem 3.4 with Corollary 2.8, we have

Corollary 3.5. *If R is a regular ring, then the following conditions are equivalent:*

- (a) *R is a direct sum of simple rings.*
- (b) *Every nonsingular quasi-injective right R -module is injective.*
- (b') *Every nonsingular quasi-injective left R -module is injective.*

Corollary 3.6. *If R is a right strongly semiprime ring, then its right socle is a direct summand of R as a ring.*

Proof. By Theorem 3.4(b), we conclude that the right socle S of R is a direct summand of R as a right R -module. Since R is a semiprime ring and S is a two-sided ideal of R , it follows that S is a direct summand of R as a ring.

Boyle and Goodearl [3] showed that every nonsingular quasi-injective right module over a semiprime right Goldie ring is injective. However, according as every essential ideal of a semiprime right Goldie ring R has a regular element, R is a right and left strongly semiprime ring. Hence Theorem 3.4 guarantees the following result.

Corollary 3.7. *If R is a semiprime right Goldie ring, then every nonsingular quasi-injective right R -module is injective and, at the same time, every nonsingular quasi-injective left R -module is also injective.*

Finally we show the following result.

Theorem 3.8. *For a given ring R , the following conditions are equivalent:*

- (a) *$R/G(R)$ is a semiprime right Goldie ring.*
- (b) *Any direct sum of nonsingular quasi-injective right R -modules is quasi-injective.*

Proof. As is well known ([13]), the following conditions are equivalent:

- (1) *$Q(R/G(R))$ is a semisimple artinian ring.*
- (2) *$R/G(R)$ is right finite dimensional.*
- (3) *Any direct sum of nonsingular injective right R -modules is injective.*

Convining this fact with Theorem 3.4 and Corollary 3.7, the proof is established.

REMARK. It seems to be also meaningful to study those rings whose nonsingular quasi-injective right modules are written as direct sums of indecomposable modules. Such rings were determined by Berry [2] as rings R such that $R/G(R)$ is right finite dimensional.

References

- [1] J. Beachy and W.D. Blair: *Rings whose faithful left ideals are cofaithful*, Pacific J. Math. **58** (1975), 1–13.
- [2] D. Berry: *Modules whose cyclic submodules have finite dimension*, Canad. Math. Bull. **19** (1976), 1–6.
- [3] Ann. K. Boyle and K.R. Goodearl: *Rings over which certain modules are injective*, Pacific J. Math. **58** (1975), 43–53.
- [4] K.R. Goodearl: *Ring theory; Nonsingular rings and modules*, New York, Marcel Dekker, 1976 (Pure and Applied Math. Series, 33).
- [5] K.R. Goodearl and Ann. K. Boyle: *Dimension theory for nonsingular injective modules*, Mem. Amer. Math. Soc. 177, 1976.
- [6] K.R. Goodearl and D. Handelman: *Simple self-injective rings*, Comm. Algebra **3** (1975), 797–834.
- [7] D. Handelman and J. Lawrence: *Strongly prime rings*, Trans. Amer. Math. Soc., **211** (1975), 209–223.
- [8] D. Handelman: *Strongly semiprime rings*, Pacific J. Math. **60** (1975), 115–122.
- [9] M. Harada: *Note on quasi-injective modules*, Osaka J. Math. **2** (1965), 351–356.
- [10] J. Kado: *Projective modules over simple regular rings*, Osaka J. Math. **16** (1979), 405–412.
- [11] R. Rubin: *Absolutely torsion-free rings*, Pacific J. Math. **46** (1973), 503–514.
- [12] F.L. Sandomierski: *Semisimple maximal quotient rings*, Trans. Amer. Math. Soc. **128** (1967), 112–120.
- [13] M.L. Teply: *Torsionfree injective modules*, Pacific J. Math. **28** (1969), 441–453.
- [14] J.M. Zelmanowitz: *The finite intersection property on annihilator right ideals*, Proc. Amer. Math. Soc. **57** (1976), 213–216.

Department of Mathematics
Yamaguchi University
Yamaguchi 753, Japan