

Title	On congruent axioms in linearly ordered spaces. I
Author(s)	Terasaka, Hidetaka; Katayama, Shigeru
Citation	Osaka Journal of Mathematics. 1966, 3(2), p. 269-292
Version Type	VoR
URL	https://doi.org/10.18910/12504
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ON CONGRUENT AXIOMS IN LINEARLY ORDERED SPACES, I

SHIGERU KATAYAMA and HIDETAKA TERASAKA

(Received September 2, 1966)

1. Introduction

In connection with the axioms of congruence of segments on a straight line given in Hilbert's *Grundlagen der Geometrie*, we will set up a group of axioms of congruence on a linearly ordered space and study their mutual dependency and independency.

In the following let L be a linearly ordered space, that is a set of points, in which for any pair of distinct points A and B either of the relations $A < B$ and $B < A$ holds, and for any three points A, B and C , if $A < B$ and $B < C$ then $A < C$.

When we write AB , it will be understood that A and B are distinct points of L such that $A < B$. AB will be called a *segment*. We write $AC \equiv AB + BC$ if and only if $A < B < C$.

The axioms we are going to study is the following:

Axiom E (UNIQUE EXISTENCE): $\forall AB \forall A' \exists_1 B' : AB = A'B'$, that is, for any segment AB and for any point A' there is one and only one point B' such that

$$AB = A'B'.$$

Axiom R (REFLEXIVITY): $AB = AB$.

Axiom S (SYMMETRICITY): $AB = A'B' \Rightarrow A'B' = AB$.


Axiom T (TRANSITIVITY): $AB = A'B', A'B' = A''B'' \Rightarrow AB = A''B''$.

Axiom A (ADDITIVITY):

$$AC \equiv AB + BC, A'C' \equiv A'B' + B'C', AB = A'B', BC = B'C' \Rightarrow AC = A'C'.$$

The following scheme will be used in application:


$$\left. \begin{array}{l} AC \equiv AB + BC, \\ A'C' \equiv A'B' \\ \quad + B'C', \\ AB = A'B', \\ BC = B'C' \end{array} \right\} \xrightarrow{(A)} AC = A'C'.$$



Axiom C (COMMUTATIVE ADDITION):

$$AC \equiv AB + BC, C'A' \equiv C'B' + B'A', AB = B'A', BC = C'B' \Rightarrow AC = C'A'.$$

In application we write:

$$\left. \begin{array}{l} AC \equiv AB + BC, \\ C'A' \equiv C'B' \\ \quad + B'A', \\ AB = B'A', \\ BC = C'B' \end{array} \right\} \xrightarrow{(C)} AC = C'A'.$$


Axiom I (INTERCHANGING): $A < B < A' < B', AB = A'B' \Rightarrow AA' = BB'$.

Under the assumption of Axiom E we studied in this paper all the relationship between the remaining axioms R, S, T, A, C, I as to their mutual dependency and independency, and obtained among others the following Main Theorem.

Main Theorem: *Under the assumption of Axiom E,*

I. *Axioms T and C are independent of each other, and Axioms R, S, A and I follow from them.* In symbol:

$$R, S, T, A, C, I$$

II. *Axioms T and I are independent of each other, and Axioms R, S, A and C follow from them.* In symbol:

$$R, S, T, A, C, I$$

III. *Axioms S, A and I are independent of one another, and Axioms R, T and C follow from them.* In symbol:

$$R, S, T, A, C, I$$

IV. *Axioms S, A and C are independent of one another, and Axioms R, T and I follow from them.* In symbol:

$$R, S, T, A, C, I$$

V. *Axioms R, A and C are independent of one another, and Axioms S, T and I follow from them.* In symbol:

$$R, S, T, A, C, I$$

2. Theorems

In the following we always assume the unique existence of Axiom E, if not otherwise stated.

To make proofs as clear as possible we introduce first some useful notations.

- (a) $X \xrightarrow{(T)} Y$ means that Y follows from the left side X by the use of T.
 (b) $A = B$ means that A coincides with B and $AB \equiv A'B'$ means that $A = A'$,

$B=B'$ at the same time.

(c) “ $\exists_1 X$:” means that “there exist one and only one X such that.”

Theorem 1. *If T is assumed, then $R \Leftrightarrow S$.*

Proof. (i) $R \Rightarrow S$.

$$\left. \begin{array}{l} \text{By Axiom E, } \exists_1 B' : AB=A'B' . \\ \text{By Axiom E, } \exists_1 B'' : A'B'=AB'' . \end{array} \right\} \xrightarrow{(T)} AB=AB'' . \quad (1)$$

Now by Axiom R,

$$\begin{aligned} AB &= AB . & (2) \\ (1), (2) & \xrightarrow{(E)^D} B''=B . \end{aligned}$$

(ii) $R \Leftarrow S$.

Assume $AB=A'B'$. Then by Axiom S, $A'B'=AB$. Hence by Axiom T, $AB=AB$.

Theorem 2. $R, C \Rightarrow I$.

Proof. Let $A < B < A' < B'$ and $AB=A'B'$.

Then we have

$$\left. \begin{array}{l} AA' \equiv AB+BA' , \\ BB' \equiv BA'+A'B' , \\ AB=A'B' , \\ BA'=BA' \text{ (by Axiom R)} \end{array} \right\} \xrightarrow{(C)} AA'=BB' .$$

Lemma 1. *Under the assumption of T: $AB=A'B' \Rightarrow A'B'=A'B'$.*

Especially: $AB=AB' \Rightarrow AB'=AB'$.

Proof. Let

$$AB=A'B' \quad (1) \quad \begin{array}{c} L \quad \overset{A}{\mid} \text{-----} \overset{B}{\mid} \\ L \quad \quad \quad \underset{A'}{\mid} \text{-----} \underset{B''}{\mid} \quad \underset{B'}{\mid} \end{array}$$

Then we have

$$\left. \begin{array}{l} AB=A'B' . \\ \text{By Axiom E, } \exists_1 B'' : A'B'=A'B'' \text{ (2).} \\ \text{By (1): } AB=A'B' . \end{array} \right\} \xrightarrow{(T)} AB=A'B'' . \left. \right\} \xrightarrow{(E)} B''=B' .$$

Therefore we have from (2) $A'B'=A'B'$.

1) If $AB=A'B'$ and $AB=A'B''$ then we have by Axiom E $B'=B''$. As a special case, if $AB=AB'$ and $AB=AB$ then $B'=B$.

Theorem 3. $T, C \Rightarrow A.$

Proof. Let $AC \equiv AB + BC$, $A'C' \equiv A'B' + B'C'$,

$$AB = A'B', \tag{1}$$

and

$$BC = B'C'. \tag{2}$$


Then by Axiom E, $\exists_1 D': A'B' = C'D'$. (3)

Then we have first from (1) and (3) by using T

$$AB = C'D'. \tag{4}$$



Further



$$\left. \begin{array}{l} AC \equiv AB + BC, \\ B'D' \equiv B'C' + C'D', \\ (4): AB = C'D', \\ (2): BC = B'C' \end{array} \right\} \xrightarrow{(C)} AC = B'D'. \tag{5}$$

By Axiom E, $\exists_1 X: C'D' = A'X$. (6)

Then by (3) and (6) we have by using T

$$A'B' = A'X. \tag{7}$$

Since by Lemma 1

$$A'B' = A'B', \tag{8}$$

we have from (7) and (8) by the use of Axiom E $X = B'$.

Hence by (6)

$$C'D' = A'B'. \tag{9}$$

Then

$$\left. \begin{array}{l} B'D' \equiv B'C' + C'D', \\ A'C' \equiv A'B' + B'C', \\ B'C' = B'C' \text{ (by Lemma 1)}, \\ (9): C'D' = A'B' \end{array} \right\} \xrightarrow{(C)} B'D' = A'C'. \tag{10}$$

From (5) and (10) we have finally by Axiom T $AC = A'C'$.


Theorem 4. $T, A \Rightarrow R.$

Proof. Let AB be a given segment. Then by Axiom E, $\exists_1 B'$:

$$AB = AB'. \tag{1}$$

(i) Suppose first $A < B' < B$.

Then we have



$$\left. \begin{array}{l} AB \equiv AB' + B'B, \\ AX \equiv AB' + B'X, \\ AB' = AB' \text{ (by Lemma 1),} \\ \text{by Axiom E, } \exists_1 X: B'B = B'X \end{array} \right\} \xrightarrow{(A)} AB = AX. \quad (2)$$

From (1) and (2) we would have by Axiom E $B' = X$, which is clearly a contradiction.

(ii) Next suppose $B < B'$.

Then we have

$$\left. \begin{array}{l} AB' \equiv AB + BB', \\ AX \equiv AB' + B'X, \\ (1): AB = AB', \\ \text{by Axiom E, } \exists_1 X: BB' = B'X \end{array} \right\} \xrightarrow{(A)} AB' = AX. \quad (3)$$

Since by Lemma 1 $AB' = AB'$, we have from (3) $X = B'$, which is clearly a contradiction.


From (i) and (ii) we conclude $AB = AB$.

Theorem 5. $T, C \Rightarrow R$.

This is an easy consequence of Theorem 3: $T, C \Rightarrow A$ and Theorem 4: $T, A \Rightarrow R$. In the following an alternative proof will be given without an intermediation of Axiom A.

Lemma 2. *Under the assumption of Axiom C*

$$\left. \begin{array}{l} AB = A'B', \quad (1) \\ A < X < B, \\ XB = A'X' \quad (2) \end{array} \right\} \implies \left\{ \begin{array}{l} A' < X' < B', \\ AX = X'B'. \end{array} \right.$$



Proof.

$$\left. \begin{array}{l} AB \equiv AX + XB, \\ A'B'' \equiv A'X' + X'B'', \\ \text{by Axiom E, } \exists_1 B'': AX = X'B'' \quad (3), \\ \text{by (2): } XB = A'X' \end{array} \right\} \xrightarrow{(C)} AB = A'B''. \quad (4)$$

Then we have

$$(1), (4) \xrightarrow{(E)} B' = B''.$$

Thus $AX = X'B'$ from (3) and clearly $A' < X' < B'' = B'$.

Proof of Theorem 5.

$$\text{By Axiom E, } \exists_1 B': AB = AB'. \tag{1}$$

(i) Suppose first $A < B' < B$.

By Lemma 2 there is an X such that


$$A < X < B', \tag{2}$$

$$B'B = AX, \tag{3}$$

$$AB' = XB'. \tag{4}$$

$$\text{Now by Axiom E, } \exists_1 X': AX = B'X' \tag{5}$$

and by Lemma 1



Thus

$$\left. \begin{array}{l} AB' \equiv AX + XB', \\ XX' \equiv XB' + B'X', \\ (5): AX = B'X', \\ (6): XB' = XB' \end{array} \right\} \xrightarrow{(C)} AB' = XX'. \tag{7}$$

Then

$$(4), (7) \xrightarrow{(E)} X' = B',$$

which is a contradiction.

(ii) Next suppose $A < B < B'$.

$$\text{By Axiom E, } \exists_1 X: BB' = AX. \tag{8}$$

$$\text{By Axiom E, } \exists_1 B'': AB' = XB''. \tag{9}$$

Then

$$(1), (9) \xrightarrow{(T)} AB = XB''. \tag{10}$$

Hence

$$\left. \begin{array}{l} AB' \equiv AB + BB', \\ AB'' \equiv AX + XB'', \\ (10): AB = XB'', \\ (8): BB' = AX \end{array} \right\} \xrightarrow{(C)} AB' = AB''. \tag{11}$$

$$\left. \begin{array}{l} \text{By Lemma 1 } AB' = AB' \end{array} \right\} \xrightarrow{(E)} B'' = B'.$$


Consequently, we have from (9)

$$AB' = XB' \tag{11}$$

Now

$$\left. \begin{aligned} AB' &\equiv AX + XB', \\ XX' &\equiv XB' + B'X', \\ \text{by Axiom E, } \exists_1 X': AX &= B'X', \\ \text{by Lemm 1, } XB' &= XB' \end{aligned} \right\} \xrightarrow{(C)} AB' = XX' \quad (12)$$

Then we have

$$(11), (12) \xrightarrow{(E)} B' = X',$$


which is a contradiction.

From (i) and (ii) we conclude $AB = AB$.


Theorem 6. $T, I \Rightarrow R$.

Proof.

$$\text{By Axiom E, } \exists_1 B': AB = AB'. \quad (1)$$

(i) Suppose first $A < B' < B$.

$$\text{By Axiom E, } \exists_1 C: AB' = BC. \quad (2)$$



Then

$$(1), (2) \xrightarrow{(T)} AB = BC. \quad (3)$$

and

$$A < B' < B < C, (2) \xrightarrow{(I)} AB = B'C. \quad (4)$$

$$\text{By Axiom E, } \exists_1 D: B'B = CD. \quad (5)$$

Then

$$B' < B < C < D, (5) \xrightarrow{(I)} B'C = BD, \quad (6)$$

and

$$(4), (6) \xrightarrow{(T)} AB = BD. \quad (7)$$

Hence

$$(3), (7) \xrightarrow{(E)} C = D.$$

which is a contradiction.

(ii) Next suppose $A < B < B'$.

$$\text{By Axiom E, } \exists_1 C: AB = B'C. \quad (8)$$

$$A < B < B' < C, (8) \xrightarrow{(I)} AB' = BC. \quad (9)$$

$$(1), (9) \xrightarrow{(T)} AB = BC. \quad (10)$$

$$\text{By Axiom E, } \exists_1 D: BB' = CD. \quad (11)$$

$$B < B' < C < D, (11) \xrightarrow{(I)} BC = B'D. \quad (12)$$

$$(10), (12) \xrightarrow{(T)} AB = B'D. \quad (13)$$

Hence

$$(8), (13) \xrightarrow{(E)} C = D,$$

which is a contradiction.

From (i) and (ii) we conclude $AB = AB$.

Lemma 3. $A < A' < B, AB = A'B' \xrightarrow{(I)} B < B', AA' = BB'$.

Proof.

$$\text{By Axiom E, } \exists_1 B'': AA' = BB''. \quad (1)$$

$$A < A' < B < B'', (1) \xrightarrow{(I)} AB = A'B''. \quad (2)$$

Let $AB = A'B'$. Then we have

$$AB = A'B', (2) \xrightarrow{(E)} B' = B''.$$

Therefore we have from (1) $B < B'$ and $AA' = BB'$.

Theorem 7. $T, I \Rightarrow A$.

Proof. (i) First let $A < B < C < A' < B' < C'$ and $AB = A'B', BC = B'C'$.

$$\left. \begin{array}{l} A < B < A' < B', AB = A'B' \xrightarrow{(I)} AA' = BB', \\ B < C < B' < C', BC = B'C' \xrightarrow{(I)} BB' = CC' \end{array} \right\} \begin{array}{l} \xrightarrow{(T)} AA' = CC', \\ A < C < A' < C' \end{array} \xrightarrow{(\text{Lem.3})} AC = A'C'.$$

(ii) Next let $A < B < C, A' < B' < C'$ with $AB = A'B', BC = B'C'$, but let $C < A'$ fail to be true.

Take points A'', B'', C'' such that $A < B < C < A'' < B'' < C''$ and $A' < B' < C' < A'' < B'' < C''$ with $A'B' = A''B'', B'C' = B''C''$.

Then by (i)

$$AC = A''C'' \quad (1)$$

and

$$A'C' = A''C''. \quad (2)$$

Now, since $T, I \Rightarrow R$ by Theorem 6 and $T, R \Rightarrow S$ by Theorem 1, Axiom S holds by our assumption of T and I .

Therefore

$$A'C' = A''C'' \xrightarrow{(S)} A''C'' = A'C'. \quad (3)$$

Hence

$$(1), (3) \xrightarrow{(T)} AC = A'C' .$$

Theorem 8. T, I \Rightarrow C.

Proof. Notice that Axiom S is a consequence of our assumption of T and I as we have shown in the proof of Theorem 7 and that Axiom A is a consequence of T and I by Theorem 7.

Let $A < B < C$, $C' < B' < A'$ and let

$$\begin{array}{ll} AB = B'A', & (1) \quad L \xrightarrow{\quad} \begin{array}{ccc} A & B & C \end{array} \\ BC = C'B', & (2) \quad L \xrightarrow{\quad} \begin{array}{cccc} C' & B' & A' & C'' \end{array} \end{array}$$

$$\left. \begin{array}{l} (2): BC = C'B', \\ \text{by Axiom E,} \\ \exists_1 C'': C'B' = A'C'' \end{array} \right\} \xrightarrow{(T)} BC = A'C''$$

$$\left. \begin{array}{l} AC \equiv AB + BC, \\ B'C'' \equiv B'A' + A'C'', \\ (1): AB = B'A', \\ (2): BC = C'B' \end{array} \right\} \xrightarrow{(A)} AC = B'C'' ,$$

$$\left. \begin{array}{l} C' < B' < A' < C'', C'B' = A'C'' \xrightarrow{(I)} C'A' = B'C'' \xrightarrow{(S)} B'C'' = C'A' \\ \xrightarrow{(T)} AC = C'A' . \end{array} \right\}$$

Theorem 9. S, A \Rightarrow R.

Proof.

$$\text{By Axiom E, } \exists_1 B': AB = AB' . \tag{1}$$

(i) Let $A < B < B'$.

$$\text{By Axiom E, } \exists_1 X: BB' = B'X \tag{2}$$

$$\left. \begin{array}{l} AB' \equiv AB + BB', \\ AX \equiv AB' + B'X, \\ (1): AB = AB', \\ (2): BB' = B'X \end{array} \right\} \xrightarrow{(A)} AB' = AX . \tag{3}$$

$$(1) \xrightarrow{(S)} AB' = AB . \tag{4}$$

$$(3), (4) \xrightarrow{(E)} X = B ,$$

which is a contradiction.

(ii) Let $A < B' < B$.

By Axiom S $AB' = AB$, $A < B' < B$ and the case (ii) reduces to that of (i).

From (i) and (ii) we conclude $AB = AB$.

Lemma 4. Under the assumption of Axioms S, A and I, if $AB=A'B'$, then

- 1) $A < A' \Rightarrow B < B', AA' = BB'$.
- 2) $A' < A \Rightarrow B' < B, A'A = B'B$.
- 3) $A = A' \Rightarrow B = B'$.

Proof. 1) follows from Lemma 3.
 2) reduces to 1) by Axiom S.
 3) follows from Theorem 9 which asserts $S, A \Rightarrow R$.

Lemma 5. $PQ = P'Q'$ (1), $P < X < Q, PX = P'X'$ (2) $\xrightarrow{(A)} P' < X' < Q', XQ = X'Q'$.

Proof.


$$\left. \begin{array}{l} PQ \equiv PX + XQ, \\ P'Q'' \equiv P'X' + X'Q'', \\ (2): PX = P'X', \\ \text{by Axiom E, } \exists_1 Q'': XQ = X'Q'' \end{array} \right\} \begin{array}{l} \xrightarrow{(A)} PQ = P'Q'', \\ (1): PQ = P'Q' \end{array} \left. \right\} \xrightarrow{(E)} Q'' = Q'.$$

Lemma 6. $PQ = P'Q', PQ = P''Q', P < P', P < P'' \xrightarrow{(A,I)} P' = P''$.

Proof. We may assume without loss of generality that $P' < P''$.

$$PQ = P'Q' \xrightarrow{(I \text{ or Lem. } 3)} PP' = QQ' \tag{1}$$

$$PQ = P''Q' \xrightarrow{(I \text{ or Lem. } 3)} PP'' = QQ' \tag{2}$$




$$\left. \begin{array}{l} PP'' \equiv PP' + P'P'', \\ QX \equiv QQ' + Q'X, \\ (1): PP' = QQ', \\ \text{by Axiom E, } \exists_1 X: P'P'' = Q'X \end{array} \right\} \begin{array}{l} \xrightarrow{(A)} PP'' = QX, \\ (2): PP'' = QQ' \end{array} \left. \right\} \xrightarrow{(E)} X = Q',$$

which is a contradiction.

Lemma 7. $\left. \begin{array}{l} PQ = P'Q', P < P', \\ P < X < Q, XQ = X'Q' \end{array} \right\} \xrightarrow{(S,A,I)} P' < X' < Q', PX = P'X'$.

Proof. By Lemma 3 we have first



From

$$XQ = X'Q' \tag{1}$$

we have $X'Q' = XQ$ by Axiom S, and combined with $Q < Q'$ we obtain by Lemma 4

$$X < X'. \tag{2}$$

Now,

$$\text{by Axiom E, } \exists X'' : PX = P'X'', \tag{3}$$

$$\text{by Lemma 5 } P' < X'' < Q' \text{ and } XQ = X''Q', \tag{4}$$

$$\text{by Lemma 4 } X < X''. \tag{5}$$

Then (1), (4), (2) and (5) yield by Lemma 6 $X' = X''$. Consequently we have $P' < X' < Q'$ and $PX = P'X'$.

Theorem 10. S, A, I \Rightarrow T.

Proof. Let $AB = A'B'$, $A'B' = A''B''$.

(i) The case where at least two of A , A' and A'' coincide:

(i)₁ $A = A'$. Since S, A \Rightarrow R by Theorem 9 we have $B = B'$ and hence $AB = A''B''$.

(i)₂ $A' = A''$. The same as (i)₁.

(i)₃ $A = A''$. $AB = A'B' \xrightarrow{(S)} A'B' = AB$, $A'B' = A''B''$, $A = A'' \Rightarrow A'B' = AB''$ } $\xrightarrow{(E)} B = B''$.

Hence

$$AB = A''B''.$$

(ii) The case where A , A' and A'' are distinct: there are six cases to be considered.

I. $A < A' < A''$, II. $A < A'' < A'$, III. $A' < A < A''$,

I'. $A'' < A' < A$, II'. $A'' < A < A'$, III'. $A' < A'' < A$.

Proof of Case I.

$$\left. \begin{array}{l} AB = A'B' \xrightarrow{(I \text{ or Lem. 3})} AA' = BB', \\ A'B' = A''B'' \xrightarrow{(I \text{ or Lem. 3})} A'A'' = B'B'', \\ A < A' < A'' \xrightarrow{(Lem. 4)} B < B' < B'' \end{array} \right\} \xrightarrow{(A)} AA'' = BB'' \xrightarrow{(I \text{ or Lem. 3})} AB = A''B''.$$

Proof of Case II.

$$AB=A'B' \stackrel{(I \text{ or Lem. } 3)}{\implies} AA'=BB' . \quad (1)$$

$$A'B'=A''B'' \stackrel{(S)}{\implies} A''B''=A'B' \stackrel{(I \text{ or Lem. } 3)}{\implies} A''A'=B''B' . \quad (2)$$

$$\left. \begin{array}{l} (1), \\ (2), \\ A < A'' < A' \end{array} \right\} \stackrel{(Lem. 7)}{\implies} \left\{ \begin{array}{l} B < B'' < B' , \\ AA''=BB'' \end{array} \right\} \stackrel{(I \text{ or Lem. } 3)}{\implies} AB=A''B'' .$$

Proof of Case III.

$$A'B'=A''B'' \stackrel{(I \text{ or Lem. } 3)}{\implies} A'A''=B'B'' . \quad (1)'$$

$$AB=A'B' \stackrel{(S)}{\implies} A'B'=AB \stackrel{(I \text{ or Lem. } 3)}{\implies} A'A=B'B . \quad (2)'$$

$$\left. \begin{array}{l} (1)', \\ (2)', \\ A' < A < A'' \end{array} \right\} \stackrel{(Lem. 5)}{\implies} \left\{ \begin{array}{l} B' < B < B'' , \\ AA''=BB'' \end{array} \right\} \stackrel{(I \text{ or Lem. } 3)}{\implies} AB=A''B'' .$$

Proof of Case I'.

$$AB=A'B' \stackrel{(S)}{\implies} A'B'=AB . \quad (1)''$$

$$A'B'=A''B'' \stackrel{(S)}{\implies} A''B''=A'B' . \quad (2)''$$

$$A'' < A' < A, (2)'', (1)'' \stackrel{(Case I)}{\implies} A''B''=AB \stackrel{(S)}{\implies} AB=A''B'' .$$

Similarly the proofs of II' and III' may be reduced to those of II and III respectively.

Theorem 11. S, A, I \Rightarrow C.

Proof. S, A, I \Rightarrow T by Theorem 10. Then by Theorem 8 T, I \Rightarrow C.

Lemma 8. Under the assumption of Axioms R and C, if $AB=A'B'$, then

- 1) $A < A' \Rightarrow B < B'$.
- 2) $A = A' \Rightarrow B = B'$.
- 3) $A' < A \Rightarrow B' < B$.

Proof. 1) $A < A'$. $B < B'$ is clear if $B < A'$ or if $B = A'$.
Let $A < A' < B$, and suppose either $B' < B$ or $B' = B$.

$$\text{By Axiom E, } \exists X: AA'=BX . \quad (1)$$

$$\left. \begin{array}{l} A < A' < B < X , \\ (1): AA'=BX \end{array} \right\} \stackrel{(I)}{\implies} AB=A'X . \left. \vphantom{\begin{array}{l} A < A' < B < X , \\ (1): AA'=BX \end{array}} \right\} \stackrel{(E)}{\implies} X=B' ,$$

By assumption $AB=A'B'$

Corollary. $AB=A'B', A' < A \xrightarrow{(R, C)} A'B' = AB.$

Theorem 13. S, C, I \Rightarrow R.

Proof.

By Axiom E, $\exists_1 B': AB = AB'.$

(i) First suppose $A < B' < B.$

By Lemma 2 there is an X such that

$$\begin{aligned} A < X < B', \quad B'B = AX, \\ AB' = XB'. \end{aligned} \tag{1}$$

By Axiom E, $\exists_1 X': AX = B'X'.$

Since $A < X < B' < X'$ we have by Axiom I

$$AB' = XX'. \tag{2}$$

From (1) and (2) we would have by Axiom E $B' = X,$ which is a contradiction.

(ii) Next suppose $B < B'.$

Since we have from (1) by Axiom S $AB' = AB,$ the argument of (i) gives again a contradiction.

Thus we conclude from (i) and (ii) $B' = B$ and then $AB = AB$ follows from (1).

3. Models

By a *model* of a geometry denoted for example by $M(S, C)$ we mean a linearly ordered space L with congruent relations which satisfy among our group of seven Axioms E, R, S, T, A, C and I Axioms S and C alone besides Axiom E but not the remaining ones.

In the following models the space L is for the most part given by the real line $-\infty < x < \infty$ or by the half line $0 \leq x < \infty.$ In these cases points denoted by A, B, A', X etc. will be those points of the real line having coordinates a, b, a', x etc. respectively. $A < B$ is defined by $a < b,$ $|AB|$ denotes the distance $b - a$ of points A and $B.$

$M(R):$ *A model of a geometry in which Axiom R alone holds besides Axiom E.*

Let L be the real line $-\infty < x < \infty.$

DEFINITION OF $AB = A'B':$

If $A = A',$ then let $AB = A'B'$ if and only if $B = B'.$

If $A \neq A',$ then let $AB = A'B'$ if and only if $|A'B'| = 1.$

This model satisfies Axioms E and R but fails to satisfy the remaining Axioms S, T, A, C, I.

M(S): *A model of a geometry in which Axiom S alone holds besides Axiom E.*
Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$:

In case $A=A'$, let $AB=A'B'$

- (i) if $|AB|=1$ and $|A'B'|=3$
- or (ii) if $|AB|=3$ and $|A'B'|=1$
- or (iii) if $|AB|$ and $|A'B'|$ are both different from 1 and 3, and $|AB| = |A'B'|$.

In case $A < A'$, let $AB=A'B'$ and $A'B'=AB$ if $2|AB|=|A'B'|$. This model satisfies Axioms E and S but fails to satisfy the remaining Axioms R, T, A, C, I.

M(T): *A model of a geometry in which Axiom T alone holds besides Axiom E.*
Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$: For any AB and for any A' , let $AB=A'B'$ if and only if $|A'B'|=1$.

This model satisfies Axioms E and T but fails to satisfy the remaining Axioms R, S, A, C, I.

M(A): *A model of a geometry in which Axiom A alone holds besides Axiom E.*
Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$:

- (i) In case $A < A'$ or $A=A'$, then let $AB=A'B'$ if and only if $2|AB| = |A'B'|$.
- (ii) In case $A' < A$, then let $AB=A'B'$ if and only if $|AB|=|A'B'|$.

This model satisfies Axioms E and A but fails to satisfy the remaining Axioms R, S, T, C, I.

M(I): *A model of a geometry in which Axiom I alone holds besides Axiom E.*
Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$:

In case $A=A'$, let $AB=A'B'$ if and only if $2|AB|=|A'B'|$.

In case $A \neq A'$, let $AB=A'B'$ if and only if $|AB|=|A'B'|$.

This model satisfies Axioms E and I but fails to satisfy the remaining Axioms R, S, T, A, C.

M(A, C): *A model of a geometry in which Axioms A and C alone hold besides Axiom E.*

Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$: Let $AB=A'B'$ if and only if $2|AB|=|A'B'|$.

This model satisfies Axioms E, A and C but fails to satisfy the remaining Axioms R, S, T, I.

$M(S, I)$: *A model of a geometry in which Axioms S and I alone hold besides Axiom E.*

Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$:

In case $A=A'$, let $AB=A'B'$ if $|AB|=1$ and $|A'B'|=2$ or if $|AB|=2$ and $|A'B'|=1$ or if $|AB|$ and $|A'B'|$ are both different from 1 and 2, and $|AB|=|A'B'|$.

In case $A \neq A'$, let $AB=A'B'$ if $|AB|=|A'B'|$.

This model satisfies Axioms E, S and I but fails to satisfy the remaining Axioms R, T, A, C.

$M(R, S, A)$: *A model of a geometry in which Axioms R, S and A alone hold besides Axiom E.*

Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$:

In case $A=A'$, let $AB=A'B'$ if $B'=B$.

In case $A < A'$, let $AB=A'B'$ and $A'B'=AB$ if $2|AB|=|A'B'|$.

This model satisfies Axioms E, R, S and A but fails to satisfy the remaining Axioms T, C, I.

$M(R, A, I)$: *A model of a geometry in which Axioms R, A and I alone hold besides Axiom E.*

Let L be the real line $-\infty < x < \infty$.

DEFINITION OF $AB=A'B'$:

In case $A=A'$ or $A < A'$, let $AB=A'B'$ if $|AB|=|A'B'|$.

In case $A' < A$, let $AB=A'B'$ if $2|A'B'|=|AB|$.

This model satisfies Axioms E, R, A and I but fails to satisfy the remaining Axioms S, T, C.


$M(R, S, T)$: *A model of a geometry in which Axioms R, S and T alone hold besides Axiom E.*

Let a point of the space L be defined as an ordered pair (x, y) of real numbers x and y such that either $x \geq 0$ and $y=0$ or $x=0$ and $y \geq 0$.

Definition of the linear order:

If $A=(x, y)$, $A'=(x', y')$, then let $A < A'$ if $x < x'$ or if $y > y'$.

DEFINITION OF $AB=A'B'$:



If $A=(x_1, y_1)$, $B=(x_2, y_2)$, $A'=(x_1', y_1')$, $B'=(x_2', y_2')$,
 then let $AB=A'B'$ if $\sqrt{(x_2-x_1)^2+(y_2-y_1)^2}=\sqrt{(x_2'-x_1')^2+(y_2'-y_1')^2}$.

This model satisfies Axioms E, R, S and T but fails to satisfy the remaining Axioms A, C, I.

M(A, C, I): *A model of a geometry in which Axioms A, C, and I alone hold besides Axiom E.*

Let L be the half real line $0 \leq x < \infty$, and let O denote the point with coordinate 0.

DEFINITION OF $AB=A'B'$:

In case $A=O$, let $AB=A'B'$ if $|AB|+1=|A'B'|$.

In case $O < A$, let $AB=A'B'$ if $|AB|=|A'B'|$.

This model satisfies Axioms E, A, C and I but fails to satisfy the remaining Axioms R, S, T.

M(R, S, I): *A model of a geometry in which Axioms R, S and I alone hold besides Axiom E.*

Let L be the half real line $0 \leq x < \infty$ with the origin O .

DEFINITION OF $AB=A'B'$: Let $f(x)=x^3$.

In case $A=O$ or $A'=O$, let $AB=A'B'$ if $f(b)-f(a)=f(b')-f(a')$.

In case $A \neq O$ and $A' \neq O$, let $AB=A'B'$ if $|AB|=|A'B'|$.

This model satisfies Axioms E, R, S and I but fails to satisfy the remaining Axioms T, A, C.

M(R, S, C, I): *A model of a geometry in which Axioms R, S, C and I alone hold besides Axiom E.*

Let L be the half real line $0 \leq x < \infty$.

For any $s > 0$ make correspond to each x with $0 \leq x \leq s$ an x' with $s \leq x' \leq 3s$ and vice versa, by the relation

$$\frac{2x+x'}{3} = s.$$

Call this correspondence σ a *skew symmetrization* with *centre* s .

$$L \quad \begin{array}{ccccccc} O & X & S & & X' & & \\ \hline 0 & x & s & & x' & & 3s \end{array}$$

It should be observed that for any pair of non negative numbers a and b there is one and only one skew symmetrization σ that interchanges a and b : $\sigma(a)=b$, $\sigma(b)=a$; indeed, if $a < b$, then we are only to set

$$\frac{2a+b}{3} = s.$$

DEFINITION OF $AB=A'B'$:

Let $AB=A'B'$, if there is a skew symmetrization σ such that $\sigma(a)=b'$, $\sigma(b)=a'$, where a, b, a' and b' are coordinates of A, B, A' and B' respectively.

Clearly Axiom E holds by the above observation. Likewise for Axioms R, S.

As for Axiom C, let $AB=B'A', BC=C'B'$. Then there must be one and only one skew symmetrization σ with centre s that carries A to A', B to B' and C to C' , hence $AC=C'A'$.

Axiom I follows then from Theorem 2.

→T: To show that Axiom T does not hold, let O, A_1, A_3, A_5 , and A_7 be points with coordinates 0, 1, 3, 5 and 7 respectively. Then $OA_1=A_1A_3$, $A_1A_3=A_3A_7$ but $OA_1 \neq A_3A_7$. Therefore $OA_1=A_3A_7$ fails to hold, as will be seen by a simple calculation.

→A: Axiom A does not hold, for otherwise T would follow by Theorem 10 which asserts $S, A, I \Rightarrow T$.

REMARK. Instead of $0 \leq x < \infty$ in our $M(R, S, C, I)$ we may take as L the real line $-\infty < x < \infty$.

In this case the skew symmetrization σ should be modified as follows, according as the centre s lies $< 0, = 0$ or > 0 , the range of symmetrization spreading along the whole line:

Case I: $s > 0$.

(i) Points x with $0 \leq x \leq s$ and x' with $s \leq x' \leq 3s$ interchange by the relation

$$\frac{2x+x'}{3} = s.$$

(ii) Points x with $x \leq 0$ and x' with $x' \geq 3s$ interchange by the relation $x+x'=3s$.

Case II: $s < 0$.

(i) Points x with $s \leq x \leq 0$ and x' with $3s \leq x' \leq s$ interchange by the same relation

$$\frac{2x+x'}{3} = s$$

as above.


(ii) Points x with $x \geq 0$ and x' with $x' \leq 3s$ interchange by the same relation $x+x'=3s$ as above.

Case III: $s=0$. For any real numbers, points x and x' interchange by the relation $x'+x=0$.

M(C): A model of a geometry in which Axiom C alone holds besides Axiom E.

Let L and \bar{L} be the half real lines $0 \leq x < \infty$ and let φ be a mapping of points X of L with coordinates x onto points \bar{X} of \bar{L} with coordinates \bar{x} such that $\bar{x}=3x$ and let i be an identical mapping $\bar{x}=x$.

DEFINITION of $AB=A'B'$: Given AB and $A'B'$ on L , let $AB=A'B'$ if and only if $i(A)i(B)=\varphi(A')\varphi(B')$ on \bar{L} in the sense of the Model $M(R, S, C, I)$.



Verification that this gives an $M(C)$ is easy.

$M(R, S, T, A)$: A model of a geometry in which Axioms R, S, T and A alone hold besides Axiom E.

Let L be the real line $-\infty < x < \infty$.


DEFINITION OF $AB=A'B'$:

For any integer n consider for a pair of real numbers x and y in $[n-1, n)$ with $x < y$ a function $d(x, y)$ defined by

$$d(x, y) = e^{1/(n-y)} - e^{1/(n-x)}.$$

In the following a, b, a', b' etc. denote the coordinates of points A, B, A', B' respectively as usual.


I. In case $a, b \in [n-1, n)$ and $a', b' \in [m-1, m)$, provided m, n denote arbitrary integers, let $AB=A'B'$ if $d(a, b) = d(a', b')$.




II. In case

$$a \in [n-1, n), \quad b \in [n+p-1, n+p), \\ a' \in [m-1, m), \quad b' \in [m+p-1, m+p)$$

for any natural number p , let $AB=A'B'$ if $d(n+p-1, b) = d(m+p-1, b')$.



Especially then, $AB=A'B'$ if $a \in [n-1, n)$, $b=n$ and $a' \in [n-1, n)$, $b'=n$ for any choice of a and a' .



E, R, S: Clearly Axioms E, R and S hold.

T: To see that Axiom T holds, let A, B, A', B', A'' and B'' be points with

coordinates, a, b, a', b', a'' and b'' respectively such that $AB=A'B', A'B'=A''B''$.

If $a, b \in [n-1, n)$ for some integer n , then by the definition of equality, $a', b' \in [n'-1, n')$ and $a'', b'' \in [n''-1, n'')$ for some integers n' and n'' . Then we have $d(a, b)=d(a', b')$ and $d(a', b')=d(a'', b'')$, hence $d(a, b)=d(a'', b'')$, therefore $AB=A''B''$.

If $a \in [n-1, n), b \in [m-1, m)$ for some integers n and m with $n < m$, then as before $a' \in [n'-1, n'), b' \in [m'-1, m'), a'' \in [n''-1, n''), b'' \in [m''-1, m'')$. Then we have by the definition of $AB=A'B'$ and $A'B'=A''B''$, $d(m-1, b)=d(m'-1, b'), d(m'-1, b')=d(m''-1, b'')$, hence $d(m-1, b)=d(m''-1, b'')$, therefore $AB=A''B''$.

A: Similarly for Axiom A.

→C, →I: To see that Axioms C and I do not hold, let A, B, A' and B' be points with coordinates a, b, a' and b' respectively such that

$$a \in [n-1, n), b = n, a' \in (n, n+1), b' = n+1.$$

Then by definition $AB=A'B'$ but not $AA'=BB'$, thus Axiom I does not hold. Axiom C fails to hold too.

Notice that this model $M(R, S, T, A)$ is non-Archimedean.

$M(S, C)$: A model of a geometry in which Axioms S and C alone hold besides Axiom E.

Let L be a linearly ordered space with points A_n^i, i, n ranging over all integers $0, \pm 1, \pm 2, \dots$, with the order relation

- (i) $A_m^i < A_n^i$, if $m < n$,
- (ii) $A_m^i < A_n^j$, if $i < j$ (for any integers m, n .)

DEFINITION OF $AB=A'B'$: let $A_m^i A_n^j = A_m^{i'} A_n^{j'}$, if

- (i) $j-i=j'-i'=0$ and $n-m=n'-m' > 0$,
- or (ii) $j-i=j'-i'$ is an even number > 0 and $m-n=m'-n'$,
- or (iii) $j-i=j'-i'$ is an odd number > 0 and $m+n+m'+n' = -1$.

E, S: Axioms E and S evidently hold.

C: To see that Axiom C holds, let

$$A_m^i A_n^j \equiv A_m^i A_q^p + A_q^p A_n^j, \tag{1}$$

$$A_m^{i'} A_n^{j'} \equiv A_m^{i'} A_q^{p'} + A_q^{p'} A_n^{j'}, \tag{2}$$

and

$$A_m^i A_q^p = A_q^{p'} A_n^{j'}, \tag{3}$$

$$A_q^p A_n^j = A_m^{i'} A_q^{p'}. \tag{4}$$

Then by the definition (i), (ii), (iii) of $=$, we have first of all from (3) and (4)

$$p-i = j'-p', \tag{5}$$

$$j-p = p'-i', \tag{6}$$

whence

$$j-i = j'-i' \tag{7}$$

follows. Next we have to consider three cases:

(i) The case: $j-i=j'-i'=0$. We have from (1) and (2):

$$p=i=j \quad \text{and} \quad p'=i'=j'.$$

From (3) and (4) we have then

$$q-m = n'-q', \quad n-q = q'-m',$$

whence

$$m-n = m'-n',$$

which is evidently different from 0 because $A_m^i < A_n^i$.

Thus in this case we have

$$A_m^i A_n^j = A_m^{i'} A_n^{j'} \tag{*}$$

(ii) The case: $j-i=j'-i'$ is an even number >0 .

Subcase 1): If $p-i$ is even, so is $j-p=(j-i)-(p-i)$ and we have from (3) and (4) by the definition of $=$,

$$m-q = q'-n',$$

$$q-n = m'-q',$$

whence

$$m-n = m'-n',$$

and (*) is proved.

Subcase 2): If $p-i$ is odd, so is $j-p=(j-i)-(p-i)$ and from (3) and (4) we obtain

$$m+q+q'+n' = -1,$$

$$q+n+m'+q' = -1,$$

whence

$$m-n = m'-n',$$

and (*) is again proved.

(iii) The case: $j-i=j'-i'$ is an odd number >0

Subcase 1): If $p-i$ is even, then $j-p=(j-i)-(p-i)$ is odd and we have from (3) and (4)

$$m-q = q'-n',$$

$$q+n+m'+q' = -1,$$

whence

$$m+n+m'+n' = -1,$$

and again (*) holds.

Subcase 2): If $p-i$ is odd, then $j-p$ is even and similarly as above we have (*).

The following examples show that Axioms R, T, A and I do not hold true.

→R: $A_0^1 A_0^2 = A_0^1 A_{-1}^2$ but not $A_0^1 A_0^2 = A_0^1 A_0^2$, so Axiom R fails to hold.

→T: $A_0^1 A_0^2 = A_0^1 A_{-1}^2$, $A_0^1 A_{-1}^2 = A_1^1 A_{-1}^2$ and $A_0^1 A_0^2 = A_1^1 A_{-2}^2$ but not $A_0^1 A_0^2 = A_1^1 A_{-1}^2$, so Axiom T fails to hold.

→A: $A_0^1 A_{-1}^2 = A_0^1 A_0^2$, $A_{-1}^2 A_0^2 = A_0^2 A_1^2$ and $A_0^1 A_0^2 = A_0^1 A_{-1}^2$ but not $A_0^1 A_0^2 = A_0^1 A_1^2$, so Axiom A fails to hold.

→I: $A_0^1 A_1^1 = A_{-1}^2 A_0^2$ ($A_0^1 < A_1^1 < A_{-1}^2 < A_0^2$) and $A_0^1 A_{-1}^2 = A_1^1 A_{-1}^2$ but not $A_0^1 A_{-1}^2 = A_1^1 A_0^2$, so Axiom I fails to hold.

A model $M(R, C, I)$ will be given in the second part of this paper.

4. Proof of Main Theorem

I. *T and C are independent, and $T, C \Rightarrow R, S, A, I$.*

Proof. (i) $T, C \Rightarrow A$ by Theorem 3.

(ii) $T, A \Rightarrow R$ by Theorem 4.

(iii) $T, R \Rightarrow S$ by Theorem 1.

(iv) $R, C \Rightarrow I$ by Theorem 2.

By Models $M(T)$ and $M(C)$ we see that T and C are independent.

II. *T and I are independent, and $T, I \Rightarrow R, S, A, C$.*

Proof. (i) $T, I \Rightarrow R$ by Theorem 6.

(ii) $T, R \Rightarrow S$ by Theorem 1.

(iii) $T, I \Rightarrow A$ by Theorem 7.

(iv) $T, I \Rightarrow C$ by Theorem 8.

By Models $M(T)$ and $M(I)$ we see that T and I are independent.

III. *S, A and I are independent, and $S, A, I \Rightarrow R, T, C$.*

Proof. (i) $S, A \Rightarrow R$ by Theorem 9.

(ii) $S, A, I \Rightarrow T$ by Theorem 10.

(iii) $S, A, I \Rightarrow C$ by Theorem 11.

1) $M(R, S, A)$ shows that S and A do not yield I.

2) $M(R, S, C, I)$ shows that S and I do not yield A.

3) $M(A, C, I)$ shows that A and I do not yield S.

Hence S, A and I are independent.

IV. *S, A and C are independent, and $S, A, C \Rightarrow R, T, I$.*

- Proof. (i) $S, A \Rightarrow R$ by Theorem 9.
 (ii) $R, C \Rightarrow I$ by Theorem 2.
 (iii) $S, A, I \Rightarrow T$ by Theorem 10.

- 1) $M(R, S, A)$ shows that S and A do not yield C .
- 2) $M(R, S, C, I)$ shows that S and C do not yield A .
- 3) $M(A, C, I)$ shows that A and C do not yield S .

Hence S, A and C are independent.

V. R, A and C are independent, and $R, A, C \Rightarrow S, T, I$.

- Proof. (i) $R, C \Rightarrow I$ by Theorem 2.
 (ii) $R, A, C \Rightarrow S$ by Theorem 12.
 (iii) $S, A, I \Rightarrow T$ by Theorem 10.

- 1) $M(R, S, A)$ shows that R and A do not yield C .
- 2) $M(R, S, C, I)$ shows that R and C do not yield A .
- 3) $M(A, C, I)$ shows that A and C do not yield R .

Hence R, A and C are independent.

REMARK: By the use of our Theorems and Models it may easily be proved that there is no further theorem of the above type I-V.

5. Tables

[1] Basic Theorems²⁾

T_1	R	S	T			
T_1	R	S	T			
T_2	R				C	I
T_3			T	A	C	
T_4	R		T	A		
T_5	R		T		C	
T_6	R		T			I
T_7			T	A		I
T_8			T		C	I
T_9	R	S		A		
T_{10}		S	T	A		I
T_{11}		S		A	C	I
T_{12}	R	S		A	C	
T_{13}	R	S			C	I

2) In the following tables **R, S, T** indicates for example that Axiom **S** follows from Axioms **R, T** and Axiom **E**. T_n means Theorem n .

[2] Models

M(R)	R	\rightarrow S	\rightarrow T	\rightarrow A	\rightarrow C	\rightarrow I
M(S)	\rightarrow R	S	\rightarrow T	\rightarrow A	\rightarrow C	\rightarrow I
M(T)	\rightarrow R	\rightarrow S	T	\rightarrow A	\rightarrow C	\rightarrow I
M(A)	\rightarrow R	\rightarrow S	\rightarrow T	A	\rightarrow C	\rightarrow I
M(C)	\rightarrow R	\rightarrow S	\rightarrow T	\rightarrow A	C	\rightarrow I
M(I)	\rightarrow R	\rightarrow S	\rightarrow T	\rightarrow A	\rightarrow C	I
M(A, C)	\rightarrow R	\rightarrow S	\rightarrow T	A	C	\rightarrow I
M(S, C)	\rightarrow R	S	\rightarrow T	\rightarrow A	C	\rightarrow I
M(S, I)	\rightarrow R	S	\rightarrow T	\rightarrow A	\rightarrow C	I
M(R, S, A)	R	S	\rightarrow T	A	\rightarrow C	\rightarrow I
M(R, A, I)	R	\rightarrow S	\rightarrow T	A	\rightarrow C	I
M(R, S, T)	R	S	T	\rightarrow A	\rightarrow C	\rightarrow I
M(A, C, I)	\rightarrow R	\rightarrow S	\rightarrow T	A	C	I
M(R, S, I)	R	S	\rightarrow T	\rightarrow A	\rightarrow C	I
M(R, S, C, I)	R	S	\rightarrow T	\rightarrow A	C	I
M(R, S, T, A)	R	S	T	A	\rightarrow C	\rightarrow I
M(R, C, I) ³⁾	R	\rightarrow S	\rightarrow T	\rightarrow A	C	I

[3] Main Theorem⁴⁾

I	R	S	T	A	C	I
II	R	S	T	A	C	I
III	R	S	T	A	C	I
IV	R	S	T	A	C	I
V	R	S	T	A	C	I

NIHAMA TECHNICAL COLLEGE, NIHAMA
SOPHIA UNIVERSITY, TOKYO

References

- [1] D. Hilbert: *Grundlagen der Geometrie*, 9te Aufl., Stuttgart, 1962.
- [2] U. Morin-F. Busulini: *Prova esistenziale della geometria generale sopra una retta*, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fiz. Mat. Natur.* **35** (1963), 269–273.
- [3] H. Terasaka: *Shotō Kikagaku* (in Japanese), Tokyo, 1952.

3) M(R, C, I) will be given in the second part of this paper.

4) For the notation, see Main Theorem, p. 270.