



Title	On a weakly unknotted 2-sphere in a simply-connected 4-manifold
Author(s)	Matumoto, Takao
Citation	Osaka Journal of Mathematics. 1984, 21(3), p. 489-492
Version Type	VoR
URL	<a href="https://doi.org/10.18910/12666">https://doi.org/10.18910/12666</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Matumoto, T.  
Osaka J. Math.  
21 (1984), 489-492

## ON A WEAKLY UNKNOTTED 2-SPHERE IN A SIMPLY-CONNECTED 4-MANIFOLD

TAKAO MATUMOTO

Dedicated to Professor Minoru Nakaoka on his sixtieth birthday

(Received May 27, 1983)

### Introduction

The purpose of this note is to present the following criterion for unknotting in a weak sense which gives us a simple geometric proof of Theorem of Kawauchi stated below.

**Theorem 1.** *Let  $M$  be a smooth 1-connected 4-manifold and  $S^2$  a smoothly embedded 2-sphere in  $M$ . Suppose that  $\pi_1(M - S^2) \cong \mathbb{Z}$  and  $S^2 \simeq 0$  in  $M$ . Then,  $S^2$  is unknotted in  $M \# (\# S^2 \times S^2)$  for some  $n \geq 0$ .*

Here  $S^2$  is called unknotted if there is a smoothly embedded  $D^3$  which is bounded by  $S^2$ . As a corollary we shall give a proof of Theorem of Kawauchi. His original proof uses the partial Poincaré duality associated to infinite cyclic covering (see [3], [4] and Suzuki [9, Th. 8.6]). Other proofs are founded in [1], [8] and [10].

**Corollary** (Theorem of Kawauchi). *Let  $S^2$  be a smoothly embedded 2-sphere in the 4-sphere  $S^4$ . Suppose that  $\pi_1(S^4 - S^2) \cong \mathbb{Z}$ . Then, it is algebraically unknotted, i.e.  $S^4 - S^2 \simeq S^1$ .*

Is a smooth 2-knot with  $\pi_1(S^4 - S^2) \cong \mathbb{Z}$  unknotted? This is a unsolved question. We stabilize the problem by making connected sum of the ambient manifold with  $\#(S^2 \times S^2)$  and another stabilization may be done by making connected sum of the embedded manifold  $S^2$  with trivially embedded  $\#(S^1 \times S^1)$ . There is a result due to [2].

**Theorem 2** (Hosokawa-Kawauchi [2]). *Under the same assumption of Theorem 1,  $S^2$  surgered by attaching  $n$  trivially embedded 1-handles is unknotted in  $M$  for some  $n \geq 0$ .*

We refer the reader to [ibid] for the precise meaning of trivial (=trivially embedded) 1-handles and unknottedness of surfaces. We shall give also a

proof of Theorem of Kawauchi using this theorem in the last section.

### 1. Proof of Theorem 1

Since  $S^2 \simeq 0$  in  $M$ , we have  $S^2 \times D^2 \subset M$ . And  $* \times \partial D^2 \subset M - S^2$  gives a generator of  $\pi_1(M - S^2) \cong \mathbb{Z}$ . This implies that there exists a map  $f: M - S^2 \times \dot{D}^2 \rightarrow S^1$  which is an extension of the projection  $S^2 \times \partial D^2 \rightarrow S^1$ . We make  $f$  transversely regular at a point of  $S^1$  and get a connected smooth 3-manifold  $N \subset M$  such that  $\partial N = S^2$  in  $M$ .

In case  $M$  has a spin structure, we can restrict the spin structure of  $M$  on  $N$  and extend it over  $N \cup D^3$ , because the spin structure is determined by a framing of the stable tangent bundle over the 2-skelton (cf. Milnor [7]). Since the 3-dimensional spin cobordism group vanishes [ibid], we have a smooth spin cobordism  $(W^4; N^3, D^3)$  relative to the boundary. We may assume that  $W^4$  is the union of the elementary cobordisms consisting of one of 1-handles, 2-handles and 3-handles in this order. The elementary cobordism  $N \times I \cup$  (1-handle) is easily embedded in  $M$  and the spin structure on the other boundary is compatible with that of  $M$ . By an inductive argument on the number of 1-handles, the level manifold  $N_1$  just above all the 1-handles is embedded in  $M$  and  $\partial N_1 = S^2$ . Remark that the spin structure of  $N_1 \subset W$  is compatible with that of  $N_1 \subset M$ . The elementary cobordism  $N_1 \times I \cup$  (2-handle) cannot be embedded in  $M$  but can be embedded in  $M \# (S^2 \times S^2)$ . In fact, we take  $S^1 \subset N_1$  which is the boundary of the axis of the 2-handle. Then,  $S^1 \simeq 0$  in  $M - S^2$ , because  $S^1$  does not link with  $S^2$  and  $\pi_1(M - S^2) \cong \mathbb{Z}$ . The framing of  $S^1 \times D^3$  is uniquely determined by the spin structure of  $N_1$  and the surgery along this framed  $S^1 \times D^3$  changes  $M - S^2$  into  $(M - S^2) \# (S^2 \times S^2)$  because of the choice of the spin structure. Of course, the spin structure on the other boundary is compatible with that of  $M \# (S^2 \times S^2)$ . The level manifold  $N_2$  just above all the 2-handles is embedded in  $M \# (\#_k S^2 \times S^2)$  and  $\partial N_2 = S^2$ , where  $k$  is equal to the number of the 2-handles of  $(W, N)$ . We note that there is a diffeomorphism  $h: (\#_l S^1 \times S^2 - \dot{D}^3, \partial) \rightarrow (N_2, \partial)$ , where  $l$  is the number of 3-handles of  $(W, N)$  i.e. 1-handles of  $(W, D^3)$ . Take the component  $S^1$  of  $S^1 \times S^2$  and consider  $h(S^1) \subset N_2 \subset M \# (\#_k S^2 \times S^2)$ . As before,  $h(S^1) \simeq 0$  in  $M \# (\#_k S^2 \times S^2) - S^2$ . The spin structure of  $N_2$  induces a framing of the tubular neighborhood of  $h(S^1)$  so that the surgery along  $h(S^1)$  changes  $M \# (\#_k S^2 \times S^2) - S^2$  into  $(M \# (\#_k S^2 \times S^2) - S^2) \# S^2 \times S^2$ . Then  $N'_2 \cong (\#_{l-1} S^1 \times S^2 - \dot{D}^3)$  is easily embedded in  $M \# (\#_{k+1} S^2 \times S^2)$  such that  $\partial N'_2 = S^2$ . By induction we get a smooth submanifold  $N_3$  of  $M \# (\#_{k+1} S^2 \times S^2)$  such that  $\partial N_3 = S^2$  and  $N_3$  is

diffeomorphic to  $D^3$ . This means that  $S^2$  is unknotted in  $M\#(\#S^2 \times S^2)$ .

In the other case that  $w_2(M) \neq 0$ , we have only to remark that the surgery along the trivial circle with any framing gives us  $M\#(S^2 \times S^2)$ . Since the closed 3-manifold  $N \cup D^3$  is orientable and the tangent bundle is trivial, there is a spin structure on  $N \cup D^3$  and any choice of the spin structure on  $N \cup D^3$  leads to the same proof as above. q.e.d.

## 2. Proof of Corollary

Let  $\tilde{E}$  be the universal covering space of  $E = S^4 - S^2$ . Then,  $E\#(\#S^2 \times S^2)$  is diffeomorphic to  $S^1 \times R^3 \#(\#S^2 \times S^2)$  by Theorem 1. Hence, we have  $H_*(\tilde{E}; Z) = 0$  for  $* \geq 3$  and there is an isomorphism as  $Z[Z]$ -modules,  $\alpha: H_2(\tilde{E}; Z) \oplus (Z[Z])^{2n} \rightarrow (Z[Z])^{2n}$ , where  $Z[Z]$  is the group ring of  $Z$  over  $Z$ . (From this fact the argument in the last paragraph of [5] completes the proof. We present here a little modified one.) Let  $\beta = p \circ \alpha^{-1}$  where  $p: H(\tilde{E}; Z) \oplus (Z[Z])^{2n} \rightarrow (Z[Z])^{2n}$  is the projection onto the 2nd factor. Since  $\beta$  is a surjection onto a free  $Z[Z]$ -module, there exists a  $Z[Z]$ -module homomorphism  $\gamma$  such that  $\beta \circ \gamma = \text{id}$ . But, since  $Z[Z]$  can be embedded in a field  $Q(t)$ , the right inverse matrix  $\gamma$  over  $Q(t)$  is also a left inverse of  $\beta$ . In particular,  $\beta$  is an injection and so is  $p = \beta \circ \alpha$ . Hence,  $H_2(\tilde{E}; Z) = 0$ , which implies that  $\tilde{E}$  is contractible and  $E$  has the homotopy type of  $S^1$ . q.e.d.

## 3. Further discussions

3.1. In Theorem 1,  $\pi_1(M - S^2) \cong Z$  implies  $S^2 \simeq 0$  in  $M$  if  $M$  is a closed manifold. In fact,  $[f(S^2)] \cap [S^2] = 0$  for any immersion  $f: S^2 \rightarrow M$ , because we can assume that  $f(S^2)$  and  $S^2$  intersect transversally and hence the algebraic intersection number times generator of  $\pi_1(M - S^2) = H_1(M - S^2)$  is zero. By the fact that  $\pi_2(M) = H_2(M)$  and the Poincaré duality this means  $S^2 \simeq 0$  in  $M$ .

3.2. Theorem of Kawauchi is valid for the locally flat topological 2-knot  $S^2$  if it has a normal micro-bundle. In this case  $S^2 \times D^2$  is embedded in  $S^4$  so that the interior of  $\tilde{E} = S^4 - S^2 \times D^2$  is homeomorphic to  $E$ . Then, Kawauchi's proof can be applied to  $\tilde{E}$  to get  $E \simeq S^1$ . Our method is also applicable. In fact, we consider an embedding of  $S^1$  parallel to  $\partial D^2$  in  $\text{Int } \tilde{E}$ . Since  $H^i(\tilde{E} - S^1, \partial \tilde{E}) = H^i(S^1 \times S^2 \times [0, +\infty), S^1 \times S^2 \times 0) = 0$  for any  $i$ , the non-compact 4-manifold  $\tilde{E} - S^1$  admits a smooth structure relative to the boundary  $\partial \tilde{E}$  (see [6, V. 1.4.1]). So we get a smooth embedding of  $S^2$  into a 1-connected smooth 4-manifold  $M$  which is homeomorphic to  $S^4 - S^1$  such that  $\pi_1(M - S^2) \cong Z$  and  $S^2 \simeq 0$  in  $M^4$ . By Theorem 1  $S^2$  is unknotted in  $M\#(\#S^2 \times S^2)$ . Then  $E\#(\#S^2 \times S^2)$  is homeomorphic to  $S^1 \times R^3 \#(\#S^2 \times S^2)$ . This implies  $\tilde{E} \simeq S^1$  by the argument of §2.

3.3. Proof of Corollary by using Theorem 2: Let  $T(n)$  be  $S^2$  surgered by attaching  $n$  trivially embedded 1-handles in  $S^4$ . Then by Theorem 2 we can assume that  $T(n)$  is unknotted. By 3.3 of [5],  $S^4 - T(n)$  has the homotopy type of  $S^1 \vee (\bigvee_{2n} S^2)$ . On the other hand we see in the same way that  $S^4 - T(n) \simeq E \vee (\bigvee_{2n} S^2)$ . Now the same argument as in §2 leads to the conclusion of Corollary.

---

### References

- [1] P. Hirshhorn and J. Ratcliff: *A simple proof of the algebraic unknotting of spheres in codimension two*, Amer. J. Math. **102** (1980), 489–491.
- [2] F. Hosokawa and A. Kawauchi: *Proposals for unknotted surfaces in four-spaces*, Osaka J. Math. **16** (1979), 233–248.
- [3] A. Kawauchi: *On partial Poincaré duality and higher dimensional knots with  $\pi_1 = \mathbb{Z}$* , Master Thesis, Kobe Univ., 1974
- [4] ———: *A partial Poincaré duality theorem for infinite cyclic coverings*, Quart. J. Math. **26** (1975), 437–458.
- [5] A. Kawauchi and T. Matumoto: *An estimate of infinite cyclic coverings and knot theory*, Pacific J. Math. **90** (1980), 99–103.
- [6] R. Kirby and L. Siebenmann: *Foundational essays on topological manifolds, smoothings, and triangulations*, Ann. Math. Studies 88 (1977).
- [7] J. Milnor: *Spin structures on manifolds*, Enseign. Math. **9** (1963), 198–203.
- [8] D. Sumners: *On asphericity of knots of  $S^2$  in  $S^4$* , Preprint.
- [9] S. Suzuki: *Knotting problems of 2-spheres in 4-space*, Math. Sem. Notes Kobe Univ. **4** (1976), 241–371.
- [10] G. Swarup: *An unknotting criterion*, J. Pure Appl. Algebra **6** (1975), 291–296.

Department of Mathematics  
 Faculty of Science  
 Hiroshima University  
 Hiroshima 730, Japan