



Title	Young diagrams and simple constituents of the Specht modules
Author(s)	Hieda, Yoshimasa; Tsushima, Yukio
Citation	Osaka Journal of Mathematics. 43(3) P.475-P.483
Issue Date	2006-09
Text Version	publisher
URL	<a href="https://doi.org/10.18910/12700">https://doi.org/10.18910/12700</a>
DOI	10.18910/12700
rights	

***Osaka University Knowledge Archive : OUKA***

<https://ir.library.osaka-u.ac.jp/repo/ouka/all/>

# YOUNG DIAGRAMS AND SIMPLE CONSTITUENTS OF THE SPECHT MODULES

YOSHIMASA HIEDA and YUKIO TSUSHIMA

(Received October 26, 2004, revised June 9, 2005)

## Abstract

We discuss the simple constituents of Specht module  $S^\lambda$  for the symmetric group  $S_n$  defined over the field of  $p$  elements. We firstly give an easier proof to the result in [6] which asserts that there exists a simple constituent of  $S^\lambda$  with the shape of “a branch” of  $\lambda$  (Theorem 3.3), and secondly give a sufficient condition for  $\lambda$  to have a particular type branch as a constituent (Proposition 3.4).

## 1. Introduction

Let  $n$  be a natural number and  $p$  a prime. Let  $S_n$  be the symmetric group on  $n$  letters and  $L$  a field of characteristic  $p$ . Given a partition  $\lambda$  of  $n$ , we have an  $LS_n$ -module  $S^\lambda$  called the Specht module corresponding to  $\lambda$ , which is not simple in general. However if the partition  $\lambda$  is  $p$ -regular, the head of  $S^\lambda$ , denoted by  $D^\lambda$ , is simple and they cover all the non-isomorphic simple modules as  $\lambda$  runs through the  $p$ -regular partitions of  $n$ .

One of the main concerns about the Specht modules is to have informations about the simple constituents of them. Especially, using information only on  $\lambda$ , we would like to describe a  $p$ -regular partition  $\mu$  for which  $D^\mu$  appears as a constituent of  $S^\lambda$ . For this purpose, it is useful to consider the operations on the partitions  $\lambda$  introduced by James and Murphy [5], each of which is roughly interpreted as a rim hook removal followed by addition on the Young diagram corresponding to  $\lambda$ . We shall call each of the resulting partitions a branch of  $\lambda$ . The Jantzen-Schaper theorem tells that if  $D^\mu$  is a constituent of  $S^\lambda$ , it follows that  $\lambda = \mu$  or  $\mu$  is obtained by making branches successively beginning with  $\lambda$  (cf. [6, Corollary 1]). One of the authors showed that if  $\lambda$  is  $p$ -regular, there is a  $p$ -regular branch  $\mu$  of  $\lambda$  such that  $D^\mu$  is a constituent of  $S^\lambda$  (cf. [6, Theorem 2]). And he gave some applications of the result in [7]. However the proof of the result cited above is rather long and complicated. In this paper we shall show a short proof to it and a result on simple constituents of the Specht modules as a byproduct of the proof.

**2. Preliminary results**

A *partition* of the integer  $n$  is a non-increasing sequence  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_m)$  of non-negative integers whose sum is  $n$ . The *Young diagram*  $[\lambda]$  associated with  $\lambda$  is the set of the ordered pairs  $(i, j)$  of integers, called the *nodes* of  $[\lambda]$ , with  $1 \leq i \leq h$  and  $1 \leq j \leq \lambda_i$ , where  $h$  denotes the largest number such that  $\lambda_h \neq 0$ . They are illustrated as arrays of squares. We denote by  $\lambda'$  the partition conjugate of  $\lambda$ , so  $[\lambda']$  is the transposed diagram of  $[\lambda]$ .

Let  $c$  be a column number of  $[\lambda]$  and  $r$  a positive integer. Then  $\lambda$  is said to be *r-singular on column  $c$*  if there is an integer  $i \geq 0$  such that  $\lambda_{i+1} = \lambda_{i+2} = \dots = \lambda_{i+r} = c$ , and is *r-regular on column  $c$*  if otherwise. We also say that  $\lambda$  is *r-singular* if it is *r-singular* on some column, and is *r-regular* if otherwise. For the convenience of later arguments, we understand that every partition is *r-regular* on column 0. We denote by  $P(n)$  and  $P(n)^0$  the sets of the partitions and *p-regular* partitions of  $n$  respectively. The *dominance order*  $\trianglelefteq$  on  $P(n)$  is defined as follows: given  $\lambda, \mu \in P(n)$ ,  $\lambda \trianglelefteq \mu$  if and only if  $\sum_{1 \leq i \leq j} \lambda_i \leq \sum_{1 \leq i \leq j} \mu_i$  for all  $j \geq 1$ .

The  $(i, j)$ -*hook* of the Young diagram  $[\lambda]$  consists of the  $(i, j)$ -node along with the  $\lambda_i - j$  nodes to the right of it (called the *arm* of the hook) and the  $\lambda'_j - i$  nodes below it. The *length* of the  $(i, j)$ -hook of  $\lambda$  is  $h_{ij}(\lambda) := \lambda_i + \lambda'_j + 1 - i - j$ . An  $(i, j)$ -*rim hook* is a connected part of the rim of  $[\lambda]$  of length  $h_{ij}(\lambda)$  beginning at the node  $(\lambda'_j, j)$ . We also call the integer  $\lambda_i - j$  the *arm length* of the node  $(i, j)$ . Moreover, a hook of  $[\lambda]$  is called a *pillar* if its arm length is zero.

Let  $(b, c)$  is a node of  $[\lambda]$  and suppose that  $a < b$ . We let  $\lambda(a, b, c)$  be the partition of  $n$  obtained from  $\lambda$  by unwrapping the  $(b, c)$ -rim hook of  $[\lambda]$  and wrapping the nodes back with the lowest nodes in the added rim hook lying on row  $a$  (if the resulting partition fails to be a non-increasing sequence of integers,  $\lambda(a, b, c)$  is not defined). We occasionally write  $\lambda(a, b, c, g)$  if the highest node in the added rim hook lies in row  $g$ . We call here each  $\lambda(a, b, c)$  a *branch* of  $\lambda$  and set

$$\Gamma_\lambda := \{ \lambda(a, b, c); v_p(h_{ac}(\lambda)) \neq v_p(h_{bc}(\lambda)) \}, \quad \Gamma_\lambda^0 := \Gamma_\lambda \cap P(n)^0,$$

where  $v_p(m)$  denotes the largest integer  $e$  such that  $p^e$  divides the integer  $m$ .

A branch  $\mu = \lambda(a, b, c)$  is called a *pillar type branch* if the rim hook which has been removed and the rim hook which has been added are both pillars. Suppose that  $\mu = \lambda(a, b, c)$  is a pillar type branch and put  $d := \lambda_a + 1$ ,  $q := h_{bc}(\lambda)$ . Then  $\mu$  is obtained by unwrapping the pillar of  $q$  nodes from column  $c$  and wrapping it back on column  $d$  (with the lowest node on row  $a$ ). Hence we sometimes write  $\mu = \lambda(c \mid d, q)$  for simplicity. For  $\lambda \in P(n)$ , let  $SC(S^\lambda)$  be the set of simple constituents of the Specht module  $S^\lambda$ .

REMARK. Let  $\lambda \in P(n)^0$ . Then if  $\mu = \lambda(a, b, c)$  is a pillar type branch of  $[\lambda]$ , we have  $h_{bc}(\lambda) \leq p - 1$ . Hence  $\mu$  lies in  $\Gamma_\lambda$  if and only if  $h_{ac}(\lambda)$  is divisible by  $p$ .

Now we list below some results for later use.

**Theorem 2.1** ([2], [3]). *Let  $\lambda \in P(n)^0$ . Then  $S^\lambda$  is simple if and only if  $v_p(h_{ac}(\lambda)) = v_p(h_{bc}(\lambda))$  for all  $a, b, c \geq 1$ .*

**Theorem 2.2** (Carter and Payne [1]). *Suppose that  $\alpha := \lambda(c \mid d, q)$  be a pillar type branch of  $\lambda$  and let  $a$  be the row index of  $[\lambda]$  such that  $d = \lambda_a + 1$ . Put  $e := v_p(h_{ac}(\lambda))$ . If  $p^e > q$ , we have*

$$\text{Hom}_G(S^\alpha, S^\lambda) \neq 0.$$

*In particular, it follows that  $D^\alpha \in \text{SC}(S^\lambda)$  if  $\alpha$  is  $p$ -regular.*

REMARK. The above statement is slightly different from the corresponding theorem in [1], but can be deduced easily from it. In fact, if  $\lambda$  and  $\alpha$  are the same as above then with the languages in [1],  $\lambda'$  is obtained from  $\alpha'$  by raising  $q$  nodes from row  $d$  to row  $c$ , whence we have  $\text{Hom}_G(S^{\lambda'}, S^{\alpha'}) \neq 0$ . The rest of the proof will be done by routine arguments, using that  $S^{\lambda'}$  is isomorphic to the  $L$ -dual of  $S^\lambda \otimes S^{(1^n)}$  ([2, Theorem 8.15]).

**Theorem 2.3** ([4, Theorem 6]). *Let  $\lambda, \mu$  be partitions of  $n$  with  $\lambda$   $p$ -regular. Suppose that there is a number  $k$  ( $1 \leq k \leq \lambda_1, \mu_1$ ) such that the subdiagrams consisting of the first  $k$  columns of  $[\lambda]$  and  $[\mu]$  are the same and that each has  $m$  nodes. Let  $[\widehat{\lambda}]$  ( $[\widehat{\mu}]$  resp.) be the subdiagram to the right of column  $k$  of  $[\lambda]$  ( $[\mu]$  resp.). Then the composition multiplicity of  $D^\lambda$  in  $S^\mu$  as  $S_n$ -modules equals the composition multiplicity of  $D^{\widehat{\lambda}}$  in  $S^{\widehat{\mu}}$  as  $S_{n-m}$ -modules.*


**Proposition 2.4** (Jantzen-Schaper, cf. [6, Corollary 1]). *Let  $\lambda \in P(n)$  and let  $\mu$  be a minimal element of  $\Gamma_\lambda$  with respect to the dominance order. If  $\mu$  is  $p$ -regular,  $D^\mu \in \text{SC}(S^\lambda)$ .*

**Proposition 2.5** ([6, Proposition 3]). *Let  $\lambda \in P(n)^0$  and let  $[\mu]$  be the diagram to the right of the first column of  $[\lambda]$ . If  $S^\mu$  is simple,  $\Gamma_\lambda$  has no  $p$ -singular partition.*

### 3. Finding simple constituents of Specht modules

We shall show a short proof to Theorem 2 of [6] and a result on simple constituents of the Specht modules. First we show

**Lemma 3.1.** *Let  $\lambda \in P(n)^0$ . If there is a pillar type branch  $\mu = \lambda(a, b, c) \in \Gamma_\lambda$  such that  $\mu$  is  $p$ -regular on column  $c - 1$ , there is a pillar type branch  $\widetilde{\lambda}$  in  $\Gamma_\lambda^0$ .*



Proof. We put  $r := h_{bc}(\lambda) (\leq p - 1)$  and  $f := \lambda_a$ . Note that  $h_{ac}(\lambda)$  is a multiple of  $p$  since  $\mu \in \Gamma_\lambda$ . We may assume that  $\mu$  is  $p$ -singular, so  $\mu$  is  $p$ -singular on column  $f + 1$  by the assumption. (In the above diagram a circle in a node indicates that the hook length at the node is divisible by  $p$ .)

Namely  $a - \lambda'_{f+2} \geq p$ , so  $a - p + 1 > \lambda'_{f+2}$ . Put  $s_1 := a - p + 1 - \lambda'_{f+2} (\geq 1)$ . Then  $r - s_1 = (p - 1) - (a - r) + \lambda'_{f+2} = (p - 1) - (\lambda'_{f+1} - \lambda'_{f+2}) \geq 0$ , so  $r \geq s_1$ . Now let  $\mu(1) = \lambda(c \mid f + 2, s_1)$ , which lies in  $\Gamma_\lambda$  since  $h_{a-p+1,c}(\lambda)$  is divisible by  $p$ . Note that  $\mu(1)$  is  $p$ -regular on column  $c - 1$ . If  $\mu(1)$  is  $p$ -regular, we may take  $\mu(1)$  as  $\tilde{\lambda}$ . Hence we may assume that  $\mu(1)$  is  $p$ -singular, so  $\lambda'_{f+2} \neq 0$  and  $\mu(1)$  is  $p$ -singular on column  $f + 2$ . Namely  $(a - p + 1) - \lambda'_{f+3} \geq p$ , so  $a - 2p + 2 > \lambda'_{f+3}$ . Put  $s_2 := a - 2p + 2 - \lambda'_{f+3} (\geq 1)$ . Then  $s_1 - s_2 = (p - 1) - (\lambda'_{f+2} - \lambda'_{f+3}) \geq 0$ , so  $s_1 \geq s_2$ . Now let  $\mu(2) = \lambda(c \mid f + 3, s_2)$ , which lies in  $\Gamma_\lambda$  since  $h_{a-2p+2,c}(\lambda)$  is divisible by  $p$ . Note that  $\mu(2)$  is also  $p$ -regular on column  $c - 1$ . By repeating similar arguments we finally obtain a  $p$ -regular pillar type branch  $\mu(i)$  for some  $i$ , completing the proof of the lemma.  $\square$

**Lemma 3.2.** *Let  $\lambda \in P(n)^0$ . If there is a branch  $\mu = \lambda(a, b, c) \in \Gamma_\lambda$  with  $c \geq 2$  such that  $\mu$  is  $p$ -singular on column  $c - 1$ , there is a pillar type branch  $\tilde{\lambda}$  in  $\Gamma_\lambda^0$ .*


Proof. We may assume that  $\mu$  is chosen so that  $c$  is the smallest and put  $r := \lambda'_{c-1} - \lambda'_c$ . Note that the  $(b, c)$ -rim hook of  $[\lambda]$  is a pillar if and only if  $b > \lambda'_{c+1}$ .

CASE I.  $r \leq p - 2$ .

As  $\mu$  is  $p$ -singular on column  $c - 1$ ,  $\lambda'_{c-1} - \lambda'_{c+1} \geq p - 1$ . Put  $x := (\lambda'_{c-1} - p + 2) - \lambda'_{c+1}$  and  $y := \lambda'_c - (\lambda'_{c-1} - p + 2)$ , so  $x \geq 1$  and  $r + y = p - 2$ .

SUBCASE (i)  $x + y \leq p - 2$ .

We have that  $x \leq r$  from  $x + y \leq p - 2 = r + y$ . Now let  $\gamma = \lambda(c - 1 \mid c + 1, x) \in \Gamma_\lambda$ . If  $\gamma$  is  $p$ -regular, we may take  $\gamma$  as  $\tilde{\lambda}$ . Hence we may assume that  $\gamma$  is  $p$ -singular.




Then by the minimality of  $c$ ,  $\gamma$  must be  $p$ -regular on column  $c - 2$  and there is a pillar type branch  $\tilde{\lambda} \in \Gamma_\lambda^0$  by Lemma 3.1, as asserted. (In the above diagrams the boldface rim hooks will be removed to make  $\mu$ .)

SUBCASE (ii)  $x + y = p - 1$ .

We have  $x = r + 1$ . As  $\mu \in \Gamma_\lambda$ , either  $h_{ac}(\lambda)$  or  $h_{bc}(\lambda)$  is divisible by  $p$ . Let  $i = a$  or  $i = b$  according as  $h_{ac}(\lambda)$  is divisible by  $p$  or not. Let furthermore  $f = \lambda_i$  and  $s_i = i - \lambda'_{f+1}$ . Then we see that  $i \leq \lambda'_{c+1}$  since  $h_{ic}(\lambda)$  is divisible by  $p$ . Since  $s_i \leq p - 1$ , we can make the pillar type branch  $\gamma = \lambda(c \mid f + 1, s_i) \in \Gamma_\lambda$ . If  $s_i + r < p$ ,  $\gamma$  is  $p$ -regular on column  $c - 1$  and the assertion follows by Lemma 3.1. Now suppose that  $s_i + r \geq p$  and put  $t_i = s_i - (p - r - 1)$ , so  $t_i \geq 1$ . Also  $\lambda'_{f+1} = i - s_i < i - (p - r - 1) = i - p + r + 1$ . Note that  $t_i \leq r$  since  $r - t_i = p - 1 - s_i \geq 0$ . Hence we can make the pillar type branch  $\delta = \lambda(c - 1 \mid f + 1, t_i)$ , which lies in  $\Gamma_\lambda$  since  $h_{i-p+r+1, c-1}(\lambda) = (r + 1) + h_{ic}(\lambda) + (p - r - 1) = h_{ic}(\lambda) + p$  is divisible by  $p$ . By the minimality of  $c$ ,  $\delta$  is  $p$ -regular on column  $c - 2$  and so there is a pillar type branch  $\tilde{\lambda} \in \Gamma_\lambda^0$  by Lemma 3.1, as asserted.

CASE II.  $r = p - 1$ .

We use the same notation as in subcase (ii). Then  $h_{i, c-1}(\lambda)$  is divisible by  $p$ , since  $h_{i, c-1}(\lambda) = h_{ic}(\lambda) + p$ . In the diagram below, we have  $r = p - 1$ , so we can make the pillar type branch  $\gamma = \lambda(c - 1 \mid f + 1, s_i) \in \Gamma_\lambda$ . By the minimality of  $c$ ,  $\gamma$  is  $p$ -regular on column  $c - 2$  and the assertion follows by Lemma 3.1. This completes the proof of the lemma.




□

Now we are ready to give an alternative proof of the following theorem:


**Theorem 3.3** ([6, Theorem 2]). *Let  $\lambda$  be a  $p$ -regular partition of  $n$ . If  $S^\lambda$  is reducible, there is a  $p$ -regular branch  $\tilde{\lambda} \in \Gamma_\lambda$  such that  $D^{\tilde{\lambda}} \in \text{SC}(S^\lambda)$ .*

*Proof.* Since  $S^\lambda$  is reducible, there is a column number  $c$  such that  $v_p(h_{a,c}(\lambda)) \neq v_p(h_{b,c}(\lambda))$  for some  $a, b$  with  $1 \leq a, b \leq \lambda'_c$ . Let  $c$  be the largest number that satisfies the condition. Let  $[\delta]$  be the subdiagram of  $[\lambda]$  with column  $c$  as the first column,  $[\gamma]$  the remaining diagram and write  $\lambda = (\gamma, \delta)$ . Then every branch in  $\Gamma_\delta$  is  $p$ -regular by Proposition 2.5. Hence, if  $\tilde{\delta}$  is a minimal element of  $\Gamma_\delta$  with respect to the dominance order,  $D^{\tilde{\delta}} \in \text{SC}(S^\delta)$  by a direct consequence of the Jantzen-Schaper theorem (see Proposition 2.4). Put  $\mu := (\gamma, \tilde{\delta}) \in \Gamma_\lambda$ . If  $\mu$  is  $p$ -singular on column  $c - 1$ , then  $c$  must be greater than 1 and there is a pillar type branch  $\tilde{\lambda} \in \Gamma_\lambda^0$  by Lemma 3.2. Thus we have  $D^{\tilde{\lambda}} \in \text{SC}(S^\lambda)$  by the Carter and Payne theorem (see Theorem 2.2). So we may assume that  $\mu$  is  $p$ -regular on column  $c - 1$ . Then  $\mu \in \Gamma_\lambda^0$  and we have  $D^\mu \in \text{SC}(S^\lambda)$  by Theorem 2.3. This completes the proof of the theorem.



□

Now a node  $(b, c)$  is called a  $\langle p, 1 \rangle$ -point of  $[\lambda]$  with arm length one, if  $h_{bc}(\lambda) = p$  and  $h_{\lambda'_{c,c}}(\lambda) = 1$ .



**Proposition 3.4.** Suppose  $p > 2$  and that  $S^\lambda$  is not simple. Let  $\lambda$  be a  $(p - 1)$ -regular partition of  $n$ . Then

- (1) If  $[\lambda]$  has no  $\langle p, 1 \rangle$ -point with arm length one, we have  $\Gamma_\lambda^0 = \Gamma_\lambda$ . Hence  $D^\mu \in \text{SC}(S^\lambda)$  for any minimal element  $\mu$  of  $\Gamma_\lambda$  with respect to the dominance order.
- (2) If  $[\lambda]$  has a  $\langle p, 1 \rangle$ -point with arm length one, there is a pillar type branch  $\mu = \lambda(c \mid d, q)$  such that  $D^\mu \in \text{SC}(S^\lambda)$  for some  $c, d, q$  with  $q \leq p - 2$ .

Proof. (1) The second half follows immediately from the first half and Proposition 2.4. So we need only prove the first half. Suppose the contrary and take a  $p$ -singular branch, say  $\mu = \lambda(a, b, c, g)$ , from  $\Gamma_\lambda$ .


CASE I.  $\mu$  is  $p$ -singular on column  $c - 1$  (hence  $c \geq 2$ ).

Since  $\lambda$  is  $(p - 1)$ -regular, it follows that  $\lambda'_{c-1} - p + 2 \leq \lambda'_c$  and so  $(\lambda'_{c-1} - p + 2, c - 1)$  is a  $\langle p, 1 \rangle$ -point of  $[\lambda]$  with arm length one, being contrary to the assumption.

CASE II.  $\mu$  is  $p$ -singular on column  $\lambda_{g-1}$  (hence  $g \geq 2$ ).

As  $\lambda$  is  $(p - 1)$ -regular, we find easily that  $\lambda_{g-1} = \lambda_g + 1$ . Let  $f = \lambda_{g-1}$ .





Then  $\lambda'_{f-1} - \lambda'_{f+1} \geq p - 1$ , and the node  $(\lambda'_{f-1} - p + 2, f - 1)$  is a  $\langle p, 1 \rangle$ -point of  $[\lambda]$  with arm length one, being contrary to the assumption. This completes the proof of (1).

(2) Let  $(i, j)$  be a  $\langle p, 1 \rangle$ -point of  $[\lambda]$  with arm length one and  $m := \lambda'_{j+1}$ . Then  $i \leq m < \lambda'_j = i + p - 2$  and  $\lambda_m - 1 = \lambda_{m+1}$ .

Now we assume that the above  $(i, j)$  is chosen so that  $j$  is the smallest. Let  $m_1 := \lambda'_{j+2}$  and  $r := h_{m+1, j}(\lambda) = i + p - 2 - m$ . Then  $m_1 < i$  since the node  $(i, j)$  has arm length one. Let  $r_1 := i - m_1$ . Then  $r - r_1 = (p - 2) - (m - m_1) \geq 0$ , so  $r_1 \leq r$ . Therefore we can make the pillar type branch  $\mu = \lambda(j \mid j + 2, r_1) \in \Gamma_\lambda$ . If  $\mu$  is  $p$ -singular on column  $j - 1$ , then  $j$  must be greater than 1 and  $(\lambda'_{j-1} - (p - 2), j - 1)$  is a  $\langle p, 1 \rangle$ -point of  $[\lambda]$  with arm length one, contradicting the minimality of  $j$ . Hence  $\mu$  is  $p$ -regular on column  $j - 1$  and by Lemma 3.1, there is a pillar type branch in  $\Gamma_\lambda^0$ , whence the assertion follows by the Carter and Payne theorem.  $\square$

ACKNOWLEDGEMENT. The authors thank the referee for his/her careful reading of the manuscript.

References

[1] R.W. Carter and M.T.J. Payne: *On homomorphisms between Weyl modules and Specht modules*, Math. Proc. Camb. Phil. Soc. **87** (1980), 419–425.  
 [2] G.D. James: *The Representation Theory of Symmetric Groups*, Lecture notes in Math. **682**, Springer-Verlag, Berlin, 1978.

- [3] G.D. James: *On a conjecture of Carter concerning irreducible Specht modules*, Math. Proc. Camb. Phil. Soc. **83** (1978), 11–17.
- [4] G.D. James: *On the decomposition matrices of the symmetric groups*, III, J. Alg. **71** (1981), 115–122.
- [5] G. James and G. Murphy: *The determinant of the Gram matrix for a Specht module*, J. Alg. **59** (1979), 222–235.
- [6] Y. Tsushima: *On some simple constituents of the Specht modules of the symmetric groups*, J. Alg. **243** (2001), 131–141.
- [7] Y. Tsushima: *On some  $S_R(H)$ -blocks for the symmetric groups*, J. Alg. **270** (2003), 281–287.

Yoshimasa Hieda  
Department of Industrial Systems Engineering: Natural Science  
Osaka Prefectural College of Technology  
Neyagawa, Osaka 572-8572  
Japan  
e-mail: hieda@las.osaka-pct.ac.jp

Yukio Tsushima  
Department of Mathematics  
Osaka City University  
Sumiyoshiku, Osaka 558-8585  
Japan