



Title	On the asymptotic behavior of solutions of certain non-autonomous differential equations
Author(s)	Hara, Tadayuki
Citation	Osaka Journal of Mathematics. 1975, 12(2), p. 267-282
Version Type	VoR
URL	https://doi.org/10.18910/12711
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

ON THE ASYMPTOTIC BEHAVIOR OF SOLUTIONS OF CERTAIN NON-AUTONOMOUS DIFFERENTIAL EQUATIONS

TADAYUKI HARA

(Received February 9, 1973)

1. Introduction

In this paper conditions are obtained under which all solutions of certain real non-autonomous nonlinear differential equations tend to zero as $t \rightarrow \infty$.

Theorem 1 is concerned with the system of differential equations;

$$(1.1) \quad \dot{x} = A(t)x + f(t, x)$$

where x, f are n -dimensional vectors, $A(t)$ is a bounded continuously differentiable $n \times n$ matrix for $t \geq 0$, and $f(t, x)$ is continuous in (t, x) for $t \geq 0$, $\|x\| < \infty$, here $\| \cdot \|$ denotes the Euclidean norm.

Theorem 2 is concerned with the differential equation of the third order;

$$(1.2) \quad \ddot{x} + a(t)f(x, \dot{x}, \ddot{x})\ddot{x} + b(t)g(x, \dot{x}) + c(t)h(x) = p(t, x, \dot{x}, \ddot{x})$$

where $a(t), b(t), c(t)$ are positive continuously differentiable and f, g, h, p are continuous real-valued functions depending only on the arguments shown, and the dots indicate the differentiation with respect to t .

The asymptotic property of solutions of third order differential equations has received a considerable amount of attention during the past two decades, particularly when (1.2) is autonomous. Many of these results are summarized in [11].

A few authors have studied non-autonomous third order differential equations. K. E. Swick [13] considered the following equations

$$(1.3) \quad \ddot{x} + p(t)\ddot{x} + q(t)g(\dot{x}) + r(t)h(x) = 0,$$

$$(1.4) \quad \ddot{x} + f(x, \dot{x}, t)\ddot{x} + q(t)g(\dot{x}) + r(t)h(x) = 0,$$

with the assumption that $q(t), r(t)$ are positive, bounded and monotone decreasing.

In [6], the author studied the asymptotic behavior of the solutions of the equation

$$(1.5) \quad \ddot{x} + a(t)f(x, \dot{x}) \dot{x} + b(t)g(x, \dot{x}) \dot{x} + c(t)h(x) = e(t)$$

under the assumptions that $|a'(t)|$, $|b'(t)|$, $|c'(t)|$ and $|e(t)|$ are integrable and suitable conditions on f, g, h . Here we assume the condition

$$\limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \{|a'(s) + b'_+(s) + c'(s)|\} ds < \gamma$$

where γ is a sufficiently small positive constant.

Conditions on $p(t, x, \dot{x}, \ddot{x})$ are also relaxed, for example $p(t, x, y, z)$ may be unbounded for $x^2 + y^2 + z^2$.

Recently H. O. Tejumola [15] established conditions under which all solutions of equations of the form

$$(1.6) \quad \ddot{x} + f(x, \dot{x}) \dot{x} + g(x, \dot{x}) \dot{x} + h(x) = p(t, x, \dot{x}, \ddot{x})$$

tend to zero as $t \rightarrow \infty$. Theorem 2 develops Tejumola's result [15; Theorem 1] to the non-autonomous equations of the form (1.2).

The main tools used in this work are Theorem A and Liapunov functions. Theorem A would be especially convenient to study the non-autonomous differential equations.

2. Auxiliary theorem

Consider a system of differential equations

$$(2.1) \quad \dot{x} = F(t, x)$$

where x and F are n -dimensional vectors.

Theorem A. Suppose that $F(t, x)$ of (2.1) is continuous in $I \times R^n$ ($I = [0, \infty)$) and that there exists a Liapunov function $V(t, x)$, continuously differentiable in $I \times R^n$, satisfying the following conditions;

(i) $a(\|x\|) \leq V(t, x) \leq b(\|x\|)$, where $a(r) \in CIP$ (the family of continuous and increasing positive definite functions), $a(r) \rightarrow \infty$ as $r \rightarrow \infty$ and $b(r) \in CIP$,

$$(ii) \quad \dot{V}_{(2.1)}(t, x) = \limsup_{h \rightarrow 0^+} \frac{1}{h} \{V(t+h, x+hF(t, x)) - V(t, x)\} \\ \leq -cV(t, x) + \lambda_1(t)V(t, x) + \lambda_2(t)(1 + V(t, x)),$$

where $c > 0$ is a constant and $\lambda_i(t) \geq 0$ ($i=1, 2$) are continuous functions satisfying

$$(2.2) \quad \limsup_{(t,t) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \lambda_1(s) ds < c,$$

$$(2.3) \quad \int_t^{t+1} \lambda_2(s) ds \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Then, any solution $x(t)$ of (2.1) is uniform-bounded and satisfies $x(t) \rightarrow 0$ as $t \rightarrow \infty$.

The following is an immediate consequence of Theorem A.

Corollary. Under the assumptions in Theorem A, if

$$(ii)' \quad \dot{V}_{(2.1)}(t, x) \leq L(t)V(t, x) + \lambda_2(t)(1 + V(t, x)),$$

where $L(t)$ is a continuous function satisfying

$$(2.2)' \quad \limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} L(\tau) d\tau < 0,$$

then any solution $x(t)$ of (2.1) is uniform bounded and satisfies $x(t) \rightarrow 0$ as $t \rightarrow \infty$.

REMARK. The condition (2.3) can be replaced by

$$(2.3)' \quad \int_t^{t+\tau_0} \lambda_2(\tau) d\tau \rightarrow 0 \quad \text{as } t \rightarrow \infty$$

where $\tau_0 > 0$ is an arbitrary constant.

3. Proof of Theorem A

Proof of Theorem A. Let $\varepsilon > 0$ be chosen such that $\frac{\hat{c}}{2} > \varepsilon$ and

$$(3.1) \quad \limsup_{(t, s-t) \rightarrow (\infty, \infty)} \frac{1}{s-t} \int_t^s \lambda_1(\tau) d\tau \leq c - 3\varepsilon.$$

Let $T_0 > 0$ such that

$$\frac{1}{s-t} \int_t^s \lambda_1(\tau) d\tau \leq \limsup_{(t, s-t) \rightarrow (\infty, \infty)} \frac{1}{s-t} \int_t^s \lambda_1(\tau) d\tau + \varepsilon \quad \text{for } t \geq T_0$$

and $s \geq T_0 + t$. T_0 does not depend on t and s .

Using (3.1) and above inequality, we have

$$(3.2) \quad \frac{1}{s-t} \int_t^s \lambda_1(\tau) d\tau \leq c - 2\varepsilon \quad \text{for } t \geq T_0 \text{ and } s \geq T_0 + t.$$

Let $K > 0$ be a constant satisfying $\exp \{cT_0 + \int_0^{T_0} \lambda_1(\tau) d\tau\} \leq K$. Then it is easy to show that for all $s \geq t \geq 0$, we have

$$(3.3) \quad e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} \leq K e^{-2\varepsilon(s-t)}.$$

Now we consider the function $U(t, x)$ defined by

$$(3.4) \quad U(t, x) = V(t, x) e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds,$$

and show that

$$(3.5) \quad e^{-c} a(\|x\|) \leq U(t, x) \leq \frac{K}{\varepsilon} b(\|x\|) \quad \text{for all } t \geq 0 \text{ and } x \in R^n.$$

To verify the right-hand side inequality in (3.5), we use the inequality (3.3). Then we obtain

$$(3.6) \quad e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds \leq e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} K e^{-2\varepsilon(s-t)} ds \\ = \frac{K}{\varepsilon}.$$

A short calculation shows the left-hand side inequality in (3.5), i.e.

$$(3.7) \quad e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds > e^{-\varepsilon t} \int_t^{t+1} e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds \\ \geq e^{-\varepsilon t} \int_t^{t+1} e^{\varepsilon s} e^{-c(t+1-t) + \int_t^t \lambda_1(\tau) d\tau} ds \\ = e^{-c}.$$

Therefore we have

$$(3.8) \quad e^{-c} V(t, x) \leq U(t, x) \leq \frac{K}{\varepsilon} V(t, x)$$

and using the hypothesis (i) of Theorem A, we obtain (3.5).

From (3.4) it follows that

$$\begin{aligned} & \dot{U}_{(2.1)}(t, x) \\ &= \dot{V}_{(2.1)}(t, x) e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds \\ & \quad - \varepsilon V(t, x) e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds \\ & \quad + \{c - \lambda_1(t)\} V(t, x) e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds - V(t, x) \\ & \leq \{-cV(t, x) + \lambda_1(t)V(t, x) + \lambda_2(t)(1 + V(t, x))\} e^{-\varepsilon t} \int_t^\infty e^{\varepsilon s} e^{-c(s-t) + \int_t^s \lambda_1(\tau) d\tau} ds \\ & \quad - \varepsilon U(t, x) + \{c - \lambda_1(t)\} U(t, x) - V(t, x). \end{aligned}$$

Using (3.6), we obtain

$$(3.9) \quad \dot{U}_{(2.1)}(t, x) \leq -\varepsilon U(t, x) + \frac{K}{\varepsilon} \lambda_2(t)(1 + V(t, x)).$$

From (3.8) and (3.9), we have

$$(3.10) \quad \dot{U}_{(2.1)}(t, x) \leq \{-\varepsilon + \frac{K}{\varepsilon} e^c \lambda_2(t)\} U(t, x) + \frac{K}{\varepsilon} \lambda_2(t).$$

Set

$$W(t) = U(t, x(t))$$

where $x(t), x(t_0) = x_0$, is any solution of (2.1). Then the inequality (3.10) implies

$$\frac{d}{dt} W(t) \leq \{-\varepsilon + \frac{K}{\varepsilon} e^c \lambda_2(t)\} W(t) + \frac{K}{\varepsilon} \lambda_2(t).$$

This immediately gives

$$W(t) \leq e^{\int_{t_0}^t g(s) ds} \left(W(t_0) + \int_{t_0}^t e^{-\int_{t_0}^s g(\tau) d\tau} h(s) ds \right)$$

where $g(t) = -\varepsilon + \frac{K}{\varepsilon} e^c \lambda_2(t)$ and $h(t) = \frac{K}{\varepsilon} \lambda_2(t)$

Using the hypothesis (2.3), we can choose a constant $\tilde{T} > 0$ so that

$$\frac{K e^c}{\delta} \cdot \frac{1}{t - t_0} \int_{t_0}^t \lambda_2(\tau) d\tau \leq \frac{\varepsilon}{2} \quad \text{for } t \geq 1 + t_0 \text{ and } t_0 \geq \tilde{T}.$$

Let $\tilde{K} > 0$ be a constant satisfying $\exp \left\{ \frac{\varepsilon}{2} (1 + \tilde{T}) + \frac{K e^c}{\varepsilon} \int_0^{\tilde{T}} \lambda_2(\tau) d\tau \right\} \leq \tilde{K}$.

Then for all $t \geq t_0 \geq 0$ we have

$$(3.11) \quad U(t, x(t)) \leq \frac{K \tilde{K}}{\delta} \{b(\|x_0\|) e^{-\varepsilon(t-t_0)/2} + \int_{t_0}^t e^{-\varepsilon(t-s)/2} \lambda_2(s) ds\}.$$

Using the left-hand side of (3.5), we find that all the solutions of (2.1) are **uniform-bounded**.

Furthermore the condition (2.3) implies that

$$U(t, x(t)) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Therefore by the inequality (3.5) we have

$$x(t) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Q.E.D.

Proof of Corollary. Let $c > 0$ be an arbitrary positive constant. By assumption (ii)', we have

$$\dot{V}_{(2,2)}(t, x) \leq -cV(t, x) + \{c + L(t)\}V(t, x) + \lambda_2(t)(1 + V(t, x)).$$

It now follows from (2.2)' that

$$\limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \{c + L(\tau)\} d\tau < c,$$

which establishes the assumption of Theorem A, and thus the proof is completed.
Q.E.D.

4. Theorems

Let $A(t)$ satisfy the hypothesis (i) of the following Theorem 1 and $P(t)$ be a solution of the matrix equation

$$(4.1) \quad A^T(t)P(t) + P(t)A(t) = -I.$$

Notice that $P(t)$ is bounded for bounded $A(t)$.

The following propositions are due to J. R. Dickerson [2].

Proposition A. $x^T P(t)x \leq C\|x\|^2$, where C is a positive constant.

Proposition B. $|x^T \dot{P}(t)x| \leq 2\|A(t)\| \|P(t)\| x^T P(t)x$, where $\dot{P}(t)$ and $A(t)$ denote the *time derivative* of matrices $P(t)$ and $A(t)$ respectively.

Theorem 1. Suppose that the following conditions are satisfied;

(i) there exists a positive constant τ_0 such that the real parts of all the eigenvalues of $A(t) \leq -\tau_0 < 0$ for all $t \geq 0$,

$$(ii) \quad \limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \|A(s)\| ds < \frac{1}{2P_1}$$

where $P_1 = \limsup_{t \rightarrow \infty} \|P(t)\|$,

$$(iii) \quad \|f(t, x)\| \leq \gamma(t)(1 + \|x\|)$$

where $\gamma(t)$ is a non-negative continuous function on $[0, \infty)$,

$$(iv) \quad \int_t^{t+1} \gamma(s) ds \rightarrow 0 \text{ as } t \rightarrow \infty \quad (i = 1, 2).$$

Then, all solutions $x(t)$ of (1.1) are uniform-bounded and satisfy $x(t) \rightarrow 0$ as $t \rightarrow \infty$.

REMARK. It may be shown by examples [16] that the smallness of $\|\dot{A}(t)\|$ is essential, even if the condition (i) is satisfied.

Next, we consider the equation (1.2) and assume that $g(x, y)$, $g_x(x, y)$, $f(x, y, z)$, $f_x(xy, z)$ and $f_z(xy, z)$ are continuous for all $(x, y, z) \in R^3$ and $h(x)$

is continuously differentiable for all $x \in R^1$.

Theorem 2. Suppose that $a(t), b(t), c(t)$ are continuously differentiable on $[0, \infty)$ and $g(x, 0) = h(0) = 0$ and the following conditions are satisfied;

- (i) $A \geq a(t) \geq a_0 > 0, B \geq b(t) \geq b_0 > 0, C \geq c(t) \geq c_0 > 0$
for $t \in I = [0, \infty)$,
- (ii) $h(x)/x \geq \delta > 0 \quad (x \neq 0),$
- (iii) $\bar{f} \geq f(x, y, z) \geq f_0 > 0$ for all (x, y, z) and $g \geq \frac{g(x, y)}{y} \geq g_0 > 0$
for all $y \neq 0$ and $x,$
- (iv) $y f_x(x, y, 0) \leq 0, y f_z(x, y, z) \geq 0, g_x(x, y) \leq 0$
for all $(x, y, z) \in R^3,$
- (v) $\frac{a_0 b_0 f_0 g_0}{C} > h_1 \geq h'(x),$
- (vi) $\delta > \frac{\mu_2}{4c_0} \left\{ A(\bar{f} - f_0) + \frac{B}{\mu_1} (\bar{g} - g_0) \right\}$

where μ_1 and μ_2 are arbitrarily fixed constants satisfying

$$\frac{Ch_1}{b_0 g_0} < \mu_1 < a_0 f_0, \quad 0 < \mu_2 < \frac{a_0 b_0 f_0 g_0 - Ch_1}{A f_0}$$

$$(vii) \quad \limsup_{(t, v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \{ |a'(s)| + b'_+(s) + |c'(s)| \} ds < \gamma,$$

where γ is a small positive constant whose magnitude depends only on the constants appeared in (i)~(vi), and $b'_+(t) = \max(b'(t), 0)$,

$$(viii) \quad |p(t, x, y, z)| \leq p(t) \{1 + (x^2 + y^2 + z^2)^{1/2}\} + \Delta (x^2 + y^2 + z^2)^{1/2}$$

where Δ is a positive constant and $p(t)$ is a non-negative continuous function,

$$(ix) \quad \int_t^{t+1} p(s) ds \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Then there exists a finite constant $\varepsilon = \varepsilon(A, a_0, B, b_0, C, c_0, \delta, \bar{f} f_0, g, g_0, h_1) > 0$ such that if $\Delta \leq \varepsilon$ then every solution $x(t)$ of (1.2) is uniform-bounded and satisfies

$$x(t) \rightarrow 0, \quad \dot{x}(t) \rightarrow 0, \quad \ddot{x}(t) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

REMARK. It should be pointed out that in the special case $f \equiv 1$ (so that the assumption (iv) is automatically satisfied) Theorem 2 reduces to the author's earlier result [7; Theorem 2]. Also in another special case in which

$a(t)f(x, y, z) = fl$, $b(t)g(x, y) \equiv by$ and $c(t)h(x) = cx$ in (1.2) (so that all the conditions (ii)~(iv) and (vi) are trivially fulfilled) the hypothesis (i) and (v) reduce to

$$a > 0, \quad b > 0, \quad c > 0, \quad ab - c > 0$$

which is the Routh-Hurwitz criterion for the asymptotic stability in the large of the zero solution of the equation

$$\ddot{x} + a\dot{x} + b\dot{x} + cx = 0.$$

5. Proof of theorems

Proof of Theorem 1. We consider the Liapunov function

$$(5.1) \quad V(t, x) = x^T P(t) x.$$

By virtue of Proposition A and the boundedness of $P(t)$, there exist positive constants C and P_2 such that

$$(5.2) \quad C\|x\|^2 \leq V(t, x) \leq P_2\|x\|^2.$$

A simple calculation shows that

$$\begin{aligned} \dot{V}_{(1.1)}(t, x) &= \dot{x}^T P(t) x + x^T \dot{P}(t) \dot{x} + x^T \dot{P}(t) x \\ &= -x^T x + f^T(t, x) P(t) x + x^T P(t) f(t, x) + x^T \dot{P}(t) x. \end{aligned}$$

Applying Proposition B to the function $x^T \dot{P}(t) x$, we obtain

$$\begin{aligned} \dot{V}_{(1.1)}(t, x) &\leq -\|x\|^2 + 2\|f(t, x)\| \cdot \|P(t)\| \cdot \|x\| \\ &\quad + 2\|\dot{A}(t)\| \cdot \|P(t)\| \cdot x^T P(t) x \end{aligned}$$

Using (5.1), (5.2) and (iii) of Theorem 1, we have

$$\begin{aligned} \dot{V}_{(1.1)}(t, x) &\leq \left\{ -\frac{1}{\|P(t)\|} + 2\|P(t)\| \|\dot{A}(t)\| \right\} V(t, x) \\ &\quad + 2\|P(t)\| \gamma(t) \left\{ \left(\frac{V(t, x)}{C} \right)^{1/2} + \frac{V(t, x)}{C} \right\}. \end{aligned}$$

We'll show that

$$\limsup_{(t, v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \left\{ -\frac{1}{\|P(\tau)\|} + 2\|P(\tau)\| \|\dot{A}(\tau)\| \right\} d\tau < 0$$

Let ρ_0 be a positive number such that

$$\rho_0 = \frac{1}{2P_1^2} - \limsup_{(t, v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \|\dot{A}(\tau)\| d\tau.$$

Given $\varepsilon > 0$, there exists a positive number T such that $\|P(\tau)\| < P_1 + \varepsilon$ for all $\tau \geq T$. This implies also

$$\frac{-1}{\|P(\tau)\|} < \frac{-1}{P_1 + \varepsilon} < -\frac{1}{P_1} + \frac{\varepsilon}{P_1^2} \quad \text{for all } \tau \geq T.$$

Using these inequalities, we have

$$\begin{aligned} \limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \left\{ -\frac{1}{\|P(\tau)\|} + 2\|P(\tau)\| \|A(\tau)\| \right\} d\tau \\ < \limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \left\{ -\frac{1}{P_1} + \frac{\varepsilon}{P_1^2} + 2(P_1 + \varepsilon) \|A(\tau)\| \right\} d\tau \\ = -2P_1\rho_0 + \varepsilon \left\{ \frac{1}{P_1^2} + \limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{2}{v} \int_t^{t+v} \|A(\tau)\| d\tau \right\}. \end{aligned}$$

If ε is chosen so that

$$\varepsilon \left\{ \frac{1}{P_1^2} + \limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{2}{v} \int_t^{t+v} \|A(\tau)\| d\tau \right\} < P_1\rho_0,$$

then we have

$$\limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{v} \int_t^{t+v} \left\{ -\frac{1}{\|P(\tau)\|} + 2\|P(\tau)\| \|A(\tau)\| \right\} d\tau < -P_1\rho_0 < 0.$$

Hence, the assumptions of Corollary hold and the proof of Theorem 1 is completed. Q.E.D.

Proof of Theorem 2. The equation (1.2) is equivalent to the system

$$(5.3) \quad \begin{cases} \dot{x} = y \\ \dot{y} = z \\ \dot{z} = -a(t)f(x, y, z)z - b(t)g(x, y) - c(t)h(x) + p(t, x, y, z). \end{cases}$$

We consider the Liapunov function

$$(5.4) \quad V(t, x, y, z) = V_1(t, x, y, z) + V_2(t, x, y, z) + V_3(t, x, y, z)$$

where V_1 , V_2 and V_3 are defined by

$$(5.5) \quad \begin{aligned} 2V_1 = & 2\mu_1 c(t) \int_0^x h(\xi) d\xi + 2c(t)h(x)y + 2b(t) \int_0^y g(x, \eta) d\eta \\ & + 2\mu_1 a(t) \int_0^y f(x, \eta, 0) \eta d\eta + 2\mu_1 yz + z^2, \end{aligned}$$

$$(5.6) \quad \begin{aligned} 2V_2 = & \mu_2 b(t) g_0 x^2 + 2a(t) f_0 c(t) \int_0^x h(\xi) d\xi + a^2(t) f_0^2 y^2 \\ & - \mu_2 y^2 + 2b(t) \int_0^y g(x, \eta) d\eta + z^2 + 2\mu_2 a(t) f_0 xy \\ & + 2\mu_2 xz + 2a(t) f_0 yz + 2c(t)h(x)y, \end{aligned}$$

$$(5.7) \quad 2V_3 = 2a^2(t)f_0^{\int_0^y} f_1 x, \quad \eta, 0) \eta d\eta - a^2(t)f_0^2 y^2,$$

and $\mu_1 > 0$, $\mu_2 > 0$ are two arbitrarily fixed constants such that

$$\frac{Ch_1}{b_0 g_0} < \mu_1 < a_0 f_0, \quad 0 < \mu_2 < \frac{a_0 b_0 f_0 g_0 - Ch_1}{A f_0}, \quad \delta > \frac{\mu_2}{4c_0} \left\{ A(\bar{f} - f_0) + \frac{B}{\mu_1} (\bar{g} - g_0) \right\}.$$

We shall prove the following two properties of V :

$$(5.8) \quad D_1 \cdot (x^2 + y^2 + z^2) \leq V(t, x, y, z) \leq D_2 \cdot (x^2 + y^2 + z^2)$$

for all $(x, y, z) \in R^3$ and

$$(5.9) \quad \dot{V}_{(5.3)} \leq -D_3 \cdot (x^2 + y^2 + z^2) + D_4 \cdot (|a'(t)| + b'_+(t) + |c'(t)|) \cdot (x^2 + y^2 + z^2) \\ + D_5 \cdot (x^2 + y^2 + z^2)^{1/2} \cdot |p(t, x, y, z)|$$

along any solution $(x(t), y(t), z(t))$ of (5.3), where $D_1 \sim D_5$ are certain positive constants.

At first we verify (5.8).

From the inequality $\frac{Ch_1}{b_0 g_0} < \mu_1 < a_0 f_0$ there exists a positive number δ_0 such that $\mu_1 b_0 g_0 (1 - \delta_0) > Ch_1$ and $a_0 f_0 (1 - \delta_0) > \mu_1$. Thus we have

$$\begin{aligned} 2V_1 &= 2\mu_1 c(t) \int_0^x \left\{ 1 - \frac{h'(\xi)}{h_1} \right\} h(\xi) d\xi + \mu_1 c(t) \left\{ \frac{h(x)}{\sqrt{h_1}} + \frac{\sqrt{h_1}}{\mu_1} y \right\}^2 \\ &\quad + (\mu_1 y + z)^2 + \frac{2}{\mu_1} \int_0^y \{ \mu_1 b(t) g(x, \eta) - c(t) h_1 \eta \} d\eta \\ &\quad + 2\mu_1 \int_0^y \{ a(t) f(x, \eta, 0) - \mu_1 \} \eta d\eta \\ &= 2\mu_1 c(t) \int_0^x \left\{ 1 - \frac{h'(\xi)}{h_1} \right\} h(\xi) d\xi + \frac{\delta_0 \mu_1 c(t)}{h_1} h^2(x) \\ &\quad + \mu_1 c(t) \left\{ \sqrt{\frac{1 - \delta_0}{h_1}} h(x) + \frac{1}{\mu_1} \sqrt{\frac{h_1}{1 - \delta_0}} y \right\}^2 + \delta_0 z^2 \\ &\quad + \left(\sqrt{1 - \delta_0} z + \frac{\mu_1}{\sqrt{1 - \delta_0}} y \right)^2 + \frac{2}{\mu_1} \int_0^y \left\{ \mu_1 b(t) g(x, \eta) - \frac{c(t) h_1}{1 - \delta_0} \eta \right\} d\eta \\ &\quad + 2\mu_1 \int_0^y \left\{ a(t) f(x, \eta, 0) - \frac{\mu_1}{1 - \delta_0} \right\} \eta d\eta \\ &\geq \frac{\delta_0 \mu_1 c_0 \delta^2}{h_1} x^2 + \left\{ b_0 g_0 - \frac{Ch_1}{\mu_1 (1 - \delta_0)} - a_0 f_0 - \frac{\mu_1^2}{1 - \delta_0} \right\} y^2 + \delta_0 z^2. \end{aligned}$$

Denoting $2D_6 = \min \left\{ \frac{\delta_0 \mu_1 c_0 \delta^2}{h_1}, b_0 g_0 - \frac{Ch_1}{\mu_1 (1 - \delta_0)} + \mu_1 a_0 f_0 - \frac{\mu_1^2}{1 - \delta_0}, \delta_0 \right\}$, we have

$$V_1(t, x, y, z) \geq D_6(x^2 + y^2 + z^2).$$

It is easy to see that there exists a positive number D_7 such that

$$V_1(t, x, y, z) \leq D_7(x^2 + y^2 + z^2)$$

Hence we obtain

$$D_6(x^2 + y^2 + z^2) \leq V_1(t, x, y, z) \leq D_7(x^2 + y^2 + z^2).$$

As before we have

$$\begin{aligned} 2V_2 &= \mu_2[g_0b(t) - \mu_2]x^2 + \mu_2^2x^2 + a^2(t)f_0^2y + z^2 \\ &\quad + 2\mu_2a(t)f_0xy + 2\mu_2xz + 2a(t)f_0yz \\ &\quad + 2a(t)f_0c(t) \int_0^x h(\xi)d\xi + 2b(t) \int_0^y g(x, \eta)d\eta - \mu_2y^2 + 2c(t)h(x)y \\ &= \mu_2[g_0b(t) - \mu_2]x^2 + [\mu_2x + a(t)f_0y + z]^2 \\ &\quad + a(t)f_0c(t) \left\{ 2 \int_0^x h(\xi)d\xi - \frac{1}{h_1}h^2(x) \right\} + \frac{h_1c(t)}{a(t)} \left\{ \frac{\sqrt{f_0}a(t)h(x)}{h_1} + \frac{y}{\sqrt{f_0}} \right\}^2 \\ &\quad + 2 \int_0^y \left\{ \frac{b(t)g(x, \eta)}{\eta} - \frac{h_1c(t)}{a(t)f_0} - \mu_2 \right\} \eta d\eta. \end{aligned}$$

We find easily that

$$\begin{aligned} [g_0b(t) - \mu_2] &> 0, \\ \left\{ 2 \int_0^x h(\xi)d\xi - \frac{1}{h_1}h^2(x) \right\} &= \frac{2}{h_1^2} \int_0^x h(\xi) \{h_1 - h'(\xi)\} d\xi \geq 0, \\ \left\{ \frac{b(t)g(x, \eta)}{\eta} - \frac{h_1c(t)}{a(t)f_0} - \mu_2 \right\} &> 0. \end{aligned}$$

Then we have a positive number D_8 such that

$$0 \leq V_2(t, x, y, z) \leq D_8(x^2 + y^2 + z^2).$$

We can see also that

$$\begin{aligned} 2V_3 &= 2a^2(t)f_0 \int_0^y \{f(x, \eta) - f_0\} \eta d\eta, \\ 0 &\leq 2V_3(t, x, y, z) \leq A^2f_0(\bar{f} - f_0)y^2. \end{aligned}$$

Therefore there exist positive numbers D_1 and D_2 such that

$$(5.8) \quad D_1(x^2 + y^2 + z^2) \leq V(t, x, y, z) \leq D_2(x^2 + y^2 + z^2).$$

Next we prove the inequality (5.9). Along any solution $(x(t), y(t), z(t))$ of (5.3), we have

$$\begin{aligned}
\dot{V}_{1(5,3)} = & - \left[\frac{\mu_1 b(t) g(x, y)}{y} - c(t) h'(x) \right] y^2 - [a(t) f(x, y, z) - \mu_1] z^2 \\
& + b(t) y \int_0^y g_x(x, \eta) d\eta + \mu_1 a(t) y \int_0^y f_x(x, \eta, 0) \eta d\eta \\
& - \mu_1 a(t) [f(x, y, z) - f(x, y, 0)] y z \\
& + \mu_1 c'(t) \int_0^x h(\xi) d\xi + c'(t) h(x) y + b'(t) \int_0^y g(x, \eta) d\eta \\
& + \mu_1 a'(t) \int_0^y f(x, \eta, 0) \eta d\eta + (\mu_1 y + z) p(t, x, y, z), \\
\dot{V}_{2(5,3)} = & - \mu_2 c(t) x h(x) - \mu_2 b(t) \left[\frac{g(x, y)}{y} - g_0 \right] x y \\
& - \left[a(t) f_0 b(t) \frac{g(x, y)}{y} - c(t) h'(x) - \mu_2 a(t) f_0 \right] y^2 \\
& - a^2(t) f_0 [f(x, y, z) - f_0] y z + b(t) y \int_0^y g_x(x, \eta) d\eta \\
& - a(t) [f(x, y, z) - f_0] z^2 - \mu_2 a(t) [f(x, y, z) - f_0] x z \\
& + \frac{1}{2} \mu_2 b'(t) g_0 x^2 + f_0 [a'(t) c(t) + a(t) c'(t)] \int_0^x h(\xi) d\xi \\
& + a(t) a'(t) f_0^2 y^2 + b'(t) \int_0^y g(x, \eta) d\eta + \mu_2 a'(t) f_0 x y \\
& + a'(t) f_0 y z + c'(t) h(x) y + [z + \mu_2 x + a(t) f_0 y] p(t, x, y, z)
\end{aligned}$$

and

$$\begin{aligned}
\dot{V}_{3(5,3)} = & a^2(t) f_0 y \int_0^y f_x(x, \eta, 0) \eta d\eta + a^2(t) f_0 [f(x, y, 0) - f_0] y z \\
& + 2a(t) a'(t) f_0 \int_0^y [f(x, \eta, 0) - f_0] \eta d\eta.
\end{aligned}$$

Thus we obtain

$$\begin{aligned}
\dot{V}_{(5,3)} = & -W(t, x, y, z) + 2b(t) y \int_0^y g_x(x, \eta) d\eta \\
& + a(t) [\mu_1 + a(t) f_0] y \int_0^y f_x(x, \eta, 0) \eta d\eta \\
& + \{ \mu_1 c'(t) + f_0 [a'(t) c(t) + a(t) c'(t)] \} \int_0^x h(\xi) d\xi \\
& + 2c'(t) h(x) y + 2b'(t) \int_0^y g(x, \eta) d\eta + \mu_1 a'(t) \int_0^y f(x, \eta, 0) \eta d\eta \\
& + \frac{1}{2} \mu_2 b'(t) g_0 x^2 + \mu_2 a'(t) f_0 x y + a'(t) f_0 y z \\
& + 2a(t) a'(t) f_0 \int_0^y f(x, \eta, 0) \eta d\eta
\end{aligned}$$

$$+ \{\mu_2 x + [\mu_1 + a(t)f_0]y + 2z\} \cdot p(t, x, y, z)$$

where

$$\begin{aligned} W = & \mu_2 c(t) x h(x) + [a(t)f(x, y, z) - \mu_1] z^2 + a(t)[f(x, y, z) - f_0] z^2 \\ & + \{[\mu_1 b(t)g_0 - c(t)h'(x)] + [a(t)b(t)f_0 g_0 - c(t)h'(x) - \mu_2 a(t)f_0]\} y^2 \\ & + a(t)b(t)f_0 \left[\frac{g(x, y)}{y} - g_0 \right] y^2 + \mu_1 b(t) \left[\frac{g(x, y)}{y} - g_0 \right] y^2 \\ & + \mu_2 b(t) \left[\frac{g(x, y)}{y} - g_0 \right] xy + \mu_1 a(t)[f(x, y, x) - f(x, y, 0)] yz \\ & + a^2(t)f_0[f(x, y, z) - f(x, y, 0)] yz \\ & + \mu_2 a(t)[f(x, y, z) - f_0] xz \\ \geq & \mu_2 c_0 \delta x^2 + [a_0 f_0 - \mu_1] z^2 \\ & + \{[\mu_1 b_0 g_0 - Ch_1] + [a_0 b_0 f_0 g_0 - Ch_1 - \mu_2 A f_0]\} y^2 \\ & + a(t)[f(x, y, z) - f_0] \{z^2 + \mu_2 xz\} \\ & + \mu_1 b(t) \left[\frac{g(x, y)}{y} - g_0 \right] \left\{ y^2 + \frac{\mu_2}{\mu_1} xy \right\} + a(t)b(t)f_0 \left[\frac{g(x, y)}{y} - g_0 \right] y^2 \\ & + a(t) \{ \mu_1 + a(t)f_0 \} [f(x, y, z) - f(x, y, 0)] yz. \end{aligned}$$

Hence

$$\begin{aligned} W \geq & \mu_2 \left\{ c_0 \delta - [f(x, y, z) - f_0] \frac{A\mu_2}{4} - \left[\frac{g(x, y)}{y} - g_0 \right] \frac{B\mu_2}{4\mu_1} \right\} x^2 \\ & + \{[\mu_1 b_0 g_0 - Ch_1] + [a_0 b_0 f_0 g_0 - Ch_1 - \mu_2 A f_0]\} y^2 \\ & + [a_0 f_0 - \mu_1] z^2 + a(t)[f(x, y, z) - f_0] \left(z + \frac{1}{2} \mu_2 x \right)^2 \\ & + \mu_1 b(t) \left[\frac{g(x, y)}{y} - g_0 \right] \left(y + \frac{1}{\mu_1} \mu_2 x \right)^2 + a(t)b(t)f_0 \left[\frac{g(x, y)}{y} - g_0 \right] y^2 \\ & + a(t) \{ \mu_1 + a(t)f_0 \} [f(x, y, z) - f(x, y, 0)] yz. \end{aligned}$$

By the assumptions

$$\begin{aligned} & \left\{ c_0 \delta - [f(x, y, z) - f_0] \frac{A\mu_2}{4} - \left[\frac{g(x, y)}{y} - g_0 \right] \frac{B\mu_2}{4\mu_1} \right\} \\ & \geq c_0 \delta - [\bar{f} - f_0] \frac{A\mu_2}{4} - [\bar{g} - g_0] \frac{B\mu_2}{4\mu_1} \\ & = c_0 \delta - \frac{\mu_2}{4} \left\{ A(\bar{f} - f_0) + \frac{B}{\mu_1} (\bar{g} - g_0) \right\} > 0, \\ & [\mu_1 b_0 g_0 - Ch_1] > \frac{Ch_1}{b_0 g_0} \cdot b_0 g_0 - Ch_1 = 0, \\ & [a_0 b_0 f_0 g_0 - Ch_1 - \mu_2 A f_0] \end{aligned}$$

$$\begin{aligned}
&> a_0 b_0 f_0 g_0 - C h_1 - \frac{a_0 b_0 f_0 g_0 - C h_1}{A f_0} \cdot A f_0 = 0, \\
&[a_0 f_0 - \mu_1] > 0.
\end{aligned}$$

Applying the Mean Value Theorem, we have for $0 < |\mathbf{z}| < |\mathbf{z}|$

$$[f(x, y, \mathbf{z}) - f(x, y, 0)] y \mathbf{z} = f_z(x, y, \mathbf{z}) y \mathbf{z}^2 \geq 0.$$

Therefore there exists a positive number D_3 such that for all $(x, y, \mathbf{z}) \in R^1$

$$W \geq D_3 \cdot (x^2 + y^2 + \mathbf{z}^2).$$

From (ii), (iii), (iv) and (v) it is easy to show that there exists a positive number D_4 such that

$$\begin{aligned}
\dot{V}_{(5.3)} &\leq -D_3(x^2 + y^2 + \mathbf{z}^2) + D_4 \cdot (|a'(t)| + b'_+(t) + |c'(t)|)(x^2 + y^2 + \mathbf{z}^2) \\
&\quad + \{\mu_2 x + [\mu_1 + a(t)f_0]y + 2\mathbf{z}\} p(t, x, y, \mathbf{z}).
\end{aligned}$$

Setting $D_9 = \max \{\mu_2, \mu_1 + A f_0, 2\}$, we have

$$\begin{aligned}
&\{\mu_2 x + [\mu_1 + a(t)f_0]y + 2\mathbf{z}\} p(t, x, y, \mathbf{z}) \\
&\leq D_9 \cdot (|x| + |y| + |\mathbf{z}|) |p(t, x, y, \mathbf{z})| \\
&\leq \sqrt{3} D_9 \cdot (x^2 + y^2 + \mathbf{z}^2)^{1/2} |p(t, x, y, \mathbf{z})|.
\end{aligned}$$

Let $D_5 = \sqrt{3} D_9$. Then we obtain the inequality (5.9).

We are now ready for the principal subsidiary results needed for the completion of the proof of Theorem 2. Application to (5.9) with the assumption (viii) leads to

$$\begin{aligned}
\dot{V}_{(5.3)} &\leq -D_3 \cdot (x^2 + y^2 + \mathbf{z}^2) + D_4 \cdot (|a'(t)| + b'_+(t) + |c'(t)|)(x^2 + y^2 + \mathbf{z}^2) \\
&\quad + D_5 p(t) [(x^2 + y^2 + \mathbf{z}^2)^{1/2} + (x^2 + y^2 + \mathbf{z}^2)] \\
&\quad + \Delta D_5 \cdot (x^2 + y^2 + \mathbf{z}^2).
\end{aligned}$$

Let Δ be fixed, in what follows, to satisfy

$$(5.10) \quad \Delta \leq \frac{D_3}{2D_5}.$$

Using the inequalities (5.8) and (5.10), we have

$$\begin{aligned}
(5.11) \quad \dot{V}_{(5.3)} &\leq -\frac{D_3}{2D_2} \cdot V + \frac{D_4}{D_1} (|a'(t)| + b'_+(t) + |c'(t)|) V \\
&\quad + D_5 p(t) \left[\left(\frac{V}{D_1} \right)^{1/2} + \frac{V}{D_1} \right].
\end{aligned}$$

Assume

$$\limsup_{(t,v) \rightarrow (\infty, \infty)} \frac{1}{V} \int_t^{t+v} \{|a'(s)| + b'_+(s) + |c'(s)|\} ds < \frac{D_1 D_3}{2D_2 D_4}.$$

Now Theorem A will be used to prove the uniform boundedness of solutions of (1.2) and that for any solution $x(t)$

$$x^2(t) + \dot{x}^2(t) + \ddot{x}^2(t) \rightarrow 0 \quad \text{as} \quad t \rightarrow \infty.$$

This completes the proof of Theorem 2.

Q.E.D.

Acknowledgement: The author is indebted to the editor and referees whose suggested revisions have improved the exposition of this paper.

OSAKA UNIVERSITY

Bibliography

- [1] F. Brauer: *Global behavior of solutions of ordinary differential equations*, J. Math. Anal. Appl. 2 (1961), 145-158.
- [2] J.R. Dickerson: *Stability of linear systems with parametric excitation*, Trans. ASME Ser.E. 37 (1970), 228-230.
- [3] J.O.C. Ezeilo: *Further results for the solution of a third order differential equations*, Proc. Cambridge Philos. Soc. 59 (1963), 111-116.
- [4] ———: *Stability results for the solutions of some third and fourth order differential equations*, Ann. Mat. Pura. Appl. 66 (1964), 233-249.
- [5] T. Hara: *On the stability of solutions of certain third order differential equations*, Proc. Japan Acad. 47 (1971), 894-902.
- [6] ———: *On the asymptotic behavior of solutions of certain third order ordinary differential equations*, Proc. Japan Acad. 47 (1971), 903-908.
- [7] ———: *Remarks on the asymptotic behavior of the solutions of certain non-autonomous differential equations*, Proc. Japan Acad. 48 (1972), 549-552.
- [8] M. Harrow: *Further results for the solution of certain third and fourth order differential equations*, Proc. Cambridge Philos. Soc. 63 (1967), 147-153.
- [9] ———: *Further results for the solution of certain third order differential equations*, J. London Math. Soc. 43 (1968), 587-592.
- [10] N. Onuchic: *Asymptotic behavior of a perturbed linear differential equation with time lag on a product space*, Ann. Mat. Pura. Appl. 86 (1970), 115-123.
- [11] R. Reissig, G. Sansone und R. Conti: *Nichtlineare Differentialgleichungen Höherer Ordnung*, Roma, 1969.
- [12] A. Strauss and J. Yorke: *Perturbing uniform asymptotically stable nonlinear systems*, J. Differential Equations 6 (1969), 452-483.
- [13] K.E. Swick: *On the boundedness and the stability of solutions of some non-autonomous differential equations of the third order*, J. London Math. Soc. 44 (1969), 347-359.
- [14] ———: *Asymptotic behavior of the solutions of certain third order differential equations*, SIAM J. Appl. Math. 19 (1970), 96-102.
- [15] H.O. Tejumola: *A note on the boundedness and the stability of solutions of certain third-order differential equations*, Ann. Mat. Pura. Appl. 92 (1972), 65-75.

- [16] R.E. Vinograd: *On a criterion of instability in the sense of **Liapunov** of the solutions of a linear system of ordinary differential equations (Russian)*, Dokl. Akad. Nauk SSSR 84 (1952), 201-204.
- [17] M. Yamamoto: *On the stability of the solutions of some non-autonomous differential equations of the third order*, Proc. Japan Acad. **47** (1971), 909-914.
- [18] ———: *Remarks on the asymptotic behavior of the solutions of certain third order non-autonomous differential equations*, Proc. Japan Acad. 47 (1971), 915-920.
- [19] T. Yoshizawa: *Stability Theory by Liapunov's Second Method*, Tokyo, 1966.