



Title	A note on relative T-nilpotency
Author(s)	Harada, Manabu
Citation	Osaka Journal of Mathematics. 1976, 13(2), p. 431-438
Version Type	VoR
URL	<a href="https://doi.org/10.18910/12896">https://doi.org/10.18910/12896</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## A NOTE ON RELATIVE T-NILPOTENCY

MANABU HARADA

(Received June 23, 1975)

This note gives some supplementary results of [6]. The first one shows an application of the idea in the proof of [6], Lemma 7 and gives a characterization of artinian rings. The second one gives a refinement of [6], Corollary 2 to Theorem A.2 and the final one is a special type of the exchange property.

Throughout we shall assume that  $R$  is a ring with identity and modules are unitary right  $R$ -modules. First, we shall recall definitions in [6].

Let  $\{P_\alpha\}_I$  and  $\{Q_\beta\}_J$  be two infinite sets of  $R$ -modules. We take a countable set  $\{M_i\}_1^\infty$  such that  $M_{2i-1}=P_{\alpha(2i-1)}\in\{P_\alpha\}_I$  and  $M_{2j}=Q_{\beta(2j)}\in\{Q_\beta\}_J$ . Further we take a set of non-isomorphisms  $f_i: M_i\rightarrow M_{i+1}$ . If for any element  $m$  in  $M_1$  there exists  $n$  such that  $f_n f_{n-1} \cdots f_1(m)=0$ , we say  $\{f_i\}_1^\infty$  is *locally T-nilpotent*. If for any countable sets  $\{M_i\}_1^\infty$  above such that  $\alpha(2i-1)\neq\alpha(2i'-1)$  ( $\beta(2j)\neq\beta(2j')$ ) if  $i\neq i'$  ( $j\neq j'$ ) any sets  $\{f_i\}$  of non-isomorphisms are always locally T-nilpotent, then we say  $\{P_\alpha\}_I$  and  $\{Q_\beta\}_J$  are *relatively and locally semi-T-nilpotent*. If we omit the assumptions  $\alpha(2i-1)\neq\alpha(2i')$  ( $\beta(2j)\neq\beta(2j')$ ) in the above, we say  $\{P_\alpha\}_I$  and  $\{Q_\beta\}_J$  are *relatively and locally T-nilpotent*. If  $\{P_\alpha\}_I=\{Q_\beta\}_J$ , we say  $\{P_\alpha\}_I$  is *locally semi-T-nilpotent* or *T-nilpotent*, corresponding to the above cases. We shall assume that the definition of relatively semi-T-nilpotency contains a case of either  $I$  or  $J$  being finite. If  $K=\sum_I \bigoplus P_\alpha=\sum_J \bigoplus Q_\beta$  and  $\{P_\alpha\}_I$ ,  $\{Q_\beta\}_J$  are locally and relatively T-nilpotent, then we say  $\sum_I \bigoplus P_\alpha$  and  $\sum_J \bigoplus Q_\beta$  are *relatively T-nilpotent decompositions* of  $K$ .

Finally, let  $M=N\oplus P$  be  $R$ -modules and  $\kappa$  a cardinal number. If for any decomposition  $M=\sum \bigoplus L_\alpha$  with  $\kappa$ -components there exist submodules  $L'_\alpha$  of  $L_\alpha$  such that  $M=N\oplus\sum \bigoplus L'_\alpha$ , then we say  $N$  has the  $\kappa$ -exchange property in  $M$ . In case  $\kappa$  is any cardinal, we say  $N$  has the exchange property in  $M$ .

### 1. T-nilpotent decompositions

First, we study a property of relative T-nilpotency. If the endomorphism ring of a module  $M$  is a local ring, then we call  $M$  *completely indecomposable*.

**Lemma 1.** *Let  $M$  be an  $R$ -module and  $f, g$  in  $\text{End}_R(M)$ . If  $fg$  is isomorphic,  $M=\text{Im } g\oplus \text{Ker } f$ .*

**Lemma 2.** *Let  $P$  be an  $R$ -module. If  $P$  is itself locally  $T$ -nilpotent,  $P$  is a completely indecomposable module.*

Proof. Put  $S = \text{End}_R(P)$ . If  $e \in S$  and  $e^2 = e$ , then  $e = 1$  or  $0$  by the assumption. Let  $x, y$  be elements in  $S$  with  $x$  non-unit. Then neither  $xy$  nor  $yx$  is unit in  $S$  from the above and Lemma 1. Furthermore, consider a sequence  $\{x^n\}_1^\infty$  of non-units in  $S$ . For any element  $p$  in  $P$ , there exists  $n = n(p)$  such that  $x^{n(p)}(p) = 0$  from the assumption. Therefore,  $X = 1 + \sum_1^\infty x^i$  is an element in  $S$  and  $((1-x)X)(p) = (1-x)(1+x+\dots+x^{n(p)-1})(p) = (1-x^{n(p)})(p) = p$ . Hence,  $1-x$  is unit in  $S$  from Lemma 1. Let  $x, y$  be non-unit in  $S$ . We assume that  $x+y$  is unit in  $S$ . Then we may assume  $x+y=1$ , which is a contradiction to the above. Therefore,  $S$  is a local ring.

**Theorem 1.** *Let  $M$  be an  $R$ -module and  $M = \sum_I \bigoplus P_\alpha = \sum_J \bigoplus Q_\beta$  two relatively  $T$ -nilpotent decompositions of  $M$ . Then all  $P_\alpha$  and  $Q_\beta$  are completely indecomposable modules and hence, those decompositions are unique up to isomorphism and every direct summand of  $M$  has the exchange property in  $M$ .*

Proof. We put  $I_1 = \{\alpha \in I \mid P_\alpha \approx Q_\beta \text{ for some } \beta \in J\}$  and  $J_1 = \{\beta \in J \mid Q_\beta \approx P_\alpha \text{ for some } \alpha \in I\}$ . We first show  $I_1 \neq \emptyset$  and so  $J_1 \neq \emptyset$ . We assume the contrary. Let  $p_\alpha, q_\beta$  be projections of  $M$  to  $P_\alpha$  and  $Q_\beta$ , respectively. Let  $x_1 \neq 0 \in P_{\alpha_1}$ . Then there exists  $\beta_2 \in J$  such that  $q_{\beta_2}(x_1) = x_2 \neq 0$ . Again there exists  $\alpha_3 \in I$  such that  $p_{\alpha_3}(x_2) = x_3 \neq 0$ . Repeating those arguments, we obtain a contradiction to the  $T$ -nilpotency, since  $I_1 = J_1 = \emptyset$  (cf. [6], Lemma 7). Hence,  $I_1 \neq \emptyset$  and so  $J_1 \neq \emptyset$ . Furthermore,  $\{P_\alpha\}_{I_1}$ ,  $\{Q_\beta\}_{J_1}$  are sets of completely indecomposable modules and locally  $T$ -nilpotent by Lemma 2 and the assumption. We put  $M = \sum_{I_1} \bigoplus P_\alpha \oplus \sum_{I_1'} \bigoplus P_{\alpha'} = \sum_{J_1} \bigoplus Q_\beta \oplus \sum_{J_1'} \bigoplus Q_{\beta'}$ , where  $I_1' = I - I_1$  and  $J_1' = J - J_1$ . Let  $\{P_{\alpha_i}\}_1^n$  be any finite subset of  $\{P_\alpha\}_{I_1}$ . Since  $\sum_1^n \bigoplus P_{\alpha_i}$  has the exchange property by [2], Lemma 3.11 and [9], Proposition 1,

$$M = \sum_1^n \bigoplus P_{\alpha_i} \oplus \sum_{J_1'} \bigoplus Q_{\beta'} \oplus \sum_{J_1'} \bigoplus Q_{\beta'}' \dots (*),$$

where  $Q_\beta = Q_{\beta'} \oplus Q_{\beta''}$ . Then  $\sum_{J_1'} \bigoplus Q_{\beta''}$  is isomorphic to a direct summand of  $\sum_1^n \bigoplus P_{\alpha_i}$ . If  $Q_{\beta''} \neq (0)$ ,  $Q_{\beta''}$  contains a direct summand isomorphic to some  $P_\alpha$ , by Krull-Remak-Schmidt's theorem, say  $Q_{\beta''} = X \oplus Y$ ;  $X \not\approx P_{\alpha_i}$ . Since  $\beta' \in J_1'$ ,  $Y \neq (0)$ . Let  $p$  be a projection of  $Q_{\beta''}$  to  $X$  and  $i$  the inclusion of  $X$  to  $Q_{\beta''}$ , then  $p$  and  $i$  are not isomorphic. However,  $\varphi p i \varphi^{-1} = 1_{P_{\alpha_i}}$  and neither  $\varphi p$  nor  $i \varphi^{-1}$  is isomorphic. Which contradicts the relative  $T$ -nilpotency. Accordingly,  $Q_{\beta''} = (0)$  and  $(\sum_1^n \bigoplus P_{\alpha_i}) \cap (\sum_{J_1'} \bigoplus Q_{\beta'}) = (0)$ . Therefore,  $(\sum_{I_1} \bigoplus P_\alpha) \cap$

$(\sum_{J_1'} \oplus Q_\beta) = (0)$ . Let  $\bar{M} = M / (\sum_{J_1'} \oplus Q_\beta)$  and  $\psi$  the natural epimorphism of  $M$  to  $\bar{M}$ . Then  $\bar{M} = \sum_{J_1} \oplus \psi(Q_\beta) \supseteq \sum_{J_1} \oplus \psi(P_\alpha)$  (note  $\psi(Q_\beta) \approx Q_\beta$  and  $\psi(P_\alpha) \approx P_\alpha$ ). On the other hand,  $\sum_{J_1} \oplus \psi(P_\alpha)$  is locally direct summand of  $\bar{M}$  from (\*) and  $\{\psi(P_\alpha)\}_{J_1}$  is locally T-nilpotent. Hence,  $\sum_{J_1} \oplus \psi(P_\alpha)$  is a direct summand of  $\bar{M}$  by [3], Theorem 9 and [7], Corollary 2 to Lemma 2 and Lemma 3, since  $\{\psi(Q_\beta)\}_{J_1}$  is a set of completely indecomposable modules. Furthermore,  $\sum_{J_1} \oplus \psi(P_\alpha)$  has the exchange property in  $\bar{M}$  by [4], Theorem 4 and so  $\bar{M} = \sum_{J_1} \oplus \psi(P_\alpha) \oplus \sum_{J_1''} \oplus \psi(Q_\beta)$ , where  $J_1'' \subseteq J_1$ . Therefore,  $M = \sum_{J_1} \oplus P_\alpha \oplus \sum_{J_1''} \oplus Q_\beta \oplus \sum_{J_1'} \oplus Q_{\beta'} = \sum_{J_1} \oplus P_\alpha \oplus \sum_{J_1'} \oplus P_{\alpha'}$ . Hence,  $\sum_{J_1''} \oplus Q_\beta \oplus \sum_{J_1'} \oplus Q_{\beta'} \approx \sum_{J_1'} \oplus P_{\alpha'}$ . However,  $\{P_{\alpha'}\}_{J_1'}$  and  $\{Q_{\beta'}\}_{J_1' \cup J_1''}$  are locally and relatively T-nilpotent and so some  $P_{\alpha'}$  is isomorphic to some  $Q_{\beta'}$  by the first part, provided  $P_{\alpha'} \neq (0)$ . Therefore,  $J_1' = J_1'' = J_1 = \phi$ . The remaining parts are clear from [1] and [4], Theorem 4.

REMARK. Theorem 1 does not remain valid if we replace the T-nilpotency by the semi-T-nilpotency in the assumption.

**Corollary 1.** *Let  $P$  be  $R$ -projective. Then  $P$  has two relatively T-nilpotent decompositions if and only if  $P$  is a perfect module.*

Proof. It is clear from Theorem 1 and [4], Theorem 6.

**Corollary 2.**  *$R$  is right artinian if and only if every projective modules and every injective modules have relatively T-nilpotent decompositions.*

Proof. It is clear from Corollary 1, [3], Corollary 1 to Proposition 1 and [8].

## 2. Exchange property

Let  $\{M_\alpha\}_I$  be a set completely indecomposable modules and  $M = \sum_I \oplus M_\alpha$ .

We consider a relation between the concept of the exchange property in  $M$  and that of the 2-exchange property in  $M$  for a direct summand  $N$  of  $M$ . If  $N$  is also a direct sum of indecomposable modules, those concepts are equivalent by [6], Theorem A.2. We do not know whether this fact is true without any assumptions.

**Lemma 3.** *Let  $A = B \oplus C \oplus D$  be  $R$ -modules. If  $B \oplus C$  has the 2-exchange property in  $A$ , then  $C$  has the same property in  $C \oplus D$ .*

Proof. We assume  $C \oplus D = K \oplus L$ . Then  $A = B \oplus C \oplus D = B \oplus K \oplus L$ . Hence,  $A = B \oplus C \oplus K' \oplus L'$  for some  $K' \subseteq K$  and  $L' \subseteq L$  from [6], Lemma A.4. Since  $C \oplus D \supseteq C \oplus K' \oplus L'$ ,  $C \oplus D = C \oplus K' \oplus L'$ .

Let  $M$  be as above and  $M=S_1 \oplus S_2$ . Let  $\{M_\alpha\}_J$  be the isomorphic representative classes of  $\{M_\alpha\}_I$  and we shall denote it by  $[M]$ . We put  $J'=\{\alpha \in J \mid M_\alpha \text{ is isomorphic to a direct summand of both } S_1 \text{ and } S_2\}$  and  $J''=\{\alpha \in J' \mid \text{directsums of any finite copies of } M_\alpha \text{ are isomorphic to direct summands of both } S_1 \text{ and } S_2\}$ .

**Theorem 2.** *Let  $M, S_i, \text{etc.}$  be as above. Then  $S_1$  has the exchange property in  $M$  if and only if  $S_1$  has the 2-exchange property in  $M$ ,  $\{M_\alpha\}_{J'}$  is locally semi-T-nilpotent and  $\{M_\alpha\}_{J''}$  is locally T-nilpotent.*

Proof. Let  $N_i$  be a dense submodule of  $S_i$  and  $N_i=\sum_{j \in I} \sum_{\beta \in I_{ij}} \oplus N_{ij\beta}$ , where  $i=1,2$  and  $N_{ij\beta}$ 's are isomorphic to some  $M_\alpha$  in  $\{M_\alpha\}_I$  and  $N_{ij\beta} \approx N_{i'j\beta} \approx N_{ij\beta'} \approx N_{i'j\beta'}$  and  $N_{ij\beta} \not\approx N_{ij'\beta}$  if  $J \neq I'$  (see [4]). We put

$$\begin{aligned} J_1 &= \{j \in J \mid I_{1j} \text{ and } I_{2j} \text{ are infinite}\} \\ J_2 &= \{j \in J \mid I_{1j} \neq \phi \text{ and } I_{2j} \text{ is finite}\} \\ J_3 &= \{j \in J \mid I_{1j} \text{ is finite and } I_{2j} \text{ is infinite}\} \\ J_4 &= \{j \in J \mid I_{1j} = \phi\} \text{ and} \\ J_5 &= \{j \in J \mid I_{2j} = \phi\}. \end{aligned}$$

Then  $J_1=J''$  and  $J_1 \cup J_2 \cup J_3=J'$ . Furthermore, we put  $N_i(k)=\sum_{j \in J} \sum_{\beta \in I_{ij}} \oplus N_{ij\beta}$ . Then  $N_1=\sum_{k \neq 4} \oplus N_1(k)$  and  $N_2=\sum_1^4 \oplus N_2(k)$ . We shall show "if" part.  $N_1(1)$ ,  $N_1(3)$  and  $N_2(2)$  are direct summands of  $M$  from the assumption and [4], Proposition 2. Since  $N_1(1) \oplus N_1(3) \subseteq S_1$  and  $N_2(2) \subseteq S_2$ ,  $M=N_1(1) \oplus N_1(3) \oplus S_1' \oplus N_2(2) \oplus S_2'$ , where  $S_1' \subseteq S_1$ . Furthermore, a dense submodule of  $S_1'$  (resp.  $S_2'$ ) is isomorphic to  $N_1(2) \oplus N_1(5)$  (resp.  $N_2(1) \oplus N_2(3) \oplus N_2(4)$ ). Since  $S_1=N_1(1) \oplus N_1(3) \oplus S_1'$ ,  $S_2=N_2(2) \oplus S_2'$  and  $S_1$  has the 2-exchange property in  $M$ ,  $S_1'$  has the same property in  $S_1' \oplus N_2(2) \oplus S_2' (=C)$  from Lemma 3. We assume  $S_1' \oplus S_2'=A \oplus B$ . Then  $C=A \oplus B \oplus N_2(2)$ . Hence,  $C=S_1' \oplus A' \oplus B' \oplus N_2(2)$  from [6], Lemma A.4, where  $A' \subseteq A$ , etc. Therefore,  $S_1' \oplus A' \oplus B'=S_1' \oplus S_2'$ , which means  $S_1'$  has the 2-exchange property in  $S_1' \oplus S_2'$ . Accordingly,  $S_1'$  has the exchange property in  $S_1' \oplus S_2'$  by [6], Corollary 2 to Theorem A.2, since  $N_1(1) \oplus N_1(3) \oplus N_2(2)$  has the exchange property in  $M$  by [5], Theorem 2 and so  $S_1' \oplus S_2'$  is a directsum of completely indecomposable modules. Therefore,  $S_1'$  is also a directsum of completely indecomposable modules and hence,  $S_1$  has the exchange property in  $M$  by [6], Theorem A.2. The converse is clear from [6], Theorem A.2.

Finally, we shall study some special properties concerning with the exchange property in  $M$  of a direct summand of  $M$ . We are interested in a relation between the exchange property in  $M$  and the relative semi-T-nilpotency. We assume  $M=N_1 \oplus N_2$  and  $N_i=\sum_k \oplus N_i(k)$  as in the proof of Theorem 2.

We know from the proof above that  $N_1$  has the exchange property in  $M$  if and only if  $N_1(2) \oplus N_1(5)$  has the same property in  $N_1(2) \oplus N_1(5) \oplus N_2(3) \oplus N_2(4)$  and  $\{N_{1ij}\}, \{N_{2ij}\}$  are locally and relatively semi-T-nilpotent (cf. [6], the proof of Corollary 1 to Theorem). Hence, we may restrict ourselves to a case of  $[N_1] \cap [N_2] = \phi$ .

In the following we shall use the category  $\underline{A}$  induced from a set of completely indecomposable modules and its factor category  $\bar{\underline{A}} = \underline{A}/J'$  studied in [3]. We refer to [3] for the notations and results on  $\underline{A}$ .

**Lemma 4.** *Let  $M$  be in  $\underline{A}$  and  $A, B$  two locally direct summands of  $M$ . If  $[A] \cap [B] = \phi$ ,  $A \cap B = (0)$  and  $A \oplus B$  is a locally direct summand of  $M$ .*

Proof. Let  $i_A, i_B$  be the inclusions of  $A$  and  $B$  into  $M$ , respectively. Since  $[A] \cap [B] = \phi$ ,  $\text{Im } i_A \cap \text{Im } i_B = (0)$  from [3], Theorem 7 and [7], Lemma 3. Let  $A \oplus B$  be the external direct sum and  $i = (i_A, i_B): A \oplus B \rightarrow M$ . Then it is clear from the above that  $i$  is monomorphic in  $\bar{\underline{A}}$ . Hence,  $\text{Im } i = A + B$  is a locally direct summand of  $M$  and  $A \cap B = (0)$ .

**Lemma 5.** *Let  $M = S \oplus T$  and  $M$  in  $\underline{A}$ . For any element  $x$  in  $S$  there exists a finite set of indecomposable modules  $S_i$  such that  $S = \sum_1^t \oplus S_i \oplus S'$ ,  $x \in \sum_1^t \oplus S_i$  and  $S_i$ 's are isomorphic to some in  $[M]$ .*

Proof. See [4], the proof of Proposition 3.

Let  $M = N_1 \oplus N_2 = \sum \oplus S_\gamma$  and  $N_i \in \underline{A}$  with  $[N_1] \cap [N_2] = \phi$ . We put  $S_\gamma(i) = \sum S_\alpha$ , where  $S_\alpha$  runs through all indecomposable direct summands of  $S_\gamma$ , which are isomorphic to some in  $[N_i]$ . By  $[S_\gamma(i)]$  we denote the representative classes of such  $S_\alpha$ 's. Then  $S_\gamma(i)$  is also the union of all locally direct summands  $A$  of  $S_\gamma$  with  $[A] \subseteq [N_i]$ . It is clear, from Lemma 5,  $S_\gamma = S_\gamma(1) + S_\gamma(2)$ . If  $N_1$  (or  $N_2$ ) has the exchange property in  $M$ ,  $S_\gamma = S_{\gamma_1} \oplus S_{\gamma_2}$  where  $S_{\gamma_i} \subseteq S_\gamma(i)$  and every indecomposable direct summand of  $S_{\gamma_i}$  is isomorphic to some in  $[N_i]$ . In the following, we shall study a case of  $S_{\gamma_1} = S_\gamma(1)$ .

The following lemma is a slight generalization of [6], Lemma 7.

**Lemma 6.** *Let  $M = N_1 \oplus N_2 = \sum_K \oplus S_\gamma$  and  $N_i = \sum_{\alpha \in I_i} \oplus N_{i\alpha}$ ;  $N_{i\alpha}$ 's are completely indecomposable modules. We assume  $\{N_{1\alpha}\}_{I_1}$  and  $\{N_{2\alpha}\}_{I_2}$  are locally and relatively semi-T-nilpotent and  $[N_1] \cap [N_2] = \phi$ . Then  $M = \sum_K \oplus S_\gamma(1) + N_2 = N_1 + \sum_K \oplus S_\gamma(2)$ .*

Proof. We shall give a sketch of the proof (cf. [6], Lemma 7). We assume  $I_1$  and  $I_2$  are infinite. Put  $M^* = \sum_K \oplus S_\gamma(1) + N_2$ . We assume  $M \neq M^*$ . Then there exists  $N_{1\alpha}$  not contained in  $M^*$ . Let  $x_1 \in N_{1\alpha} - M^*$  and  $x_{1\alpha} = \sum x_{\gamma_i}$ ;  $x_{\gamma_i} \in S_{\gamma_i}$ .

We may assume  $x_{\gamma_1} \notin M^*$ . Then from Lemma 5 we have  $x_{\gamma_1} = \sum_{\beta} y_{i\beta}$ ;  $y_{i\beta} \in A_{\beta} \setminus \bigoplus S_{\gamma_1}$  and  $A_{\beta}$ 's are indecomposable. Since  $x_{\gamma_1} \notin M^*$ , there exists  $y_{2\beta} \in A_{\beta} - M^*$  and so  $[A_{\beta}] \in [N_2]$ . Now, we can find  $S_{\gamma_1'}$  which contains a direct summand  $S_{\gamma_1'\alpha'}$  isomorphic to  $N_{1\alpha}$ . Since  $S_{\gamma_1'\alpha'}$  has the exchange property by [9], Proposition 1,

$$M = S_{\gamma_1'\alpha'} \oplus N_1' \oplus N_2; N_1' \subseteq N_1,$$

since  $[N_1] \cap [N_2] = \emptyset$ . Let  $y_{2\beta} = a + b + c$ ;  $a \in S_{\gamma_1'\alpha'}$ ,  $b \in N_1'$  and  $c \in N_2$ . Then  $b \in N_1' - M^*$ . Hence, we can find an indecomposable direct summand  $N_{1\delta}$  of  $N_1'$  such that  $b = x_{1\delta} + \dots$ ,  $x_{1\delta} \in N_{1\delta} - M^*$ . On the other hand, since  $[A_{\beta}] \in [N_2]$ , there exists  $N_{2\epsilon}$  isomorphic to  $A_{\beta}$  and

$$M = N_{2\epsilon} \oplus \sum_{\gamma} \bigoplus S_{\gamma}'; S_{\gamma}' \subseteq S_{\gamma}.$$

Let  $x_{1\delta} = d + \sum f_{\gamma}$ ;  $d \in N_{2\epsilon}$ ,  $f_{\gamma} \in S_{\gamma}'$ . Again there exists  $f_{\gamma_2} \in S_{\gamma_2}' - M^*$ . Similarly to the above, we can find a direct summand  $A_{\gamma}$  of  $S_{\gamma_2}'$  such that  $[A_{\gamma}] \in [N_2]$ ,  $f_{\gamma_2} = \sum y_{i\gamma}$  and  $y_{2\gamma} \in A_{\gamma} - M^*$ . Furthermore, since  $N_{1\delta}$  is a direct summand of  $N_1'$ , there exists  $S_{\gamma_2'}$  which contains a direct summand  $S_{\gamma_2'\delta'}$  isomorphic to  $N_{1\delta}$  such that

$$M = S_{\gamma_1''\alpha''} \oplus S_{\gamma_2'\delta'} \oplus N_1'' \oplus N_2; N_1'' \subseteq N_1'$$

and  $S_{\gamma_1''\alpha''} \approx S_{\gamma_1'\alpha'}$ . Repeating those arguments we obtain a sequence  $\{x_{1\alpha}, y_{2\beta}, x_{1\delta}, y_{2\gamma}, \dots\}$ , which contradicts the assumption of relative semi-T-nilpotency.

**Theorem 3.** *Let  $M$  be a direct sum of completely indecomposable modules and  $M = N_1 \oplus N_2 = \sum_{\gamma} \bigoplus S_{\gamma}$ . We assume  $N_i$  is a direct sum of indecomposable modules  $\{N_{i\alpha}\}_{I_i}$  such that  $N_{1\alpha} \not\approx N_{2\beta}$  for any  $\alpha \in I_1$ ,  $\beta \in I_2$  and  $\{N_{1\alpha}\}_{I_1}$ ,  $\{N_{2\beta}\}_{I_2}$  are locally and relatively semi-T-nilpotent. Then the following conditions are equivalent.*

- 1)  $M = \sum_{\gamma} \bigoplus S_{\gamma}(1) \oplus N_2$
- 2)  $\text{Hom}_R([S_{\gamma}(1)], [S_{\gamma}(2)]) = (0)$  for every  $\gamma$
- 3)  $S_{\gamma}(1)$  is a direct summand of  $S_{\gamma}$  and  $[A] \in [N_1]$  for every indecomposable direct summands  $A$  of  $S_{\gamma}(1)$ , where  $\text{Hom}_R([S_{\gamma}(1)], [S_{\gamma}(2)]) = (0)$  means  $\text{Hom}_R(B, C) = (0)$  for all  $B \in [S_{\gamma}(1)]$  and all  $C \in [S_{\gamma}(2)]$ .

**Proof.** 1)  $\rightarrow$  2) and 3). 1) implies  $S_{\gamma} = S_{\gamma}(1) \oplus S_{\gamma}'$  and  $\sum_{\gamma} \bigoplus S_{\gamma}(1) \approx N_1$ ,  $\sum_{\gamma} \bigoplus S_{\gamma}' \approx N_2$ . Hence, we obtain 3). Furthermore, every  $C$  in  $[S_{\gamma}(2)]$  is isomorphic to a direct summand  $C'$  of  $S_{\gamma}'$ . Let  $B$  be in  $[S_{\gamma}(1)]$  and  $f' \in \text{Hom}_R(B, C) \not\approx \text{Hom}_R(B, C')$ . Then  $f = \phi(f')$  is not isomorphic from the assumption. Put  $B' = \{x + f(x) \mid x \in B\}$ . Then  $B' \approx B$  and  $B'$  is a direct summand of  $S_{\gamma}$  from [7],

Lemma 3. Hence,  $B' \subseteq S_\gamma(1)$ . Therefore,  $\text{Im } f \subseteq (B + B') \cap S_\gamma' \subseteq S_\gamma(1) \cap S_\gamma' = (0)$ .

2)  $\rightarrow$  1). Let  $x$  be in  $S_\gamma(1)$  and  $x = \sum x_i; x_i \in A_i \subset \bigoplus S_\gamma$  with  $[A_i] \in [N_i]$ . On the other hand, there exists, from Lemma 5, a direct summand  $B = \sum_{j=1}^t \bigoplus B_j$  of  $S_\gamma$  such that  $x \in B$  and  $[B_j] \in [M]$ . Let  $p_j$  be the projection of  $S_\gamma$  onto  $B_j$ . If  $0 \neq p_j(x) = \sum p_j(x_i), [B_j] \in [N_1]$  from 2). Hence,  $x \in \sum \bigoplus B_{j,s}$  with  $[B_{j,s}] \in [N_1]$ . Now, let  $y$  be in  $(\sum \bigoplus S_\gamma(1)) \cap N_2$  and  $y = \sum y_i; y_i \in S_{\gamma_i}(1)$ . Then there exists a direct summand  $\sum_{\gamma_i} \bigoplus B_{j,s}$  containing  $y$  as above. Hence,  $y \in (\sum \bigoplus B_{j,s}) \cap N_2 = (0)$  by Lemma 4. Therefore,  $M = \sum \bigoplus S_\gamma(1) \oplus N_2$  from Lemma 6.

3)  $\rightarrow$  1). Let  $S_\gamma = S_\gamma(1) \oplus S_\gamma'$  and  $C$  an indecomposable direct summand of  $S_\gamma'$ . Then  $C \in [N_2]$  otherwise  $C \subseteq S_\gamma(1)$ . Hence, for dense submodules  $B_{\gamma_1}, B_{\gamma_2}$  in  $S_\gamma(1)$  and  $S_\gamma'$ , respectively, we have  $[B_{\gamma_i}] \subseteq [N_i]$ . Since  $\sum \bigoplus B_{\gamma_1}$  and  $\sum \bigoplus B_{\gamma_2}$  are dense submodules of  $\sum \bigoplus S_\gamma(1)$  and  $\sum \bigoplus S_\gamma'$ , respectively by [4], Theorem 1,  $M = \sum \bigoplus S_\gamma(1) \oplus N_2$  by [6], Lemma 7.

**Corollary 1.** *Let  $M$  be as above and  $M = N_1 \oplus N_2$ . If either  $\text{Hom}_R(N_1, N_2) = 0$  or  $\text{Hom}_R(N_1, N_2) = (0)$ , then  $N_1$  and  $N_2$  have the exchange property in  $M$ , (cf. [6], Corollary 5 to Theorem).*

Proof.  $N_i$  is in  $\underline{A}$  by [6], Corollary 5 to Theorem. The condition 2) in the theorem is satisfied for any decompositions  $M = \sum \bigoplus S_\gamma$ . Hence,  $N_1$  and  $N_2$  have the exchange property in  $M$  from Theorem 3 and [6], Theorem A.2.

**Corollary 2.** *Let  $M, N_i$  and  $S_\gamma$  be as in Theorem 3. Then the following conditions are equivalent.*

- 1)  $M = (\sum \bigoplus S_\gamma(1)) \oplus N_2 = N_1 \oplus (\sum \bigoplus S_\gamma(2))$
- 2)  $\text{Hom}_R([S_\gamma(1)], [S_\gamma(2)]) = (0) = \text{Hom}_R([S_\gamma(2)], [S_\gamma(1)])$
- 3)  $S_\gamma = S_\gamma(1) \oplus S_\gamma(2)$ .

Proof. 1)  $\rightarrow$  3). 1) implies  $S_\gamma = S_\gamma(1) \oplus S_\gamma' = S_\gamma(2) \oplus S_\gamma''$ . Hence,  $S_\gamma(2) = S_\gamma' \oplus S_\gamma(1) \cap S_\gamma(2)$ , since  $S_\gamma' \subseteq S_\gamma(2)$ . If  $S_\gamma(1) \cap S_\gamma(2) = T \neq (0)$ ,  $T$  contains an indecomposable direct summand  $A$  of  $S_\gamma$  from Lemma 5. Then  $[A] \in [N_1] \cap [N_2]$  from the first decompositions and 1). Therefore,  $S_\gamma(1) \cap S_\gamma(2) = (0)$  and  $S_\gamma = S_\gamma(1) \oplus S_\gamma(2)$ . Other implications are clear from Theorem 3.

## References

- [1] G. Azumaya: *Correction and supplementaries to my paper concerning Krull-Remak-Schmidt's theorem*, Nagoya Math. J. **1** (1950), 117–124.
- [2] P. Crawley and B. Jónnsson: *Refinements for infinite direct decomposition of algebraic systems*, Pacific J. Math. **14** (1964), 797–855.
- [3] M. Harada and Y. Sai: *On categories of indecomposable modules I*, Osaka J. Math. **7** (1970), 323–344.
- [4] M. Harada: *On categories of indecomposable modules II*, ibid. **8** (1971), 309–321.
- [5] \_\_\_\_\_: *Supplementary remarks on categories of indecomposable modules*, ibid. **9** (1972), 49–55.
- [6] \_\_\_\_\_: *On the exchange property in a directsum of indecomposable modules*, Osaka J. Math. **12** (1975), 719–736.
- [7] T. Ishii: *On locally direct summands of modules*, Osaka J. Math. (1975), 473–482.
- [8] Z. Papp: *On algebraically closed modules*, Publ. Math. Debrecen **6** (1959), 311–323.
- [9] R.B. Warfield Jr.: *A Krull-Schmidt theorem for infinite sums of modules*, Proc. Amer. Math. Soc. **22** (1969), 460–465.