

Title	ISOLATION AND SELECTION OF LIPASE-PRODUCING BACTERIA IN VIETNAM
Author(s)	Lai, Thuy Hien; Le, Phi Nga; Nguyen, Thi Thu Huyen et al.
Citation	Annual Report of FY 2001, The Core University Program between Japan Society for the Promotion of Science(JSPS) and National Centre for Natural Science and Technology(NCST). 2003, p. 234-238
Version Type	VoR
URL	https://hdl.handle.net/11094/12946
rights	
Note	

# Osaka University Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

Osaka University

# ISOLATION AND SELECTION OF LIPASE - PRODUCING BACTERIA IN VIETNAM

Lai Thuy Hien<sup>1</sup>, Le Phi Nga<sup>1</sup>, Nguyen Thi Thu Huyen<sup>1</sup>, Pham Thi Hang<sup>1</sup>, Vuong Thi Nga<sup>1</sup>, Nguyen Van Long<sup>1</sup>, M. Tateda<sup>2</sup>, M. Fujita<sup>2</sup>

1 Institute of Biotechnology, NCST, Hanoi 3 Osaka University

# Abstract

From deferent sources of Vietnam such as petrol product reservoirs, oil polluted sand, airplane fuel, food wastewater were isolated decades of lipase-producing bacteria. Six strains with high lipase-producing activity were selected. The effect of olive oil on lipase production was investigated. Olive oil concentration was optimal for lipase producing from 4% to 7%. In particularly, all strain was capable to produce lipase on medium containing 10% olive oil. Using API 20E kit identified three of strains. They belong to *Pseudomonas* genus. One of which is *Pseudomonas aeruginosa* N9. This strain produced high lipase (2.53 u/ml) therefore it could have great potential for application in the detergent industry.

Key words: API 20E kit, Characterization, Lipase activities, Lipase - producing bacteria, *Pseudomonas* sp., Olive oil, RhodaminB.

## Introduction

A large number of surface-active agents are used in the detergent industry. In particular, lipases that catalyze syntheses of a wide variety of esters have been studied by a large number of researcher [7].

Lipases can be obtained from many species of plants, animals and microorganisms. However, most attention has been paid to microbial source [3].

Lipases are a class of triglyceride-hydrolyzing enzymes whose act at oil-water interfaces to catalyze the hydrolysis of long-chain triglycrides. Lipases not only hydrolyze ester bonds, trans-esterify triglycerides, and resolve racemic mixture, but also, in the reverse mode, to synthesize ester and peptide bonds depending on their sources with respect to positional, specificity, fatty acid specificity, thermo-stability, pH and temperature optimum, etc. Since lipases can evidently catalyze numerous different reactions, they have been widely used in industrial applications such as in food, chemical, pharmaceutical, and detergent industries [5,6,8,9,11].

In this paper we report the production of lipase by some bacteria strains. The aim of work is isolation, sellection high lipase-producing bacteria, investigation the effect of olive oil concentration on lipase production, and identification some species producing high lipase.

## Materials and methods

## Microorganisms

Some lipase-producing strains: TL-N<sub>1</sub>A, TL-N<sub>1</sub>B, TL-N<sub>1</sub>C were isolated from Thuongly petrol product reservoirs. Strain ASB was isolated from S7-A320 airplane fuel. Another strain, P.HQV4 was isolated from food oil wastewater. The other strain, *Pseudomonas aeruginosa* N9 was isolated from expanse of stand containing oil polluted on Nhatrang beach.

# Medium and cultivation

Medium A for isolation and selection of the strains contained 0.5%(w/v) peptone; 0.3%(w/v) yeast extract; 1%(v/v) olive oil; and 0.2%(v/v) TritonX-100 in running water. The pH was initially adjusted to 6.0 by

PBS buffer 0.1M(pH 6.0). For plate cultivation, 1.5%(w/v) agar and 0.01%(v/v) RhodaminB were added to this medium. Cultivation on agar plate for isolation of strains was carried out at  $30^{\circ}$ C for 3 days [8].

## Isolation of strains

All strains were pre-cultivated with shaking (140 rpm) in medium A for 1 day, and then were inoculated on agar plates. After 3 days, the strains growing on the plates were picked up, and single-colony isolation on medium agar A was done and observed on fluorescence microscope.

## Characterization of bacteria

The morphological properties and taxonomical characteristics of bacteria were determined by Gram staining methods, were observed on JEM 1010 electron microscope and was identified by using API 20E kit.

## Preparation of extracellular lipases

Medium B for lipase production is the same to medium A without agar. However, ASB strain was inoculated in mineral medium supplemented 1% NaCl, 0.5% yeast extract, and 0.2% TritonX-100 and olive oil. For determination of extracellular lipase activity, shake culture were carried out in 20 ml of medium B with shaking (140 rpm) in 100 ml flasks at  $30^{\circ}$ C. After 2 days, the cultures were centrifuged at 10000 rpm for 10 min at  $4^{\circ}$ C, and the supernatants were used as extracellular lipase fractions.

## Assay of lipase activity

The reaction mixture contained 1ml olive oil, 8ml PBS buffer (100 mM) pH 7.6, 1ml CaCl<sub>2</sub> 0.1M and 1ml enzyme solution. The reaction was carried out at  $37^{\circ}$ C for 15 min with stirring at 500 rpm. Amount of liberated fatty acid was titrated with 10mM KOH by a pH meter. One unit of lipase activity was defined as the amount of enzyme required to release 1µmol fatty acid per min at  $37^{\circ}$ C and pH7.6 [1,3,5,8,9,].

## **Results and discussion**

## Isolation of bacteria strains

We isolated decade's strains from water samples from various environmental sites that were screened as potential sources of lipase-producing bacteria. Among 10 trains, we selected 6 strains showing better produce lipase than the other did.

## Characterization of strains

6 strains were Gram negative, rod shapes, flagella. 4 of strains were observed on electron microscope (Fig1). TL-N<sub>1</sub>C strain had  $0.55\mu$ m×  $1.15\mu$ m, rod shape, and polar flagella. ASB strain had  $0.5\mu$ m×  $1\mu$ m, short, rod shape, polar flagella. *Pseudomonas aeruginosa* N9 had  $0.83\mu$ m× $3.08\mu$ m, rod shape, and single polar flagella. P.HQV4 had  $0.6\mu$ m× $1.25\mu$ m, polar flagella. After 3 days incubating on Rhodamin B agar medium, ASB and *Pseudomonas aeruginosa* N9 were observed on fluorescence microscope. Both two strains tinted fluorescence (Fig 2).

# Dentification of strains

3 of strains were identified using API 20E test as *Pseudomonas*. One strain was *Pseudomonas aeruginosa* N9. TL-N<sub>1</sub>B and P.HQV4 strains were *Pseudomonas* sp.

Fig. 1. Cells of lipase-producing bacteria on electron microscope (× 20000): A - cell of TL-N<sub>1</sub>C; B - cell of ASB; C - cell of *Pseudomonas aeruginosa* N9.



Fig 2. Colony of lipase producing bacteria on fluorescence microscope (×330): E - ASB; F - Pseudomonas aeruginosa N9.

## Effect of olive oil on the production of lipase

Because olive oil is suitable natural substrate for lipase producing bacteria. It is a limiting factor in lipase production. In this work, we cultivated 6 strains in medium B containing different olive oil concentration such as 1%, 4%, 7% and 10%. The result was showed on Fig 3 to Fig 8.



Fig. 3. Production of lipase by TL-N1A





Fig. 5. Production of lipase by TL-N<sub>1</sub>C



Fig. 7. Production of lipase by Pseudomonas





Fig. 8. Production of lipase by P.HQV4

The result showed that all strains were able to produce lipase in 10% olive oil. In particular, each strain produced lipase optimal in different olive oil concentration. 4% olive oil was optimal for TL-N<sub>1</sub>C. 7% olive oil was optimal for Pseudomonas aeruginosa N9, TL-N<sub>1</sub>A, TL-N<sub>1</sub>B, ASB strains. 10% olive oil was optimal for P.HQV4 strain and this strain was capable to produce lipase in high oil concentration. It was very interesting that *Pseudomonas aeruginosa* N9 produced high lipase in all different olive oil concentration. Pseudomonas strain may be applied in detergent industry.

## Conclusion

From different sources of Vietnam, 6 strains of lipase producing bacteria were isolated and selected. The highest lipase activity was produced by Pseudomonas aeruginosa N9 (2,53 u/ml). The lipase activity of this strain could be applied in detergent industry.

## References

- 1. Bornscheuer U., Reif O-W. Lausch R., Freitag R., Scheper T., Kolisis N.F., Menge U. Lipase of Pseudomonas cepacia for biotechnological purposes: purification, crystallization and characterization. Biochimica et biophysica acta. 1201, pp. 55-60. (1994)
- Chen J., Shimura S., Kirimura K., Usami S. Enhancement of lipase production from hydrocarbons by mutation of Trichosporon fermentans. Applied and microbiology biotechnology. 38:pp. 714-718.1993
- Chen S-J. Cheng C-Y., Chen T-L. Production of alkaline lipase by Acinetobacter radioresistens. Journal of fermentation and bioengineering. Vol.86. No.3., pp. 308-312. (1998).
- 4. Dunhaupt A., Lang B., Wagner P. Pseudomonas cepacia lipase: studied on aggregation purification and on the cleavage of olive oil. Biotechnology letters. Vol.14. No.10. pp. 953-958. (1992).
- 5. Gilbert J.E., Drozd W.J., Jones W.C. Physiological regulation and optimization of lipase activity in Pseudomonas aeruginosa EF2. Journal of general microbiology. 137, pp. 2215-2221. (1991)

- 6. Ishimoto R., Sugimoto M., Kawai F. Screening and characterization of trehalose-oleate hydrolyzing lipase. FEMS microbiology letters. pp. 231-23. (2001).
- 7. Kawakami K., Yoshida S., *Thermal stabilization of lipase by sol-gel entrapment in organically modified silicates formed on kieselguhr*. Journal of fermentation and bioengineering. Vol. 82. No.3. pp.239-245.96
- 8. Lin S-F. Production and stabilization of a solvent-tolerant alkaline lipase from Pseudomonas pseudoalcaligenes *F-111*. Journal of fermentation and bioengineering. Vol.82. No.5. pp. 448-451.1996
- 9. Sugihara A., Tani T., Tominaga Y. Purification and characterization of a novel thermostable lipase from Bacillus sp. Journal of biochemistry.109, pp. 211-216. (1991).
- Taipa M.A., Liebeton K., Costa J.V., Cabral J.M.S., Jaeger K-E. Lipase from Chromobacterium viscosum: biochemical characterization indicating homology to the lipase from Pseudamonas glumae. Biochimica et biophysica acta. 1256, pp. 396-402. (1995).
- 11. Takahashi S., Ueda M., Atomi H., Beer H.D., Bornscheuer U.B., Schmid R.D., Tanaka A., *Extracellar production of active Rhizopus oryzae lipase by Saccharomyces cerevisiae*. Journal of fermentation and bioengineering. Vol.86. No.2. pp. 164-168. (1998).