

Title	Unified Theory of Elementary Particles : in Search of Extra Dimensions
Author(s)	Hosotani, Yutaka
Citation	
Issue Date	2005-11-05
oaire:version	VoR
URL	https://hdl.handle.net/11094/13056
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Unified Theory of Elementary Particles – in Search of Extra Dimensions –

Yutaka Hosotani

Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

Even though the unified theory of electroweak interactions is very successful at low energies, there remains one part to be confirmed. It is the sector involving Higgs particles. Those Higgs particles are expected to be discovered. It has been shown recently that Higgs particles can be viewed as gauge fields in higher dimensional gauge theory. The mass of a Higgs particle and its coupling to other particles are constrained by the gauge principle. In this scenario the mass of a Higgs particle is predicted to be in the range of 120 GeV - 290 GeV, exactly in the range explored at LHC, provided that the extra dimension is curved and warped. Thus physics of extra dimensions can manifest itself in collider experiments at the LHC energies.

1 Unification in extra dimensions

At the most fundamental level quarks and leptons interact with each other by exchanging gauge bosons. Strong interactions are described by $SU(3)_C$ color gauge interactions whereas electroweak interactions by $SU(2)_L \times U(1)_Y$ gauge interactions. The associated gauge bosons are gluons, W bosons, Z bosons, and photons. There is one more field necessary to make the standard model of elementary particles to work. It is the Higgs field which not only spontaneously breaks the electroweak symmetry $SU(2)_L \times U(1)_Y$ to the electromagnetic symmetry $U(1)_{EM}$, but also gives fermions finite masses. There appear many parameters whose values are chosen to fit the observed data. There is no principle regulating the Higgs sector of the standard model.

This seemingly awkward dilemma is resolved in higher dimensional gauge theory. Long time ago Kaluza and Klein proposed an intriguing scenario in which we are living in five-dimensional spacetime.[1] They assumed that our spacetime is close to the product of four-dimensional Minkowski spacetime (M_4) and a circle (S^1) with a radius R . The metric in the five-dimensional space, g_{MN} ($M, N = 0, \dots, 4$), decomposes into the four-dimensional metric $g_{\mu\nu}$ ($\mu, \nu = 0, \dots, 3$), the off-diagonal components $g_{\mu 4}$, and g_{44} . The general coordinate invariance in the fifth

dimension implies that the $g_{\mu 4}$ components behave as four-dimensional electromagnetic gauge potential A_μ . In this manner the four-dimensional gravity and electromagnetism are unified in the five-dimensional gravity.

Motivated by Kaluza and Klein's idea, we consider non-Abelian gauge theory in five-dimensional spacetime. Gauge potential decomposes into two parts;

$$A_M = (A_\mu , A_y) . \quad (1)$$

On $M_4 \times S^1$, for instance, fields are expanded in Fourier series in the fifth coordinate y ;

$$A_M(x, y) = \sum_{n=-\infty}^{\infty} A_M^{(n)}(x) e^{iny/R} . \quad (2)$$

$A_\mu^{(0)}(x)$ describes four-dimensional gauge fields. $A_y^{(0)}(x)$ transforms as a four-dimensional scalar. It is our contention that $A_y^{(0)}(x)$ contains the four-dimensional Higgs scalar field. Thus the Higgs field is a part of gauge fields and the unification of gauge fields and Higgs fields is achieved. The scenario is called the gauge-Higgs unification.[2,3]

2 Dynamical gauge-Higgs unification

To apply the gauge-Higgs unification scenario to the electroweak interactions, several ingredients must be implemented.

(i) In the electroweak theory $SU(2)_L \times U(1)_Y$ breaks down to $U(1)_{EM}$ and the Higgs fields

transform as a doublet of the $SU(2)_L$ group. On the other hand, the extra-dimensional component of the gauge fields in the decomposition in (1) belongs to the adjoint representation of the gauge group. This means that one must begin with a larger group to achieve gauge-Higgs unification, as first clarified by Fairlie and by Manton.[2]

(ii) The electroweak symmetry is spontaneously broken by the 4-d Higgs fields, which are a part of the 5-d gauge fields. Dynamical electroweak symmetry breaking is induced by the Hosotani mechanism.[3,4] When the extra-dimensional space is non-simply connected, there appear non-Abelian generalization of the Aharonov-Bohm phases. Those non-Abelian Aharonov-Bohm phases, $\{\theta_j\}$, become dynamical degrees of freedom, even though they give vanishing field strengths at the classical level. At the quantum level those θ_j , in general, develop nonvanishing expectation values, thus breaking the gauge symmetry.

(iii) Quarks and leptons are chiral in the electroweak theory. Left-handed and right-handed fermions interact with other particles differently. The most natural and powerful way of incorporating chiral fermions in higher dimensional theory is to have an orbifold in extra dimensions.[5] As a typical example, consider $M_4 \times S^1$. An orbifold is obtained by identifying two points on S^1 :

$$(x^\mu, y) \sim (x^\mu, -y) .$$

The resultant spacetime is $M_4 \times (S^1/Z_2)$.

(iv) As is seen below, phenomenology emerging from dynamical gauge-Higgs unification in flat space contradicts with the observation. To have realistic phenomenology in Higgs particles, quarks, and leptons, extra-dimensional space should be curved. In particular, dynamical gauge-Higgs unification in the Randall-Sundrum warped space yields intriguing consequences which can be tested in the experiments at LHC.[6]

3 Extra dimensions : flat or curved?

One example in flat space with dynamical gauge-Higgs unification in electroweak

interactions is given by the $U(3)_S \times U(3)_W$ model.[7] There are two relevant non-Abelian AB phases (θ_1, θ_2) . The effective potential $V_{\text{eff}}(\theta_1, \theta_2)$ is depicted in fig. 1. The absolute minimum is located at $(\theta_1, \theta_2) = (0, \pm 0.269\pi)$ so that the electroweak symmetry $SU(2)_L \times U(1)_Y$ breaks down to the electromagnetic symmetry $U(1)_{EM}$.




Fig. 1. The effective potential for the non-Abelian AB phases $(\theta_1 = \pi a, \theta_2 = \pi b)$ in the $U(3)_S \times U(3)_W$ model in flat space.

Although the symmetry is dynamically broken, there are two major problems. The W boson mass is predicted to be $0.135/R$ where R is the radius of extra-dimensions. It implies that the Kaluza-Klein mass scale $M_{KK} = 1/R$ is about 600 GeV, which is too low. The mass of the Higgs particle is estimated from the curvature of the effective potential at its global minimum. One finds that $m_H \sim 0.871 \times \sqrt{\alpha_W} m_W$ where $\alpha_W = g_4^2/4\pi$ is the weak fine structure constant. It leads to $m_H \sim 10$ GeV, contradicting with experimental data.

These two features are generic in flat space. The observational fact that the Higgs mass should be much bigger than m_W indicates that the extra-dimensional space, if it exists, must be curved.

The most promising spacetime in the context of dynamical gauge-Higgs unification is the Randall-Sundrum (RS) warped spacetime.[8,9] It has the same topology as $M_4 \times (S^1/Z_2)$. The metric is given by

$$\begin{aligned} ds^2 &= e^{-2\sigma(y)} dx_\mu dx^\mu + dy^2 , \\ \sigma(y) &= k|y| \quad \text{for } |y| \leq \pi R , \\ \sigma(y + 2\pi R) &= \sigma(y) . \end{aligned} \tag{3}$$

As in $M_4 \times (S^1/Z_2)$, (x^μ, y) and $(x^\mu, -y)$ are identified. It is the anti-de Sitter space with a cosmological constant $-1/k^2$ sandwiched by two branes at $y = 0$ and $y = \pi R$. It has been speculated that five-dimensional anti-de Sitter space naturally emerges from such a more fundamental theory as superstring theory. Further reduction to four dimensions yields approximately conformal theory with gauge fields and light fermions.

The RS spacetime is specified with two parameters, k and R . It is natural to suppose that the structure of spacetime is determined at the Planck scale $M_{pl} = 1.2 \times 10^{19}$ GeV. As a consequence it is expected that $k = O(M_{pl})$. The size R is determined such that the theory predicts $m_W = 80.4$ GeV. As is shown below, it implies that $kR = 12 \pm 0.3$.

4 W bosons and the Kaluza-Klein mass

Consider $SU(3)$ gauge group which contains $SU(2)_L \times U(1)_Y$. Boundary conditions for the gauge fields in the RS spacetime (3) are given by

$$\begin{aligned} \begin{pmatrix} A_\mu \\ A_y \end{pmatrix} (x, y_j - y) &= P_j \begin{pmatrix} A_\mu \\ -A_y \end{pmatrix} (x, y_j + y) P_j^\dagger, \\ y_0 = 0, \quad y_1 &= \pi R, \\ P_0 = P_1 &= \begin{pmatrix} -1 & & \\ & -1 & \\ & & +1 \end{pmatrix}. \end{aligned} \quad (4)$$

With these boundary conditions zero modes (massless modes) of A_M in four dimensions exist only for components

$$A_\mu = \begin{pmatrix} \circ & \circ & \\ \circ & \circ & \\ & & \circ \end{pmatrix}, \quad A_y = \begin{pmatrix} & & \circ \\ & & \circ \\ \circ & \circ & \end{pmatrix} \quad (5)$$

where \circ is marked. The zero modes of A_μ are W bosons, Z bosons, and photons, whereas those of A_y constitute the Higgs doublet. In particular, the zero mode of A_y gives rise to non-Abelian Aharonov-Bohm phase θ_W :

$$\exp \left\{ \frac{i}{2} \theta_W \Lambda \right\} = P \exp \left\{ ig \int_0^{\pi R} dy \langle A_y \rangle \right\},$$

$$\langle A_y \rangle = ce^{2ky} \Lambda, \quad \Lambda = \begin{pmatrix} & & \\ & & 1 \\ 1 & & \end{pmatrix}. \quad (6)$$

When $\theta_W \neq 0 \pmod{2\pi}$, the $SU(2)_L$ symmetry breaks down and W bosons acquire a mass m_W given by

$$m_W \sim \sqrt{\frac{2k}{\pi R}} e^{-\pi k R} \sin \frac{\theta_W}{2}. \quad (7)$$

In the RS warped space the Kaluza-Klein mass scale M_{KK} , characterizing a mass spectrum $m_n \sim nM_{KK}$ is given by

$$\begin{aligned} M_{KK} &\sim \frac{\pi k}{e^{\pi k R} - 1} \\ &= \begin{cases} 1/R & \text{for } k \rightarrow 0 \\ \pi k e^{-\pi k R} & \text{for } kR > 2. \end{cases} \end{aligned} \quad (8)$$

In a typical model θ_W takes a value ($0.2\pi \sim 0.5\pi$). To yield $k = M_{pl}$ in (7), kR must be ($11.75 \sim 12.0$). Note that $kR = 6$ and 24 yield $k = 10^{11}$ GeV and 10^{36} GeV, respectively. Thus the value of kR is determined to be 12 ± 0.3 .

Combining (7) and (8), one obtains

$$M_{KK} \sim \frac{\pi}{\sin \frac{1}{2} \theta_W} \left(\frac{\pi k R}{2} \right)^{1/2} m_W. \quad (9)$$

This should be compared with the formula in flat space; $M_{KK} \sim (2\pi/\theta_W)m_W$. There appears an enhancement factor $\frac{1}{2}\pi k R \sim 20$ in the RS warped space. Inserting the value for kR , one finds that

$$M_{KK} = 1.6 \text{ TeV} \sim 3.6 \text{ TeV}. \quad (10)$$

At LHC, Kaluza-Klein excited states can be produced in intermediate processes so that their existence can be indirectly checked for the value in (10).

5 Higgs particles

The Higgs field ϕ in four dimensions corresponds to fluctuations of the non-Abelian Aharonov-Bohm phase θ_W . More explicitly

$$\theta_W \rightarrow g_4 \left(\frac{\pi k R}{2} \right)^{1/2} \frac{v + \phi}{M_{KK}} \quad (11)$$

where g_4 is the four-dimensional gauge coupling constant.

At the quantum level the effective potential for θ_W becomes nontrivial. Expanding it around its global minimum, one finds

$$V_{\text{eff}}(\theta_W) = \text{const.} + \frac{m_H^2}{2}\phi^2 + \frac{\eta}{3}\phi^3 + \frac{\lambda}{4}\phi^4 + \dots \quad (12)$$

The effective potential is determined, once the mass spectrum $m_n(\theta_W)$ is found for each field. It is shown that

$$V_{\text{eff}}(\theta_W) = \frac{3}{128\pi^6} M_{KK}^4 f(\theta_W) \quad (13)$$

where $f(\theta_W + 2\pi) = f(\theta_W)$ is a periodic function with an amplitude of $O(1)$.

It follows from (12) and (13) that

$$m_H = \sqrt{\frac{3\alpha_W}{32\pi} f^{(2)}(\theta_W)} \frac{\pi k R}{2} \frac{m_W}{\sin \frac{1}{2}\theta_W} \quad (14)$$

and

$$\lambda = \frac{\alpha_W^2}{16} f^{(4)}(\theta_W) \left(\frac{\pi k R}{2}\right)^2 \quad (15)$$

where $\alpha_W = g_4^2/4\pi \sim 0.03$. Notice the appearance of the enhancement factor $\frac{1}{2}\pi k R$ in (14) and (15), which distinguishes the formulas in the warped space from those in flat space. In a typical model we have found that $f^{(2)}(\theta_W)$ and $f^{(4)}(\theta_W)$ are about 4. For $\theta_W = (0.2 \sim 0.5)\pi$, one finds $m_H = (125 \sim 286)$ GeV! (The experimental bound is $m_H > 116$ GeV.[10]) It is remarkable that the dynamical gauge-Higgs unification predicts the mass of the Higgs particle exactly in the range where experiments at LHC will explore. The quartic coupling constant λ is predicted to be ~ 0.09 , though there is ambiguity in the value of $f^{(4)}(\theta_W)$.

We summarize the prediction for M_{KK} , m_H , and λ in Table 1. The values for M_{KK} and m_H predicted in flat space are inconsistent with the observation, but the situation drastically changes for the better in the Randall-Sundrum warped space.

	flat space	RS space
M_{KK}	320 \sim 800 GeV	1.6 \sim 3.5 TeV
m_H	6 \sim 15 GeV	125 \sim 286 GeV
λ	0.00025	0.09

Table 1

Prediction in the dynamical gauge-Higgs unification scenario in flat space and in the Randall-Sundrum warped space. θ_W is chosen to be $0.2\pi \sim 0.5\pi$.

6 Quarks and leptons

Another magic in the RS warped space appears in the fermion sector.[11] Each multiplet of fermions enters as a 5-d Dirac fermion in a triplet representation of $SU(3)$. For instance, a lepton multiplet ψ in the first generation consists of

$$\psi_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}, \quad \psi_R = \begin{pmatrix} \tilde{\nu}_R \\ \tilde{e}_R \\ e_R \end{pmatrix}. \quad (16)$$

The components ν_L , e_L , and e_R have zero modes which appear as 4-d ν_L , e_L , and e_R . On the other hand $\tilde{\nu}_R$, \tilde{e}_R , and \tilde{e}_L have no zero modes so that they drop from the particle spectrum in four dimensions at low energies.

The Lagrangian density for a fermion multiplet in the RS space is given by

$$\mathcal{L} = \bar{\psi} i\Gamma^a e_a^M D_M \psi - c k \bar{\psi} \psi, \quad D_M = \partial_M + \frac{1}{8} \omega_{bcM} [\Gamma^b, \Gamma^c] - ig A_M. \quad (17)$$

The covariant derivative D_M is dictated by the general coordinate invariance and the gauge invariance. $ck\bar{\psi}\psi$ is called the bulk kink mass term.[12] c is a dimensionless parameter.

Although c is called a bulk mass parameter, quarks and leptons remain massless even with $c \neq 0$ unless the electroweak symmetry breaks down. Their wave functions in the fifth dimension, however, depend on the value c . When the electroweak symmetry breaks down with nonvanishing θ_W , those quarks and lep-

tons acquire finite masses given by

$$m_f = \sqrt{\frac{\pi k R}{2} \frac{(1 - 4c^2)(z_1^2 - 1)}{(z_1^{1-2c} - 1)(z_1^{1+2c} - 1)}} m_W \quad (18)$$

where $z_1 = e^{\pi k R}$. A fermion mass is determined by c , or vice versa. See fig. 2.

$c = \pm \frac{1}{2}$ corresponds to $m_f = m_W$. Except for top quarks all fermions have $|c| > \frac{1}{2}$. As shown in Table 2, the top quark mass corresponds to $c = 0.43$ whereas the electron mass to $c = 0.87$. Although $m_t/m_e \sim 10^5$, there appears no hierarchical structure in the c space. This gives us a good hint to understand the hierarchy in the quark-lepton mass spectrum.




Fig. 2. The bulk mass parameter c v.s. the fermion mass m_f in (18).

particle	mass (GeV)	c
e	0.5110×10^{-3}	0.87
μ	0.105	0.72
τ	1.777	0.63
u	4×10^{-3}	0.81
c	1.3	0.64
t	175	0.43

Table 2
The bulk kink mass parameter c for each quark or lepton, following from (18)

7 Outlook

The results obtained in the dynamical gauge-Higgs unification in the Randall-Sundrum warped spacetime are surprising. The mass of the Higgs particle is predicted

in the range 125 GeV to 285 GeV. We have determined the fermion wave functions in terms of their masses, with which couplings of quarks and leptons to the KK excited states of W bosons etc. can be determined. In the LHC experiments we might be able to see the trace of the extra dimension, directly or indirectly.

8 Acknowledgement

This work was financially supported by the Japanese Ministry of Education and the 21st Century COE Program at Osaka University, “Towards a New Basic Science: Depth and Synthesis”.

References

- [1] Th. Kaluza, Sitzungsber, Preuss. Akad. Wiss. Berlin, Phys. Math. Klasse (1921) 966; O. Klein, Z. Phys. 37 (1926) 895.
- [2] D.B. Fairlie, *Phys. Lett.* B82 (1979) 97; N. Manton, *Nucl. Phys.* B158 (1979) 141.
- [3] Y. Hosotani, *Phys. Lett.* B126 (1983) 309; *Ann. Phys. (N. Y.)* 190 (1989) 233.
- [4] Y. Hosotani, in the Proceedings of “Dynamical Symmetry Breaking”, Nagoya University, December 2004, page 17. (arXiv:hep-ph/0504272)
- [5] A. Pomarol and M. Quiros, *Phys. Lett.* B438 (1998) 255.
- [6] Y. Hosotani and M. Mabe, *Phys. Lett.* B615 (2005) 257.
- [7] Y. Hosotani, S. Noda and K. Takenaga, *Phys. Lett.* B607 (2005) 276.
- [8] L. Randall and R. Sundrum, *Phys. Rev. Lett.* 83 (1999) 3370.
- [9] K. Oda and A. Weiler, *Phys. Lett.* B606 (2005) 408.
- [10] LEP Electroweak Working Group, <http://lepewwg.web.cern.ch/lepewwg/>.
- [11] Y. Hosotani, S. Noda and Y. Sakamura, and S. Shimasaki, in preparation.
- [12] T. Gherghetta and A. Pomarol, *Nucl. Phys.* B586 (2000) 141.