



Title	FUNGAL BIODEGRADATION OF BISPHENOL A AND BENZOPHENONE
Author(s)	Takita, O; Tanaka, T; Ooi, Toshiko et al.
Citation	Annual Report of FY 2004, The Core University Program between Japan Society for the Promotion of Science (JSPS) and Vietnamese Academy of Science and Technology (VAST). 2005, p. 185-188
Version Type	VoR
URL	<a href="https://hdl.handle.net/11094/13057">https://hdl.handle.net/11094/13057</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

# FUNGAL BIODEGRADATION OF BISPHENOL A AND BENZOPHENONE

O. Takita\*, T. Tanaka\*, T. Ooi\*, S. Kinoshita \*, L. T. T. Ai\*\* and T. L. Thuoc\*\*

\* Division of Molecular Chemistry, Graduate School of Engineering, Hokkaido University, Sapporo, 060-8628, Japan

\*\* Faculty of Biology, University of Natural Sciences, VNU-HCMC, HCMC, Vietnam

## ABSTRACT

Sixteen fungal strains from our stock culture were tested the ability to degrade endocrine disrupters, bisphenol A and benzophenone. More than 79 % of bisphenol A could degrade by the culture of *Aspergillus oryzae*, *Cheatomium globosum*, *Penicillium janthillum* LM12, and *Rhizopus oryzae*, respectively. Degradation of benzophenone was archived to less than detection limit by the culture of *Cladosporium* sp. DD618, *Geotrichum candidum*, and *Rhizopus oryzae*, respectively.

## KEYWORDS

Bisphenol, Benzophenone, Biogradation, Endocrine disrupting chemicals, Fungi,

## INTRODUCTION

Bisphenol A and benzophenone have been listed among “chemicals suspected of having endocrine disrupting effects” by the World Wildlife Fund and Japanese Environment agency.

Bisphenol A is widely used in industry and dentistry to make polycarbonate plastics and epoxy resins (Staples et al., 1998) and is one of the most common environmental endocrine disruptors with an oestrogenic activity (Krishnan et al, 1993) and an acute toxicity to aquatic organisms (Alexander et al. 1988).

Benzophenone has been used as an ingredient of pharmaceuticals, insecticides, agricultural chemicals and fragrance in medicine and industry and as an ultraviolet light-absorber in plastic and polymers. Its derivatives are widely used in cosmetic products such as photostabilizers to protect the skin and hair from ultraviolet irradiation. Benzophenone has also weak estrogenic activity as well as bisphenol A (Nakagawa et al. 2000).

There is an increasing interest in the microbial degradation of such endocrine disruptors. The white rot fungi and their enzymes can degrade bisphenol A (Hirano et al. 2000; Tsutsumi et al. 2001) and remove oestrogenic activity. Some gram-negative bacteria and *Streptomyces* sp. strain also reported to degrade bisphenol A (Lobos et al. 1992; Kang et al. 2004).

The aim of this study was to identify bisphenol A and benzophenone biodegradability by 16 fungi from our stock culture. A HPLC system was used for analysis of these compounds as the economical methods.

## MATERIALS AND METHODS

### Chemicals, microorganisms, and culture condition

Bisphenol A and benzophenone (Fig. 1) were purchased from Wako pure chemical (Osaka). Sixteen fungal strains from our stock culture were used in this study as listed in Table 1. The medium used for the fungal degradation contained 4.5 g/l glycerol, 1.0 g/l yeast extract, 0.4 g/l ammonium

sulfate, 0.2 g/l  $\text{KH}_2\text{PO}_4$ , 0.3 g/l  $\text{K}_2\text{HPO}_4$ , and 0.1 g/l  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ . Initial pH was adjusted to 5.6. The fungal strains were cultivated in a reciprocal shaker at 25°C. Bisphenol A and benzophenone were solubilized in ethanol and then added to the medium at final concentration of 15 ppm and 7.5 ppm, respectively.




Fig. 1. Chemical structure of bisphenol A and benzophenone

Table 1 List of fungi used for degradation of bisphenol A and benzophenone

<i>Aspergillus niger</i>	<i>Penicillium roquefortii</i>
<i>Aspergillus oryzae</i>	<i>Penicillium SL413</i>
<i>Chaetomium globosum</i>	<i>Rhizopus oryzae</i>
<i>Cladosporium</i> DD618	<i>Schizophyllum commune</i>
<i>Cladosporium herbarum</i>	<i>Scopulariopsis brevicaulis</i>
<i>Geotrichum candidum</i>	<i>Sporotrichum cellulophilum</i>
<i>Mucor javahicus</i>	<i>Trametes hirsuta</i>
<i>Penicillium janthillum</i> LM12	<i>Trichoderma viride</i>

### HPLC analysis

After the cultivation at appropriate day, 2 volume of methanol was added to cultured medium and filtrate with a glass filter (GS25, Advantec) to remove insoluble materials and mycelium. Bisphenol A and benzophenone concentration in the filtrate was assayed by HPLC on a Microsorb MV 86 203-C5 reversed phase column (4.6 ID X 25 cm, Rainin, USA) in a Hitachi L6200 apparatus. The solvent system was 50% acetonitrile. The flow rate was 1.0 ml/min and elution was monitored at 279 nm for bisphenol A and 252 nm for benzophenone with Hitachi L4000H UV detector. Concentration of bisphenol A and benzophenone were calculated from peak area of standard curve. Detection limit by the HPLC was 0.05 ppm for bisphenol A and 0.01 ppm for benzophenone, respectively.

## RESULTS AND DISCUSSION

### Degradation of bisphenol A

Fungal strains were cultivated in bisphenol A (15 ppm) containing medium for 7 days. More than 80 % of bisphenol A were degraded by the culture of *Aspergillus oryzae* (80.7 % for 7 days), *Cheatomium globosum* (100 % for 6 days), *Penicillium janthillum* LM12 (80 % for 7 days), and *Rhizopus oryzae* (100 % for 5 days) as shown in Fig. 2.

### Degradation of benzophenone

Sixteen fungal strains were cultivated in the medium containing 15 ppm of benzophenone at 25°C for 7 days. Three strains, *Cladosporium* sp. DD618 (5 days), *Geotricum candidum* (6 days), and *Rhizopus oryzae* (3 days) could degrade efficiently benzophenone to below detection limit. Time course for benzophenone degradation of the three strains are shown in Fig. 3.




Fig. 2 Time course of bisphenol A degradation by fungi




Fig. 3 Time course of benzophenone degradation by fungi

## CONCLUSIONS

Biodegradability of bisphenol and benzophenone were tested by using 16 fungal strains. Bisphenol A was degraded by the culture of *Aspergillus oryzae* (79.7 % for 7 days), *Cheatomium globosum* (100 % for 6 days), *Penicillium janthillum LM12* (79.6 % for 7 days), and *Rhizopus oryzae* (100 % for 6 days). Benzophenone could degrade by the culture of *Cladosporium sp. DD618*, *Geotrichum candidum*, and *Rhizopus oryzae* could degrade benzophenone to less than detection limit. .

## REFERENCES

Alexander H.C., Dill D.C., Smith L.W., Guiney P.D., Dorn P.B. (1988) Bisphenol A: acute aquatic toxicity. *Environmental Toxicology and Chemistry* **7**, 19-26.

Hirano T., Honda Y., Watanabe T., Kuwahara M. (2000) degradation of bisphenol A by the lignin-degrading enzyme, manganese peroxidase, produced by the white-rot basidiomycete. *Bioscience, Biotechnology and Biochemistry*. **64**, 1958-1962.

Kang J. -H., Ri N., Kondo F. (2004) *Streptomyces* sp. strain isolated from river water has high bisphenol A degradability. *Letters in Applied microbiology* **39**, 178-180.

Krishnan A.V., Stathis P., Permuth S.F., Tokes L., Feldman D. (1993) Bisphenol A: an estrogenic substance is released from polycarbonate flasks during autoclaving. *Endocrinology* **132**, 2279-2286.

Lobos J.H., Leib T.K., Su T.M. (1992) Biodegradation of bisphenol A and other bisphenols by gram-negative aerobic bacterium. *Applied and Environmental Microbiology* **58**, 1823-1831.

Nakagawa Y., Suzuki T., Toyama S. Metabolism and toxicity of benzophenone in isolated rat hepatocytes and estrogenic activity of its metabolites in MCF-7 cells. *Toxicology* **156**, 27-36.

Staples C.A., Dorn P.B., Klecka G.M., O'Block S.T., Harris L.R. (1998) A review of the environmental fate, effects, and exposures of bisphenol A. *Chemosphere* **36**, 2149-2173.

Tsutumi Y., Haneda T., Nishida T. (2001) Removal of estrogenic activity of bisphenol A and nonylphenol by oxidative enzymes from lignin-degrading basidiomycetes. *Chemosphere* **42**, 271-276.