

Title	THE AR- PROPERTY AND THE FIXED POINT PROPERTY FOR COMPACT MAPS OF A SOME CONVEX SUBSET IN THE SPACE $LP(0 < p < 1)$
Author(s)	Le, Hoang Tri
Citation	Annual Report of FY 2007, The Core University Program between Japan Society for the Promotion of Science (JSPS) and Vietnamese Academy of Science and Technology (VAST). p590-p.595
Issue Date	2008
oaire:version	VoR
URL	https://hdl.handle.net/11094/13105
DOI	
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

THE AR- PROPERTY AND THE FIXED POINT PROPERTY FOR COMPACT MAPS OF A SOME CONVEX SUBSET IN THE SPACE $l_p(0 < p < 1)$

Le Hoang Tri

Abstract. We know that every convex subset in locally convex linear metric space is an absolute retract and it has the fixed point property for compact maps. However it is not know whether a convex subset of a non-locally convex space has the property? The aim of this paper is to prove the AR- property of bound convex subsets in the space $l_p(0 < p < 1)$ (non-locally convex linear metric space) and prove that normal simple in the space l_p has the fixed point property for compact maps.

1 Introduction

A topological space Y is called an absolute retract whether

- a) Y is metrizable and
- b) for an metriczable X and closed $A \subset X$ each $f : A \mapsto Y$ is extendable over X .

The class of absolute retracts is denoted by AR.

In 1951 Dugundji proved the following theorem, see [1]

Theorem 1.1 (Dugundji). Every convex subset of a locally convex linear metric space is a absolute retract.

It is not know that whether a convex subset of a non-locally convex space has the property?

For every $p \in (0, 1)$, we will denote

$$l_p = \{x = (x_n) \subset \mathbb{R} : \sum_{n=1}^{\infty} |x_n|^p < +\infty\}$$

$\forall x = (x_n), y = (y_n)$; let $\|x\| = \sum_{n=1}^{\infty} |x_n|^p, d(x, y) = \sum_{n=1}^{\infty} |x_n - y_n|^p$. The space l_p is a non-locally convex linear metric space.

In 2, we will prove that every bound convex subset in the space l_p is a absolute retract.

Let \mathbf{X}, \mathbf{Y} be topological spaces. A continous map $F : \mathbf{X} \rightarrow \mathbf{Y}$ is called compact if $F(\mathbf{X})$ is contained in a compact subset of \mathbf{Y} .

A topological \mathbf{X} is called has the fixed point property for compact maps iff every compact map $F : \mathbf{X} \rightarrow \mathbf{X}$, there exits a point $x_0 \in \mathbf{X}$:

$$F(x_0) = x_0.$$

We know that every convex subset in locally convex linear topological space has the fixed point property for compact maps (see [2], page 71).

In [3], we will prove that normal simplex in the space l_p (non-locally convex linear metric space) has the fixed point property for compact maps.

We know that every AR has the fixed point property for compact maps(Borsuk Theorem, see [2],p.94). Therefore every convex subset in locally convex linear metric space has the fixed point property for compact maps; where normal simplex in the space l_p is unbound convex set.

2 The AR - property of bound convex subset in the space $l_p(0 < p < 1)$

The main result of 2 is

Theorem 2.1. Every bound convex subset in the space l_p is an AR.

To prove this theorem, we have to use following lemmas

Lemma 2.2. Every bound convex subset in the space l_p is totally bounded

Proof. See [3]. □

Let K is a convex subset in the linear space \mathbf{X} , a map f : from K into the linear space \mathbf{Y} is called affine map if

$$\forall n \in \mathbb{N}; \forall x_1, x_2, \dots, x_n \in K; \forall \alpha_1, \alpha_2, \dots, \alpha_n \geq 0 : \alpha_1 + \alpha_2 + \dots + \alpha_n = 1,$$

$$f(\alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n) = \alpha_1 f(x_1) + \alpha_2 f(x_2) + \dots + \alpha_n f(x_n).$$

We have

Lemma 2.3. Every compact convex subset in the space l_p ($0 < p < 1$) is homeomorphic to a compact convex subset in the locally convex linear metric space $\mathbb{R}^\infty = \mathbb{R} \times \mathbb{R} \times \dots$.

Proof. The space $\mathbb{R}^\infty = \mathbb{R} \times \mathbb{R} \times \dots$ with metric d is defined by

$$d(x, y) = \sum_{n=1}^{\infty} \frac{2^{-n} |x_n - y_n|}{1 + |x_n - y_n|}, \quad \forall x = (x_n), y = (y_n) \in \mathbb{R}^\infty$$

(see [1], p.36)

$\forall r > 0, \exists n_0 \in \mathbb{N}$ such that $\sum_{n=n_0+1}^{\infty} 2^{-n} < \frac{r}{2}$. We have $V = \{x = (x_n) \in \mathbb{R}^\infty : |x_n| < \frac{r}{2 \cdot 2^n}, \forall n \in \{1, 2, \dots, n_0\}\}$ is open convex neighbourhood of zero in \mathbb{R}^∞ and $V \subset B(0, r)$. Therefore \mathbb{R}^∞ is the locally convex linear metric space.

$\forall n \in \mathbb{N}$, let $p_n : l_p \mapsto \mathbb{R}$ is defined by

$$p_n(x) = x_n$$

where $x = (x_1, x_2, \dots, x_n, \dots) \in l_p$.

We have $\forall x = (x_1, x_2, \dots, x_n, \dots), y = (y_1, y_2, \dots, y_n, \dots) \in l_p :$

$$d(x, y) = \sum_{i=1}^{\infty} |x_i - y_i|^p \geq |x_n - y_n|^p.$$

Hence $p_n : l_p \mapsto \mathbb{R}$ is continuous and

$$P : l_p \mapsto \mathbb{R}^\infty$$

is defined by

$$P(x) = (p_1(x), p_2(x), \dots, p_n(x), \dots), \forall x = (x_1, x_2, \dots, x_n, \dots) \in l_p$$

is continuous linear map.

Now, let K is a arbitrary compact convex subset in the l_p , then restrict of $P \circ K$ is a continuous affine map and injection. We see that K is compact. Therefore this map is a homeomorphic from K on compact convex subset $f(K)$ in \mathbb{R}^∞ . \square

Proof of Theorem 2.1. Let A is a bound convex subset in the space l_p , by Lemma 2.3, A is total bound subset $\Rightarrow K = clA$ is compact convex subset of l_p ; let B is a closed subset of a metric space \mathbf{X} and $g : B \rightarrow A$ is a continuous map $\Rightarrow P \circ g : B \rightarrow \mathbb{R}^\infty$ is continuous and $P \circ g(B) \subset P(A)$ (where P is defined in Lemma 2.3)

Using Dugundji Theorem, there exists $H : \mathbf{X} \rightarrow \mathbb{R}^\infty$ is continuous extension of $P \circ g$ and $H(\mathbf{X}) \subset conv P \circ g(B) \subset conv P(A) = P(A)$.

Let $P^* : K \mapsto P(K)$ is homeomorphic defined by $P^*(x) = P(x), \forall x \in K$. Therefore $(P^*)^{-1} \circ H : \mathbf{X} \mapsto A$ is a continuous extension of $g \Rightarrow A$ is a AR \square

3 The fixed point property for compact maps of normal simplex in the space $l_p(0 < p < 1)$

In $l_p(0 < p < 1)$. Let

$$e_1 = (1, 0, 0, \dots, 0, \dots)$$

$$e_2 = (0, 1, 0, \dots, 0, \dots)$$

$$e_3 = (0, 0, 1, 0, \dots, 0, \dots)$$

.....

$$e_n = (0, 0, \dots, 1, 0, \dots)$$

.....

and $A = \overline{conv\{e_1, e_2, \dots, e_n, \dots\}}$, A is the normal simplex in the space $l_p(0 < p < 1)$. We see that A is a unbound convex subset in the space $l_p(0 < p < 1)$. We will prove that A has fixed point property for compact maps.

Theorem 3.1. A has fixed point property for compact maps.

We need prove following lemmas.

Lemma 3.2. Let (X, d) is a metric space, $f : X \rightarrow X$ is a compact map without fixed point. Then $\exists \epsilon_0 > 0 : d(x, f(x)) \geq \epsilon_0; \forall x \in X$

Proof. Let K is a compact in X such that $f(X) \subset K$. We need prove that $\exists \epsilon_0 > 0 : d(x, f(x)) \geq \epsilon_0; \forall x \in X$. Suppose contrary to our claim, that $\forall \epsilon > 0, \exists x \in X : d(x, f(x)) < \epsilon$.

$\forall n \in \mathbb{N}$, choose $\epsilon = \frac{1}{n} \Rightarrow \exists x_n \in X : d(x_n, f(x_n)) < \frac{1}{n}; \forall n \in \mathbb{N}$. We have $\{f(x_n)\} \subset K$ and K is compact \Rightarrow there exists a subsequence $\{f(x_{m_n})\}$ of

$\{f(x_n)\}$ and $\exists y_0 \in K : \lim_{n \rightarrow \infty} d(y_0, f(x_{m_n})) = 0(1)$.

We have $d(x_n, f(x_n)) < \frac{1}{n}; \forall n \in \mathbb{N} \Rightarrow d(x_{m_n}, f(x_{m_n})) < \frac{1}{m_n} \leq \frac{1}{n}; \forall n \in \mathbb{N}$
 $\Rightarrow \lim_{n \rightarrow \infty} d(x_{m_n}, f(x_{m_n})) = 0(2)$.

(1), (2) $\Rightarrow \lim_{n \rightarrow \infty} d(y_0, x_{m_n}) = 0 \Rightarrow \lim_{n \rightarrow \infty} d(f(y_0), f(x_{m_n})) = 0$.

$\Rightarrow f(y_0) = y_0 \Rightarrow$ This contradicts our assumption. □

Definition 3.3. Let A is a convex subset in a linear metric space (X, d) , A is called admissible if $\forall \epsilon > 0$, every compact subset K in A , there exists continuous function $h : K \rightarrow A$ such that $d(x, f(x)) < \epsilon; \forall x \in K$ and $h(K)$ is contained a finite dimensional linear subspace of X

Lemma 3.4. Let A is a convex subset in a linear metric space (X, d) . If A is admissible, the A has fixed point property for compact maps.

Proof. Suppose contrary, that there exists compact map $f : A \rightarrow A$ and $f(x) \neq x; \forall x \in A$. Using lemma 3.2, then $\exists \epsilon_0 > 0 : d(x, f(x)) \geq \epsilon_0; \forall x \in A$ Let K is a compact map in A such that $f(K) \subset K$. We see that A is admissible, there exists continuous function $g : K \rightarrow A : d(g(x), x) < \frac{\epsilon_0}{4}; \forall x \in K$ and $g(K)$ is contained in a finite dimensional linear subspace of X . $\Rightarrow g(K) \subset L \cap A \Rightarrow L \cap A$ is a convex in a finite dimensional linear metric space L .

Consider $g \circ f|_{L \cap A} : L \cap A \rightarrow L \cap A$. We know that every finite dimensional linear metric space is a locally convex linear metric space $\Rightarrow L \cap A$ is an AR $\Rightarrow L \cap A$ has fixed point property for compact maps and $g \circ f|_{L \cap A}(L \cap A) \subset g(K)$, $g(K)$ is a compact set $\Rightarrow \exists x_0 \in L \cap A : g \circ f(x_0) = x_0 \Rightarrow d(g(f(x_0)), x_0) < \frac{\epsilon_0}{4} \Rightarrow d(f(x_0), x_0) < \frac{\epsilon_0}{4}$.

This contradicts our assumption. □

We only give main ideas of the proof of theorem 3.1. By Lemma 3.4, we only prove that A is admissible.

$\forall \epsilon > 0$. Let K is a arbitray compact in A . $\forall n \in \mathbb{N}, \forall x = (x_1, x_2, \dots, x_n, \dots) \in K$. Let $f_n(x) = (x_1, x_2, \dots, x_n, 1 - x_1 - x_2 - \dots - x_n, 0, 0, \dots)$. We see that $f_n(K)$ is contained in a finite dimensional linear subspace of l_p and when n is sufficient large, we have: $d(f_n(x), x) < \epsilon; \forall x \in K$.

Therefore A has fixed point property for compact maps.

References

- [1] C. Bessage and A. Pelczynsky, Selected topics in infinite dimensional topology, PWN, Warszawa, 1975.

- [2] J. Dugundji and A. Granas, Fixed point property, I, Warszawa, 1982.
- [3] Le Hoang Tri, The AR - property of bound convex in the space $l_p(0 < p < 1)$, Journal of science and technology, University of Danang, pp.59-64, No 1(13), April 2006
- [4] Le Hoang Tri, The fixed point property for compact maps of normal simplex in the space $l_p(0 < p < 1)$, Journal of science and technology, University of Danang, preprint