

Title	Arsenic contamination in groundwater and residents from Ha Nam and Ha Tay Provinces of Red River Delta, Vietnam
Author(s)	Agusa, Tetsuro; Kunito, Takashi; Inoue, Suguru; Tu, Binh Minh; P, T. K. Trang; Subramanian, A.; Iwata, Hisato; Pham, Hung Viet; Tanabe, Shinsuke
Citation	Annual Report of FY 2005, The Core University Program between Japan Society for the Promotion of Science (JSPS) and Vietnamese Academy of Science and Technology (VAST). P.33-P.36
Issue Date	2006
Text Version	publisher
URL	http://hdl.handle.net/11094/13171
DOI	
rights	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/repo/ouka/all/>

Arsenic contamination in groundwater and residents from Ha Nam and Ha Tay Provinces of Red River Delta, Vietnam

T. Agusa¹, T. Kunito², S. Inoue¹, T.B. Minh¹, P.T.K. Trang³, A. Subramanian¹,
H. Iwata¹, P.H. Viet³, S. Tanabe¹

¹ Center for Marine Environmental Studies (CMES), Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan

(E-mail: ax@agr.ehime-u.ac.jp)

² Department of Environmental Sciences, Faculty of Science, Shinshu University, 3-1-1 Asahi, Matsumoto 390-8621, Japan

³ Center for Environmental Technology and Sustainable Development (CETASD), Hanoi National University, Hanoi, 334 Nguyen Trai, Thanh Xuan, Hanoi, Vietnam

Abstract To make clear the distribution of As contamination in the Red River Delta, Vietnam, As concentrations were measured in groundwater and human urine from Ha Nam and Ha Tay Provinces. Concentrations of As in groundwater were from 3.0 µg/l to 486 µg/l (median; 256 µg/l) in Ha Nam Province and from 132 µg/l to 344 µg/l (median; 194 µg/l) in Ha Tay Province. Most of groundwater samples had As more than the WHO guideline value (10 µg/l) for drinking. These results suggest that human health risk in both provinces is of concern and that As pollution may be widely distributed in the Red River Delta. Dimethylarsinic acid (DMA) was the most predominant compound in almost all the residents. Monomethylarsonic acid (MMA), arsenite (As[III]) and arsenate (As[V]) were also detected in the urine samples. A significant positive correlation between concentrations of As in groundwater and in urine of residents was observed. DMA, As[III] and As[V] concentrations in human urine were also positively correlated with As concentrations in groundwater. These results suggest that residents in Ha Nam and Ha Tay are exposed to As through drinking groundwater and human health risks by As exposure are of great concern.

Keywords: arsenic; groundwater; human urine; Red River Delta; Vietnam

Introduction

Numerous occurrences of As contamination in groundwater and its related health issues have been reported in the world (Nordstrom, 2002). Especially, this problem in Bangladesh and West Bengal, India is known as one of the most serious cases (Nordstrom, 2002). In Vietnam, Berg et al. (2001) reported for the first time that there is As pollution in groundwater collected from Hanoi and its maximum level (up to 3,050 µg/l) was comparable to those in Bengal Delta. A significant positive

correlation between As concentrations in groundwater and human hair in this region was observed, suggesting that people are exposed to As through groundwater consumption (Agusa et al., 2004; Agusa et al., 2006). Because population using groundwater in the Red River Delta is expected over 11 million, As contamination in this region and hence human health risk with As exposure are of great concern. However, there is little information on As contamination in the Red River Delta. This study was conducted to make clear the contamination status of As in groundwater and residents in Ha Nam Province and Ha Tay Province of the Red River Delta, Vietnam.

Materials and Methods

Sampling was conducted in Ha Nam and Ha Tay Provinces in the Red River Delta during December 2004. From these regions, 23 groundwater and 97 human urine samples were collected and all the samples were stored at $-20\text{ }^{\circ}\text{C}$. Informed consent was obtained from all the donors and the samples were collected in an ethical manner.

The groundwater samples were acidified with sulfuric acid. Analysis of As in groundwater was performed with a hydride generation atomic absorption spectrometer (Shimadzu HVG-1 hydride system coupled to a Shimadzu-AA680 AAS) (Agusa et al., 2004). The urine samples were diluted by Milli-Q water without chemical treatment. Arsenic compounds (arsenobetaine (AB), dimethylarsinic acid (DMA), monomethylarsonic acid (MMA), arsenite (As[III]) and arsenate (As[V])) were determined in urine samples with a high performance liquid chromatograph (Shimadzu, LC10A Series) – inductively coupled mass spectrometer (Hewlett-Packard, HP-4500) using an anion exchange column (Shodex Asahipak ES-502N 7C) (Mandal et al., 2001). Total As concentration in urine was represented as sum of As compounds detected.

Results and Discussion

Arsenic was detected in all the groundwater samples and their levels were from $3.0\text{ }\mu\text{g/l}$ to $486\text{ }\mu\text{g/l}$ (median; $256\text{ }\mu\text{g/l}$) in Ha Nam Province and from $132\text{ }\mu\text{g/l}$ to $344\text{ }\mu\text{g/l}$ (median; $194\text{ }\mu\text{g/l}$) in Ha Tay Province. There is no significant difference between As concentrations in groundwater of Ha Nam and Ha Tay ($p > 0.05$). Interestingly, As concentrations in most of the groundwater samples (22/23) were over the WHO guideline level ($10\text{ }\mu\text{g/l}$) for drinking (Fig. 1)(WHO, 2004), suggesting possible human health risk in both provinces and that As pollution may be widely distributed in the Red River Delta.

Concentrations of As in groundwater have been reported in other As-contaminated areas in Vietnam (Berg et al., 2001; Agusa et al., 2004; Iwata et al., 2004; Agusa et al., 2006). In the Red River Delta, North Vietnam, high concentrations of As were observed in groundwater of Gia Lam District (up to $3,050\text{ }\mu\text{g/l}$, mean; $127\text{ }\mu\text{g/l}$) and Thanh Tri District (up to $3,010\text{ }\mu\text{g/l}$, mean; $432\text{ }\mu\text{g/l}$) in Hanoi (Berg et al., 2001). Our preliminary study found high concentrations of As (up to $411\text{ }\mu\text{g/l}$) in groundwater of the Mekong Delta, South Vietnam (Iwata et al., 2004). Median levels of As in groundwater in the present study were comparable to mean values in the previous study by Berg et al.




Fig. 1. Concentrations of As in groundwater in Ha Nam and Ha Tay Provinces in Red River Delta, Vietnam. Sample numbers (HN01-10 and HT01-13) indicate the well in each house. The vertical bold line denote the guideline value (10 µg/l) established by WHO (WHO, 2004).

(2001).

To assess the As exposure, concentrations of As compounds in urine of residents in Ha Nam and Ha Tay were quantified. Although high abundance of AB, which may be derived from consumption of seafood, was observed in urine of some donors, DMA was the most predominant compound in almost all the residents. MMA, As[III] and As[V] were also observed in the urine samples. A significant positive correlation between concentrations of As in groundwater and in urine of residents was observed ($p < 0.01$). DMA ($p < 0.01$), As[III] ($p < 0.01$) and As[V] ($p < 0.001$) concentrations in human urine were also positively correlated with As concentrations in groundwater (Fig. 2). On the other hand, no significant correlation was observed between concentrations of As in groundwater and AB in urine ($p > 0.05$) (Fig. 2). These results suggest that residents in Ha Nam and Ha Tay are exposed to As through drinking groundwater and hence human health risks by As exposure




Fig. 2. Relationships between As concentrations in groundwater and concentrations of dimethylarsinic acid (DMA), arsenite (As[III]), arsenate (As[V]) and arsenobetaine (AB) in urine of residents in Ha Nam and Ha Tay Provinces in Red River Delta, Vietnam.

are great concern. Further investigation is needed to evaluate the toxic effects by As to the residents in these areas.

Acknowledgments

The authors wish to thank the staff of the Research Center for Environmental Technology and Sustainable Development, Hanoi University of Science, Hanoi. This study was supported (in part) by Japan Society for the Promotion of Science (JSPS) for the cooperative research program under the Core University Program between JSPS and Vietnamese Academy of Science and Technology, and grants from Research Revolution 2002 (RR2002) Project for Sustain-able Coexistence of Human, Nature and the Earth (FY2002) and “21st Century COE Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- Agusa, T., Kunito, T., Fujihara, J., Kubota, R., Minh, T.B., Trang, P.T.K., Subramanian, A., Iwata, H., Viet, P.H. and Tanabe, S. (2004) Contamination by trace elements in groundwater of Vietnam. *Biomedical Research on Trace Elements*, 15, 339-341.
- Agusa, T., Kunito, T., Fujihara, J., Kubota, R., Minh, T.B., Trang, P.T.K., Iwata, H., Subramanian, A., Viet, P.H. and Tanabe, S. (2006) Contamination by arsenic and other trace elements in tube-well water and its risk assessment to humans in Hanoi, Vietnam. *Environmental Pollution*, 139, 95-106.
- Berg, M., Tran, H.C., Nguyen, T.C., Pham, H.V., Schertenleib, R. and Giger, W. (2001) Arsenic contamination of groundwater and drinking water in Vietnam: a human health threat. *Environmental Science and Technology*, 35, 2621-2626.
- Iwata, H., Agusa, T., Inoue, S., Kubota, R., Minh, N.H., Minh, T.B., Tu, N.P.C., Kajiwara, N., Kunisue, T., Subramanian, A., Tanabe, S., Viet, P.H. and Tuyen, B.C. (2004) Contamination of trace elements in groundwater and persistent organochlorines in sediment from Mekong Delta, South Vietnam. *Proceeding on International Symposium on the Development of Water Resource Management System in Mekong Watershed, Hanoi, Vietnam*, 25-31.
- Mandal, B.K., Ogra, Y. and Suzuki, K.T. (2001) Identification of dimethylarsinous and monomethylarsonous acids in human urine of the arsenic-affected areas in West Bengal, India. *Chemical Research in Toxicology*, 14, 371-378.
- Nordstrom, D.K. (2002) Public health. Worldwide occurrences of arsenic in ground water. *Science*, 296, 2143-2145.
- WHO (2004) *Guidelines for Drinking Water Quality*. 3rd edition, World Health Organization, Geneva, Switzerland.