



Title	Quantized enveloping algebras associated to simple Lie su-peralgebras and universal R-matrices
Author(s)	山根, 宏之
Citation	大阪大学, 1992, 博士論文
Version Type	VoR
URL	https://doi.org/10.11501/3060137
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

博士論文 : Quantized enveloping algebras associated to simple Lie
superalgebras and universal R-matrices
(複素単純リー超代数の量子包絡環と普遍R-行列)

山根宏之 (大阪府)

平成4年2月3日 提出

Quantized Enveloping Algebras
Associated To Simple Lie Superalgebras
And Universal R-matrices

Hiroyuki Yamane

Introduction.

0.1. In [2], Drinfeld introduced the notion of *quasi-triangular Hopf-algebras* from the point of view of the quantum inverse scattering method.

Let \mathbb{H} be a (topological) Hopf algebra with coproduct Δ . Let $\mathcal{R} = \sum_i a_i \otimes b_i \in \mathbb{H} \otimes \mathbb{H}$ be an invertible element. We say that $(\mathbb{H}, \Delta, \mathcal{R})$ is a quasi-triangular Hopf algebra if it satisfies the following properties:

$$\bar{\Delta}(x) = \mathcal{R} \cdot \Delta(x) \cdot \mathcal{R}^{-1} \quad (x \in \mathbb{H}),$$

$$(\Delta \otimes 1)(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{23}, \quad (1 \otimes \Delta)(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{12},$$

where $\bar{\Delta} = \tau \circ \Delta$, $\tau(x \otimes y) = y \otimes x$ and $\mathcal{R}_{12} = \sum_i a_i \otimes b_i \otimes 1$

$$\mathcal{R}_{13} = \sum_i a_i \otimes 1 \otimes b_i, \quad \mathcal{R}_{23} = \sum_i 1 \otimes a_i \otimes b_i.$$

The above \mathcal{R} is called the *universal R-matrix* of \mathbb{H} . The importance of \mathcal{R} lies in the fact that it satisfies the

Yang-Baxter equation:

$$\mathcal{R}_{12}\mathcal{R}_{13}\mathcal{R}_{23} = \mathcal{R}_{23}\mathcal{R}_{13}\mathcal{R}_{12} .$$

0.2. Drinfeld [2] (and Jimbo [3]) introduced a family of quasi-triangular Hopf algebras $U_h(G)$ associated to complex simple Lie algebras G . The Hopf algebras $U_h(G)$ are called *quantum groups* or *quantized enveloping algebras*. Moreover Drinfeld [2] gave a method of constructing the universal R-matrix of $U_h(G)$, which is called the *quantum double construction*. Several authors gave explicit formulas for the universal R-matrices of the Hopf algebras $U_h(G)$ by using the quantum double construction. See [6], [8], [13].

In this paper, we introduce a new family of quasi-triangular Hopf algebras $U_h^\sigma = U_h(\mathcal{G})^\sigma$ associated to complex simple Lie superalgebras \mathcal{G} of types A - G. We shall also give explicit formulas for the universal R-matrices \mathcal{R} of the Hopf algebras U_h^σ by using the quantum double construction.

0.3. Let \mathcal{G} be a complex Lie superalgebras of type A - G and $U(\mathcal{G})$ the enveloping superalgebra of \mathcal{G} . Let (Φ, Π, p) be a root system of \mathcal{G} , i.e., $\Phi, \Pi = \{\alpha_1, \dots, \alpha_n\}$ and $p : \Pi \rightarrow \{0, 1\}$ are the set of roots, the set of simple roots and the parity function respectively. In this paper, we assume that (Φ, Π, p) is of distinguished type (see [4]) if \mathcal{G} is of type F_4 or G_3 . For each such (Φ, Π, p) , we introduce an h -adic topologically free $\mathbb{C}[[h]]$ -Hopf superalgebra $U_h = U_h(\mathcal{G}) = U_h(\Pi, p)$ such that

U_h/hU_h is isomorphic to $U(\mathcal{G})$ as an \mathbb{C} -Hopf superalgebra. The Hopf superalgebra structure of $U_h(\Pi, p)$ depends on the choice of (Φ, Π, p) . (Note that two root systems of a simple Lie superalgebra are not necessarily isomorphic.)

0.4. Let $\mathfrak{H} = \mathfrak{H}_0 \oplus \mathfrak{H}_1$ be any Hopf superalgebra. Let $\sigma : \mathfrak{H} \rightarrow \mathfrak{H}$ be an involution defined by $\sigma(x) = (-1)^i x$ for $x \in \mathfrak{H}_i$. In the end of §1, we remark that $\mathfrak{H}^\sigma = \mathfrak{H} \rtimes \langle \sigma \rangle (\simeq \mathfrak{H} \oplus \mathfrak{H}\sigma)$ has a Hopf algebra structure.

In this paper, we give an explicit formula for the universal R-matrix \mathcal{R} of U_h^σ by using the quantum double construction. In particular, we can show that U_h^σ is quasi-triangular. As in "non-super" cases, our \mathcal{R} is described in terms of q-root vectors and the q-exponential.

0.5. Here we explain the content of this paper in more detail. In §1, we explain the quantum double construction applied to h-adic topological $\mathbb{C}[[h]]$ -Hopf algebras. In §2, for any pair (Π, p) of a set of simple roots Π and a parity function p of any symmetrizable Kac-Moody type Lie superalgebra, we define an h-adic topologically free $\mathbb{C}[[h]]$ -Hopf superalgebra $U_h(\Pi, p)$. We also show that, if $p(\alpha) = 0$ for any $\alpha \in \Pi$, then $U_h(\Pi, p)$ coincides with Drinfeld - Jimbo quantized enveloping algebra $U_h(\mathcal{G})$ defined for the Kac-Moody Lie algebra \mathcal{G} with simple roots Π .

In §3-10, for the pair (Π, p) satisfying the assumption in 0.3, we give the defining relations of $U_h(\Pi, p)$ and show $U_h(\Pi, p)/hU_h(\Pi, p) = U(\mathcal{G})$. We remark that the defining relations are not just q-analogue of Serre relations; additional relations

are also needed. In the end of §10, we give an explicit formula for the universal R-matrix \mathcal{R} of $U_h(\pi, p)^\sigma$.

In doing these, our basic references are Lusztig's papers [9] and [10]. By imitating the definition of q-root vectors of $U_h(G)$ in [9] and [10], we introduce q-root vectors of $U_h(\pi, p)$.

The main result of this paper has been announced in [17].

§1. Quantum double construction

1.1. Let $R = \mathbb{C}[[h]]$ be the \mathbb{C} -algebra of formal power series. We explain briefly elementary facts concerning the h -adic topological R -modules. For details, see [12].

Let V be an R -module. Let $v_V = v : V \rightarrow \mathbb{Z}_+ \cup \{+\infty\}$ be the h -valuation defined as follows; if $v \in h^i V \setminus h^{i+1} V$, put $v(v) = i$, and, if $v \in \bigcap_{i \in \mathbb{Z}_+} h^i V$, put $v(v) = +\infty$. We can regard V as a topological space such that a fundamental system of neighborhoods of $v \in V$ is given by $v + h^i V$ ($i \in \mathbb{Z}_+$). This topology is called the h -adic topology. For $v, w \in V$, we put

$$d_V(v, w) = 2^{-v(v-w)}.$$

Then $d_V(,)$ is a quasi-metric for the topological space V . For a subset S of V , the symbol \bar{S} denotes the closure of S . Note that any R -module homomorphism is continuous with

respect to the h-adic topology.

If any Cauchy sequence (with respect to $d_V(\cdot, \cdot)$) has a limit, then V is called *complete*. If $\{0\} = \overline{\{0\}}$, then V is called *separated*. If V is separated, then $d_V(\cdot, \cdot)$ is a metric on V . Note that, for a submodule W , the quotient topology of V/W coincides with the h-adic topology of it. If V is complete, then V/W is also complete. If V is separated and W is closed, then V/W is separated.

It is well known that, for any h-adic topological R -module V , there exists a pair (\hat{V}, i) of an h-adic topological R -module \hat{V} and an R -homomorphism $i : V \rightarrow \hat{V}$ satisfying: For any h-adic topological complete separated R -module W , and, for any R -homomorphism $\varphi : V \rightarrow W$, there exists uniquely an R -homomorphism $\hat{\varphi} : \hat{V} \rightarrow W$ such that $\varphi \circ i = \hat{\varphi}$. We note that \hat{V} is complete and that $i(V)$ is dense in \hat{V} . It is also well known that, if V is separated, then i is injective and the induced topology of V ($\subset \hat{V}$) coincides with the h-adic topology of V .

If a complete separated h-adic R -module V has a submodule W such that W is a free R -module and V is the completion of W , then we say that V is *topologically free*. A basis of W is called a *topological basis* of V .

Example 1.1.1. Let V_0 be a \mathbb{C} -vector space and $V = R \otimes V_0$. Let \hat{V} be the completion of V . Then we have the following natural identifications:

$$\hat{V} = \left\{ \sum_{i=0}^{\infty} h^i a_i \mid a_i \in V_0 \right\},$$

$$V = \left\{ \sum_{i=0}^{\infty} h^i a_i \mid a_i \in V_0 \quad \dim(\sum_{i=0}^{\infty} \mathbb{C}a_i) < \infty \right\}$$

where $\sum_{i=0}^{\infty} h^i a_i$ is a formal infinite sum.

Definition 1.1.2. We say that an R -module V has a *handy basis* $\{v_i\}_{i \in I}$ if (i) V is a topologically free R -module with a topological basis $\{v_i\}_{i \in I}$, (ii) I is a partially ordered set and (iii) there is an order homomorphism $p : I \rightarrow \mathbb{Z}_+$ such that, for each $n \in \mathbb{Z}_+$, $p^{-1}(n)$ is a finite set.

Example 1.1.3. Retainig the notation in Definition 1.1.2. We have natural identifications:

$$\hat{V} = \left\{ \sum_{i=0}^{\infty} \alpha_i v_i \mid \alpha_i \in R, \lim_{i \rightarrow \infty} v(\alpha_i) = +\infty \right\},$$

$$V = \left\{ \sum_{i=0}^{\infty} \alpha_i v_i \mid \alpha_i \in R, v(\alpha_i) \neq 0 \text{ for finitely many } i's \right\}$$

where $\sum_{i=0}^{\infty} \alpha_i v_i$ is a formal infinite sum.

For R -modules V and W , we denote the completion of

$V \otimes W$ by $\hat{V \otimes W}$. If V and W have handy bases $\{v_i\}_{i \in I}$ and $\{w_j\}_{j \in J}$ respectively, then $\{v_i \otimes w_j\}_{(i,j) \in I \times J}$ is also a handy basis of $\hat{V \otimes W}$. In particular, $V \otimes W$ and $\hat{V \otimes W}$ are separated.

1.2. Let $A = (A, m, \eta, \Delta, S, \varepsilon)$ be an h -adic topological R -Hopf algebra. Namely, the h -adic topological R -module A has a topological R -Hopf algebra structure with the product $m : A \hat{\otimes} A \rightarrow A$, the unit $\eta : R \rightarrow A$, the coproduct $\Delta : A \rightarrow A \hat{\otimes} A$, the antipode $S : A \rightarrow A$ and the counit $\varepsilon : A \rightarrow R$. Here the definition of the h -adic topological Hopf algebras is given by replacing $A \otimes A$, $A \otimes A \otimes A$ in the definition of the Hopf algebras by their completions $A \hat{\otimes} A$, $A \hat{\otimes} A \hat{\otimes} A$. For the definition of the ordinary Hopf algebras, see [1].

Define $\tau : A \hat{\otimes} A \rightarrow A \hat{\otimes} A$ by $\tau(a \otimes b) = b \otimes a$. It is well known that $A^{op} = (A, m, \eta, \tau \circ \Delta, S^{-1}, \varepsilon)$ is also an h -adic topological R -Hopf algebra. We call A^{op} the *opposite Hopf algebra* of A .

Let \mathcal{J} be an ideal of the R -algebra A . We say that \mathcal{J} is a bi-ideal if \mathcal{J} satisfies: $\Delta(\mathcal{J}) \subseteq \mathcal{J} \otimes A + A \otimes \mathcal{J}$ and $\varepsilon(\mathcal{J}) = 0$. Moreover, if \mathcal{J} satisfies $S(\mathcal{J}) = \mathcal{J}$, we say that \mathcal{J} is a Hopf ideal.

1.3. Let $A = (A, m, \eta, \Delta, S, \varepsilon)$ be an h -adic topological R -Hopf

algebra. In this subsection, we assume that A has a handy basis $\{a_i\}_{i \in I}$. Let $A^* = \text{Hom}_R(A, R)$ be the dual space of the R -module A . Define $a_i^* \in A^*$ ($i \in I$) by $a_i^*(a_j) = \delta_{ij}$. Then we have a natural identification:

$$A^* = \left\{ \sum_{i \in I} \alpha_i a_i^* \mid \alpha_i \in R \right\}$$

where $\sum_{i=0}^{\infty} \alpha_i a_i^*$ is a formal infinite sum. Then A^* is a torsion free complete separated R -module.

The R -module A^* has a two-sided A -module structure defined by:

$$a.f.\delta(c) = f(\delta ca) \quad (f \in A^*, a, \delta, c \in A).$$

Put

$$A^O = \{f \in A^* \mid A.f.A \text{ is a finitely generated free } R\text{-module}\}.$$

Similarly to the case of ordinary Hopf algebras (see [1]), it can be shown easily that

$$\begin{aligned} A^O &= \{f \in A^* \mid A.f \text{ is a finitely generated free } R\text{-module}\} \\ &= \{f \in A^* \mid f.A \text{ is a finitely generated free } R\text{-module}\}, \end{aligned}$$

and that $A^O = (A^O, t_\Delta, t_\varepsilon, t_m, t_S, t_\eta)$ is a non-topological R -Hopf

algebra where t denotes the transpose. We call A^O the *dual Hopf algebra* of A . Let \hat{A}^O be the completion of A^O . It is obvious that $(\hat{A}^O, t_\Delta, t_\varepsilon, t_m, t_S, t_\eta)$ is an h -adic R -Hopf algebra. We note that \hat{A}^O (resp. $\hat{A}^O \otimes \hat{A}^O$, $\hat{A}^O \otimes \hat{A}^O \otimes \hat{A}^O$) is naturally identified with the closure $\overline{\hat{A}^O}$ of \hat{A}^O (resp.

$\overline{A^O \otimes A^O}$ of $A^O \otimes A^O$, $\overline{A^O \otimes A^O \otimes A^O}$ of $A^O \otimes A^O \otimes A^O$) in A^* (resp. $(\hat{A} \otimes \hat{A})^*$, $(\hat{A} \otimes \hat{A} \otimes \hat{A})^*$). Hence we shall denote $(\hat{A}^O, t_\Delta, t_\varepsilon, t_m, t_S, t_\eta)$ by $\overline{\hat{A}^O}$.

1.4. Let $A = (A, m, \eta, \Delta, S, \varepsilon)$ be an h -adic R -Hopf algebra. For $i \geq 2$, $\Delta^{(i)}$ and $m^{(i)}$ denote $(\Delta^{(i-1)} \otimes \text{id}) \circ \Delta$ and $m \circ (m^{(i-1)} \otimes \text{id})$ respectively.

Let $A = (A, m_A, \eta_A, \Delta_A, S_A, \varepsilon_A)$ and $B = (B, m_B, \eta_B, \Delta_B, S_B, \varepsilon_B)$ be h -adic topological R -Hopf algebras. Let $\langle , \rangle : A \otimes B \rightarrow R$ be an R -bilinear form. We say that \langle , \rangle is a Hopf pairing if \langle , \rangle satisfies:

- (i) $\langle a_1 a_2, b \rangle = \langle a_1 \otimes a_2, \Delta_B(b) \rangle$,
 $\langle a, b_1 b_2 \rangle = \langle \Delta_A(a), b_1 \otimes b_2 \rangle$,
- (ii) $\langle a, S_B(b) \rangle = \langle S_A(a), b \rangle$,
- (iii) $\langle \eta_A(1), b \rangle = \varepsilon_B(b)$, $\langle a, \eta_B(1) \rangle = \varepsilon_A(a)$,

where $a, a_i \in A$ and $b, b_i \in B$.

Define an R -module homomorphism $\Phi : B \hat{\otimes} A \rightarrow A \hat{\otimes} B$ by

$$\Phi(\delta \otimes a) = \sum_{i,j} \langle a_i^{(3)}, \delta_j^{(1)} \rangle \langle s_A^{-1}(a_i^{(1)}), \delta_j^{(3)} \rangle \cdot a_i^{(2)} \otimes \delta_j^{(2)}$$

$$\text{where } \Delta_A^{(2)}(a) = \sum_i a_i^{(1)} \otimes a_i^{(2)} \otimes a_i^{(3)} \text{ and}$$

$$\Delta_B^{(2)}(\delta) = \sum_j \delta_j^{(1)} \otimes \delta_j^{(2)} \otimes \delta_j^{(3)}.$$

In the following lemma, we define an h-adic topological R -Hopf algebra $D(A, B^{op})$ which is called the *quantum double* of A and B . The notion of quantum double was introduced by Drinfeld [2].

Proposition 1.4.1. Let A and B be h-adic R -Hopf algebras with a (possibly degenerate) Hopf pairing $\langle , \rangle : A \hat{\otimes} B \rightarrow R$. Then there exists uniquely an h-adic topological R -Hopf algebra $D(A, B^{op}) = (D(A, B^{op}), m_D, \eta_D, \Delta_D, S_D, \varepsilon_D)$ satisfying:

- (i) As an h-adic topological R -module, $D(A, B^{op})$ is isomorphic to $A \hat{\otimes} B$,
- (ii) The R -module maps $A \rightarrow D(A, B^{op})$ ($a \mapsto a \otimes 1$), $B^{op} \rightarrow D(A, B^{op})$ ($\delta \mapsto 1 \otimes \delta$) are h-adic topological R -Hopf algebra homomorphisms,
- (iii) The multiplication m_D is given by $m_D = (m_A \otimes m_B) \circ (\text{id}_A \otimes \Phi \otimes \text{id}_B)$.

Proof. Here we prove the associativity of the multiplication m_D of $D(A, B^{op})$ only.

Let $a \otimes b, c \otimes d, e \otimes f \in D(A, B^{op})$. Put

$$\begin{aligned}\Delta_B^{(2)}(b) &= \sum_u b_u^{(1)} \otimes b_u^{(2)} \otimes b_u^{(3)}, \\ \Delta_A^{(2)}(c) &= \sum_v c_v^{(1)} \otimes c_v^{(2)} \otimes c_v^{(3)}.\end{aligned}$$

By (iii), we have:

$$\begin{aligned}&(a \otimes b) \cdot (c \otimes d) \\ &= \sum_{u,v} \langle s_A^{-1}(c_v^{(1)}), b_u^{(3)} \rangle \langle c_v^{(3)}, b_u^{(1)} \rangle a c_v^{(2)} \otimes b_u^{(2)} d.\end{aligned}$$

Put

$$\begin{aligned}\Delta_B^{(2)}(b) &= \sum_\xi b_\xi^{(1)} \otimes b_\xi^{(2)} \otimes b_\xi^{(3)} \otimes b_\xi^{(4)} \otimes b_\xi^{(5)}, \\ \Delta_B^{(2)}(d) &= \sum_w d_w^{(1)} \otimes d_w^{(2)} \otimes d_w^{(3)}, \\ \Delta_A^{(2)}(e) &= \sum_x e_x^{(1)} \otimes e_x^{(2)} \otimes e_x^{(3)}.\end{aligned}$$

By (iii), we have:

$$\begin{aligned}&((a \otimes b) \cdot (c \otimes d)) \cdot (e \otimes f) \\ (1.4.2) \quad &= \sum_{v,w,x,\xi} \langle s_A^{-1}(c_v^{(1)}), b_\xi^{(5)} \rangle \langle c_v^{(3)}, b_\xi^{(1)} \rangle \\ &\quad \langle s_A^{-1}(e_x^{(1)}), b_\xi^{(4)} d_w^{(3)} \rangle \langle e_x^{(3)}, b_\xi^{(2)} d_w^{(1)} \rangle a c_v^{(2)} e_x^{(2)} \otimes b_\xi^{(3)} d_w^{(2)} f\end{aligned}$$

Putting

$$\Delta_A^{(4)}(e) = \sum_\xi e_\xi^{(1)} \otimes e_\xi^{(2)} \otimes e_\xi^{(3)} \otimes e_\xi^{(4)} \otimes e_\xi^{(5)},$$

we have that (1.4.2) is equal to

$$(1.4.3) \quad \sum_{v,w,x,\xi,\zeta} \langle s_A^{-1}(c_v^{(1)}), b_\xi^{(5)} \rangle \langle c_v^{(3)}, b_\xi^{(1)} \rangle \cdot \\ \langle s_A^{-1}(e_\xi^{(1)}), d_w^{(3)} \rangle \langle s_A^{-1}(e_\xi^{(2)}), b_\xi^{(4)} \rangle \cdot \langle e_x^{(4)}, b_\xi^{(2)} \rangle \langle e_x^{(5)}, d_w^{(1)} \rangle \\ ac_v^{(2)} e_\xi^{(3)} \otimes b_\xi^{(3)} d_w^{(2)} f .$$

Similarly, we can show that $(a \otimes c) \cdot ((b \otimes c) \cdot (d \otimes f))$ is equal to (1.4.3). Hence, we have:

$$((a \otimes c) \cdot (b \otimes c)) \cdot (d \otimes f) = (a \otimes c) \cdot ((b \otimes c) \cdot (d \otimes f)) .$$

1.5. Let V be a complete and separated R -module. Here, we define the notion the convergence of a multi-series in V .

Let $\{a_{i_1 \dots i_u}\}_{i_1, \dots, i_u \in \mathbb{Z}_+}$ be a multi-sequence in V . If there exists an element $\alpha \in V$ such that, for any $M \in \mathbb{Z}_+$, there exists $N \in \mathbb{Z}_+$ satisfying that $v(\alpha - a_{i_1 \dots i_u}) > M$ for all $i_1, \dots, i_u > N$, then we say that $\{a_{i_1 \dots i_u}\}$ converge to α as a multi-sequence. The element α is denoted by

$\lim_{i_1 \dots i_u} \alpha_{i_1 \dots i_u}$. The uniqueness of α follows from the separatedness of V . The following lemma is obvious.

Lemma 1.5.1. Let V be a complete separated R -module. Let $\{b_{i_1 \dots i_u} \mid (i_1, \dots, i_u \in \mathbb{Z}_+)\}$ be a subset of V . Assume that, for any $M \in \mathbb{Z}_+$, there exists $N \in \mathbb{Z}_+$ such that

$v(b_{i_1 \dots i_u}) > M$ if $i_1 > N$ or ... or $i_u > N$. Then there exists the limit $\beta = \lim_{i_1 \dots i_u} \sum b_{i_1 \dots i_u}$. Moreover, for any permutation ρ of $\{1, 2, \dots, u\}$, it holds that

$$\lim_{i_{\rho(1)}} (\lim_{i_{\rho(2)}} \dots (\lim_{i_{\rho(u)}} \sum b_{i_1 \dots i_u}) \dots) = \beta.$$

1.6. Let V be a complete separated h -adic topological R -module with a handy basis $\{v_i\}_{i \in I}$. Let

$K = \mathbb{C}((h)) (= \{\sum_{i=n}^{\infty} a_i h^i \mid n \in \mathbb{Z}\})$ be the fraction field of

$R = \mathbb{C}[[h]]$. The h -valuation $v_K : K \rightarrow \mathbb{R}_+$ is defined by putting $v_K(\sum a_i h^i) = \min\{i \mid a_i \neq 0\}$. Let $V^K = K \otimes V$ be the scalar extention. Then we have the following identification:

$$V^K = \{ \sum_{i \in I} \alpha_i v_i \mid \alpha_i \in K, \lim_{i \rightarrow +\infty} v(\alpha_i) = +\infty \}.$$

The h -valuation v_{V^K} on V^K is given by putting

$$v_{V^K}(\sum \alpha_i v_i) = \min\{v_K(\alpha_i) \mid i \in I\}.$$

Let A and B be complete separated h -adic topological R -Hopf algebras with a non-degenerate Hopf pairing $\langle , \rangle : A \hat{\otimes} B \rightarrow R$. We assume that A (resp. B) has a handy basis $\{a_i\}_{i \in I}$ (resp. $\{\ell_i\}_{i \in I}$). Moreover we assume that $\langle a_i, \ell_j \rangle = \delta_{ij} c_i$ for some $c_i \in R \setminus \{0\}$. Let $D = D(A, B^{op})$ be

the quantum double.

Let A^K, B^K, D^K be h-adic topological K -Hopf algebras which are obtained by the scalar extenstions of R -modules A, B, D respectively. Then $D^K \cong A^K \hat{\otimes} B^K$ as an h-adic topological K -vector spaces. Moreover we consider A^K (resp. $(B^K)^{op}$) as a topological K -Hopf subalgebra of D^K by the embedding $A^K \rightarrow D^K$ ($a \rightarrow a \otimes 1$) (resp. $(B^K)^{op} \rightarrow D^K$ ($b \rightarrow 1 \otimes b$)). We denote the Hopf pairing $K \otimes \langle , \rangle : A^K \hat{\otimes} B^K \rightarrow R$ simply by \langle , \rangle .

Let $\{e_i\}_{i \in I}$ and $\{e^i\}_{i \in I}$ be the subsets of A^K and B^K respectively such that $e_i \in K e_i$, $e^i \in K e_i$ and $\langle e_i, e^j \rangle = \delta_{ij}$. Let us define $m_j^{i_1 \dots i_u}, \mu^i_{j_1 \dots j_v}, \gamma^i_j \in K$ by

$$\begin{aligned} m_{A^K}^{(u-1)}(e_{i_1} \otimes \dots \otimes e_{i_u}) &= \sum m_j^{i_1 \dots i_u} e_j, \\ \Delta_{A^K}^{(v-1)}(e_j) &= \sum \mu^i_{j_1 \dots j_v} e_{j_1} \otimes \dots \otimes e_{j_v}, \\ s_{A^K}^{-1}(e_i) &= \sum \gamma^i_j e_j. \end{aligned}$$

Using Hopf pairings \langle , \rangle , we have:

$$\begin{aligned} \Delta_{B^K}^{(u-1)}(e^j) &= \sum m_j^{i_1 \dots i_u} e_{i_1} \otimes \dots \otimes e_{i_u}, \\ m_{B^K}^{(v-1)}(e^{j_1} \otimes \dots \otimes e^{j_v}) &= \sum \mu^i_{j_1 \dots j_v} e^i, \\ s_{B^K}^{-1}(e^i) &= \sum \gamma^i_j e^j. \end{aligned}$$

From Proposition 1.4.1, we obtain the following lemma. We

omit the proof.

Lemma 1.6.1. In D^K , the following equations hold.

$$(i) \quad e^t e_s = \sum \mu_{njk}^s m_{t}^{k\ell p} \gamma_p^n e_j e^\ell .$$

$$(ii) \quad e_s e^t = \sum \mu_{kjn}^s m_{t}^{p\ell k} \gamma_p^n e^\ell e_j .$$

1.7. In [2], Drinfeld introduced the following construction of a universal R-matrix, which are called the *quantum double construction*.

Proposition 1.7.1, (*the quantum double construction*).

Retaining the notation in 1.6. Let C be a complete separated h -adic topological R -algebras with 1 . Let $\Omega : D \rightarrow C$ be an R -algebra homomorphism. Denote the scalar extention

$\text{id}_K \otimes \Omega : D^K \rightarrow C^K$ by again Ω . Assume that

$\mathcal{R} = \sum_{i \in I} \Omega(e_i) \otimes \Omega(e^i)$ converges in $C^K \hat{\otimes} C^K$. Then \mathcal{R}

satisfies:

(i) The element \mathcal{R} is invertible. The inverse \mathcal{R}^{-1} is given by $\mathcal{R}^{-1} = \sum_{i \in I} \Omega(S(e_i)) \otimes \Omega(e^i)$,

(ii) $\mathcal{R}(\Omega \otimes \Omega(\Delta(x)))\mathcal{R}^{-1} = \Omega \otimes \Omega(\tau \circ \Delta(x))$ for all $x \in D^K$,

(iii) $\sum_{i \in I} (\Omega \otimes \Omega \otimes \Omega)((\Delta \otimes \text{id})(e_i \otimes e^i)) = \mathcal{R}_{13}\mathcal{R}_{23}$,

$\sum_{i \in I} (\Omega \otimes \Omega \otimes \Omega)((\text{id} \otimes \Delta)(e_i \otimes e^i)) = \mathcal{R}_{13}\mathcal{R}_{12}$.

Proof. Note that any multi-series below satisfies the assumption in Lemma 1.5.1. In the proof, we simply write $a \otimes b$ for $(\Omega \otimes \Omega)(a \otimes b)$. For all $i \in I$, we have:

$$\begin{aligned}\mathcal{R} \cdot \Delta(e_i) &= \sum \mu_{rs}^i e_r \otimes e_s \\ &= \sum \mu_{rs}^i m_x^r \mu_{njk}^s m_t^{kl} p_{\gamma_p^n} e_x \otimes e_j e^l = \sum \mu_{rnjk}^i m_x^k p_{\gamma_p^n} e_x \otimes e_j e^l \\ &= \sum \mu_{jkl}^i m_x^{kl} e_x \otimes e_j e^l = \sum \mu_{jkl}^i e_k e_l \otimes e_j e^l = (\tau \circ \Delta(e_i)) \cdot \mathcal{R}\end{aligned}$$

Similarly, we have $\mathcal{R} \cdot \Delta(e^i) = \tau \circ \Delta(e^i) \cdot \mathcal{R}$ for all $i \in I$. Since $\{a_i \cdot e_j = a_i \otimes e_j \in R e_i \cdot e^j\}_{(i,j) \in I \times I}$ is a topological basis of D^K , we obtain (i). The equations (ii), (iii) can be proved similarly.

1.8. Here, we comment on the definition of an h-adic topological algebra with generators and relations.

Let V be a free R -module with a basis $\{x_\ell\}_{\ell \in L}$. Let $\mathcal{F} = R\langle x_\ell | \ell \in L \rangle$ be the tensor algebra $T(V)$ of V and let $\hat{\mathcal{F}}$ be the completion of \mathcal{F} . Let P_λ ($\lambda \in \Lambda$) be the elements of $\hat{\mathcal{F}}$. Put $\mathcal{G} = \hat{\mathcal{F}} / (\overline{\sum_\lambda \mathcal{F} \cdot P_\lambda \cdot \mathcal{F}})$. We say that the h-adic topological R -algebra \mathcal{G} is h-adically generated by x_ℓ ($\ell \in L$) with the relations $\{P_\lambda | \lambda \in \Lambda\}$.

1.9. Let $\mathfrak{H} = \mathfrak{H}_0 \oplus \mathfrak{H}_1$ be a (topological) R -superalgebra. For $i \in \{0,1\}$, define $p_i : \mathfrak{H} \rightarrow \mathfrak{H}_i$ by $p_i(x_0 + x_1) = x_i$ where $x_k \in \mathfrak{H}_k$ ($k \in \{0,1\}$). Let $\langle \sigma \rangle$ be the cyclic group of order two with a generator σ . Let $R\langle \sigma \rangle$ be the group ring of $\langle \sigma \rangle$ over R . We define an R -algebra structure on an R -module $\mathfrak{H}^\sigma = \mathfrak{H} \otimes_R R\langle \sigma \rangle$ by

$$(x \otimes \sigma^c)(y \otimes \sigma^d) = x(p_0(y) + (-1)^c p_1(y)) \otimes \sigma^{c+d}.$$

We write $x\sigma^c$ for $x \otimes \sigma^c$. Define $r_\sigma : \mathfrak{H} \rightarrow \mathfrak{H}^\sigma$ (resp. $\ell_\sigma : \mathfrak{H} \rightarrow \mathfrak{H}^\sigma$) by $r_\sigma(x) = x\sigma$ (resp. $\ell_\sigma(x) = x\sigma$) ($x \in \mathfrak{H}$). From the axiom of Hopf superalgebras (see [14]), we can easily show:

Proposition 1.9.1. Let $(\mathfrak{H} = \mathfrak{H}_0 \oplus \mathfrak{H}_1, \dot{\Delta}, \dot{\varepsilon}, \dot{S})$ be a (topological) R -Hopf superalgebra. Then the R -algebra \mathfrak{H}^σ has a (topological) R -Hopf algebraic structure $(\mathfrak{H}^\sigma, \Delta, \varepsilon, S)$ such that

- (i) The coproduct $\Delta : \mathfrak{H}^\sigma \rightarrow \mathfrak{H}^\sigma \otimes \mathfrak{H}^\sigma$ is defined by $\Delta(x) = ((\text{id} \otimes p_0 + r_\sigma \otimes p_1) \circ \dot{\Delta})(x)$ ($x \in \mathfrak{H}$) and $\Delta(\sigma) = \sigma \otimes \sigma$,
- (ii) The counit $\varepsilon : \mathfrak{H}^\sigma \rightarrow R$ is defined by $\varepsilon(x) = \dot{\varepsilon}(x)$ ($x \in \mathfrak{H}$) and $\varepsilon(\sigma) = 1$,
- (iii) The antipode $S : \mathfrak{H}^\sigma \rightarrow \mathfrak{H}^\sigma$ is defined by $S(x) = ((p_0 + \ell_\sigma \circ p_1) \circ \dot{S})(x)$ ($x \in \mathfrak{H}$) and $S(\sigma) = \sigma$.

Conversely, we can also show:

Proposition 1.9.2. Let \mathfrak{H} be a (topological) R -superalgebra. Assume that \mathfrak{H}^σ has a (topological) R -Hopf algebra structure $(\mathfrak{H}^\sigma, \Delta, \varepsilon, S)$ satisfying:

- (i) $\Delta(\mathfrak{H}_0) \subset \mathfrak{H}_0 \otimes \mathfrak{H}_0 + \mathfrak{H}_1 \sigma \otimes \mathfrak{H}_1$,
 $\Delta(\mathfrak{H}_1) \subset \mathfrak{H}_1 \otimes \mathfrak{H}_0 + \mathfrak{H}_0 \sigma \otimes \mathfrak{H}_1$ and $\Delta(\sigma) = \sigma \otimes \sigma$,
- (ii) $\varepsilon(\mathfrak{H}_1) = \{0\}$ and $\varepsilon(\sigma) = 1$,
- (iii) $S(\mathfrak{H}_0) \subset \mathfrak{H}_0$, $S(\mathfrak{H}_1) \subset \sigma \mathfrak{H}_1$ and $S(\sigma) = \sigma$.

Then there uniquely exists a (topological) R -Hopf superalgebra structure $(\mathfrak{H}, \dot{\Delta}, \dot{\varepsilon}, \dot{S})$ such that $(\mathfrak{H}^\sigma, \Delta, \varepsilon, S)$ coincides with the Hopf algebra defined in Proposition 1.9.1 for $(\mathfrak{H}, \dot{\Delta}, \dot{\varepsilon}, \dot{S})$.

§2. Quantized enveloping (super) algebras

Notation. In §2-§10, the following notation will be used:

- (\mathfrak{E}, Π, p) := a triple of an N -dimensional \mathbb{C} -vector space \mathfrak{E} with a non-degenerate symmetric bilinear form
- $(\cdot, \cdot) : \mathfrak{E} \times \mathfrak{E} \rightarrow \mathbb{C}$, a linearly independent finite subset
- $\Pi = \{\alpha_1, \dots, \alpha_n\}$ of \mathfrak{E} and a function $p : \Pi \rightarrow \{0, 1\}$ (see 2.1)
- $P_+ := \mathbb{Z}_+ \alpha_1 \oplus \mathbb{Z}_+ \alpha_2 \oplus \dots \oplus \mathbb{Z}_+ \alpha_n (\subset \mathfrak{E})$ (see 2.3)
- $D = \text{diag}(d_1, \dots, d_n) :=$ a diagonal matrix of degree n whose matrix elements are half integers (see 2.1)
- $\mathcal{K} := \mathfrak{E}^*$ (see 2.1)
- H_λ ($\lambda \in \mathfrak{E}$) := the element of \mathcal{K} defined by $\mu(H_\lambda) = (\mu, \lambda)$ ($\mu \in \mathfrak{E}$) (see 2.1)
- $R := \mathbb{C}[[h]]$, the ring of formal power series in an indeterminate h
- $q := e^h \in R$ (see 2.9)
- $\tilde{N}_+ :=$ a free R -algebra with generators $\{E_i \mid 1 \leq i \leq n\}$ (see 2.1)
- $\tilde{N}_- :=$ a free R -algebra with generators $\{F_i \mid 1 \leq i \leq n\}$ (see 2.1)
- $S[\mathcal{K}^R] :=$ a symmetric R -algebra generated by $\mathcal{K}^R = \mathcal{K} \otimes R$ (see 2.1)
- $R<\sigma> :=$ a group ring over R of a cyclic group $\langle\sigma\rangle$ of order two (see 2.1)
- $\tilde{U}_h^\sigma = \tilde{U}_h^\sigma((\mathfrak{E}, \Pi, p), D) :=$ an h -adic R -Hopf algebra, which is, as an R -module, isomorphic to $\tilde{N}_+ \otimes S[\mathcal{K}^R] \otimes R<\sigma> \otimes \tilde{N}_-$ (see Lemma 2.1.4)
- $R' := \mathbb{C}[[\sqrt{h}]]$ (see 2.2)

$\tilde{N}'_+ := \tilde{N}_+ \otimes R'$ (see 2.2)

$S[\mathcal{R}^{R'}] := S[\mathcal{R}^R] \otimes R'$ (see 2.2)

$R' \langle \sigma \rangle := R \langle \sigma \rangle \otimes R'$ (see 2.2)

$\tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma = \tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma(\mathcal{E}, \pi, p) :=$ an \sqrt{h} -adic R' -Hopf algebra, which is as an R' -module, isomorphic to $\tilde{N}'_+ \otimes S[\mathcal{R}^{R'}] \otimes R' \langle \sigma \rangle$; We put $E'_i = E_i \otimes 1 \otimes 1$, $\sigma' = 1 \otimes 1 \otimes \sigma$, $H' = 1 \otimes H \otimes 1$ ($H \in \mathcal{H}$); usually we identify E'_i and σ' with E_i and σ (but not H' with H) (see 2.2)

E_i° , H_λ° , $\sigma^\circ :=$ elements of $\overline{(\tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma)^\circ}$ (see (2.3.1-3))

$\langle , \rangle : \tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma \times \tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma \rightarrow R'$:= a Hopf paring (see 2.4)

$\mathfrak{D}' := D(\tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma, (\tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma)^{op})$ the quantum double defined with respect to \langle , \rangle (see 2.5)

Since, as R' -modules, $\mathfrak{D}' \cong \tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma \otimes \tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma$, we write $X \in \mathfrak{D}'$ for $X \otimes 1$ and $X^\circ \in \mathfrak{D}'$ for $1 \otimes X$ where $X \in \tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma$ (see 2.5)

$I'_\mathcal{C}_+ := \text{Ker } \langle , \rangle$ (see 2.6)

$U'_\sqrt{h} \mathcal{C}_+^\sigma := \tilde{U}'_{\sqrt{h}} \mathcal{C}_+^\sigma / I'_\mathcal{C}_+$ (see 2.7)

$I'_+ := \text{Ker } \langle , \rangle|_{\tilde{N}'_+ \times \tilde{N}'_+}$ (see 2.6)

$N'_+ := \tilde{N}'_+ / I'_+$ (see 2.7)

$\mathfrak{D}' := D(U'_\sqrt{h} \mathcal{C}_+^\sigma, (U'_\sqrt{h} \mathcal{C}_+^\sigma)^{op})$ the quantum double defined with respect to \langle , \rangle (see 2.8)

Since, as R' -modules, $\mathfrak{D}' \cong U'_\sqrt{h} \mathcal{C}_+^\sigma \otimes U'_\sqrt{h} \mathcal{C}_+^\sigma$, we write $X \in \mathfrak{D}'$ for $X \otimes 1$ and $X^\circ \in \mathfrak{D}'$ for $1 \otimes X$ where $X \in U'_\sqrt{h} \mathcal{C}_+^\sigma$ (see 2.5)

$N_+ :=$ a unital R -subalgebra of N'_+ such that $N'_+ = N_+ \oplus \sqrt{h}N_+$ (see Lemma 2.9.1)

$I_+ := I'_+ \cap \tilde{N}_+ ; I'_+ = I_+ \oplus \sqrt{h}I_+$, $N_+ = \tilde{N}_+ / I_+$ (see Lemma 2.9.1)

$U_h^\sigma = U_h^\sigma((\mathcal{E}, \Pi, p), D) :=$ a topologically free R -Hopf algebra
 (see Theorem 2.9.4); if Π is a set of simple roots of a
 Kac-Moody Lie algebra G (resp. a simple Lie superalgebra \mathfrak{G}
 in 3.1) and $p(\alpha_i) = 0$ for all $\alpha_i \in \Pi$, then, as a \mathbb{C} -Hopf
 algebra, $U_h^\sigma/hU_h^\sigma \simeq U(G)^\sigma$ (resp. $U_h^\sigma/hU_h^\sigma \simeq U(\mathfrak{G})^\sigma$)
 (see Theorem 2.10.1 (resp. Theorem 10.5.1))

$\mathcal{I}_+ :=$ an ideal of \tilde{N}_+ generated by q-Serre relations and
 additional relations (see Definition 4.2)

$N_+ := \tilde{N}_+/\mathcal{I}_+$ (see 4.3)
 $\mathcal{I}_{\ell_+} :=$ an ideal of $\tilde{U}_{\sqrt{h}}\ell_+^\sigma$ generated by elements of \mathcal{I}_+ (see
 4.3)

$\tilde{U}_{\sqrt{h}}\ell_+^\sigma := \tilde{U}_{\sqrt{h}}\ell_+^\sigma/\mathcal{I}_{\ell_+}$ (see 4.3)

In fact, it will be shown that $\mathcal{I}_+ = I_+$, $\mathcal{I}_{\ell_+} = I_{\ell_+}$,
 $\tilde{U}_{\sqrt{h}}\ell_+^\sigma = U_{\sqrt{h}}\ell_+^\sigma$ (see Proposition 10.4.1)

$N_{+,v} :=$ a weight space of N_+ of a weight $v \in P_+$ (see 4.4)

2.1. In § 2, we construct quantized enveloping algebras
 associated with generalized symmetric Cartan matrices of
 Kac-Moody type Lie superalgebras.

Let \mathcal{E} be an N -dimensional complex linear space with a
 non-degenerate symmetric bilinear form $(,) : \mathcal{E} \times \mathcal{E} \rightarrow \mathbb{C}$. Let
 $\Pi = \{\alpha_1, \dots, \alpha_n\}$ be a finite linearly independent subset of \mathcal{E} .

We call a function $p : \Pi \rightarrow \{0,1\}$ the *parity function*. We call (\mathcal{E}, Π, p) a *triple system*. Let $d_i \in \frac{1}{2}\mathbb{Z} \setminus \{0\}$ ($1 \leq i \leq n$). Define the diagonal matrix D by $\text{diag}(d_1, \dots, d_n)$. Put $\mathcal{K} = \mathcal{E}^*$. For $\lambda \in \mathcal{E}$, let us define $H_\lambda \in \mathcal{K}$ by $\mu(H_\lambda) = (\mu, \lambda)$ for all $\mu \in \mathcal{E}$.

Let $\tilde{U}_h^\sigma = \tilde{U}_h^\sigma((\mathcal{E}, \Pi, p), D)$ be an h -adic topological R -algebra h -adically defined with generators E_i, F_i ($1 \leq i \leq n$), $H \in \mathcal{K}$, σ and relations: (Here $[X, Y]$ denotes $XY - YX$.)

$$(2.1.1) \quad \sigma^2 = 1, \quad \sigma H \sigma = H \quad (H \in \mathcal{K}), \quad \sigma E_i \sigma = (-1)^{p(\alpha_i)} E_i,$$

$$\sigma F_i \sigma = (-1)^{p(\alpha_i)} F_i,$$

$$(2.1.2) \quad [H_1, H_2] = 0 \quad (H_1, H_2 \in \mathcal{K}),$$

$$[H, E_i] = \alpha_i(H) E_i, \quad [H, F_i] = -\alpha_i(H) F_i \quad (H \in \mathcal{K}),$$

$$(2.1.3) \quad E_i F_j - (-1)^{p(\alpha_i)p(\alpha_j)} F_j E_i = \delta_{ij} \frac{sh(hH_{\alpha_i})}{sh(hd_i)}.$$

Similarly to [15], we have:

Lemma 2.1.4. (the triangular decomposition of \tilde{U}_h^σ) Let \tilde{N}_+ , (resp. $U(\mathcal{K}^R)$, $R<\sigma>$ or \tilde{N}_-) be the unital R -subalgebra of \tilde{U}_h^σ algebraically generated by the elements $\{E_1, \dots, E_n\}$ (resp. $\{H_1, \dots, H_N\}$, $\{\sigma\}$ or $\{F_1, \dots, F_n\}$). Then \tilde{N}_+ (resp. \tilde{N}_-) is isomorphic to the free algebra $R<E_1, \dots, E_n>$ (resp. $R<F_1, \dots, F_n>$). The algebra $U(\mathcal{K}^R)$ is isomorphic to the symmetric algebra $S[\mathcal{K}^R]$ of the R -module $\mathcal{K}^R = R \otimes \mathcal{K}$. The

R -module $R<\sigma>$ is isomorphic to $R\sigma \oplus R$. Moreover we have an isomorphism of h -adic topological R -modules:

$$\hat{N}_+ \otimes S[\mathcal{R}^R] \otimes R<\sigma> \otimes \hat{N}_- \rightarrow \hat{U}_h^\sigma \quad (X \otimes Z \otimes \sigma^c \otimes Y \mapsto X \cdot Z \cdot \sigma^c \cdot Y)$$

($c = 0, 1$).

Proof. Let $R<x_1, \dots, x_n>$ and $R<y_1, \dots, y_n>$ be the tensor algebras of the free R -modules with bases x_1, \dots, x_n and y_1, \dots, y_n respectively. Let z_1, \dots, z_N be a basis of \mathcal{R} . Put $V = \hat{N}_+ \otimes S[\mathcal{R}^R] \otimes R<\sigma> \otimes \hat{N}_-$. Note that the topological basis $\{x_{i_1} \cdots x_{i_u} \otimes z_1^{a_1} \cdots z_N^{a_N} \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v}\}$ is a handy basis with $I = \{(i_1, \dots, i_u, a_1, \dots, a_N, c, j_1, \dots, j_v)\}$ and $p((i_1, \dots, i_u, a_1, \dots, a_N, c, j_1, \dots, j_v)) = i_1 + \dots + i_u + a_1 + \dots + a_N + c + j_1 + \dots + j_v$. We can define a \hat{U}_h^σ -module structure on V by the following formulas: (Here $p(i)$ denotes $p(\alpha_i)$.)

$$\begin{aligned} & F_i \cdot x_{i_1} \cdots x_{i_u} \otimes z_1^{a_1} \cdots z_N^{a_N} \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v} \\ &= - \sum_{s=1}^n \delta_{i_s, i} (-1)^{p(i)(p(i_1) + \dots + p(i_{s-1}))} \\ & \quad x_{i_1} \cdots \hat{x}_{i_s} \cdots x_{i_u} \otimes \frac{sh(h(H_{\alpha_i} - (\alpha_{i_{s+1}} + \dots + \alpha_{i_u}(H_{\alpha_i}))))}{sh(hd_i)} z_1^{a_1} \cdots z_N^{a_N} \\ & \quad \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v} \\ &+ (-1)^{p(i)(p(i_1) + \dots + p(i_u) + c)} \\ & \quad x_{i_1} \cdots x_{i_u} \otimes (z_1 + \alpha_i(z_1))^{a_1} \cdots (z_N + \alpha_i(z_N))^{a_N} \sigma^c \otimes y_i y_{j_1} \cdots y_{j_v} \end{aligned}$$

($1 \leq i \leq n$) ,

$$\begin{aligned} & H \cdot x_{i_1} \cdots x_{i_u} \otimes z_1^{a_1} \cdots z_N^{a_N} \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v} \\ &= x_{i_1} \cdots x_{i_u} \otimes (H + (\alpha_{i_1} + \cdots + \alpha_{i_u})(H)) z_1^{a_1} \cdots z_N^{a_N} \\ & \quad \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v} \quad (H \in \mathcal{R}) , \end{aligned}$$

$$\begin{aligned} & \sigma \cdot x_{i_1} \cdots x_{i_u} \otimes z_1^{a_1} \cdots z_N^{a_N} \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v} \\ &= (-1)^{p(i_1) + \cdots + p(i_u)} x_{i_1} \cdots x_{i_u} \otimes z_1^{a_1} \cdots z_N^{a_N} \otimes \\ & \quad \sigma^{c+1} \otimes y_{j_1} \cdots y_{j_v} , \end{aligned}$$

$$\begin{aligned} & E_i \cdot x_{i_1} \cdots x_{i_u} \otimes z_1^{a_1} \cdots z_N^{a_N} \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v} \\ &= x_i x_{i_1} \cdots x_{i_u} \otimes z_1^{a_1} \cdots z_N^{a_N} \otimes \sigma^c \otimes y_{j_1} \cdots y_{j_v} \quad (1 \leq i \leq n) . \end{aligned}$$

On the other hand, by using (2.1.1) - (2.1.3) , we can show that \tilde{U}_h^σ is generated by the elements

$$E_{i_1} \cdots E_{i_u} \cdot z_1^{a_1} \cdots z_N^{a_N} \cdot \sigma^c \cdot F_{j_1} \cdots F_{j_v} \quad (a_1, \dots, a_n \in \mathbb{Z}_+, c \in \{0, 1\})$$

as an h -adic topological R -module.

Hence we see that the R -module homomorphism $\tilde{U}_h^\sigma \rightarrow V$ ($x \mapsto x \cdot 1_V$) is isomorphism. This completes the proof.

Defining the coproduct Δ , the antipode S and the counit ε by

$$\begin{aligned}\Delta(E_i) &= E_i \otimes 1 + \exp(hH_{\alpha_i}) \cdot \sigma^{p(\alpha_i)} \otimes E_i, \\ \Delta(F_i) &= F_i \otimes \exp(-hH_{\alpha_i}) + \sigma^{p(\alpha_i)} \otimes F_i, \\ \Delta(H) &= H \otimes 1 + 1 \otimes H \quad (H \in \mathcal{H}), \quad \Delta(\sigma) = \sigma \otimes \sigma, \\ S(E_i) &= -\exp(-hH_{\alpha_i}) \cdot \sigma^{p(\alpha_i)} E_i, \quad S(F_i) = -F_i \exp(hH_{\alpha_i}) \sigma^{p(\alpha_i)}, \\ S(H) &= -H \quad (H \in \mathcal{H}), \quad S(\sigma) = \sigma, \\ \varepsilon(E_i) &= \varepsilon(F_i) = \varepsilon(H) = 0, \quad \varepsilon(\sigma) = 1,\end{aligned}$$

the algebra $\tilde{\mathbb{U}}_h^\sigma$ becomes an Hopf algebra.

2.2. Let $(\mathcal{E}, \Pi = \{\alpha_1, \dots, \alpha_n\}, p)$ be a triple system. Put $R' = \mathbb{C}[[\sqrt{h}]]$. Then $R' = R \oplus \sqrt{h}R$. Let $\tilde{\mathbb{U}}_{\sqrt{h}}^{\sigma\sigma} = \tilde{\mathbb{U}}_{\sqrt{h}}^{\sigma\sigma}(\mathcal{E}, \Pi, p)$ be a \sqrt{h} -adic topological R' -algebra with generators E_i ($1 \leq i \leq n$), $H' \in \mathcal{H}$, σ and relations: (In $\tilde{\mathbb{U}}_{\sqrt{h}}^{\sigma\sigma}$, we write H' for $H \in \mathcal{H}$).

$$(2.2.1) \quad \sigma^2 = 1, \quad \sigma H' \sigma = H' \quad (H' \in \mathcal{H}), \quad \sigma E_i \sigma = (-1)^{p(\alpha_i)} E_i,$$

$$(2.2.2) \quad [H'_1, H'_2] = 0 \quad (H'_1, H'_2 \in \mathcal{H}),$$

$$(2.2.3) \quad [H', E_i] = \sqrt{h}\alpha_i(H')E_i \quad (H' \in \mathcal{H}).$$

A topological Hopf algebra structure of $\tilde{U}_{\sqrt{h}}^{\sigma}$ introduced by defining the coproduct Δ' , the antipode S' and the counit ε' defined as follows: (Here H' denotes an arbitrary element of \mathcal{H} .)

$$(2.2.4) \Delta'(E_i) = E_i \otimes 1 + \exp(\sqrt{h}H_{\alpha_i}) \cdot \sigma^{p(\alpha_i)} \otimes E_i ,$$

$$\Delta'(H') = H' \otimes 1 + 1 \otimes H' \quad (H' \in \mathcal{H}), \quad \Delta'(\sigma) = \sigma \otimes \sigma ,$$

$$(2.2.5) S'(E_i) = -\exp(-\sqrt{h}H_{\alpha_i}) \cdot \sigma^{p(\alpha_i)} E_i ,$$

$$S'(H') = -H' \quad (H' \in \mathcal{H}), \quad S'(\sigma) = \sigma ,$$

$$(2.2.6) \varepsilon'(E_i) = \varepsilon'(F_i) = \varepsilon'(H') = 0 , \quad \varepsilon'(\sigma) = 1 .$$

Let $\tilde{\Omega}'_+ : \tilde{U}_{\sqrt{h}}^{\sigma} \rightarrow R' \otimes \tilde{U}_h^{\sigma}$ be a continuous R' -algebra homomorphism defined by putting $\tilde{\Omega}'_+(E_i) = E_i$, $\tilde{\Omega}'_+(H') = \sqrt{h} H'$ ($H' \in \mathcal{H}$), $\tilde{\Omega}'_+(\sigma) = \sigma$. Then $\tilde{\Omega}'_+$ is an h -adic topological Hopf algebra homomorphism. Similarly to the proof of Lemma 2.1.4, we have:

Lemma 2.2.7. (i) $\tilde{\Omega}'_+$ is injective. (ii) Put $\tilde{N}'_+ = R' \otimes \tilde{N}_+ = R' \langle E_1, \dots, E_n \rangle$, $S[\mathcal{H}^{R'}] = R' \otimes S[\mathcal{H}^R]$ and $R' \langle \sigma \rangle = R' \otimes R \langle \sigma \rangle$. Then we have an isomorphism of \sqrt{h} -adic topological R' -modules:

$$\hat{\tilde{N}'_+} \otimes \hat{S[\mathcal{H}^{R'}]} \otimes \hat{R' \langle \sigma \rangle} \rightarrow \tilde{U}_{\sqrt{h}}^{\sigma}$$

$$(X \otimes Z' \otimes \sigma^c \rightarrow X \cdot Z' \cdot \sigma^c \quad (c = 0, 1)) .$$

In particular, if H_1, \dots, H_N are \mathbb{C} -basis of \mathcal{H} , the elements

$$(2.2.8) \quad E_{i_1} \cdots E_{i_u} \cdot H_1^{a_1} \cdots H_N^{a_N} \cdot \sigma^c$$

$$(1 \leq i_1, \dots, i_u \leq n, a_1, \dots, a_N \in \mathbb{Z}_+, c \in \{0, 1\})$$

form a topological basis of $\tilde{\cup}_{\sqrt{h}} \ell_+^\sigma$.

2.3. For $a \in \mathbb{Z}_+$, let S_a' be the submodule of homogeneous elements of degree a of $S[\mathcal{H}^{R'}]$. Then $S[\mathcal{H}^{R'}] = \bigoplus_{a \in \mathbb{Z}_+} S_a'$.

Put $P_+ = \mathbb{Z}_+ \alpha_1 \oplus \cdots \oplus \mathbb{Z}_+ \alpha_n \in \mathcal{E}$. For $\lambda \in P_+$, we put $\tilde{N}_{+\lambda}' = \bigoplus_{\alpha_{i_1} + \cdots + \alpha_{i_v} = \lambda} R' E_{i_1} \cdots E_{i_v}$. Then $\tilde{N}_+' = \bigoplus_{\lambda \in P_+} \tilde{N}_{+\lambda}'$.

We define the elements H_λ° ($\lambda \in \mathcal{E}$), E_i° ($1 \leq i \leq n$), $\sigma^\circ \in (\tilde{\cup}_{\sqrt{h}} \ell_+^\sigma)^*$ as follows: (Here X (resp. Z') denotes a non-zero element of $\tilde{N}_{+\mu}'$ (resp. $S[\mathcal{H}]_a'$) and $c \in \{0, 1\}$.)

$$(2.3.1) \quad H_\lambda^\circ (X \cdot Z' \cdot \sigma^c) = \begin{cases} \lambda(Z') & \text{if } X = 1 \text{ and } a = 1 , \\ 0 & \text{if } \mu \neq 0 \text{ or } a \neq 1 , \end{cases}$$

$$(2.3.2) \quad E_i^\circ (X \cdot Z' \cdot \sigma^c) = \begin{cases} 1 & \text{if } X = E_i \text{ and } Z' = 1 , \\ 0 & \text{if } \mu \neq \alpha_i \text{ or } a \neq 0 , \end{cases}$$

$$(2.3.3) \quad \sigma^\circ (X \cdot Z' \cdot \sigma^c) = \begin{cases} (-1)^c & \text{if } X = Z' = 1 , \\ 0 & \text{if } \mu \neq 0 \text{ or } a \neq 0 . \end{cases}$$

Lemma 2.3.4. (i) $E_i^\circ, H_\lambda^\circ, \sigma^\circ \in (\tilde{U}_{\sqrt{h}}^{\ell^\sigma_+})^\circ$.

(ii) There is a topological R' -Hopf algebra homomorphism

$\theta: \tilde{U}_{\sqrt{h}}^{\ell^\sigma_+} \rightarrow \overline{(\tilde{U}_{\sqrt{h}}^{\ell^\sigma_+})^\circ}$ such that $\theta(H_\lambda^\circ) = H_\lambda^\circ$, $\theta(E_i^\circ) = E_i^\circ$, $\theta(\sigma^\circ) = \sigma^\circ$.

Proof. (i) We show $E_i^\circ \in (\tilde{U}_{\sqrt{h}}^{\ell^\sigma_+})^\circ$ only. It can be easily shown that the basis elements x of (2.2.8) satisfying $x \cdot E_i^\circ \neq 0$ are $1, \sigma, E_i$ and $E_i \sigma$. Hence we have $\text{rank } (\tilde{U}_{\sqrt{h}}^{\ell^\sigma_+}) \cdot E_i^\circ \leq 4$, which implies $E_i^\circ \in (\tilde{U}_{\sqrt{h}}^{\ell^\sigma_+})^\circ$. Similarly, we have $H_\lambda^\circ, \sigma^\circ \in (\tilde{U}_{\sqrt{h}}^{\ell^\sigma_+})^\circ$.

(ii) First we show that θ is an algebra map. We show that the elements $E_i^\circ, H_\lambda^\circ$ satisfy (2.2.3). By the definition of the coproduct Δ' and the definitions of $E_i^\circ, H_\lambda^\circ, \sigma^\circ$, for the basis elements x of (2.2.8), we have:

$$E_i^\circ \cdot H_\lambda^\circ (x) = \begin{cases} (\lambda, \mu) & \text{if } x = E_i \cdot H_\mu^\circ \cdot \sigma^c, \\ 0 & \text{otherwise,} \end{cases}$$

$$H_\lambda^\circ \cdot E_i^\circ (x) = \begin{cases} \sqrt{h}(\lambda, \alpha_i) & \text{if } x = E_i \cdot \sigma^c \\ (\lambda, \mu) & \text{if } x = E_i \cdot H_\mu^\circ \cdot \sigma \\ 0 & \text{otherwise.} \end{cases}$$

Hence it follows that $H_\lambda^\circ \cdot E_i^\circ = E_i^\circ \cdot H_\lambda^\circ + \sqrt{h}(\lambda, \alpha_i) E_i^\circ$, which is the relation (2.2.3). Similarly, we can show that the elements $E_i^\circ, H_\lambda^\circ, \sigma^\circ$ satisfy the relations (2.2.1-2).

Next we show that θ is a Hopf algebra map. Let m denote

the multiplication of $\tilde{U}_{\sqrt{h}}^{\sigma} \ell_+^\sigma$. We are going to show that

$$(2.3.5) \quad t_m(E_i^\circ) = E_i^\circ \otimes \varepsilon + \exp(\sqrt{h} H_{\alpha_i}^\circ) \cdot (\sigma^\circ)^{p(\alpha_i)} \otimes E_i^\circ,$$

which is the one of (2.2.4). Let x and y be a pair of the basis elements (2.2.8). By (2.2.1-3), we have:

$$\begin{aligned} & t_m(E_i^\circ)(x \otimes y) \\ &= \begin{cases} 1 & \text{if } x = E_i \sigma^c \ (c = 0, 1) \text{ and } y = \sigma^d \ (d = 0, 1), \\ (-1)^{p(\alpha_i)c} \cdot (\sqrt{h})^{a_1 + \dots + a_N} \cdot \alpha_i(H_1) \alpha_i(H_2) \dots \alpha_i(H_N) & \text{if } x = H_1^{a_1} \dots H_N^{a_N} \sigma^c \ (a_1, \dots, a_N \in \mathbb{Z}_+, c = 0, 1), y = E_i \sigma^d, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Here we note that, for the basis elements x of (2.2.8),

$$\begin{aligned} (2.3.6) \quad & (H_\lambda^\circ)^a \cdot (\sigma^\circ)^c(x) = \\ &= \begin{cases} a! \lambda(H_{i_1}^\circ) \dots \lambda(H_{i_a}^\circ) (-1)^{cd} & \text{if } x = H_{i_1}^\circ \dots H_{i_a}^\circ \cdot \sigma^d \\ & \quad (1 \leq i_1 \leq \dots \leq i_a \leq N, d = 0, 1), \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Hence we have:

$$t_m(E_i^\circ) = E_i^\circ \otimes \varepsilon + \sum_{a \in \mathbb{Z}_+} \frac{1}{a!} (\sqrt{h})^a (H_{\alpha_i}^\circ)^a \cdot (\sigma^\circ)^{p(\alpha_i)} \otimes E_i^\circ$$

in $\overline{(\tilde{U}_{\sqrt{h}}^{\sigma} \ell_+^\sigma)^0} \hat{\otimes} \overline{(\tilde{U}_{\sqrt{h}}^{\sigma} \ell_+^\sigma)^0}$. The above is nothing else but (2.3.5).

Similarly, we can prove that E_i° , H_λ° , σ° satisfy (2.2.4-6). This completes the proof.

2.4. We define a symmetric topological R' -Hopf algebra pairing $\langle \cdot, \cdot \rangle : \tilde{U}_{\sqrt{h}}^{\sigma} \times \tilde{U}_{\sqrt{h}}^{\sigma} \rightarrow R'$ by $\langle x, y \rangle = \theta(x)(y)$. By (2.3.1-3) and (2.3.6), we have:

Lemma 2.4.1. Let ε_i ($1 \leq i \leq N$) be a \mathbb{C} -basis of \mathfrak{e} such that $(\varepsilon_i, \varepsilon_j) = \delta_{ij}$. Let $X \in \tilde{N}_{+, \lambda}$ and $Y \in \tilde{N}_{+, \mu}$. Then we have:

$$\begin{aligned} & \langle X \cdot H_{\varepsilon_1}^{a_1} \cdots H_{\varepsilon_N}^{a_N} \cdot \sigma^c, Y \cdot H_{\varepsilon_1}^{b_1} \cdots H_{\varepsilon_N}^{b_N} \cdot \sigma^d \rangle \\ &= (-1)^{cd} \cdot \prod_{i=1}^N (\delta_{a_i b_i} \cdot a_i !) \cdot \delta_{\lambda \mu} \langle X, Y \rangle. \end{aligned}$$

2.5. Let $\mathfrak{D}' = D(\tilde{U}_{\sqrt{h}}^{\sigma}, (\tilde{U}_{\sqrt{h}}^{\sigma})^{op})$ be the quantum double defined in §1. Then $\mathfrak{D}' \simeq \tilde{U}_{\sqrt{h}}^{\sigma} \otimes \tilde{U}_{\sqrt{h}}^{\sigma}$ as R' -modules. For an element $X \in \tilde{U}_{\sqrt{h}}^{\sigma}$, we write X for $X \otimes 1$ and X° for $1 \otimes X$.

Lemma 2.5.1. The h -adic topological R' -algebra \mathfrak{D}' is h -adically defined with the generators E_i , E_i° ($1 \leq i \leq n$), H , H° ($H \in \mathcal{H}$), σ , σ° and the relations:

(2.5.2) The elements σ , σ° , H , H° are mutually commutative.

$$(2.5.3) \sigma E_i \sigma = (-1)^{p(\alpha_i)} E_i, \quad \sigma E_i^{\circ} \sigma = (-1)^{p(\alpha_i)} E_i^{\circ},$$

$$\sigma^{\circ} E_i \sigma^{\circ} = (-1)^{p(\alpha_i)} E_i, \quad \sigma^{\circ} E_i^{\circ} \sigma^{\circ} = (-1)^{p(\alpha_i)} E_i^{\circ},$$

$$(2.5.4) [H^-, E_i] = \sqrt{h}\alpha_i(H^-)E_i. \quad [H^{-\circ}, E_i] = -\sqrt{h}\alpha_i(H^{-\circ})E_i,$$

$$[H^{-\circ}, E_i^{\circ}] = \sqrt{h}\alpha_i(H^{-\circ})E_i^{\circ}. \quad [H^-, E_i^{\circ}] = -\sqrt{h}\alpha_i(H^-)E_i^{\circ},$$

$$(2.5.5) E_i E_j^{\circ} - E_j^{\circ} E_i = \delta_{ij} (\exp(\sqrt{h} H_{\alpha_i}^{-\circ}) \sigma^{\circ} E_i^{\circ} - \exp(\sqrt{h} H_{\alpha_i}^-) \sigma E_i).$$

Proof. Put $L_i = \exp(\sqrt{h} H_{\alpha_i}^-) \sigma^{p(\alpha_i)}$ and

$L_i^{\circ} = \exp(\sqrt{h} H_{\alpha_i}^{-\circ}) \sigma^{p(\alpha_i)}$. First we show that (2.5.5) holds in

\mathfrak{D}' . By Proposition 1.4.1 and (2.3.1-3), we have:

$$\begin{aligned} E_j^{\circ} E_i &= \Phi(E_j^{\circ} \otimes E_i) \\ &= \langle L_j^{\circ}, 1 \rangle \langle 1, S'^{-1}(L_i) \rangle E_i E_j^{\circ} + \langle E_j^{\circ}, E_i \rangle \langle 1, S'^{-1}(L_i) \rangle L_i \\ &\quad + \langle L_j^{\circ}, L_i \rangle \langle E_j^{\circ}, S'^{-1}(E_i) \rangle L_j^{\circ} \\ &= -E_i E_j^{\circ} + \delta_{ij} L_i - \delta_{ij} L_j^{\circ}. \end{aligned}$$

Similarly, we can show (2.5.2-4). On the other hand, it can be easily shown that the R' -algebra defined by the relations (2.5.2-5) is generated by the elements

$$E_{i_1} \cdots E_{i_u} H_1^{a_1} \cdots H_N^{a_N} \sigma^c \cdot E_{j_1}^{\circ} \cdots E_{j_v}^{\circ} H_1^{b_1} \cdots H_N^{b_N} \sigma^d$$

as an R' -module where $\{H_1, \dots, H_N\}$ is a basis of \mathcal{H} and

$1 \leq i_1, \dots, i_u, j_1, \dots, j_v \leq N, \quad a_1, \dots, a_N, b_1, \dots, b_N \in \mathbb{Z}_+$,

$c, d \in \{0, 1\}$. Since $\mathfrak{D}' \cong \widetilde{\mathfrak{U}}_h^{\sigma} \otimes \widetilde{\mathfrak{U}}_h^{\sigma}$, this completes the

proof.

2.6. Let $I_{\ell_+}' (\subset \tilde{U}_{\sqrt{h}}^{\ell_+}{}^\sigma)$ (resp. $I_+' (\subset \tilde{N}_+')$) be the kernel of \langle , \rangle (resp. $\langle , \rangle|_{\tilde{N}_+'} \times \tilde{N}_+')$. The following lemma is useful.

Lemma 2.6.1. If J' is an bi-ideal of $\tilde{U}_{\sqrt{h}}^{\ell_+}{}^\sigma$ such that $J' \subset \sum_{i,j} E_i E_j \tilde{U}_{\sqrt{h}}^{\ell_+}{}^\sigma$, then $J' \subset I_{\ell_+}'$.

Proof. By (2.3.1-3), the generators E_i ($1 \leq i \leq n$), H ($H \in \mathcal{H}$), σ are orthogonal to J' . Hence the lemma follows.

2.7. Put $U_{\sqrt{h}}^{\ell_+}{}^\sigma = \tilde{U}_{\sqrt{h}}^{\ell_+}{}^\sigma / I_{\ell_+}'$ and $N_+'' = \tilde{N}_+'' / I_+'$.

Put $I_{+, \lambda}' = I_+ \cap \tilde{N}_{+, \lambda}$ and $N_{+, \lambda}'' = \tilde{N}_{+, \lambda}'' / I_{+, \lambda}'$ for $\lambda \in P_+$.

Lemma 2.7.1. (i) $N_{+, \lambda}''$ is a free R' -module of finite rank.

$N_+'' = \bigoplus_{\lambda \in P_+} N_{+, \lambda}''$. (ii) $I_{\ell_+}' = I_+ \cdot S[\mathcal{H}^{R'}] \cdot R' \langle \sigma \rangle$. (iii) As

topological R' -modules, $U_{\sqrt{h}}^{\ell_+}{}^\sigma \simeq N_+'' \otimes S[\mathcal{H}^{R'}] \otimes R' \langle \sigma \rangle$ ($X \cdot Z' \cdot \sigma^c \leftarrow X \otimes Z' \otimes \sigma^c$). In particular, $U_{\sqrt{h}}^{\ell_+}{}^\sigma$ is a topologically free R' -Hopf algebra.

Proof. Let $c \in R' \setminus \{0\}$ and $x \in N_{+, \lambda}''$. If $cx \in I_{+, \lambda}'$ then $c\langle x, y \rangle = \langle cx, y \rangle = 0$ for all $y \in N_{+, \lambda}''$. Hence $\langle x, y \rangle = 0$ which

implies $x \in I_{+, \lambda}^{\perp}$. Hence the freeness of N_+^{\perp} follows.

From Lemma 2.4.1, we obtain:

$\langle , \rangle \cong$

$$\langle , \rangle |_{N_+^{\perp} \otimes N_+^{\perp}} \otimes \langle , \rangle |_{S[\mathbb{X}^{R'}] \otimes S[\mathbb{X}^{R'}]} \otimes \langle , \rangle |_{R' \langle \sigma \rangle \otimes R' \langle \sigma \rangle}$$

and $\tilde{N}_+^{\perp} = \bigoplus_{\lambda \in P_+^{\perp}} \tilde{N}_{+, \lambda}^{\perp}$ (Here \oplus^{\perp} denotes the orthogonal direct sum).

Then we have (i) and (ii). We immediately obtain (iii) from (ii).

2.8. Let $D' = D(U_{\sqrt{h}}^{\perp}, (U_{\sqrt{h}}^{\perp})^{op})$ ($\cong U_{\sqrt{h}}^{\perp} \otimes (U_{\sqrt{h}}^{\perp})^{op}$) be the quantum double defined in §1. For an element $X \in U_{\sqrt{h}}^{\perp}$, we write X for $X \otimes 1 \in D'$ and X° for $1 \otimes X \in D'$. For a subset M of $U_{\sqrt{h}}^{\perp}$ (resp. $\tilde{U}_{\sqrt{h}}^{\perp}$), we write M for $\{m \otimes 1 \in D' \text{ (resp. } \tilde{D}') \mid m \in M\}$ and M° for $(1 \otimes m \in D' \text{ (resp. } \tilde{D}') \mid m \in M\}$.

By the definitions of the quantum doubles \tilde{D}' and D' (see Proposition 1.4.1), we have:

Lemma 2.8.1. (i) As R' -modules,

$$(2.8.2) \quad \tilde{D}' \cong \tilde{N}' \otimes S[\mathbb{X}^{R'}] \otimes R' \langle \sigma \rangle \otimes \tilde{N}'^\circ \otimes S[\mathbb{X}^{R'}]^\circ \otimes R' \langle \sigma^\circ \rangle,$$

$$(2.8.3) \quad D' \cong N' \otimes S[\mathbb{X}^{R'}] \otimes R' \langle \sigma \rangle \otimes N'^\circ \otimes S[\mathbb{X}^{R'}]^\circ \otimes R' \langle \sigma^\circ \rangle.$$

(ii) Let $\Psi : \mathfrak{D}' \rightarrow \mathfrak{D}'$ be a natural epimorphism defined by $\Psi(\tilde{X} \otimes \tilde{Y}) = X \otimes Y$ where $X = \tilde{X} + I_{\ell_+}'$ and $Y = \tilde{Y} + I_{\ell_+}'$. Then

$\text{Ker } \Psi =$

$$I_+^{\wedge} \cdot S[\mathcal{R}^{\wedge}] \cdot R^{\wedge} < \sigma > \oplus (\tilde{U}_{\sqrt{h}} \ell_+^{\sigma})^{\circ} + \tilde{U}_{\sqrt{h}} \ell_+^{\sigma} \otimes I_+^{\wedge \circ} \cdot S[\mathcal{R}^{\wedge}]^{\circ} \cdot R^{\wedge} < \sigma^{\circ} > .$$

2.9. Let N_+ (resp. \tilde{N}_+) be the unital R -subalgebra of N_+^{\wedge} (resp. \tilde{N}_+^{\wedge}) generated by the elements E_i ($1 \leq i \leq n$). Since N_+^{\wedge} is free (see 2.7.1), N_+ is a free R -module. Let $I_+ = I_+^{\wedge} \cap \tilde{N}_+$. For $\lambda \in P_+$, we put $\tilde{N}_{+, \lambda} = \tilde{N}_+^{\wedge, \lambda} \cap \tilde{N}_+$, $N_{+, \lambda} = N_+^{\wedge, \lambda} \cap N_+$ and $I_{+, \lambda} = I_+^{\wedge, \lambda} \cap N_+$.

Lemma 2.9.1. (i)

$$\begin{aligned} N_+^{\wedge} &= N_+ \oplus \sqrt{h} N_+ , \quad I_+^{\wedge} = I_+ \oplus \sqrt{h} I_+ , \quad N_+ = \tilde{N}_+ / I_+ , \\ N_{+, \lambda}^{\wedge} &= N_{+, \lambda} \oplus \sqrt{h} N_{+, \lambda} , \quad I_{+, \lambda}^{\wedge} = I_{+, \lambda} \oplus \sqrt{h} I_{+, \lambda} , \\ N_{+, \lambda} &= \tilde{N}_{+, \lambda} / I_{+, \lambda} . \end{aligned}$$

where $\lambda \in P_+$.

(ii) For $\lambda \in P_+$, there exists a free R -module $L_{+, \lambda}$ such that $\tilde{N}_{+, \lambda} = I_{+, \lambda} \oplus L_{+, \lambda}$.

Proof. (i) We have $I_{+, \lambda}^{\wedge} = I_{+, \lambda} \oplus \sqrt{h} I_{+, \lambda}$, since $\langle X, Y \rangle \in R$ for all $X, Y \in \tilde{N}_+$. The rests follow easily from this.

(ii) By Lemma 2.7.1, $N_{+, \lambda}^-$ is a free R' -module of finite rank. Hence, by (i), $N_{+, \lambda}$ is a free R -module of finite rank. Since $N_{+, \lambda} = \tilde{N}_{+, \lambda}/I_{+, \lambda}$, choosing representatives $x_1, \dots, x_u \in \tilde{N}_{+, \lambda}$ such that x_1, \dots, x_u form a basis of $N_{+, \lambda}$ (modulo $I_{+, \lambda}$), we have $\tilde{N}_{+, \lambda} = I_{+, \lambda} \oplus (Rx_1 \oplus \dots \oplus Rx_u)$. Hence, putting $L_{+, \lambda} = Rx_1 \oplus \dots \oplus Rx_u$, the part (ii) follows.

Lemma 2.9.2. Put $K_i = \exp(\sqrt{h} H_{\alpha_i}^-)$, $K_i^\circ = \exp(\sqrt{h} H_{\alpha_i}^{\circ -}) \in D'$ ($1 \leq i \leq n$). Let T (resp. T°) be the R -subalgebra of D' generated by the elements $K_i^{\pm 1}$ (resp. $K_i^{\circ \pm 1}$) ($1 \leq i \leq n$). Let U be the R -subalgebra of D' algebraically generated by the elements $\sigma^{\pm 1}$, $\sigma^{\circ \pm 1}$ and $K_i^{\pm 1}$, $K_i^{\circ \pm 1}$, E_i , E_i° ($1 \leq i \leq n$). Then

- (i) U is a (non-topological) R -Hopf subalgebra of D' .
- (ii) As R -modules, $U \cong N_+ \otimes T \otimes R<\sigma> \otimes N_+^\circ \otimes T^\circ \otimes R<\sigma^\circ>$ ($Xt\sigma^c Y^\circ s^\circ \sigma^d \leftarrow X \otimes t \otimes \sigma^c \otimes Y^\circ \otimes s^\circ \otimes \sigma^d$) ($X \in N_+$, $t \in T$, $c \in \{0, 1\}$, $Y^\circ \in N_+^\circ$, $s^\circ \in T^\circ$, $d \in \{0, 1\}$). The elements $K_1^{\ell_1} \cdots K_n^{\ell_n}$ (resp. $K_1^{\circ \ell_1} \cdots K_n^{\circ \ell_n}$) ($\ell_1, \dots, \ell_n \in \mathbb{Z}$) form an R -basis of T (resp. T°).

Proof. (i) By (2.2.4-6) and Proposition 1.4.1, we have (i).

(ii) By (2.2.1-3), (2.5.2-5), we have

$$U = N_+ \cdot T \cdot R<\sigma> \cdot N_+^\circ \cdot T^\circ \cdot R<\sigma^\circ>.$$

By Lemma 2.8.1 (i) , we can easily show that

$$\begin{aligned} & N_+ \cdot K_1^{\ell_1} \cdots K_n^{\ell_n} \cdot R<\sigma> \cdot N_+^\circ \cdot K_1^{\circ m_1} \cdots K_n^{\circ m_n} \cdot R<\sigma^\circ> \\ & \simeq N_+ \otimes K_1^{\ell_1} \cdots K_n^{\ell_n} \otimes R<\sigma> \otimes N_+^\circ \otimes K_1^{\circ m_1} \cdots K_n^{\circ m_n} \otimes R<\sigma^\circ> \end{aligned}$$

holds in D' for each $\ell_1, \dots, \ell_n, m_1, \dots, m_n \in \mathbb{Z}$. Therefore, it is enough to show

$$(2.9.3) \quad U \simeq \bigoplus_{\ell_i, \ell'_i \in \mathbb{Z}} N_+ \cdot K_1^{\ell_1} \cdots K_n^{\ell_n} \cdot R<\sigma> \cdot N_+^\circ \cdot K_1^{\circ m_1} \cdots K_n^{\circ m_n} \cdot R<\sigma^\circ>$$

Take $\alpha_{n+1}, \dots, \alpha_N \in \mathcal{E}$ such that $\alpha_1, \dots, \alpha_n, \alpha_{n+1}, \dots, \alpha_N$ form a basis of \mathcal{E} . For $1 \leq i \leq n$, define R' -module maps

$\theta_i : D' \rightarrow D'$ and $\theta_i^\circ : D' \rightarrow D'$ by:

$$\begin{aligned} & \theta_i(X \otimes H_{\alpha_1}^{a_1} \cdots H_{\alpha_N}^{a_N} \otimes \sigma^c \otimes Y^\circ \otimes H_{\alpha_1}^{\circ b_1} \cdots H_{\alpha_N}^{\circ b_N} \otimes \sigma^\circ d) \\ & = a_i \cdot X \otimes H_{\alpha_1}^{a_1} \cdots H_{\alpha_i}^{a_i-1} \cdots H_{\alpha_N}^{a_N} \otimes \sigma^c \otimes Y^\circ \otimes H_{\alpha_1}^{\circ b_1} \cdots H_{\alpha_N}^{\circ b_N} \otimes \sigma^\circ d, \end{aligned}$$

$$\begin{aligned} & \theta_i^\circ(X \otimes H_{\alpha_1}^{a_1} \cdots H_{\alpha_N}^{a_N} \otimes \sigma^c \otimes Y^\circ \otimes H_{\alpha_1}^{\circ b_1} \cdots H_{\alpha_N}^{\circ b_N} \otimes \sigma^\circ d) \\ & = b_i \cdot X \otimes H_{\alpha_1}^{a_1} \cdots H_{\alpha_N}^{a_N} \otimes \sigma^c \otimes Y^\circ \otimes H_{\alpha_1}^{\circ b_1} \cdots H_{\alpha_i}^{b_i-1} \cdots H_{\alpha_N}^{\circ b_N} \otimes \sigma^\circ d \end{aligned}$$

where the elements

$$X \otimes H_{\alpha_1}^{a_1} \cdots H_{\alpha_N}^{a_N} \otimes \sigma^c \otimes Y^\circ \otimes H_{\alpha_1}^{\circ b_1} \cdots H_{\alpha_N}^{\circ b_N} \otimes \sigma^\circ d$$

are basis elements of

$$D' \cong N' \otimes S[\mathbb{R}^R] \otimes R' \langle \sigma \rangle \otimes N'^\circ \otimes S[\mathbb{R}^R]^\circ \otimes R' \langle \sigma^\circ \rangle$$

(see (2.8.3)). Then we have:

$$\begin{aligned} & \theta_i(X \otimes K_1^{\ell_1} \cdots K_n^{\ell_n} \otimes \sigma^c \otimes Y^\circ \otimes K_1^{m_1} \cdots K_n^{m_n} \otimes \sigma^{\circ d}) \\ &= \ell_i \sqrt{h} \cdot (X \otimes K_1^{\ell_1} \cdots K_n^{\ell_n} \otimes \sigma^c \otimes Y^\circ \otimes K_1^{m_1} \cdots K_n^{m_n} \otimes \sigma^{\circ d}), \end{aligned}$$

$$\begin{aligned} & \theta_i^\circ(X \otimes K_1^{\ell_1} \cdots K_n^{\ell_n} \otimes \sigma^c \otimes Y^\circ \otimes K_1^{m_1} \cdots K_n^{m_n} \otimes \sigma^{\circ d}) \\ &= m_i \sqrt{h} \cdot (X \otimes K_1^{\ell_1} \cdots K_n^{\ell_n} \otimes \sigma^c \otimes Y^\circ \otimes K_1^{m_1} \cdots K_n^{m_n} \otimes \sigma^{\circ d}) \end{aligned}$$

Hence (2.9.3) is the eigenspace decomposition of U with respect to θ_i and θ_i° .

Now we state the main theorem of §2. We put $q = e^h$.

Theorem 2.9.4. Let $(\mathcal{E}, \Pi = \{\alpha_1, \dots, \alpha_n\}, p)$ be a triple system and $D = \text{diag}(d_1, \dots, d_n)$ ($d_i \in \frac{1}{2}\mathbb{Z} \setminus \{0\}$). Put $q_i = q^{d_i} \in R$. Then there exists a unique topologically free R -Hopf algebra $U_h^\sigma = U_h^\sigma((\mathcal{E}, \Pi, p), D)$ satisfying the following conditions:

(i) The R -algebra U_h^σ contains $S[\mathbb{R}^R]$, $R\langle \sigma \rangle$, N_+ , N_+° as R -subalgebras. Here N_+° is another algebra isomorphic to N_+ . For $Y \in N_+$, we denote the corresponding element of N_+°

by Y° . As topological R -modules,

$$U_h^\sigma \cong N_+ \otimes S[\mathbb{K}^R] \otimes R<\sigma> \otimes N_+^\circ \quad (X \cdot P \cdot \sigma^C \cdot Y^\circ \leftarrow X \otimes P \otimes \sigma^C \otimes Y^\circ).$$

(ii) There is a topological R' -Hopf algebra homomorphism $\Omega' : D' \rightarrow U_h^\sigma \otimes R'$ such that

$$(2.9.5) \quad \Omega'(E_i) = E_i \quad (1 \leq i \leq n), \quad \Omega'(H') = -\Omega'(H'^\circ) = \sqrt{h} H \quad (H \in \mathbb{K}) .$$

$$\Omega'(\sigma) = \Omega'(\sigma^\circ) = \sigma, \quad \Omega'(E_i^\circ) = (q_i^{-1} - q_i) E_i^\circ \quad (1 \leq i \leq n) .$$

Proof. Step I. Put $E_i^\circ = F_i \sigma^{p(\alpha_i)} \in \tilde{U}_h^\sigma \quad (1 \leq i \leq n)$. By Lemma 2.5.1, we see that there exists a topological R' -Hopf algebra homomorphism $\tilde{\Omega}' : D' \rightarrow \tilde{U}_h^\sigma \otimes R'$ satisfying the conditions (2.9.5). Let \tilde{N}_+° be the unital subalgebra of \tilde{U}_h^σ algebraically generated by the elements $E_i^\circ \quad (1 \leq i \leq n)$. By Lemma 2.1.4, it is clear that

$$(2.9.6) \quad \tilde{U}_h^\sigma \cong \tilde{N}_+ \otimes S[\mathbb{K}^R] \otimes R<\sigma> \otimes \tilde{N}_+^\circ .$$

Step II. We construct U_h^σ as a quotient of \tilde{U}_h^σ . Set $J_1 = \overline{I_+ \cdot S[\mathbb{K}^R] \cdot R<\sigma> \cdot \tilde{N}_+^\circ}$ and $J_2 = \overline{\tilde{N}_+ \cdot S[\mathbb{K}^R] \cdot R<\sigma> \cdot I_+^\circ}$ where $I_+^\circ = \{Y^\circ \mid Y \in I_+\}$. We claim:

$$(2.9.7) \quad J_1 \text{ and } J_2 \text{ are Hopf ideals of } \tilde{U}_h^\sigma .$$

Assume this fact for a moment. We define the Hopf algebra U_h^σ by $\tilde{U}_h^\sigma / (J_1 + J_2)$. By Lemma 2.8.1, there exists an R' -Hopf algebra homomorphism $\Omega' : \mathfrak{D}' \rightarrow U_h^\sigma \otimes R'$ naturally induced from $\tilde{\Omega}' : \tilde{\mathfrak{D}'} \rightarrow \tilde{U}_h^\sigma \otimes R'$ of Step 1. By (2.9.6) and the definition of U_h^σ , it is clear that $U_h^\sigma \simeq N_+ \otimes S[\mathbb{R}] \otimes R<\sigma> \otimes N_+$ as R -modules. In particular, U_h^σ is topologically free. Since $h(U_h^\sigma \otimes R') \subset \text{Im } \Omega'$, the product in U_h^σ is uniquely determined by Ω' . Hence the uniqueness of U_h^σ follows.

It remains to prove (2.9.7). We shall prove this only for J_1 ; J_2 can be treated similarly.

First we prove that J_1 is a two-sided ideal of the algebra \tilde{U}_h^σ . Evidently, J_1 is a right ideal and $E_i \cdot J_1 \subset J_1$ ($1 \leq i \leq n$), $H \cdot J_1 \subset J_1$ ($H \in \mathcal{H}$), $\sigma \cdot J_1 \subset J_1$. Recall $I_+ = \bigoplus_{\lambda \in P_+} I_{+, \lambda}$.

By Lemma 2.9.1, J_1 is a direct summand of \tilde{U}_h^σ . Hence it is enough to show that

$$(2.9.8) \quad (q_i^{-1} - q_i) E_i^\circ \cdot I_{+, \lambda} \subset I_{+, \lambda} \cdot E_i^\circ \\ + I_{+, \lambda - \alpha_i} \cdot \exp(-hH_{\alpha_i}) \sigma^{p(\alpha_i)} + I_{+, \lambda - \alpha_i} \cdot \exp(hH_{\alpha_i}) \sigma^{p(\alpha_i)}$$

Let X be an element of $I_{+, \lambda}$. By Lemma 2.5.1, we have that

$$(2.9.9) \quad E_i^\circ \cdot X = \\ X \cdot E_i^\circ + X^{(1)} \cdot \exp(\sqrt{h} H_{\alpha_i}^\circ) \sigma^{p(\alpha_i)} + X^{(2)} \cdot \exp(\sqrt{h} H_{\alpha_i}^\circ) \sigma^{p(\alpha_i)}$$

with some $X^{(1)}, X^{(2)} \in \tilde{N}_{+, \lambda - \alpha_i}$ in the algebra \mathfrak{D}' . Let

$\Psi : \mathfrak{D}' \rightarrow \mathfrak{D}'$ be the Hopf algebra homomorphism defined in Lemma 2.8.1. By Lemma 2.9.2, $X^{(1)}, X^{(2)} \in \text{Ker } \Psi$. Hence, by Lemma 2.8.1, $X^{(1)}, X^{(2)} \in I_{+, \lambda-\alpha_i}$. If we let operate $\tilde{\Omega}'$ on the left and right hand sides of (2.9.9), we obtain:

$$(q_i^{-1} - q_i) E_i^\circ \cdot X = (q_i^{-1} - q_i) X \cdot E_i^\circ \\ + X^{(1)} \cdot \exp(-hH_{\alpha_i}) \sigma^{p(\alpha_i)} + X^{(2)} \cdot \exp(hH_{\alpha_i}) \sigma^{p(\alpha_i)}.$$

Hence (2.9.9) follows. Thus we showed that J_1 is a two-sided ideal.

Noting that $\text{Ker } \Psi$ is a Hopf ideal, and using an argument similar to the above, we have

$$\Delta(X) = \sum_{\mu, \nu \in P_+, \mu+\nu=\lambda} \{ X_\mu^{(1)} \cdot \exp(hH_\nu) \sigma^{p(\nu)} \otimes Y_\nu^{(1)} \\ + Y_\mu^{(2)} \cdot \exp(hH_\nu) \sigma^{p(\nu)} \otimes X_\nu^{(2)} \}, \quad X \in I_{+, \lambda},$$

with some $X_\mu^{(1)} \in I_{+, \mu}$, $X_\nu^{(2)} \in I_{+, \nu}$, $Y_\nu^{(1)} \in N_{+, \nu}$, $Y_\mu^{(2)} \in N_{+, \mu}$. Hence $\Delta(J_1) \subset J_1 \otimes \tilde{U}_h^\sigma + \tilde{U}_h^\sigma \otimes J_1$. Similarly we have:

$$S(I_{+, \lambda}) \subset I_{+, \lambda} \exp(hH_\lambda) \cdot \sigma^{p(\lambda)}.$$

Hence $S(J_1) \subset J_1$. It is clear that $\varepsilon(J_1) = 0$. Hence (2.9.7) is proved. This completes the proof.

By Proposition 1.7.1 and Lemma 2.4.1, we have:

Lemma 2.9.10. Let $\Omega' : D' \rightarrow U_h^\sigma \otimes R'$ be the Hopf algebra homomorphism defined in Theorem 2.9.4. Let K' (resp. K) be the fraction field of R' (resp. R). Denote the scalar extention $\Omega' \otimes \text{id}_{K'} : D' \otimes K' \rightarrow U_h^\sigma \otimes K'$ of Ω' again by Ω' . For $\lambda \in P_+$, put $p_\lambda = \text{rank } N_{+, \lambda}$, and let $\{e_{(\lambda, i)}\}_{1 \leq i \leq p_\lambda}$, $\{e^{(\lambda, i)}\}_{1 \leq i \leq p_\lambda}$ be bases of $N_{+, \lambda} \otimes K$ such that $\langle e_{(\lambda, i)}, e^{(\lambda, j)} \rangle = \delta_{ij}$. Let $\{\varepsilon_i\}_{1 \leq i \leq N}$ be a basis of \mathcal{E} such that $\langle \varepsilon_i, \varepsilon_j \rangle = \delta_{ij}$. Put $t_0 = \sum_{i=1}^N h_{\varepsilon_i} \otimes h_{\varepsilon_i} \in \mathcal{H} \otimes \mathcal{H}$. If the element

$$\mathcal{R} = \left(\sum_{\lambda \in P_+} \sum_{i=1}^{p_\lambda} \Omega'(e_{(\lambda, i)}) \otimes \Omega'(e^{(\lambda, i)}) \right) \cdot \exp(-ht_0) \cdot \left(\frac{1}{2} \sum_{c, d=0}^1 (-1)^{cd} \sigma^c \otimes \sigma^d \right)$$

converges in $U_h^\sigma \otimes U_h^\sigma \otimes K'$ and $\mathcal{R} \in U_h^\sigma \otimes U_h^\sigma$, then $(U_h^\sigma, \Delta, \mathcal{R})$ is a quasi-triangular Hopf algebra.

Remark 2.9.11. By Proposition 1.9.2, for U_h^σ , we define an h -adic topologically free R -Hopf superalgebra U_h . In Introduction, we wrote $U_h(\pi, p)$ for this U_h .

2.10. Here we show that, if $\Pi = \{\alpha_1, \dots, \alpha_n\}$ is the set of simple roots of a symmetrizable Kac-Moody Lie algebra G and $p(\alpha_i) = 0$ ($1 \leq i \leq n$), then $U_h = U_h^\sigma / U_h^{\sigma(\sigma-1)}$ is isomorphic to the quantized enveloping algebra $U_h(G)$ introduced by Drinfeld [2] and Jimbo [3]. More precisely, we obtain the following theorem.

Theorem 2.10.1. Suppose $p(\alpha_i) = 0$ for all i . Assume that $(\alpha_i, \alpha_i) > 0$ ($1 \leq i \leq n$), $(\alpha_i, \alpha_j) \leq 0$ ($i \neq j$) and $a_{ij} = 2(\alpha_i, \alpha_j)/(\alpha_i, \alpha_i) \in \mathbb{Z}$. Let $d_i = \frac{(\alpha_i, \alpha_i)}{2}$ ($1 \leq i \leq n$) and $D = \text{diag}(d_1, \dots, d_n)$. Then I_+ is the ideal of \tilde{N}_+ generated by the elements

$$(2.10.2) \quad \sum_{v=0}^{1-a_{ij}} (-1)^v \left[\begin{matrix} 1-a_{ij} \\ v \end{matrix} \right]_{q_i} E_i^v E_j E_i^{1-a_{ij}-v} \quad (i \neq j)$$

where $q_i = \exp(hd_i)$ and

$$\left[\begin{matrix} 1-a_{ij} \\ v \end{matrix} \right]_{q_i} = \frac{v}{\prod_{s=1}^{2-a_{ij}-v} \frac{q_i^{a_{ij}+s-2} - q_i^{-s}}{q_i^s - q_i^{-s}}} \in R.$$

Proof. Let us denote $y_{ij} \in \tilde{N}_+$ the element (2.10.2). By a direct computation, we see that

$\Delta'(y_{ij}) = y_{ij} \otimes 1 + \exp(\sqrt{h}H'((1-a_{ij})\alpha_i + \alpha_j)) \otimes y_{ij}$. Hence the ideal of $\tilde{U}_{\sqrt{h}}^\sigma$ generated by the elements y_{ij} is the bi-ideal.

Hence, by Lemma 2.6.1, $y_{ij} \in I_+$. Put $\mathbb{U} = U_h/hU_h$ where $U_h = U_h^\sigma/U_h^{\sigma-1}$. Since $U(G)$ is defined by the Serre relations, there exists a natural epimorphism $\psi : U(G) \rightarrow \mathbb{U}$.

Let $U(G) = U(n_+) \otimes U(\mathfrak{k}) \otimes U(n_-)$ be the triangular decomposition. Put $N_+ = N_+/hN_+$. Let $U(n_+)_\gamma$ and $N_{+, \gamma}$ denote the weight space of a weight $\gamma \in P_+$. Let $V(\lambda)$ and $V(\lambda)$ denote the irreducible highest weight module with highest weight λ of $U(G)$ and \mathbb{U} respectively. Let $V(\lambda)_{\lambda-\gamma} \subset V(\lambda)$ and $V(\lambda)_{\lambda-\gamma} \subset V(\lambda)$ be the weight spaces of weight $\lambda-\gamma$. From a well-known fact in the representation theory of G (see the formula (10.4.6) in [4]), we can see that, if λ is sufficiently large as compared with $\gamma \in P_+$, $\dim U(n_+)_\gamma = \dim V(\lambda)_{\lambda-\gamma}$. On the other hand, using $\psi : U(G) \rightarrow \mathbb{U}$, we can regard $V(\lambda)$ as an irreducible $U(G)$ -module isomorphic to $V(\lambda)$. Hence we have:

$$\dim U(n_+)_\gamma = \dim V(\lambda)_{\lambda-\gamma} = \dim V(\lambda)_{\lambda-\gamma} \leq \dim N_{+, \gamma}.$$

Since ψ is an epimorphism, $\dim U(n_+)_\gamma \geq \dim N_{+, \gamma}$. Hence we have

$$(2.10.3) \quad \dim U(n_+)_\gamma = \dim N_{+, \gamma}.$$

Note that $\tilde{N}_{+, \lambda}$ is a free R -module of finite rank and $I_{+, \lambda}$ is a direct summand of $\tilde{N}_{+, \lambda}$ (see Lemma 2.9.1). Hence, if w_u ($1 \leq u \leq \text{rank } I_{+, \lambda}$) are elements of $I_{+, \lambda}$ such that $\{w_u + hI_{+, \lambda}\}$ is a \mathbb{C} -basis of $I_{+, \lambda}/hI_{+, \lambda}$, then $\{w_u\}$ is an

R -basis of $I_{+, \lambda}$. In particular, by (2.10.3), we can put

$$w_u = E_{u_1} \cdots E_{u_{k-1}} y_{u_k, u_{K+1}} E_{u_{k+2}} \cdots E_{u_p} \in I_{+, \lambda}$$
$$(\lambda = -a_{u_k, u_{k+1}} \alpha_{u_k} + \sum_{t=1}^p \alpha_{u_t}) .$$

Hence the theorem follows.

Remark 2.10.3. We remark that the above theorem can also be proved by using Tanisaki's result; as an immediate consequence of Proposition 2.4.1 in [16], we can show the non-degeneracy of the Hopf pairing $\langle , \rangle : U_h(n_+) \times U_h(n_+) \rightarrow R$ where $U_h(n_+) = \tilde{N}_+ / (y_{ij} \ (1 \leq i \neq j \leq n))$. Hence $I_+ = (y_{ij})$.

§3. Root systems of simple Lie superalgebras.

3.1. Let $\mathfrak{G} = \mathfrak{G}_0 \oplus \mathfrak{G}_1$ be a finite dimensional complex simple Lie superalgebra of type A-G. Let $(\Phi, \mathcal{E}_{\mathbb{R}})$ be a root system of \mathfrak{G} . For terminologies related to simple Lie superalgebras, see [5]. Here $\mathcal{E}_{\mathbb{R}}$ is an N-dimensional real vector space with a non-degenerate symmetric bilinear form $(,) : \mathcal{E}_{\mathbb{R}} \times \mathcal{E}_{\mathbb{R}} \rightarrow \mathbb{R}$ and $\Phi (\subset \mathcal{E}_{\mathbb{R}})$ is the set of roots. Put $\mathcal{E} = \mathbb{C} \otimes \mathcal{E}_{\mathbb{R}}$. Let $\Pi = \{\alpha_1, \dots, \alpha_n\}$ be a set of simple roots and $p : \Pi \rightarrow \{0,1\} = \mathbb{Z}/2\mathbb{Z}$ the parity function. Put $P = \mathbb{Z}\alpha_1 + \dots + \mathbb{Z}\alpha_n (\subset \mathcal{E})$. We extend p the function $p : P \rightarrow \mathbb{Z}/2\mathbb{Z}$ additively. We assume that the triple (\mathcal{E}, Π, p) is of distinguished type if \mathfrak{G} is of type F_4 or G_3 . We do not treat (\mathcal{E}, Π, p) of type $D(2,1;\alpha)$. That case is easy. However we need some unpleasant notation for type $D(2,1;\alpha)$. To fix notation, we list below Dynkin diagrams, systems of simple roots $\Pi = \{\alpha_1, \dots, \alpha_n\}$, the set of positive roots Φ_+ and parity function $p : \Pi \rightarrow \{0,1\}$ of triples (\mathcal{E}, Π, p) of type A - G. We put $N = n + 1$ if (Φ, Π, p) is of type A, and $N = n$ otherwise. Let $\{\bar{\varepsilon}_i\}_{1 \leq i \leq N}$ be a fixed orthogonal basis of $\mathcal{E}_{\mathbb{R}}$; the values of $(\bar{\varepsilon}_i, \bar{\varepsilon}_i)$ are given below. The element of $\mathcal{E}_{\mathbb{R}}$ written under the dot with the i -th label is the simple root α_i . Note that the numbering of α_i 's is not the standard one for types F_4 and G_3 . In the following diagram, the parity function $p : \Pi \rightarrow \{0,1\}$ is defined as follows. The dot \times at the i -th label stands for the dot \circ (resp. \otimes) if $(\alpha_i, \alpha_i) \neq 0$ (resp. $(\alpha_i, \alpha_i) = 0$). If the i -th dot is \circ , \otimes or \bullet , then we define $p(\alpha_i) = 0, 1, 1$ respectively. We also give the diagonal

matrix $D = \text{diag}(d_1, \dots, d_n)$ such that $A = D^{-1}[(\alpha_i, \alpha_j)]$ is the Cartan matrix of (Φ, Π) .

(i) Types A, B, C or D. For types A-D, we put

$(\bar{\varepsilon}_i, \bar{\varepsilon}_j) = \pm \delta_{ij}$ ($1 \leq i, j \leq N$), where we can arbitrarily choose the signs of $(\bar{\varepsilon}_i, \bar{\varepsilon}_i)$.

$$(A_{N-1}) \quad \begin{array}{ccccccccc} 1 & & 2 & & & & N-1 & & \\ \overline{\varepsilon}_1 - \overline{\varepsilon}_2 & \times & \overline{\varepsilon}_2 - \overline{\varepsilon}_3 & \times & \cdots & \times & \overline{\varepsilon}_{N-1} - \overline{\varepsilon}_N & & \end{array}, \quad D = \text{diag}(1, \dots, 1),$$

$$\Phi_+ = \{\bar{\varepsilon}_i - \bar{\varepsilon}_j \mid 1 \leq i < j \leq N\}.$$

$$(B_N) \quad \begin{array}{ccccccccc} 1 & & 2 & & & & N-1 & & N \\ \overline{\varepsilon}_1 - \overline{\varepsilon}_2 & \times & \overline{\varepsilon}_2 - \overline{\varepsilon}_3 & \times & \cdots & \times & \overline{\varepsilon}_{N-1} - \overline{\varepsilon}_N & \xrightarrow{O} & \overline{\varepsilon}_N \end{array} \quad \text{or} \quad \begin{array}{ccccccccc} 1 & & 2 & & & & N-1 & & N \\ \overline{\varepsilon}_1 - \overline{\varepsilon}_2 & \times & \overline{\varepsilon}_2 - \overline{\varepsilon}_3 & \times & \cdots & \times & \overline{\varepsilon}_{N-1} - \overline{\varepsilon}_N & \xrightarrow{\bullet} & \overline{\varepsilon}_N \end{array},$$

$$D = \text{diag}(1, \dots, 1, \frac{1}{2}),$$

$$\Phi_+ = \{\bar{\varepsilon}_i \pm \bar{\varepsilon}_j \mid 1 \leq i < j \leq N\}, \quad \bar{\varepsilon}_i \quad (1 \leq i \leq N) \quad 2\bar{\varepsilon}_i \quad (1 \leq i \leq N), \quad p(\bar{\varepsilon}_i) = 0\}.$$

$$(C_N) \quad \begin{array}{ccccccccc} 1 & & 2 & & & & N-1 & & N \\ \overline{\varepsilon}_1 - \overline{\varepsilon}_2 & \times & \overline{\varepsilon}_2 - \overline{\varepsilon}_3 & \times & \cdots & \times & \overline{\varepsilon}_{N-1} - \overline{\varepsilon}_N & \xleftarrow{O} & 2\bar{\varepsilon}_N \end{array}, \quad D = \text{diag}(1, \dots, 1, 2),$$

$$\Phi_+ = \{\bar{\varepsilon}_i \pm \bar{\varepsilon}_j \mid 1 \leq i < j \leq N\}, \quad 2\bar{\varepsilon}_i \quad (1 \leq i \leq N), \quad p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 0\}.$$

$$(D_N) \quad \begin{array}{ccccccccc} 1 & & 2 & & & & N-2 & & N-1 \\ \overline{\varepsilon}_1 - \overline{\varepsilon}_2 & \times & \overline{\varepsilon}_2 - \overline{\varepsilon}_3 & \times & \cdots & \times & \overline{\varepsilon}_{N-2} - \overline{\varepsilon}_{N-1} & \xrightarrow{O} & \overline{\varepsilon}_{N-1} - \overline{\varepsilon}_N \end{array} \quad \text{if } (\bar{\varepsilon}_{N-1}, \bar{\varepsilon}_{N-1}) = (\bar{\varepsilon}_N, \bar{\varepsilon}_N),$$

$$\begin{array}{ccccccccc} 1 & & 2 & & & & N-2 & & N-1 \\ \overline{\varepsilon}_1 - \overline{\varepsilon}_2 & \times & \overline{\varepsilon}_2 - \overline{\varepsilon}_3 & \times & \cdots & \times & \overline{\varepsilon}_{N-2} - \overline{\varepsilon}_{N-1} & \xrightarrow{O} & \overline{\varepsilon}_{N-1} - \overline{\varepsilon}_N \end{array} \quad \text{if } (\bar{\varepsilon}_{N-1}, \bar{\varepsilon}_{N-1}) = -(\bar{\varepsilon}_N, \bar{\varepsilon}_N),$$

$$D = \text{diag}(1, \dots, 1).$$

$$\Phi_+ = \{\bar{\varepsilon}_i \pm \bar{\varepsilon}_j \mid 1 \leq i < j \leq N\}, \quad 2\bar{\varepsilon}_i \quad (1 \leq i \leq N), \quad p(\bar{\varepsilon}_i - \bar{\varepsilon}_{N+1}) = 0\}.$$

(ii) Types F_4 and G_3 .

$$(F_4) \quad \begin{array}{ccccccccc} & 1 & & 4 & & 3 & & 2 & \\ \textcircled{1} & \xrightarrow{\varepsilon_2 - \bar{\varepsilon}_3} & \textcircled{4} & \xrightarrow{\varepsilon_3 - \bar{\varepsilon}_4} & \textcircled{3} & \xrightleftharpoons{\varepsilon_4} & \textcircled{2} & \xrightarrow{\frac{1}{2}(\varepsilon_1 - \bar{\varepsilon}_2 - \bar{\varepsilon}_3 - \bar{\varepsilon}_4)} & \end{array},$$

$$[(\bar{\varepsilon}_i, \bar{\varepsilon}_j)] = \text{diag}(6, -2, -2, -2), \quad D = \text{diag}(2, 1, 1, 2).$$

$$\Phi_+ = \{n_1\alpha_1 + n_4\alpha_4 + n_3\alpha_3 + n_2\alpha_2 \mid (n_1, n_4, n_3, n_2) = (1, 0, 0, 0), (1, 1, 0, 0), (1, 1, 1, 0), (1, 1, 2, 0), (1, 1, 1, 1), (1, 2, 2, 0), (1, 1, 2, 1), (1, 2, 2, 1), (1, 2, 3, 1), (1, 2, 3, 2), (0, 0, 0, 1), (0, 0, 1, 1), (0, 1, 1, 1), (0, 1, 2, 1), (0, 0, 1, 0), (0, 1, 2, 0), (0, 1, 1, 0), (0, 1, 0, 0)\}.$$

$$(G_3) \quad \begin{array}{ccccccccc} & 1 & & 3 & & 2 & & & \\ \textcircled{1} & \xrightleftharpoons{\varepsilon_1 - \bar{\varepsilon}_2} & \textcircled{3} & \xrightleftharpoons{\frac{1}{2}(\varepsilon_2 - \bar{\varepsilon}_3)} & \textcircled{2} & \xrightleftharpoons{\varepsilon_3} & & & \end{array},$$

$$[(\bar{\varepsilon}_i, \bar{\varepsilon}_j)] = \text{diag}(-2, 2, 6), \quad D = \text{diag}(1, 3, 1).$$

$$\Phi_+ = \{n_1\alpha_1 + n_3\alpha_3 + n_2\alpha_2 \mid (n_1, n_4, n_3) = (1, 0, 0), (1, 1, 0), (1, 1, 1), (1, 2, 1), (1, 3, 1), (1, 3, 2), (1, 4, 2), (2, 4, 2), (0, 0, 1), (0, 1, 1), (0, 3, 2), (0, 2, 1), (0, 3, 1), (0, 1, 0)\}.$$

3.2. Let Φ_+ be the set of positive roots with respect to Π . Put $\Phi_+^{\text{red}} = \{\beta \in \Phi_+ \mid \frac{1}{2}\beta \notin \Phi_+\}$. We define the partial order $<$ on Φ_+^{red} as follows. Given $\beta = c_1\alpha_1 + \cdots + c_n\alpha_n \in \Phi_+^{\text{red}}$, we define integers $ht(\beta)$, $g(\beta)$, $c_\beta \in \mathbb{Z}_+$ by $ht(\beta) = c_1 + \cdots + c_n$, $g(\beta) = \min\{i \mid c_i \neq 0\}$, $c_\beta = c_{g(\beta)}$ respectively. Define a half integer $ht'(\beta)$ by $ht'(\beta) = ht(\beta)/c_\beta$. Let α, β be elements of Φ_+^{red} . We say

that $\alpha < \beta$ if they satisfy one of the following

$$(i) \quad g(\alpha) < g(\beta) ,$$

$$(ii) \quad g(\alpha) = g(\beta) \text{ and } ht'(\alpha) < ht'(\beta)$$

or

$$(iii) \Phi_+^{\text{red}} \text{ is of type } D_N, \quad p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 0 \quad \text{and} \quad \alpha = \bar{\varepsilon}_i - \bar{\varepsilon}_N , \\ \beta = 2\bar{\varepsilon}_i \quad \text{or} \quad \alpha = 2\bar{\varepsilon}_i, \quad \beta = \bar{\varepsilon}_i + \bar{\varepsilon}_N, \quad \text{or}, \quad \alpha = \bar{\varepsilon}_i - \bar{\varepsilon}_N, \quad \beta = \bar{\varepsilon}_i + \bar{\varepsilon}_N .$$

For $\alpha, \beta \in \Phi_+^{\text{red}}$, let $\Phi_+^{\text{red}}(<\alpha) = \{\gamma \in \Phi_+^{\text{red}} | \gamma < \alpha\}$,
 $\Phi_+^{\text{red}}(\alpha < \beta) = \{\gamma \in \Phi_+^{\text{red}} | \alpha < \gamma < \beta\}$, $\Phi_+^{\text{red}}(\beta <) = \{\gamma \in \Phi_+^{\text{red}} | \beta < \gamma\}$.
Let $\Phi_{+,i}^{\text{red}} = \{\beta \in \Phi_+^{\text{red}} | g(\beta) = i\}$. For $\alpha, \beta \in \Phi_{+,i}^{\text{red}}$, let
 $\Phi_{+,i}^{\text{red}}(<\alpha) = \Phi_{+,i}^{\text{red}} \cap \Phi_+^{\text{red}}(<\alpha)$, $\Phi_{+,i}^{\text{red}}(\alpha < \beta) = \Phi_{+,i}^{\text{red}} \cap \Phi_+^{\text{red}}(\alpha < \beta)$,
 $\Phi_{+,i}^{\text{red}}(\beta <) = \Phi_{+,i}^{\text{red}} \cap \Phi_+^{\text{red}}(\beta <)$.

§4. Defining relations of N_+ .

4.1. Keep the notation in §2-3. Assume that the triple system $(\mathcal{E}, \Pi = \{\alpha_1, \dots, \alpha_n\}, p)$ satisfies the assumption in 3.1.
Let $A = (a_{ij})_{1 \leq i, j \leq n} = D^{-1}((\alpha_i, \alpha_j))$ be the corresponding Cartan matrix. Let $\tilde{U}_{\sqrt{h}} \mathcal{E}_+^\sigma = \tilde{N}_+ \otimes S[\mathbb{X}^{R'}] \otimes R'^{\langle \sigma \rangle}$ be the topologically free R' -Hopf algebra defined in 2.2 for the triple (\mathcal{E}, Π, p) . The purpose of this section is to define an ideal $\mathcal{I}_+ (\subseteq I_+)$ of \tilde{N}_+ with an explicit set of generators. In §10, we will show $\mathcal{I}_+ = I_+$.

4.2. We define the \mathbb{Z}_2 -graded algebra structure on $\tilde{\mathbb{N}}_+$ such that the parity of E_i is $p(\alpha_i)$. Denote the parity of $X \in \tilde{\mathbb{N}}_+$ by $p(X)$. Put $[X, Y]_v = XY - (-1)^{p(X)p(Y)} v Y X$ and $[X, Y] = [X, Y]_1$ where $p(X)$ and $p(Y)$ are the parities of X and Y . Set $\left[\begin{smallmatrix} m+n \\ n \end{smallmatrix} \right]_t = \prod_{i=0}^{n-1} ((t^{m+n-i} - t^{-m-n+i}) / (t^{i+1} - t^{-i-1})) \in \mathbb{C}[t]$. Put $q = e^h$, $v_i = q^{(\bar{\epsilon}_i, \bar{\epsilon}_i)}$ and $q_i = q^{d_i}$.

Definition 4.2.1. Let \mathcal{I}_+ be the ideal of $\tilde{\mathbb{N}}_+$ generated by the following elements:

- (i) $[E_i, E_j]$ for $1 \leq i, j \leq n$ such that $a_{ij} = 0$,
- (ii) $\sum_{\nu=0}^{1+|a_{ij}|} (-1)^\nu \left[\begin{smallmatrix} 1+|a_{ij}| \\ \nu \end{smallmatrix} \right]_{q_i} E_i^{1+|a_{ij}|-\nu} E_j^\nu E_i$ for $1 \leq i \neq j \leq n$
such that $p(\alpha_i) = 0$,
- (iii) $[[[E_i, E_j]_{v_j}, E_k]_{v_{j+1}}, E_j]$ with $\overset{i}{x} \otimes \overset{j}{x} \otimes \overset{k}{x}$ ($i < j < k$),
(resp. $\overset{i}{x} \otimes \overset{j}{x} \otimes \overset{k}{\bullet}$ or $\overset{i}{x} \otimes \overset{j}{\bullet} \otimes \overset{k}{x}$),
- (iv) $[[[E_{N-1}, E_N]_{v_N}, E_N], E_N]_{v_N-1}$ with $x \Rightarrow \bullet$
if A is of type B_N ,
- (v) $[[E_{N-2}, E_{N-1}]_{v_{N-1}}, E_N]_{v_N} - [[E_{N-2}, E_N]_{v_{N-1}}, E_{N-1}]_{v_N}$
with $\overset{N-2}{x} \otimes \overset{N-1}{x} \otimes \bullet$ if A is of type D_N ,
- (vi) $[[[E_{N-2}, E_{N-1}]_{v_{N-1}}, E_N]_{2v_N}, [E_{N-2}, E_{N-1}]_{v_{N-1}}], E_{N-1}]$
(resp.

$$[[[[[E_{N-3}, E_{N-2}]_{v_{N-2}}, E_{N-1}]_{v_{N-1}}, E_N]_{2v_N}, E_{N-1}]_{v_N}, E_{N-2}]_{v_{N-1}}, E_{N-1}]$$

for $\begin{array}{c} N-2 & N-1 & N \\ \otimes & \otimes & \longleftarrow \end{array}$ (resp. $\begin{array}{c} N-3 & N-2 & N-1 & N \\ \times & \circ & \otimes & \longleftarrow \end{array}$) if A is of type C_N .

4.3. The following proposition will be used in proving

$$\mathcal{I}_+ = I_+$$

Proposition 4.3.1. The R' -submodule

$$\mathcal{I}'_{\mathcal{B}_+} = (\mathcal{I}_+ \otimes R') \cdot S[\mathcal{R}^{R'}] \cdot R' \langle \sigma \rangle \text{ is a bi-ideal of } \mathcal{U}_{\sqrt{h}} \mathcal{B}_+^\sigma.$$

Proof. Here we consider the case of the triple (\mathcal{E}, Π, p)

given by $\begin{array}{ccccc} 1 & 2 & 3 & & \\ \overline{\varepsilon}_1 - \overline{\varepsilon}_2 & \overline{\varepsilon}_2 - \overline{\varepsilon}_3 & \overline{\varepsilon}_3 - \overline{\varepsilon}_4 & \dots & \end{array}$. In this case, the ideal \mathcal{I}_+ is

generated by the elements

$$\begin{aligned} s_{\alpha_1 + \alpha_3} &= [E_1, E_3], \quad s_{2\alpha_2} = E_2^2, \\ s_{2\alpha_1 + \alpha_2} &= E_1^2 E_2 - (q+q^{-1}) E_1 E_2 E_1 + E_2 E_1^2, \\ s_{\alpha_2 + 2\alpha_3} &= E_2 E_3^2 - (q+q^{-1}) E_3 E_2 E_3 + E_3^2 E_2 \end{aligned}$$

and

$$t_{1223} = [w_{123}, E_2]$$

$$\text{where } w_{123} = [[E_1, E_2]_{v_2}, E_3]_{v_3}.$$

By an easy computation, it follows

$$\Delta'(\sigma_v) = \sigma_v \otimes 1 + \exp(\sqrt{h} H_v^-) \cdot \sigma^{p(v)} \otimes \sigma_v$$

for any σ_v . We also have:

$$\begin{aligned} \Delta'(w_{123}) &= w_{123} \otimes 1 \\ &+ (v_2^{-1} - v_2)[E_2, E_3]_{v_3} \cdot \exp(\sqrt{h} H^- \bar{\epsilon}_1 - \bar{\epsilon}_2) \otimes E_1 \\ &+ (v_3^{-1} - v_3)E_3 \cdot \exp(\sqrt{h} H^- \bar{\epsilon}_1 - \bar{\epsilon}_2) \sigma \otimes [E_1, E_2]_{v_2} \\ &\quad \exp(\sqrt{h} H^- \bar{\epsilon}_1 - \bar{\epsilon}_4) \cdot \sigma \otimes w_{123}. \end{aligned}$$

Since the ideal \mathcal{G} generated by the element $\sigma_{2\alpha_2}$ is a bi-ideal, we have:

$$[[E_1, E_2]_{v_2}, E_2]_{v_2} \equiv [E_2, [E_2, E_3]_{v_3}]_{v_3} \equiv 0 \pmod{\mathcal{G}}.$$

Therefore we have:

$$\begin{aligned} \Delta'(t_{1223}) &= \Delta'([w_{123}, E_2]) \\ &= [\Delta'(w_{123}), E_2 \otimes 1 + \exp(\sqrt{h} H^- \bar{\epsilon}_2 - \bar{\epsilon}_3) \cdot \sigma \otimes E_2] \\ &\equiv \{t_{1223} \otimes 1 + 0\} + \\ &\quad \{0 + (v_2^{-1} - v_2)[E_2, E_3]_{v_3} \cdot \exp(\sqrt{h} H^- \bar{\epsilon}_1 - \bar{\epsilon}_3) \cdot \sigma \otimes [E_1, E_2]_{v_2}\} \\ &+ \{(v_3^{-1} - v_3)[E_2, E_3]_{v_3} \cdot \exp(\sqrt{h} H^- \bar{\epsilon}_1 - \bar{\epsilon}_2) \cdot \sigma \otimes [E_1, E_2]_{v_2} + 0\} \\ &+ \{0 + \exp(\sqrt{h} H^- \bar{\epsilon}_1 - \bar{\epsilon}_4) \cdot \sigma \otimes t_{1223}\} \\ &\pmod{\mathcal{G} \otimes \tilde{U}_{\sqrt{h}}^{\sigma} \ell_+^{\sigma} + \tilde{U}_{\sqrt{h}}^{\sigma} \ell_+^{\sigma} \otimes \mathcal{G}} \\ &= t_{1223} \otimes 1 + \exp(\sqrt{h} H^- \bar{\epsilon}_1 - \bar{\epsilon}_4) \cdot \sigma \otimes t_{1223}. \end{aligned}$$

Hence $\Delta'(\mathcal{I}_+) \subset \mathcal{I}_+ \otimes \tilde{U}_{\sqrt{h}}^{\sigma} + \tilde{U}_{\sqrt{h}}^{\sigma} \otimes \mathcal{I}_+$. This implies that $\mathcal{I}_{\tilde{U}_{\sqrt{h}}^{\sigma}}$ is a bi-ideal of $\tilde{U}_{\sqrt{h}}^{\sigma}$. The other cases can be proved similarly.

Denote the h -adic topological R' -bialgebra $\tilde{U}_{\sqrt{h}}^{\sigma}/\mathcal{I}_{\tilde{U}_{\sqrt{h}}^{\sigma}}$ by $\tilde{U}_{\sqrt{h}}^{\sigma}$. Put $\mathcal{N}_+ = \tilde{N}_+/\mathcal{I}_+$ and $\mathcal{N}'_+ = \mathcal{N}_+ \otimes R'$ ($= \tilde{N}'_+/\mathcal{I}'_+$). Then we have:

$$\tilde{U}_{\sqrt{h}}^{\sigma} \simeq \mathcal{N}'_+ \hat{\otimes} S[\mathbb{X}^{R'}] \hat{\otimes} R'<\sigma>.$$

4.4. Put $\mathcal{N}_{+,v} = \tilde{N}_{+,v}/(\mathcal{I}_+ \cap \tilde{N}_{+,v})$ ($\subset \mathcal{N}_+$) for $v \in P_+$.

For $v, \mu \in P_+$ and $x_v \in \mathcal{N}_{+,v}$, $x_\mu \in \mathcal{N}_{+,\mu}$, we put

$$[x_v, x_\mu] = [x_v, x_\mu]_{q^{-(v, \mu)}}.$$

Let $x_v \in \mathcal{N}_{+,v}$, $x_\mu \in \mathcal{N}_{+,\mu}$, $x_\eta \in \mathcal{N}_{+,\eta}$ ($v, \mu, \eta \in P_+$). In §6-§9, we shall frequently use the following identities:

$$(4.4.1) \quad [x_v \cdot x_\mu, x_\eta] \\ = x_v \cdot [x_\mu, x_\eta] + (-1)^{p(\mu)p(\eta)} q^{-(\mu, \eta)} [x_v, x_\mu] \cdot x_\eta.$$

$$(4.4.2) \quad [[X_\nu, X_\mu], X_\eta] - [X_\nu, [X_\mu, X_\eta]] \\ = (-1)^{p(\mu)p(\eta)} q^{-(\mu, \eta)} [X_\nu, X_\eta] \cdot X_\mu \\ - (-1)^{p(\nu)p(\mu)} q^{-(\nu, \mu)} X_\mu \cdot [X_\nu, X_\eta].$$

If $X_\mu^2 = 0$, we have:

$$(4.4.3) \quad [[X_\nu, X_\mu], X_\mu] = 0, \\ [X_\mu, [X_\mu, X_\nu]] = 0.$$

$$(4.4.4) \quad [[[X_\nu, X_\mu], X_\eta], X_\chi] \\ = [X_\nu, [[X_\mu, X_\eta], X_\chi]] \\ + (-1)^{p(\mu+\eta)p(\chi)} q^{-(\mu+\eta, \chi)} [X_\nu, X_\chi] [X_\mu, X_\eta] \\ - (-1)^{p(\nu)p(\mu+\eta)} q^{-(\nu, \mu+\eta)} [X_\mu, X_\eta] [X_\nu, X_\chi] \\ + (-1)^{p(\mu)p(\eta)} q^{-(\mu, \eta)} [X_\nu, X_\eta] [X_\mu, X_\chi] \\ + (-1)^{p(\mu)p(\eta+\chi)} q^{-(\mu, \eta+\chi)} [[[X_\nu, X_\eta], X_\chi], X_\mu] \\ - (-1)^{p(\nu)p(\mu)} q^{-(\nu, \mu)} [X_\mu, [[X_\nu, X_\eta], X_\chi]] \\ - (-1)^{p(\nu)p(\mu)} q^{-(\nu, \mu)} (-1)^{p(\nu+\eta)p(\chi)} q^{-(\nu+\eta, \chi)} [X_\mu, X_\chi] [X_\nu, X_\eta].$$

In particular, if $\chi = \eta$, we have:

$$(4.4.5) \quad [[[X_\nu, X_\mu], X_\eta], X_\eta] \\ = [X_\nu, [[X_\mu, X_\eta], X_\eta]] \\ + (1+(-1)^{p(\eta)} q^{-(\eta, \eta)}) \{ (-1)^{p(\mu)p(\eta)} q^{-(\mu, \eta)} [X_\nu, X_\eta] [X_\mu, X_\eta] \\ - (-1)^{p(\nu)p(\mu+\eta)} q^{-(\nu, \mu+\eta)} [X_\mu, X_\eta] [X_\nu, X_\eta] \} \\ + q^{-(\mu, 2\eta)} [[[X_\nu, X_\eta], X_\eta], X_\mu]$$

$$- (-1)^{p(\nu)p(\mu)} q^{-(\nu, \mu)} [X_\mu, [[X_\nu, X_\eta], X_\eta]] .$$

§5. Root vectors of \mathcal{N}_+ .

5.1. Here we define q -root vectors E_α ($\alpha \in \Phi_+^{\text{red}}$) of \mathcal{N}_+ .

Definition 5.1. For $\beta \in \Phi_+^{\text{red}}$, we define the element

$E_\beta \in \mathcal{N}_+$ as follows. (For type F_4 (resp. G_3), we write E_{abcd} and E'_{abcd} (resp. E_{abc} and E'_{abc}) for $E_{a\alpha_1+b\alpha_4+c\alpha_3+d\alpha_2}$ and $E'_{a\alpha_1+b\alpha_4+c\alpha_3+d\alpha_2}$ (resp. $E_{a\alpha_1+b\alpha_3+c\alpha_2}$ and $E'_{a\alpha_1+b\alpha_3+c\alpha_2}$).

(i) We put $E_{\alpha_i} = E_i$ ($1 \leq i \leq n$).

(ii) Let $\alpha \in \Phi_+^{\text{red}}$ and $1 \leq i \leq n$ be such that $g(\alpha) < i$ (see 3.2 for the definition of $g(\alpha)$) and $\alpha + \alpha_i \in \Phi$. We put $E'_{\alpha+\alpha_i} = [E_\alpha, E_{\alpha_i}] q^{-(\alpha, \alpha_i)}$. If A is of type B_N , $i = N$ and $\alpha = \bar{\epsilon}_j$ ($1 \leq j \leq N-1$), let $E_{\alpha+\alpha_N} = (q^{1/2} + q^{-1/2})^{-1} E'_{\alpha+\alpha_N}$. If A is of type D_N , $i = N$ and $\alpha = \alpha_{N-1}$, let $E_{\alpha+\alpha_N} = (q + q^{-1})^{-1} E'_{\alpha+\alpha_N}$.

If A is of type F_4 , let $E_{1120} = (q + q^{-1})^{-1} E'_{1120}$ and $E_{1232} = (q^2 + 1 + q^{-2})^{-1} E'_{1232}$. If A is of type G_3 , let $E_{121} = (q + q^{-1})^{-1} E'_{121}$, $E_{021} = (q + q^{-1})^{-1} E'_{021}$ and $E_{031} = (q^2 + 1 + q^{-2})^{-1} E'_{031}$. Otherwise, put $E_{\alpha+\alpha_i} = E'_{\alpha+\alpha_i}$.

(iii) For $\alpha, \beta \in \Phi_+^{\text{red}}$ such that $g(\alpha) = g(\beta)$, $\alpha < \beta$, $ht(\beta) - ht(\alpha) \leq 1$ and $\alpha + \beta \in \Phi_+^{\text{red}}$, we put

$E'_{\alpha+\beta} = [E_\alpha, E_\beta] q^{-(\alpha, \beta)}$. If A is of type C_N (resp. D_N , F_4 or G_3), then $E_{\alpha+\beta}$ is defined by $(q + q^{-1})^{-1} E'_{\alpha+\beta}$ (resp.

$$(q+q^{-1})^{-1}E_{\alpha+\beta}^-, (q^2+q^{-2})^{-1}E_{\alpha+\beta}^- \text{ or } (q^2+1+q^{-2})^{-1}E_{\alpha+\beta}^-.$$

5.2. The following lemma will play key role in proving our main results (Theorem 10.6.1).

Lemma 5.2.1. (i) Let $\alpha \in \Phi_{+, i}^{\text{red}}$ (see 3.2 for this definition) and $j > i$. Then we have:

$$[E_\alpha, E_j]_q - (\alpha, \alpha_j) = \sum_{\gamma_1, \dots, \gamma_u \in \Phi_{+, i}^{\text{red}}(\alpha <)} c_{\gamma_1, \dots, \gamma_u} E_{\gamma_1} \cdots E_{\gamma_u}$$

for some $c_{\gamma_1, \dots, \gamma_u} \in R$.

(ii) Let $\alpha, \beta \in \Phi_{+, i}^{\text{red}}$. If $\alpha < \beta$, then we have:

$$[E_\alpha, E_\beta]_q - (\alpha, \beta) = \sum_{\gamma_1, \dots, \gamma_u \in \Phi_{+, i}^{\text{red}}(\alpha < \beta)} c_{\gamma_1, \dots, \gamma_u} E_{\gamma_1} \cdots E_{\gamma_u}$$

for some $c_{\gamma_1, \dots, \gamma_u} \in R$.

(iii) $E_\alpha^2 = 0$ if $(\alpha, \alpha) = 0$.

Remark 5.2.2. In some special cases, we can show more detailed results than Lemma 5.2.1.

(i) Let $\alpha \in \Phi_+^{\text{red}}$ satisfy $c_\alpha = 1$. Take $\alpha_i \in \Pi$ such that $i > g(\alpha)$ and $\alpha + \alpha_i \notin \Phi_+^{\text{red}}$. Then we have:

$$[E_\alpha, E_i]_{q^{-(\alpha, \alpha_i)}} = 0.$$

(ii) Assume that $\alpha, \beta \in \Phi_+^{\text{red}}$ satisfies $\alpha \nless \beta$ and $\beta \nless \alpha$.

Then we have:

$$[E_\alpha, E_\beta] = 0.$$

5.3. Let $\prod_{\alpha \in \Phi_+^{\text{red}}}^<$ denote the product taken with respect to a total order on Φ_+^{red} compatible with the partial order $<$.

As an immediate consequence of Lemma 5.2.1, we have:

Proposition 5.3.1. The R -module N_+ is generated by the elements

$$\prod_{\alpha \in \Phi_+^{\text{red}}}^< E_\alpha^{n_\alpha} \quad (n_\alpha \in \mathbb{Z}_+, \text{ if } (\alpha, \alpha) \neq 0, n_\alpha = 0, 1 \text{ if } (\alpha, \alpha) = 0).$$

5.4. The purpose of §6-9 below, is to prove Lemma 5.2.1 and Remark 5.2.2.

In §10, we shall prove that the monomials $\prod_{\alpha \in \Phi_+^{\text{red}}} E_\alpha^{n_\alpha}$

form an orthogonal basis of \mathcal{N}_+ , which implies $\mathcal{S}_+ = I_+$.

§6. Commutator relations of root vectors of \mathcal{N}_+

(type A_{N-1} , B_N , C_N or D_N).

6.1. In this section, we assume that \mathcal{N}_+ is of type A_{N-1} , B_N , C_N or D_N . Put $\bar{d}_i = (\bar{\varepsilon}_i, \bar{\varepsilon}_i) \in \{1, -1\}$ ($1 \leq i \leq N$).

Lemma 6.1.1. The following identities hold in \mathcal{N}_+ .

$$(i) [E_{\alpha_i + \alpha_j + \alpha_k}, E_j] = 0 \quad \text{for } \begin{matrix} i & j & k \\ \times & \times & \times \end{matrix} \quad (i < j < k),$$

$$\begin{matrix} i & j & k \\ \times & \times & \Rightarrow O \end{matrix} \quad \text{or} \quad \begin{matrix} i & j & k \\ \times & \times & \Rightarrow \bullet \end{matrix}.$$

$$(ii) [E_{\alpha_i + \alpha_j}, E_j] q^{-\bar{d}_j} = [E_i, E_{\alpha_i + \alpha_j}] q^{-\bar{d}_i} = 0 \quad \text{for } \begin{matrix} i & j \\ \times & \times \end{matrix} \quad (i < j).$$

$$(iii) [E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N}, E_{N-1}] q^{-\bar{d}_{N-1}} = [E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N}, E_N] q^{-\bar{d}_{N-1}} = 0 \quad \text{for type } D_N.$$

$$(iv) [E_{\alpha_{N-1} + \alpha_N}, E_{N-1}] q^{-\bar{d}_{N-1}} = [E_{\alpha_{N-1} + 2\alpha_N}, E_N] q^{-\bar{d}_N} = 0 \quad \text{for type } B_N.$$

$$(v) [E_{\alpha_{N-1} + \alpha_N}, E_N] q^{-2\bar{d}_N} = 0,$$

$$[E_{N-1}, E_{2\alpha_{N-1} + \alpha_N}] q^{-2\bar{d}_{N-1}} = 0 \quad \text{for type } C_N.$$

$$(vi) [E_{\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}, E_{N-1}] q^{-\bar{d}_{N-1}} = 0 \quad \text{for type } C_N.$$

$$(vii) [E_{2\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}, E_{N-1}] = 0 \quad (p(\alpha_{N-2} + \alpha_{N-1}) = 0) \quad \text{for type } D_N.$$

C_N .

$$(viii) [E_{\alpha_{N-3}+2\alpha_{N-2}+2\alpha_{N-1}+\alpha_N}, E_{N-1}] = 0 \text{ for type } C_N.$$

Proof. (ii)-(v) These can be proved by direct computations.

We omit the details.

(i) This is the defining relation (iii) of Definition 4.2.1 if $p(\alpha_j) = 1$. If $p(\alpha_j) = 0$, by Definition 4.2.1 (ii), and the formula (ii), we have:

$$\begin{aligned} 0 &= [E_i, E_j^2 E_k - (q+q^{-1}) E_j E_k E_j + E_k E_j^2] \\ &= \{E_{\alpha_i+\alpha_j} E_j + q^{\bar{d}_j} E_j E_{\alpha_i+\alpha_j}\} E_k - (q+q^{-1}) E_{\alpha_i+\alpha_j} E_k E_j \\ &\quad - (q+q^{-1}) q^{\bar{d}_j} E_j E_k E_{\alpha_i+\alpha_j} + E_k \{E_{\alpha_i+\alpha_j} E_j + q^{\bar{d}_j} E_j E_{\alpha_i+\alpha_j}\} \\ &= (q+q^{-1}) E_j E_{\alpha_i+\alpha_j} E_k - (q+q^{-1}) E_{\alpha_i+\alpha_j} E_k E_j \\ &\quad - (q+q^{-1}) q^{\bar{d}_j} E_j E_k E_{\alpha_i+\alpha_j} + q^{\bar{d}_j} (q+q^{-1}) E_k E_{\alpha_i+\alpha_j} E_j \\ &= - (q+q^{-1}) [E_{\alpha_i+\alpha_j+\alpha_k}, E_j]. \end{aligned}$$

(vi) The case $p(\alpha_{N-1}) = 1$ follows from (4.4.3) since, in this case, we have $E_{N-1}^2 = 0$. Assume $p(\alpha_{N-1}) = 0$. By using the facts $p(\alpha_{N-1}) = p(\alpha_N) = 0$ and $\bar{d}_{N-1} = \bar{d}_N$, and, by using Definition 4.2.1 (i-ii), (4.4.1) and (ii), we have:

$$\begin{aligned} 0 &= [E_{N-2}, \{E_{N-1}^3 E_N - (q^2+1+q^{-2}) E_{N-1}^2 E_N E_{N-1} \\ &\quad + (q^2+1+q^{-2}) E_{N-1} E_N E_{N-1}^2 - E_N E_{N-1}^3\}] \end{aligned}$$

$$\begin{aligned}
&= (q^2 + 1 + q^{-2}) \{ E_{N-1}^2 E_{\alpha_{N-2} + \alpha_{N-1}} E_N - q^{2\bar{d}_{N-1}} E_{N-1}^2 E_N E_{\alpha_{N-2} + \alpha_{N-1}} \\
&- (q+q^{-1}) E_{N-1} E_{\alpha_{N-2} + \alpha_{N-1}} E_N E_{N-1} + q^{2\bar{d}_{N-1}} (q+q^{-1}) E_{N-1} E_N E_{\alpha_{N-2} + \alpha_{N-1}} E_{N-1} \\
&+ E_{\alpha_{N-2} + \alpha_{N-1}} E_N E_{N-1}^2 - q^{2\bar{d}_{N-1}} E_N E_{\alpha_{N-2} + \alpha_{N-1}} E_{N-1}^2 \} \\
&= (q^2 + 1 + q^{-2}) \{ E_{N-1}^2 E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} \\
&- (q+q^{-1}) E_{N-1} E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{N-1} + E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{N-1}^2 \} \\
&= (q^2 + 1 + q^{-2}) [E_{\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}, E_{N-1}] .
\end{aligned}$$

(vii) This is the defining relation (vi) of Definition 4.2.1 if $p(\alpha_{N-1}) = 1$. Assume $p(\alpha_{N-1}) = 0$. By (ii), we easily see $[E_{N-2}, E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N}] = 0$. By (vi), similarly to the proof of (i), we have:

$$\begin{aligned}
0 &= [E_{N-2}, [E_{\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}, E_{N-1}]] \\
&= [E_{N-2}, \{ E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{N-1}^2 - (q+q^{-1}) E_{N-1} E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{N-1} \\
&\quad + E_{N-1}^2 E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} \}] \\
&= (q+q^{-1}) \{ E_{N-1} E_{\alpha_{N-2} + \alpha_{N-1}} E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} \\
&- E_{N-1} E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{\alpha_{N-2} + \alpha_{N-1}} - E_{\alpha_{N-2} + \alpha_{N-1}} E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{N-1} \\
&+ E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{\alpha_{N-2} + \alpha_{N-1}} E_{N-1} \} \\
&= -(q+q^{-1})^2 [E_{\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}, E_{N-1}] .
\end{aligned}$$

(viii) If $p(\alpha_{N-2}) = 0$ and $p(\alpha_{N-1}) = 1$, then (viii) is

defining relation (vi) of Definition 4.2.1. Next, we assume $p(\alpha_{N-1}) = 0$. Note that, by (i) and (vi), we have

$$(6.1.2) \quad [\![E_{\alpha_{N-3}+\alpha_{N-2}+\alpha_{N-1}+\alpha_N}, E_{N-2}]\!] = 0,$$

$$(6.1.3) \quad [\![E_{\alpha_{N-3}+\alpha_{N-2}+2\alpha_{N-1}+\alpha_N}, E_{N-1}]\!] = 0$$

respectively. Hence, by (4.4.5) and (ii), we have:

$$\begin{aligned} 0 &= [\![[\![E_{\alpha_{N-3}+\alpha_{N-2}+\alpha_{N-1}+\alpha_N}, E_{N-2}]\!], E_{N-1}]\!] \\ &= (1+(-1))^{\frac{p(\alpha_{N-1})}{q}} \bar{d}_{N-1} - \bar{d}_N \\ &\quad ((-1)^{\frac{p(\alpha_{N-1})p(\alpha_{N-2})}{q}} \bar{d}_{N-1} E_{\alpha_{N-3}+\alpha_{N-2}+2\alpha_{N-1}+\alpha_N} E_{\alpha_{N-2}+\alpha_{N-1}} \\ &\quad - (-1)^{\frac{p(\alpha_{N-3}+\alpha_{N-2}+\alpha_{N-1}+\alpha_N)p(\alpha_{N-2}+\alpha_{N-1})}{q}} \bar{d}_N \\ &\quad E_{\alpha_{N-2}+\alpha_{N-1}} E_{\alpha_{N-3}+\alpha_{N-2}+2\alpha_{N-1}+\alpha_N}) . \end{aligned}$$

Using $p(\alpha_{N-1}) = 0$ and $\bar{d}_{N-1} = \bar{d}_N$, the right hand side equals

$$(q+q^{-1}) [\![E_{\alpha_{N-3}+\alpha_{N-2}+2\alpha_{N-1}+\alpha_N}, [\![E_{N-2}, E_{N-1}]\!]] .$$

Here we used the identity $E_{\alpha_{N-2}+\alpha_{N-1}} = [\![E_{N-2}, E_{N-1}]\!]$. Hence, by (4.4.2) and (6.1.3), this equals

$$(6.1.4) \quad (q+q^{-1}) [\![E_{\alpha_{N-3}+2\alpha_{N-2}+2\alpha_{N-1}+\alpha_N}, E_{N-1}]\!] .$$

Finally, we assume $p(\alpha_{N-2}) = p(\alpha_{N-1}) = 1$. By (i) and (ii), we have:

$$[E_{\alpha_{N-3} + \alpha_{N-2} + \alpha_{N-1}}, E_{\alpha_{N-2} + \alpha_{N-1}}] = 0.$$

From this, we have:

$$\begin{aligned} & [E_{N-3}, E_{2\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}] \\ &= (q+q^{-1})^{-1} [E_{N-3}, (E_{\alpha_{N-2} + \alpha_{N-1}})^2 E_N \\ &\quad - (1+q^{2\bar{d}_N}) E_{\alpha_{N-2} + \alpha_{N-1}} E_N E_{\alpha_{N-2} + \alpha_{N-1}} + q^{2\bar{d}_N} E_N E_{\alpha_{N-2} + \alpha_{N-1}}^2] \\ &= (q+q^{-1})^{-1} \{ (q+q^{-1}) E_{\alpha_{N-2} + \alpha_{N-1}} E_{\alpha_{N-3} + \alpha_{N-2} + \alpha_{N-1}} E_N \\ &\quad - (q+q^{-1}) q^{2\bar{d}_N} E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_N E_{\alpha_{N-2} + \alpha_{N-1}} \\ &\quad - (q+q^{-1}) q^{2\bar{d}_{N-2} + \bar{d}_N} E_{\alpha_{N-2} + \alpha_{N-1}} E_N E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} \\ &\quad + (q+q^{-1}) q^{2\bar{d}_{N-2} + \bar{d}_N} E_N E_{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} E_{\alpha_{N-2} + \alpha_{N-1}} \} \\ &= [E_{\alpha_{N-2} + \alpha_{N-1}}, E_{\alpha_{N-3} + \alpha_{N-2} + \alpha_{N-1} + \alpha_N}] \end{aligned}$$

Hence, by Definition 4.2.1 (vi), Lemma 6.1.1 (iii) and (4.4.2), we have:

$$\begin{aligned} 0 &= [E_{N-3}, [E_{2\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}, E_{N-1}]] \\ &= [[E_{N-3}, E_{2\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}], E_{N-1}] \end{aligned}$$

$$\begin{aligned}
&= [[E_{\alpha_{N-2}+\alpha_{N-1}}, E_{\alpha_{N-3}+\alpha_{N-2}+\alpha_{N-1}+\alpha_N}], E_{N-1}] \\
&= - [E_{\alpha_{N-3}+\alpha_{N-2}+\alpha_{N-1}+\alpha_N}, E_{\alpha_{N-2}+\alpha_{N-1}}]
\end{aligned}$$

But, by (6.1.3) and (4.4.2), this equals

$$[E_{\alpha_{N-3}+2\alpha_{N-2}+2\alpha_{N-1}+\alpha_N}, E_{N-1}]$$

Hence we get the desired formula.

6.2. Similarly to the proof of the formula (i) of Lemma 6.1.1, we obtain the following lemma.

Lemma 6.2.1. For $\overset{i}{x} \overset{j}{x} \overset{k}{x}$ ($i < j < k$) , $\overset{i}{x} \overset{j}{x} \overset{k}{o}$ or $\overset{i}{x} \overset{j}{x} \overset{k}{\bullet}$, the following identities hold:

$$[[[E_i, E_j]_q(\alpha_i, \alpha_j), E_k]_q(\alpha_i + \alpha_j, \alpha_k), E_j] = 0,$$

$$[E_{\alpha_i + \alpha_j}, [E_j, E_k]_q(\alpha_j, \alpha_k)] = 0.$$

6.3. Lemma 6.3.1. For $1 \leq i \leq N-1$, we have the following identities:

(i) For type B_N , we have:

$$[\bar{E}_{\bar{\varepsilon}_i}, \bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}]_{q^{-\bar{d}_i}} = 0 .$$

(ii) For type C_N , we have:

$$[\bar{E}_{2\bar{\varepsilon}_i}, \bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}]_{q^{-2\bar{d}_i}} = 0 \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 0) .$$

In the formulas (iii) - (ix) below, we assume that A is of type D_N .

$$(iii) [\bar{E}_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, \bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}] = 0 \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 0),$$

$$(iv) [\bar{E}_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, \bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-1}}]_{q^{-\bar{d}_i}} = 0 ,$$

$$(v) [\bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}, \bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-1}}]_{q^{-\bar{d}_i}} = 0 ,$$

$$(vi) [\bar{E}_{2\bar{\varepsilon}_i}, \bar{E}_{N-1}] = 0 \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 1),$$

$$(vii) [\bar{E}_{2\bar{\varepsilon}_i}, \bar{E}_N] = \bar{d}_N(q-q^{-1})\bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-1}}\bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_N} \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 1),$$

$$(viii) [\bar{E}_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, \bar{E}_{2\bar{\varepsilon}_i}]_{q^{-2\bar{d}_i}} = 0 \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 1),$$

$$(ix) [\bar{E}_{2\bar{\varepsilon}_i}, \bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}]_{q^{-2\bar{d}_i}} = 0 \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 1) .$$

Proof. (i) By Lemma 6.1.1 (i), (ii), (iv), we can easily show

$$[\bar{E}_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, \bar{E}_{\bar{\varepsilon}_i}] = 0 ,$$

$$[\bar{E}_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}, \bar{E}_N] = 0 .$$

By (4.4.5), we have:

$$\begin{aligned}
0 &= [[[[E_{\bar{\varepsilon}_i} - \bar{\varepsilon}_N, E_{\bar{\varepsilon}_i}], E_N], E_N]] \\
&= (q^{1/2} + q^{-1/2}) \left((1 + (-1)^{\frac{p(\bar{\varepsilon}_N)}{q} - \bar{d}_N}) \{ (-1)^{\frac{p(\bar{\varepsilon}_i)p(\bar{\varepsilon}_N)}{E_{\bar{\varepsilon}_i}E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}}} \right. \\
&\quad - (-1)^{\frac{p(\bar{\varepsilon}_i-\bar{\varepsilon}_N)p(\bar{\varepsilon}_i+\bar{\varepsilon}_N)}{q} - \bar{d}_i + \bar{d}_N} \frac{E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}E_{\bar{\varepsilon}_i}}{E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}E_{\bar{\varepsilon}_i}} \} \\
&\quad + E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}E_{\bar{\varepsilon}_i} - (-1)^{\frac{p(\bar{\varepsilon}_i-\bar{\varepsilon}_N)p(\bar{\varepsilon}_i)}{q} - \bar{d}_i} \frac{E_{\bar{\varepsilon}_i}E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}}{E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}E_{\bar{\varepsilon}_i}} \right) \\
&= (q^{1/2} + q^{-1/2}) \left(-((-1)^{\frac{p(\bar{\varepsilon}_i-\bar{\varepsilon}_N)}{q} - \bar{d}_i} - (-1)^{\frac{p(\bar{\varepsilon}_N)}{-q} - \bar{d}_N}) \right. \\
&\quad \left. (-1)^{\frac{p(\bar{\varepsilon}_i-\bar{\varepsilon}_N)p(\bar{\varepsilon}_N)}{E_{\bar{\varepsilon}_i}E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}}} \right. \\
&\quad + ((-1)^{\frac{p(\bar{\varepsilon}_i-\bar{\varepsilon}_N)}{-q} - \bar{d}_i + \bar{d}_N} - (-1)^{\frac{p(\bar{\varepsilon}_N)}{q} - \bar{d}_i} (-1)^{\frac{p(\bar{\varepsilon}_i-\bar{\varepsilon}_N)}{E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}E_{\bar{\varepsilon}_i}}}) \\
&= - (q^{1/2} + q^{-1/2}) ((-1)^{\frac{p(\bar{\varepsilon}_i-\bar{\varepsilon}_N)}{q} - \bar{d}_i} - (-1)^{\frac{p(\bar{\varepsilon}_N)}{-q} - \bar{d}_N}) \\
&\quad \left. (-1)^{\frac{p(\bar{\varepsilon}_i)p(\bar{\varepsilon}_N)}{E_{\bar{\varepsilon}_i}, E_{\bar{\varepsilon}_i+\bar{\varepsilon}_N}}} \right].
\end{aligned}$$

(ii) This can be proved similarly to (i).

(iii) By Lemma 6.1.1 (i)-(ii), it can be easily proved that

$$(6.3.2) \quad 0 = [[E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1}}, E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}]],$$

$$(6.3.3) \quad 0 = [[E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1}}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}]].$$

If $p(\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1}) = 0$ (resp. $p(\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1}) = 1$) , by

Definition 4.2.1. (i)-(ii) (resp. (v)), we have:

$$0 = [\![E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_N]\!] - [\![E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}, E_{N-1}]\!] .$$

Hence, by (6.3.2-3) and (4.4.2), we have:

$$\begin{aligned} 0 &= [\![E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1}}, ([\![E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_N]\!] - [\![E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}, E_{N-1}]\!])]\!] \\ &= ((-1)^{\frac{p(\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1})p(\bar{\varepsilon}_i - \bar{\varepsilon}_N)}{q}} - \bar{d}_i) \\ &\quad + (-1)^{\frac{p(\bar{\varepsilon}_i + \bar{\varepsilon}_N)p(\bar{\varepsilon}_{N-1} - \bar{\varepsilon}_N)}{q}} \bar{d}_N \} E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N} E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N} \\ &\quad - ((-1)^{\frac{p(\bar{\varepsilon}_i - \bar{\varepsilon}_N)p(\bar{\varepsilon}_{N-1} + \bar{\varepsilon}_N)}{q}} \bar{d}_N \\ &\quad + (-1)^{\frac{p(\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1})p(\bar{\varepsilon}_i + \bar{\varepsilon}_N)}{q}} - \bar{d}_i) \} E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N} E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N} . \end{aligned}$$

Since $p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = p(\bar{\varepsilon}_i + \bar{\varepsilon}_N) = 0$ and $\bar{d}_i = \bar{d}_N$, this equals $(q+q^{-1})[\![E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}]\!]$, which implies (iii).

(iv) By Lemma 6.1.1 (ii), we can easily show:

$$(6.3.4) \quad [\![E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_{N-1}]\!] = 0 .$$

If $p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 0$, by (iii), (6.3.4) and (4.4.2), we have:

$$0 = [\![[\![E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}]\!], E_{N-1}]\!] = [\![E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-1}}]\!] .$$

By (6.3.2) and (6.3.4), we have:

$$0 = [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_N}] .$$

If $p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 1$, this equals $(1+q)^{2\bar{d}_N} E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}^2$. Hence, by (4.4.3), we have (iv).

The formula (v) can be proved quite similarly to (iv).

The formula (vi)-(ix) can be easily proved by using (iii)-(v).

6.4. Lemma 6.4.1. The following identities hold in \mathcal{N}_+ .

- (i) $[E_{\alpha_{N-3} + 2\alpha_{N-2} + \alpha_{N-1} + \alpha_N}, E_N] = 0$ for type D_N .
- (ii) $[E_{\alpha_{N-3} + 2\alpha_{N-2} + \alpha_{N-1} + \alpha_N}, E_{N-1}] = 0$ for type D_N .
- (iii) $[E_{\alpha_{N-2} + 2\alpha_{N-1} + 2\alpha_N}, E_N] = 0$ for type B_N .
- (iv) $[E_{\alpha_{N-2} + 2\alpha_{N-1} + \alpha_N}, E_N] = 0$ for type C_N .

Proof. In the proof, we may assume that $N = 4$. Put

$$E_{abcd} = E_{a\alpha_1 + b\alpha_2 + c\alpha_3 + d\alpha_4}. \text{ Denote } p(\alpha_i) \text{ by } p(i).$$

(i) We are in one of the following three cases.

- (1) $p(3) = p(4) = 0$,
- (2) $p(2) = 0$,
- (3) $p(2) = p(3) = p(4)$.

First consider the case (1). From (4.4.5), we have:

$$\begin{aligned}
 0 &= [[E_{1110}, E_2], E_4] \\
 &= [E_{1110}, [E_{0101}, E_4]] \\
 &+ (1 + (-1)^{p(4)} q^{-\bar{d}_3 - \bar{d}_4}) \{ (-1)^{p(2)p(4)} q^{\bar{d}_3} E_{1111} E_{0101} \\
 &- (-1)^{p(123)p(24)} q^{\bar{d}_4} E_{0101} E_{1111} \} \\
 &+ q^{2\bar{d}_4} [E_{1111}, E_4] E_2 - (-1)^{p(123)p(2)} E_2 [E_{1111}, E_4] .
 \end{aligned}$$

By Definition 4.2.1, we can easily show $[E_{1111}, E_4] = 0$. Since $p(3) = p(4) = 0$ and $\bar{d}_3 = \bar{d}_4$, the right hand side equals

$$(q + q^{-1}) [E_{1111}, E_{0101}] .$$

Using (4.4.2) and Lemma 6.1.1 (iii), we have:

$$(6.4.2) \quad [E_{1111}, E_{0101}] = [E_{1111}, [E_2, E_4] q^{\bar{d}_3}] = [E_{1211}, E_4] .$$

Hence we have $[E_{1211}, E_4] = 0$.

The case (2); by Lemma 6.3.1 (v), $[E_{0101}, E_{0111}] q^{-\bar{d}_2} = 0$.

Similarly to the case (1), by (4.4.2) and (4.4.4), we have:

$$\begin{aligned}
0 &= [[[[E_{1101}, E_2], E_3], E_4] - [E_1, [E_{0101}, E_{0111}]] \\
&= \{ [[E_{1101}, E_{0111}] + (-1)^{p(23)p(4)} q^{\bar{d}_4} [E_{1101}, E_4] E_{0110} \\
&\quad - (-1)^{p(124)p(23)} q^{\bar{d}_4} E_{0110} [E_{1101}, E_4] \\
&\quad + (-1)^{p(2)p(3)} q^{\bar{d}_3} E_{1111} E_{0101} + (-1)^{p(2)p(34)} q^{2\bar{d}_3} [E_{1111}, E_4] E_2 \\
&\quad - (-1)^{p(124)p(2)} E_2 [E_{1111}, E_4] \\
&\quad - (-1)^{p(124)p(2)} (-1)^{p(1234)p(4)} q^{-\bar{d}_3} E_{0101} E_{1111} \} \\
&\quad - \{ [[E_{1101}, E_{0111}] - (-1)^{p(24)p(234)} q^{-\bar{d}_2} E_{1111} E_{0101} \\
&\quad + (-1)^{p(1)p(24)} q^{\bar{d}_2} E_{0101} E_{1111} \}.
\end{aligned}$$

By Definition 4.2.1, it can be easily shown that

$$[[E_{1101}, E_4]] = [[E_{1111}, E_4]] = 0.$$

Since $p(2) = 0$, $p(3) = p(4)$ and $\bar{d}_3 = \bar{d}_2$, the above equals

$$\begin{aligned}
&\{ q^{\bar{d}_2} E_{1111} E_{0101} - (-1)^{p(1)p(4)} q^{-\bar{d}_2} E_{0101} E_{1111} \} \\
&+ \{ q^{-\bar{d}_2} E_{1111} E_{0101} + (-1)^{p(1)p(4)} q^{\bar{d}_2} E_{0101} E_{1111} \} \\
&= (q+q^{-1}) [[E_{1111}, E_{0101}]].
\end{aligned}$$

Hence, similarly to (6.4.2), we have $[E_{1211}, E_4] = 0$.

Finally assume that we are in the case (3). We can easily show that $E_{0101}^2 E_1 - (q+q^{-1}) E_{0101} E_1 E_{0101} + E_1 E_{0101}^2 = 0$ and

$E_{0101}^2 E_3 - (q+q^{-1})E_{0101} E_3 E_{0101} + E_3 E_{0101}^2 = 0$. Then, similarly to Lemma 6.1.1 (i), we can prove our formula.

(ii) The proof of (ii) is quite similar to that of (i).

(iii) The proof is similarly to the case (2) in the proof of (i). By Lemma 6.4.1 (i), we have $[E_{0011}, E_{0012}] = 0$. Hence, by (4.4.2) and (4.4.5), we have:

$$\begin{aligned}
 0 &= [[E_{0111}, E_3], E_4] \frac{-\bar{d}_4}{q^{d_4}}, E_4 \\
 &\quad - (q^{1/2} + q^{-1/2}) [E_2, [E_{0011}, E_{0012}]] \\
 &= \{ [E_{0111}, (q^{1/2} + q^{-1/2}) E_{0012}] \\
 &\quad + (1 + (-1)^{p(4)} q^{-\bar{d}_4}) ((-1)^{p(3)p(4)} q^{\bar{d}_4} (q^{1/2} + q^{-1/2}) E_{0112} E_{0011} \\
 &\quad - (-1)^{p(234)p(34)} (q^{1/2} + q^{-1/2}) E_{0011} E_{0112}) \\
 &\quad + q^{2\bar{d}_4} (q^{1/2} + q^{-1/2}) [E_{0112}, E_4] E_3 \\
 &\quad + (-1)^{p(234)p(3)} (q^{1/2} + q^{-1/2}) [E_{0112}, E_4] E_3 \} \\
 &\quad - (q^{1/2} + q^{-1/2}) \{ [E_{0111}, E_{0012}] - (-1)^{p(34)p(344)} q^{-\bar{d}_3} E_{0112} E_{0011} \\
 &\quad + (-1)^{p(2)p(34)} q^{\bar{d}_3} E_{0011} E_{0112} \}
 \end{aligned}$$

By Lemma 6.1.1 (iv), we can easily show $[E_{0112}, E_4] = 0$. Hence the above equals

$$\begin{aligned}
 &\{ (1 + (-1)^{p(4)} q^{-\bar{d}_4}) ((-1)^{p(3)p(4)} q^{\bar{d}_4} (q^{1/2} + q^{-1/2}) E_{0112} E_{0011} \\
 &\quad - (-1)^{p(234)p(34)} (q^{1/2} + q^{-1/2}) E_{0011} E_{0112}) \}
 \end{aligned}$$

$$\begin{aligned}
& - (q^{1/2} + q^{-1/2}) \{ -(-1)^{p(34)p(3)} q^{-\bar{d}_3} E_{0112} E_{0011} \\
& + (-1)^{p(2)p(34)} q^{\bar{d}_3} E_{0011} E_{0112} \} \\
& = (q^{1/2} + q^{-1/2}) \{ (-1)^{p(3)p(4)} (q^{\bar{d}_4} + (-1)^{p(4)} + (-1)^{p(3)} q^{-\bar{d}_3}) E_{0112} E_{0011} \\
& - (-1)^{p(234)p(34)} (1 + (-1)^{p(4)} q^{-\bar{d}_4} + (-1)^{p(34)} q^{\bar{d}_3}) E_{0011} E_{0112} \\
& = (q^{1/2} + q^{-1/2}) (q^{\bar{d}_4} + (-1)^{p(4)} + (-1)^{p(3)} q^{-\bar{d}_3}) \\
& (-1)^{p(3)p(4)} [E_{0112}, E_{0011}] .
\end{aligned}$$

Hence, similarly to (6.4.2), we have $[E_{0122}, E_4] = 0$.

(iv) By Lemma 6.1.1 (v), we have

$[E_{0011}, E_4]_{q^{-2\bar{d}_4}} = [E_{0111}, E_4]_{q^{-2\bar{d}_4}} = 0$. Therefore, using (4.4.5), we have:

$$\begin{aligned}
0 &= [[E_{0110}, E_3], E_4], E_4] \\
&= (1 + (-1)^{p(4)} q^{-4\bar{d}_4}) ((-1)^{p(3)p(4)} q^{2\bar{d}_4} E_{0111} E_{0011} \\
&- (-1)^{p(23)p(34)} q^{\bar{d}_4} E_{0011} E_{0111}) .
\end{aligned}$$

Since $p(4) = 0$, this equals

$$(q^2 + q^{-2}) [E_{0111}, E_{0011}] .$$

Hence, by Lemma 6.1.1 (v) and (4.4.2), we have

$$0 = [E_{0111}, E_{0011}] = [E_{0111}, [E_3, E_4]] = [E_{0121}, E_4] .$$

This completes the proof of Lemma 6.4.1.

6.5. Lemma 6.5.1. Let \mathcal{M}_+ be of type C_N . Let $i \in \{1, \dots, N-1\}$. We have:

$$(i) \quad [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-1}}, E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}] = 0 \quad (1 \leq i \leq N-3) .$$

$$(ii) \quad [E_{2\bar{\varepsilon}_i}, E_{N-1}] = 0 \quad (1 \leq i \leq N-2) \text{ if } p(\bar{\varepsilon}_i - \bar{\varepsilon}_{N-1}) = 0 .$$

Proof. (i) If $i = N-2$, this is proved easily by using Lemma 6.1.1 (ii), (vii). Assume $i \leq N-3$. By Lemma 6.1.1 (i)-(ii) (resp. (viii)), we can easily show that

$$(6.5.2) \quad 0 = [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{N-2}}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-1}}]$$

(resp.

$$(6.5.3) \quad 0 = [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-2}}, E_{N-1}] .$$

Using (4.4.2), from (6.5.3), we obtain:

$$0 = [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{N-1}}, E_{\bar{\varepsilon}_{N-2} - \bar{\varepsilon}_N}] .$$

Hence, using this and (6.5.2), we have (i) .

(ii) By Lemma 6.1.1 (ii), we can easily show that

$[\bar{E}_{\bar{\varepsilon}_i} - \bar{E}_{\bar{\varepsilon}_N}, E_{N-1}] = 0$. From this and the formula (i) , we can immediately prove (ii).

6.6. Lemma 5.2.1 (and Remark 5.2.2) for Φ_+^{red} of type A_{N-1} , B_N , C_N or D_N can be proved using lemmas in 6.1 - 6.5. In 6.7 - 6.9 below, this will be done only in some special cases. In the remaining cases, the proof can be done similarly and more easily.

6.7. Proof of Lemma 5.2 (i) for Φ_+^{red} of type A_{N-1} , B_N , C_N or D_N . Here we give a proof in the case when Φ_+^{red} is of type B_N and $\alpha = \bar{\varepsilon}_i + \bar{\varepsilon}_k$ ($i+1 < k < N-1$) . The other case can be treated similarly.

Since $[E_i, E_j] = 0$ if $|i-j| \geq 2$, by (4.4.2) , we have

$$(6.7.1) \quad \bar{E}_{\bar{\varepsilon}_i} = [[\bar{E}_{\bar{\varepsilon}_i} - \bar{E}_{\bar{\varepsilon}_{u-1}}, \bar{E}_{\bar{\varepsilon}_{u-1}} - \bar{E}_{\bar{\varepsilon}_{u+2}}]_{q \bar{d}_{u-1}}, \bar{E}_{\bar{\varepsilon}_{u+2}}]_{q \bar{d}_{u+2}}$$

if $i < u < N$.

Hence, by Lemma 6.1.1 (i), we have:

$$(6.7.2) \quad [\bar{E}_{\bar{\varepsilon}_i}, E_u] = 0 \quad \text{if } i < u < N .$$

Hence, putting $\bar{E}_{\bar{\varepsilon}_k}^V = [\dots [E_N, E_{N-1}]_{q \bar{d}_N} \dots, E_{k+1}]_{q \bar{d}_{k+2}}, E_k]_{q \bar{d}_{k+1}}$,

we have:

$$(6.7.3) \quad E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k} = (q^{1/2} + q^{-1/2})^{-1} [E_{\bar{\varepsilon}_i}, E_{\bar{\varepsilon}_k}^\vee].$$

Similarly to (6.7.2), using Lemma 6.2.1 (i), we have:

$$(6.7.4) \quad [E_{\bar{\varepsilon}_k}^\vee, E_u] = 0 \quad \text{if } k < u < N.$$

Hence, we have:

$$(6.7.5) \quad [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}, E_u] = 0 \quad \text{if } i < u < k-1 \quad \text{or} \quad k < u < N.$$

By Definition 5.1.1, we have:

$$[E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}, E_{k-1}]_{q\bar{d}_k} = E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{k-1}}.$$

By Definition 4.2.1 (i), (ii), it can be easily shown that

$$[E_{\bar{\varepsilon}_k}^\vee, E_k]_{q^{-\bar{d}_k}} = 0.$$

Hence, we have:

$$(6.7.6) \quad [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}, E_k]_{q^{-\bar{d}_k}} = 0.$$

By (6.7.2), putting

$$E_{\bar{\varepsilon}_k - \bar{\varepsilon}_{N-1}}^V = [\dots [E_{N-2}, E_{N-3}]_{q \bar{d}_{N-3}}, \dots, E_{k+1}]_{q \bar{d}_{k+2}}, E_k]_{q \bar{d}_{k+1}},$$

we have:

$$E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k} = [[E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{k-1}}, E_{\bar{\varepsilon}_{k-1} + \bar{\varepsilon}_{N-1}}]_{q \bar{d}_{k-1}}, E_{\bar{\varepsilon}_k - \bar{\varepsilon}_{N-1}}]_{q \bar{d}_{N-1}}.$$

Hence, by Lemma 6.4.1 (iii), we have:

$$(6.7.7) \quad [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}, E_N] = 0,$$

as required. Remark 5.2.2 (i) also follows from this and (6.7.5).

6.8. *Proof of Lemma 5.2 (ii) for Φ_+^{red} of type A_{N-1} , B_N , C_N or D_N .* Here we give a proof in the case when Φ_+^{red} is of type B_N and $\beta = \bar{\varepsilon}_i + \bar{\varepsilon}_k$ ($i+1 < k < N-1$) .

By Lemma 6.1.1 (ii) and (6.7.2), we have:

$$(6.8.1) \quad [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_u}, E_{\bar{\varepsilon}_i}]_{q \bar{d}_i} = 0 \quad \text{if } i < u \leq N.$$

Similarly to (6.7.1), we have

$$\begin{aligned} E_{\bar{\varepsilon}_i - \bar{\varepsilon}_u} &= [[E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{k-1}}, E_{\bar{\varepsilon}_{k-1} - \bar{\varepsilon}_{k+1}}]_{q \bar{d}_{k-1}}, E_{\bar{\varepsilon}_{k+1} - \bar{\varepsilon}_u}]_{q \bar{d}_{k+1}}, \\ E_{\bar{\varepsilon}_k}^V &= [E_{\bar{\varepsilon}_{k+2}}, [E_{k+1}, E_k]_{q \bar{d}_k}]_{q \bar{d}_{k+2}} \end{aligned}$$

if $i < k < u \leq N$.

Hence, by Lemma 6.2.1 (ii) and (6.7.4), we have:

$$(6.8.2) \quad [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_u}, E_{\bar{\varepsilon}_k}^\vee] = 0 \quad \text{if } k < u \leq N.$$

By (6.7.3) and (6.8.1-2), we have:

$$(6.8.3) \quad [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_u}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}]_{q^{-d_i}} = 0$$

if $i < u < k-1$ or $k < u \leq N$.

If $\alpha = \bar{\varepsilon}_i - \bar{\varepsilon}_k$, then we can inductively show the formula by using the following fact.

$$\begin{aligned} & [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_k}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}]_{q^{-d_k - d_i}} \\ &= [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_k}, [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{k+1}}, E_{\bar{\varepsilon}_k}]_{q^{-d_{k+1}}}]_{q^{-d_k - d_i}} \\ &= (-1)^{p(\bar{\varepsilon}_i + \bar{\varepsilon}_{k+1})p(\bar{\varepsilon}_k - \bar{\varepsilon}_{k+1})} (q^{-d_{k+1} - q^{-d_{k+1}}}) E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{k+1}} E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{k+1}} \\ &\quad - (-1)^{p(\bar{\varepsilon}_i + \bar{\varepsilon}_{k+1})p(\bar{\varepsilon}_k - \bar{\varepsilon}_{k+1})} q^{-d_{k+1}} [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_{k+1}}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{k+1}}]_{q^{-d_{k+1} - d_i}}. \end{aligned}$$

Since $E_{\bar{\varepsilon}_i} = [E_{\bar{\varepsilon}_i - \bar{\varepsilon}_N}, E_N]_{q^{-d_N}}$, by (6.7.7) and (6.8.3), we have:

$$(6.8.4) \quad [E_{\bar{\varepsilon}_i}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}]_{q^{-d_i}} = 0.$$

By (6.7.5), (6.7.7) and (6.8.4), we have

$$(6.8.5) \quad [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_u}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}]_{q^{-\bar{d}_i}} = 0 \quad \text{if } k < u \leq N.$$

Other cases can be shown similarly.

6.9. *Proof of Lemma 5.2 (iii) for Φ_+^{red} of type A_{N-1} , B_N , C_N or D_N .* Here we give a proof in the case when Φ_+^{red} is of type B_N and $\alpha = \bar{\varepsilon}_i + \bar{\varepsilon}_k$ ($i+1 < k < N-1$) .

By Definition 5.1.1 and (6.7.6), (6.8.5), we have:

$$\begin{aligned} 0 &= [[E_{\bar{\varepsilon}_i + \bar{\varepsilon}_{K+1}}, E_k]_{q^{-\bar{d}_k}}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}]_{q^{-\bar{d}_k - \bar{d}_i}} \\ &= [E_{\bar{\varepsilon}_i + \bar{\varepsilon}_K}, E_{\bar{\varepsilon}_i + \bar{\varepsilon}_k}]_{q^{-\bar{d}_k - \bar{d}_i}} \\ &= (1 - (-1)^{p(\bar{\varepsilon}_i + \bar{\varepsilon}_k)} q^{-\bar{d}_k - \bar{d}_i}) E_{\bar{\varepsilon}_i + \bar{\varepsilon}_K}^2. \end{aligned}$$

Since $p(\bar{\varepsilon}_i + \bar{\varepsilon}_k) = 1$, we have $E_{\bar{\varepsilon}_i + \bar{\varepsilon}_K}^2 = 0$.

Other cases can be proved similarly.

§7. Braid group action on the quantized enveloping algebra
 $U_h(G)$ of a simple Lie algebra G .

7.1. In this section, we briefly explain the braid group action on $U_h(G)$ introduced by Lusztig [9] and [10].

Let $(\mathcal{E}, \Pi = \{\alpha_1, \dots, \alpha_n\}, p)$ be a triple system. Assume that Π is the set of the simple roots of a complex simple Lie algebra G and that $p(\alpha_i) = 0$ for any α_i . Put

$$d_i = (\alpha_i, \alpha_i)/2 \quad (1 \leq i \leq n) \quad \text{and} \quad D = \text{diag}(d_1, \dots, d_n).$$

$$s_i \in GL(\mathcal{E}) \quad \text{be such that} \quad s_i(x) = x - \frac{2(\alpha_i, x)}{(\alpha_i, \alpha_i)} \alpha_i \quad (x \in \mathcal{E}). \quad \text{Let}$$

\mathbb{W} be the Weyl group, i.e., the group generated by the elements s_i ($1 \leq i \leq n$). Let $U_h^\sigma = U_h^\sigma((\mathcal{E}, \Pi, p), D)$ be the h -adic R -Hopf algebra defined in Theorem 2.9.4. Put $F_i = E_i^\circ$. Then

Drinfeld's [2] $U_h(G)$ is equal to the unital subalgebra of U_h^σ h -adically generated by the elements E_i, F_i ($1 \leq i \leq n$), $H \in \mathcal{H}$ (see Theorem 2.10.1). We have $U_h^\sigma = U_h(G) \oplus R<\sigma>$. We put

$$U_h(\mathcal{E}, \Pi) = U_h(G).$$

Let $d_i = \frac{(\alpha_i, \alpha_i)}{2}$ ($1 \leq i \leq n$) and $q_i = \exp(hd_i)$. Put $K_i = \exp(hH_{\alpha_i})$ and $E_i^{(r)} = E_i^r / [r]_{q_i}!$, $F_i^{(r)} = F_i^r / [r]_{q_i}!$

where $[r]_{q_i}! = \prod_{v=1}^r \frac{q_i^v - q_i^{-v}}{q_i - q_i^{-1}}$. In [9] and [10], Lusztig

introduced a braid group action on $U_h(\mathcal{E}, \Pi)$.

Proposition 7.1.1. ([9] and [10]) (i) For any $1 \leq i \leq n$, there is a unique algebra automorphism T_i (resp. T_i^{-1}) on $U_h(G)$ such that

$$\begin{aligned}
T_i(E_i) &= -F_i K_i, \quad (\text{resp. } T_i^{-1}(E_i) = -K_i^{-1} F_i), \\
T_i(F_i) &= -K_i^{-1} E_i, \quad (\text{resp. } T_i^{-1}(F_i) = -E_i K_i), \\
T_i(E_j) &= \sum_{r+s=-a_{ij}} (-1)^r q_i^{-s} E_i^{(r)} E_j E_i^{(s)} = 0, \\
(\text{resp. } T_i^{-1}(E_j) &= \sum_{r+s=-a_{ij}} (-1)^r q_i^{-s} E_i^{(s)} E_j E_i^{(r)} = 0) \quad (i \neq j), \\
T_i(F_j) &= \sum_{r+s=-a_{ij}} (-1)^r q_i^s F_i^{(s)} E_j E_i^{(r)} = 0, \\
(\text{resp. } T_i^{-1}(F_j) &= \sum_{r+s=-a_{ij}} (-1)^r q_i^s E_i^{(r)} E_j E_i^{(s)} = 0) \quad (i \neq j). \\
T_i(H_\lambda) &= H_{s_i(\lambda)} \quad (\text{resp. } T_i^{-1}(H_\lambda) = H_{s_i(\lambda)}) \quad (\lambda \in \delta).
\end{aligned}$$

(ii) T_i, s satisfy the braid relations:

$$\begin{array}{c}
T_i T_j T_i \cdots = T_i T_j T_i \cdots \quad (1 \leq i \neq j \leq n) \\
(\mathfrak{m}_{ij}) \quad (\mathfrak{m}_{ij})
\end{array}$$

where $\mathfrak{m}_{ij} = 2 + 4(\alpha_i, \alpha_j)^2 / (\alpha_i, \alpha_i)(\alpha_j, \alpha_j)$. In particular, for any $w \in W$, there is a unique element T_w such that

$$T_w = T_{i_1} \cdots T_{i_r} \quad \text{for any reduced expression } w = s_{i_1} \cdots s_{i_r}.$$

7.2. Let Φ ($\subset \delta$) be the set of the roots of G and Φ_+ the set of the positive roots with respect to π . Put $N_- = N_+^\circ$. Lusztig proved:

Proposition 7.2.1. ([9] and [10]) (i) If $w \in W$ and


$\alpha_i \in \Pi$ satisfies $w(\alpha_i) \in \Phi_+$, then $T_w(E_i) \in N_+$ and $T_w(F_i) \in N_-$.

(ii) If $w \in W$ and $\alpha_i, \alpha_j \in \Pi$ satisfies $w(\alpha_i) = \alpha_j$, then $T_w(E_i) = E_j$ and $T_w(F_i) = F_j$.

§8. Commutator relations for root vectors of N_+ (type F_4).

8.1. First, in the subsections 8.1-3, we treat

N_+ ($\subset U_h(G(F_4))$) associated to the complex simple Lie algebra $G(F_4)$ of type F_4 . We denote this N_+ by N_+ . In §8, the symbols $\Pi, \alpha_i, I_+, E_i, \dots$ respectively mean $\Pi, \alpha_i, I_+, E_i, \dots$ defined for the simple Lie algebra $G(F_4)$. So, for example, Π is the set of simple roots of $G(F_4)$ in an Euclidean space \mathcal{E} . Namely $\Pi = \{\ddot{\alpha}_1, \ddot{\alpha}_2, \ddot{\alpha}_3, \ddot{\alpha}_4\}$ with $\ddot{\alpha}_1 = \varepsilon_2 - \varepsilon_3, \ddot{\alpha}_2 = \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 - \varepsilon_4), \ddot{\alpha}_3 = \varepsilon_4,$ $\ddot{\alpha}_4 = \varepsilon_3 - \varepsilon_4$ where ε_i ($1 \leq i \leq 4$) is a basis of \mathcal{E} satisfying $(\varepsilon_i, \varepsilon_j) = \delta_{ij}$. The Dynkin diagram of (\mathcal{E}, Π) is given by:



8.2. Let Φ_+ be the set of positive roots of $G(F_4)$. Put $\Phi_{+,1} = \{\beta = \Phi_+ \mid \beta = \ddot{\alpha}_1 + n_4 \ddot{\alpha}_4 + n_3 \ddot{\alpha}_3 + n_2 \ddot{\alpha}_2\}$. Then the number of elements $\Phi_{+,1}$ is equal to 15. Define $w_1 \in W$ by the following reduced expression:

$$(8.2.1) \quad w_1 = s_1 s_2 s_3 \cdot s_2 s_4 \cdot s_1 s_3 \cdot s_2 \cdot s_3 s_1 \cdot s_4 s_2 \cdot s_3 s_2 s_1 \quad .$$

The following lemma can be verified directly.

Lemma 8.2.2. For $1 \leq t \leq 15$, let s_{i_t} be the t -th generator in the reduced expression (8.2.1) of w_1 . We put $\beta_t = s_{i_1} \cdots s_{i_{t-1}} (\alpha_{i_t})$. Then $\Phi_{+,1} = \{\beta_t \ (1 \leq i \leq 15)\}$.

For $\beta_t = s_{i_1} \cdots s_{i_{t-1}} (\alpha_{i_t}) \in \Phi_{+,1}$, put $e_{\beta_t} = T_{i_1} \cdots T_{i_{t-1}} (E_{i_t})$. By Proposition 7.2.1, we see that $e_{\beta_t} \in N_+$.

8.3. By using Proposition 7.2.1, and reducing to rank 2 cases, we can obtain the following identities. Here we put $e_{abcd} = e_{a\alpha_1 + b\alpha_4 + c\alpha_3 + d\alpha_2}$. See also [10].

Lemma 8.3.1. The following identities hold.

$$\begin{aligned} -[e_{1110}, E_2]_q^{-1} &= e_{1111}, \quad [e_{1111}, e_{1120}] = 0, \\ -[e_{1120}, E_2]_q^{-2} &= e_{1121}, \\ -[e_{1111}, E_4] &= 0, \quad -[e_{1111}, E_3]_q^{-1} = e_{1121}, \end{aligned}$$

$$\begin{aligned} -[e_{1220}, E_2]_q^{-2} &= e_{1221}, [e_{1121}, e_{1220}] = 0, \\ -[e_{1121}, E_4]_q^{-2} &= e_{1221}, [e_{1121}, E_3]_q = 0, \\ [e_{1221}, E_4]_q^{-2} &= 0, -[e_{1221}, E_3]_q = e_{1231}, \\ [e_{1231}, E_4] &= 0, -[e_{1231}, E_3]_q = 0. \end{aligned}$$

8.4. For type F_4 , we use the following fact.

Lemma 8.4.1. Let \mathcal{N}_+ be the R -algebra defined for the distinguished triple system (δ, π, p) of type F_4 (see §4). Let $v = n_1\alpha_1 + n_4\alpha_4 + n_3\alpha_3 + n_2\alpha_2 \in P_+$ be such that $n_2 = 0$ or 1. Then there exists an R -module isomorphism $j_v : N_{+,v} \rightarrow \mathcal{N}_{+,v}$ such that $j_v(E_{i_1} \cdots E_{i_u}) = E_{i_1} \cdots E_{i_u}$ for any monomial $E_{i_1} \cdots E_{i_u}$ ($\alpha_{i_1} + \cdots + \alpha_{i_u} = v$) in $N_{+,v}$.

Proof. By Theorem 2.10.1, $I_+ = (y_{ij} \ (i \neq j))$ where $y_{ij} \ (i \neq j)$ are elements given in (2.10.2). Let \mathcal{I}_+ be the ideal of \tilde{N}_+ defined in Definition 4.2.1 for type F_4 . Then, for $v \in P_+$, $N_{+,v} = \tilde{N}_{+,v}/(I_+ \cap \tilde{N}_{+,v})$, $\mathcal{N}_{+,v} = \tilde{N}_{+,v}/(\mathcal{I}_+ \cap \tilde{N}_{+,v})$. The lemma now follows by observing that $I_+ \cap \tilde{N}_{+,v} = \mathcal{I}_+ \cap \tilde{N}_{+,v}$ if $v = n_1\alpha_1 + n_4\alpha_4 + n_3\alpha_3 + n_2\alpha_2 \in P_+$ with $n_2 = 0$ or 1.

8.5. By Lemma 8.3.1, we can easily show:

Lemma 8.5.1. Let

$\alpha = a\alpha_1 + b\alpha_4 + c\alpha_3 + d\alpha_2 \in \Phi_{+,1}^{\text{red}} \setminus \{\alpha_1 + 2\alpha_4 + 3\alpha_3 + 2\alpha_2\}$. Then we have

$$j_v(e_{a\alpha_1 + b\alpha_4 + c\alpha_3 + d\alpha_2}) = -(-1)^{a+b+c+d} E_{a\alpha_1 + b\alpha_4 + c\alpha_3 + d\alpha_2}.$$

8.6. Proof of Lemma 5.2.1 (i) for Φ_{+}^{red} of type F_4 . Here we put $E_{abcd} = E_{a\alpha_1 + b\alpha_4 + c\alpha_3 + d\alpha_2}$. By Lemma 8.3.1, Lemma 8.4.1 and Lemma 5.2.1 (i) for type B_3 and C_3 (see §6), it is enough to show:

$$(8.6.1) [E_\alpha, E_2]_{q^{-(\alpha, \alpha_2)}} = 0$$

for $E_\alpha = E_{1111}, E_{1121}, E_{1221}, E_{1232}$,

and

$$(8.6.2) [E_{1232}, E_4] = [E_{1232}, E_3] = 0.$$

Since $E_2^2 = 0$ and $E_\alpha \in R[E_{\alpha-\alpha_2}, E_2]_{q^{-(\alpha-\alpha_2, \alpha_2)}}$, the formulas (8.6.1) follow from (4.4.3).

Since $E_{1232} = (q^2 + 1 + q^{-2})^{-1} [E_{1231}, E_2]_{q^{-3}}$, by Lemma 8.3.1 and Lemma 8.4.1, we have $[E_{1232}, E_4] = 0$. By Lemma 8.3.1, Lemma 8.4.1, (8.6.1) and (4.4.5), we have:

$$\begin{aligned} 0 &= [[[E_{1221}, E_2]_{q^{-2}}, E_3]_{q^{-2}}, E_3] \\ &= (q + q^2) \{ q^{-1} E_{1231} E_{0011} + q^{-3} E_{0011} E_{1231} \} \end{aligned}$$

$$= (q+q^{-1}) [E_{1231}, E_{0011}]_{q^{-2}} = (q+q^{-1}) [E_{1232}, E_3] .$$

8.7. *Proof of Lemma 5.2.1 (iii) for type F_4* . By Lemma 5.2.1 (iii) for type C_3 (see §6), it is enough to show:

$$(8.7.1) \quad E_\alpha^2 = 0 \quad \text{if} \quad \alpha \in \Phi_{+,1}^{\text{red}} \quad \text{and} \quad (\alpha, \alpha) = 0 .$$

We show (8.7.1) by the induction on $\text{ht}(\alpha)$.

Since $E_2^2 = 0$ and $E_{1111} = [E_{1110}, E_2]_{q^{-1}}$,

$[E_{1111}, E_2]_{q^{-1}} = 0$. Hence, by Lemma 8.3.1 and Lemma 8.4.1,

$$0 = [[E_{1110}, E_2]_{q^{-1}}, E_{1111}]_{q^2} = (1+q^2) E_{1111}^2 .$$

If $\text{ht}(\alpha) > 4$, $E_\alpha = [E_\beta, E_i]_{q^{-(\alpha, \alpha_i)}}$ for some $i \in \{2, 3, 4\}$

and $\beta \in \Phi_{+,1}^{\text{red}}$ such that $(\beta, \beta) = 0$. In this case,

$[E_\alpha, E_i]_{q^{-(\alpha, \alpha_i)}} = 0$ by Lemma 8.3.1 and Lemma 8.4.1, and

$[E_\beta, E_\alpha]_{q^{-(\alpha, \beta)}} = 0$ since $E_\beta^2 = 0$ and (4.4.3). Hence we have:

$$\begin{aligned} 0 &= [[E_\beta, E_i]_{q^{-(\beta, \alpha_i)}}, E_\alpha]_{q^{(-(\alpha, \beta) + (\alpha, \alpha_i))}} \\ &= (1 + q^{(-(\alpha, \beta) + (\alpha, \alpha_i))}) E_\alpha^2 . \end{aligned}$$

8.8. *Proof of Lemma 5.2.1 (ii) for type F_4* . By Lemma 5.2.1 (ii) for type B_3 and C_3 (see §6), it is enough to show:

$$(8.8.1) \quad [E_\alpha, E_\beta]_{q-(\alpha, \beta)} = 0 \quad \text{if } \alpha, \beta \in \Phi_{+,1}^{\text{red}}.$$

By Lemma 8.3.1, Lemma 8.4.1 and (8.7.1), (4.4.3), we can easily show:

$$[E_\alpha, E_\beta] = 0 \quad \text{if } \alpha, \beta \in \Phi_{+,1}^{\text{red}} \text{ and } \text{ht}(\beta) = \text{ht}(\alpha),$$

and

$$[E_\alpha, E_\beta]_{q-(\alpha, \beta)} = 0 \quad \text{if } \alpha, \beta \in \Phi_{+,1}^{\text{red}} \text{ and } \text{ht}(\beta) - \text{ht}(\alpha) = 1.$$

In the case of $\text{ht}(\beta) - \text{ht}(\alpha) \geq 2$, we can choose the elements $\gamma \in \Phi_{+,1}^{\text{red}}$ and $\alpha_i \in \Pi$ in such a way that $E_\beta = x_\alpha [E_\gamma, E_i]_{q-(\gamma, \alpha_i)}$ holds for some $x_\alpha \in R^\times$. By (4.4.2), we see:

$$\begin{aligned} & [E_\alpha, [E_\gamma, E_i]_{q-(\gamma, \alpha_i)}]_{q-(\alpha, \gamma+\alpha_i)} \\ &= [[E_\alpha, E_\gamma]_{q-(\alpha, \gamma)}, E_i]_{q-(\alpha+\gamma, \alpha_i)} \\ &+ (-1)^{p(\alpha)p(\gamma)} E_\gamma [E_\alpha, E_i]_{q-(\alpha, \alpha_i)} \\ &- (-1)^{p(\gamma)p(\alpha_i)} [E_\alpha, E_i]_{q-(\alpha, \alpha_i)} E_\gamma. \end{aligned}$$

Since $\alpha < \gamma < \beta$, we finish the proof using part (i) of Lemma 5.2.1.

§9. Commutator relations for root vectors of \mathcal{N}_+ (type G_3)

9.1. Let $(\mathcal{E}, \Pi = \{\alpha_1, \alpha_2, \alpha_3\}, p)$ be the distinguished triple system of type G_3 (see §3). Let $\mathfrak{U}_h = \mathfrak{U}_h(\mathcal{G}(G_3))$ be an h -adic R -algebra with generators E_i, F_i ($1 \leq i \leq 3$), $H \in \mathcal{H}$ and relations:

$$(9.1.1) [H_1, H_2] = 0 \quad (H_1, H_2 \in \mathcal{H}),$$

$$(9.1.2) [H, E_i] = \alpha_i(H)E_i, \quad [H, F_i] = -\alpha_i(H)F_i \quad (H \in \mathcal{H}),$$

$$(9.1.3) E_i F_j - (-1)^{p(\alpha_i)p(\alpha_j)} F_j E_i = \delta_{ij} \frac{sh(hH\alpha_i)}{sh(hd_i)},$$

$$(9.1.4) E_1^2 = 0,$$

$$\sum_{v=0}^{1+|a_{ij}|} (-1)^v \left[\begin{matrix} 1+|a_{ij}| \\ v \end{matrix} \right] q^{d_i} E_i^{1+|a_{ij}|-v} E_j E_i = 0 \quad \text{for } i \neq j \text{ and}$$

$$p(\alpha_i) = 0,$$

$$(9.1.5) F_1^2 = 0,$$

$$\sum_{v=0}^{1+|a_{ij}|} (-1)^v \left[\begin{matrix} 1+|a_{ij}| \\ v \end{matrix} \right] q^{d_i} F_i^{1+|a_{ij}|-v} F_j F_i = 0 \quad \text{for } i \neq j \text{ and}$$

$$p(\alpha_i) = 0.$$

Let \mathcal{N}_+ be the R -algebra with generators E_1, E_2, E_3 which was defined in §4 for the distinguished triple system (\mathcal{E}, Π, p) of type G_3 . Then it is obvious that there exists an R -algebra map $i_+ : \mathcal{N}_+ \rightarrow \mathfrak{U}_h$ (resp. $i_- : \mathcal{N}_+ \rightarrow \mathfrak{U}_h$) such that $i_+(E_i) = E_i$ (resp. $i_-(E_i) = F_i$) ($i = 1, 2, 3$). Let $\mathcal{N}_- = i_-(\mathcal{N}_+)$ and $\mathcal{N}_{-, \alpha} = i_-(\mathcal{N}_+, \alpha)$ ($\alpha \in P_+$). Put $U_h(\mathcal{G}(G_2)) = U_h(\mathbb{C}\alpha_3 \oplus \mathbb{C}\alpha_2, \{\alpha_3, \alpha_2\})$ (see 7.1).

Similarly to Theorem 1 (iii)-(iv) of [15], we have:

Lemma 9.1.1. (*The triangular decomposition of $\mathfrak{U}_h(\mathcal{G}(G_3))$*)

(i) The maps i_+ and i_- are injective. As h -adic topological R -modules,

$$\mathfrak{U}_h \simeq \mathcal{N}_+ \hat{\otimes} S[\mathcal{R}] \hat{\otimes} \mathcal{N}_-.$$

(ii) There exists an injective h -adic topological algebra map $j : U_h(G(G_2)) \rightarrow \mathfrak{U}_h$ such that $j(E_i) = E_i$, $j(F_i) = F_i$ ($i = 3, 2$) and $j(H_\lambda) = H_\lambda$ ($\lambda \in \delta$).

We omit the proof.

9.2. We shall extend the braid group action on $U_h((G(G_2))$ in §7 to the one on $U_h((\mathcal{G}(G_3)))$ ($\supset \mathfrak{U}_h(G(G_2))$). By direct computations, we can show the following lemma. We omit the proof.

Lemma 9.2.1. Let $T_i, T_i^{-1} \in Aut(U_h((G(G_2)))$ ($i = 3, 2$) be of Proposition 7.2.1, Then T_i, T_i^{-1} ($i = 3, 2$) can be extended to automorphisms of $\mathfrak{U}_h(\mathcal{G}(G_3))$ such that

$$T_3(E_1) = -E_3E_1 + q^{-1}E_1E_3, \quad T_2(E_1) = E_1,$$

$$T_3^{-1}(E_1) = -E_1E_3 + q^{-1}E_3E_1, \quad T_2^{-1}(E_1) = E_1,$$

$$T_3(F_1) = -qF_3F_1 + F_1F_3, \quad T_2(F_1) = F_1,$$

$$T_3^{-1}(F_1) = -qF_1F_3 + F_3F_1, \quad T_2^{-1}(F_1) = F_1,$$

$$T_i(H_\lambda) = H_{S_i(\lambda)} \quad (\text{resp. } T_i^{-1}(H_\lambda) = H_{S_i(\lambda)}) \quad (\lambda \in \delta)$$

(i = 3, 2).

9.3. Put

$$(9.3.1) \quad e_{110} = T_3(E_1), \quad e_{111} = T_2 T_3(E_1), \quad e_{131} = T_3 T_2 T_3(E_1),$$

$$(9.3.2) \quad e_{121} = (q+q^{-1})^{-1} [e_{111}, E_3] \quad (\text{see 4.4 for the notation } [,]).$$

$$(9.3.3) \quad e_{132} = T_2 T_3 T_2 T_3(E_1), \quad e_{142} = T_2 T_3 T_2 T_3 T_2 T_3(E_1).$$

Lemme 9.3.4. We have:

$$[e_{abc}, E_i] = 0$$

if $i \in \{3, 2\}$ and $a\alpha_1 + b\alpha_3 + c\alpha_2 + \alpha_i \notin \Phi_+^{\text{red}}$.

Proof. We are in one of the following three cases.

- (i) $(a, b, c, i) = (1, 1, 0, 3), (1, 1, 1, 2), (1, 3, 1, 3), (1, 4, 2, 2),$
- (ii) $(a, b, c, i) = (1, 2, 1, 1),$
- (iii) $(a, b, c, i) = (1, 4, 2, 3).$

(i) In this case, if we write $e_{abc} = T_{i_1} T_{i_2} \cdots T_{i_u}(E_1)$ as in

(9.3.1-3), then, by Proposition 7.1.1 and Proposition 7.2.1, we have:

$$(T_{i_1} T_{i_2} \cdots T_{i_u})^{-1}(E_i) = -\exp(-hH_\gamma)X$$

where $\gamma = (s_{i_2} \cdots s_{i_u})^{-1}(\alpha_i)$ and
 $X = (T_{i_2} \cdots T_{i_u})^{-1}(F_i) \in \mathcal{N}_{-, \gamma}$. Hence we have (9.3.5) in this case.

(ii) By (4.4.2) and (i), we have:

$$\begin{aligned} [e_{121}, E_2] &= (q+q^{-1})^{-1} [[e_{111}, E_3], E_2] \\ &= (q+q^{-1})^{-1} [e_{111}, [E_3, E_2]] = (q+q^{-1})^{-1} [T_2 T_3(E_1), -T_2(E_3)] \\ &= -(q+q^{-1})^{-1} [e_{110}, E_3] = 0. \end{aligned}$$

(iii) By Proposition 7.2.1 and (i), we have:

$$\begin{aligned} [e_{142}, E_3] &= [T_2 T_3 T_2 T_3 T_2 T_3(E_1), T_2 T_3 T_2 T_3 T_2(E_3)] \\ &= [e_{110}, E_3] = 0 \end{aligned}$$

This completes the proof.

Lemme 9.3.5. We have:

$$(i) e_{110} = q^{-1} [E_1, E_3].$$

$$(ii) e_{142} = q^{-1} [e_{132}, E_3] .$$

(iii)

$$(9.3.6) \quad e_{111} = q^{-3} [e_{110}, E_2] , \quad e_{131} = q^{-2} [e_{121}, E_3] , \\ e_{132} = q^{-3} [e_{131}, E_2] .$$

In particular, $e_{abc} \in \mathcal{N}_+$.

Proof. (i) Clear.

(ii)

$$e_{142} = T_2 T_3 T_2 T_3 T_2 T_3 (E_1) = T_2 T_3 T_2 T_3 T_2 (e_{110}) \\ = q^{-1} [T_2 T_3 T_2 T_3 (E_1), T_2 T_3 T_2 T_3 T_2 (E_3)] .$$

By Proposition 7.2.1, $T_2 T_3 T_2 T_3 T_2 (E_3) = E_3$. Hence

$$e_{142} = q^{-1} [e_{132}, E_3] .$$

(iii) The formulas (9.3.6) can be verified by direct computations. We sketch the proof. We write

$$e_{1bc} = T_{i_1} \cdots T_{i_{u-1}} T_{i_u} (E_1) \text{ as in (9.3.1-3). Then}$$

$$e_{1bc} = q^{-1} T_{i_1} \cdots T_{i_{u-1}} ([E_1, E_3]) = q^{-1} [e_{1yz}, T_{i_1} \cdots T_{i_{u-1}} (E_3)] .$$

Here, if $(b, c) = (1, 1)$ (resp. $(3, 1)$, $(3, 2)$), then

$(y, z) = (0, 0)$ (resp. $(1, 0)$, $(1, 1)$). By Proposition 7.2.1 (ii),

$$T_{i_1} \cdots T_{i_{u-1}} (E_3) \in \mathcal{N}_{+, (b-y)\alpha_3 + (c-z)\alpha_2} .$$

In fact, by direct computations, we can show that
 $T_2(E_3) = q^{-3}[[E_3, E_2]]$ (resp. $T_3T_2(E_3) = q^{-4}(q+q^{-1})^{-1}[[[E_2, E_3], E_3]]$,
 $T_2T_3T_2(E_3) = q^{-4}(q+q^{-1})^{-1}[[[E_3, E_2], E_2]]$) if $(b, c) = (1, 1)$ (resp.
 $(3, 1), (3, 2)$). By the formulas in Lemma 9.3.1 and the formulas
which we have already shown in this lemma, and by using the
formula (4.4.2) repeatedly, we have the formulas (iii). For
example,

$$\begin{aligned} e_{111} &= T_2T_3(E_1) = T_2(e_{110}) = q^{-1}[[E_1, T_2(E_3)]] \\ &= q^{-4}[[[E_1, E_3], E_2]] = q^{-3}[[e_{110}, E_2]]. \end{aligned}$$

This proves the first formula. The second (resp. third) formula
can be proved similarly using the first one (resp. the first and
the second ones).

Lemma 9.3.7. We have:

- (i) $e_{1bc}^2 = 0$ if $(b, c) \neq (2, 1)$.
- (ii) $[[e_{1bc}, e_{1yz}], e_{1yz}] = 0$ if $b+c-y-z = 1$.

Proof. (i) This is obvious from (9.3.1) and (9.3.3).

(ii) If $(b, c) \neq (3, 1)$, then, by Lemma 9.3.6,

$e_{1bc} \in R[[e_{1yz}, E_i]]$ for some $i \in \{3, 2\}$. By (i), $e_{1yz}^2 = 0$.

Hence, by (4.4.3), we have (ii). If $(b, c) = (3, 1)$, then
 $(y, z) = (2, 1)$.

Since $[e_{111}, e_{121}] = 0$, we have

$$\begin{aligned} 0 &= [[e_{111}, e_{121}], E_3], E_3] \\ &= (1+q^{-2})\{e_{121}((q^2 e_{131}) - (-1)q^4((q^2 e_{131})e_{121}) \\ &\quad + (q^2 e_{131})e_{121} - (-1)q^2 e_{121}(q^2 e_{131}) \\ &= q^2(q^{-2}+1+q^2)[e_{121}, e_{131}] \end{aligned}$$

by (4.4.5). Hence we get (ii).

For $\alpha = a\alpha_1 + b\alpha_3 + c\alpha_2 \in \Phi_+^{\text{red}}$, put $E_{abc} = E_{a\alpha_1 + b\alpha_3 + c\alpha_2}$.

The next lemma easily follows from Lemma 9.3.6.

Lemma 9.3.8. We have:

$$\begin{aligned} E_{110} &= q e_{110}, E_{111} = q^4 e_{111}, E_{121} = q^4 e_{121}, \\ E_{131} &= q^6 e_{131}, E_{132} = q^9 e_{132}, E_{142} = q^{10} e_{142}. \end{aligned}$$

By Lemma 9.3.7 and Lemma 9.3.8, and an argument similar to that in 8.8, we have:

Lemma 9.3.9. Let $\alpha, \beta \in \Phi_{+,1}^{\text{red}}$ be such that $\alpha < \beta$. Then we have:

$$[E_\alpha, E_\beta]_{q-(\alpha, \beta)} = \sum_{\gamma_1, \dots, \gamma_u \in \Phi_{+, 1}^{\text{red}}(\alpha < \beta)} c_{\gamma_1, \dots, \gamma_u} E_{\gamma_1} \cdots E_{\gamma_u}$$

for some $c_{\gamma_1, \dots, \gamma_u} \in R$.

9.4. By the definition of \mathfrak{U}_h (see 9.1), we can easily see that there is a \mathbb{C} -algebra isomorphism $r : \mathfrak{U}_h \rightarrow \mathfrak{U}_h$ such that

$$r(E_i) = E_i, \quad r(F_i) = F_i, \quad r(H) = H \quad (H \in \mathfrak{X}), \quad r(h) = -h.$$

Put

$$\begin{aligned} e_{010} &= E_3, \quad e_{011} = T_2(E_3), \quad e_{032} = T_2 T_3(E_2), \\ e_{021} &= T_2 T_3 T_2(E_3), \quad e_{031} = T_2 T_3 T_2 T_3(E_2), \quad e_{001} = E_2. \end{aligned}$$

By Proposition 7.2.1, we see that the above elements belong to \mathfrak{N}_+ .

By direct computations, we can get commutator relations for the above elements. For example, such commutator relations are found in Section 5 in [10]. From them, we have:

Lemma 9.4.1. We have:

$$\begin{aligned} (i) \quad E_{010} &= E_3, \quad E_{011} = -r(e_{011}), \quad E_{032} = -r(e_{032}), \\ E_{021} &= r(e_{021}), \quad E_{031} = -r(e_{031}), \quad E_{001} = E_2. \end{aligned}$$

(ii) The q-root vectors $\{E_{010}, E_{011}, E_{032}, E_{021}, E_{031}, E_{001}\}$ satisfy the commutator relations in Lemma 5.2.1 (i)-(ii).

9.5. By lemmas in 9.3-4, we can prove Lemma 5.2.1 and Remark 5.2.2 for N_+ of type G_3 . We omit the details.

§10. Main results.

10.1. Let $(\delta, \Pi = \{\alpha_1, \dots, \alpha_n\}, p)$ be the triple system satisfying the assumption in 3.1.

Lemma 10.1.1. Let $\alpha \in \Phi_+^{\text{red}}$. Then, in the h -adic topological R' -Hopf algebra $\mathcal{U}_{\sqrt{h}}^{\sigma}$, we have:

$$(10.1.2) \quad \Delta'(E_\alpha) = \{E_\alpha \otimes 1 + \exp(\sqrt{h} H'_\alpha) \sigma^{p(\alpha)} \otimes E_\alpha\} \\ \in \sum_{\gamma_1, \dots, \gamma_u \in \Phi_{+, g(\alpha)}^{\text{red}} (< \alpha)} \mathcal{U}_{\sqrt{h}}^{\sigma} \otimes E_{\gamma_1} \cdots E_{\gamma_u}.$$

Proof. We use the induction with respect to the order $<$ on $\Phi_{+, g(\alpha)}^{\text{red}}$. Then, by using Definition 5.1.1 and Lemma 5.2.1, we can show that

$$(10.1.3) \quad \Delta'(E_\alpha) = \{E_\alpha \otimes 1 + \exp(\sqrt{h} H'_\alpha) \sigma^{p(\alpha)} \otimes E_\alpha\} \\ \in \sum_{\gamma_1, \dots, \gamma_u \in \Phi_{+, g(\alpha)}^{\text{red}} (< \alpha)}$$

$$x_{\alpha-\gamma_1-\dots-\gamma_u} \exp(\sqrt{h}H_{\gamma_1+\dots+\gamma_u})^{\sigma^{p(\gamma_1)+\dots+p(\gamma_u)}} \otimes e_{\gamma_1} \cdots e_{\gamma_u}.$$

for some $x_{\alpha-\gamma_1-\dots-\gamma_u} \in N_{+, \alpha-\gamma_1-\dots-\gamma_u}$ where $u = c_\alpha$ (see 3.2 for the notation c_α).

10.2. Put $\Psi_n(t) = \prod_{i=1}^n \frac{t^i - 1}{t - 1} \in \mathbb{C}[t]$. Let $q = e^h$. As an

immediate consequence of Lemma 5.2.1 and Lemma 10.1.1, we have:

Lemma 10.2.1

$$(10.2.2) \quad \begin{aligned} & \left\langle \prod_{\alpha \in \Phi_+^{\text{red}}} E_\alpha^{m_\alpha}, \prod_{\alpha \in \Phi_+^{\text{red}}} E_\alpha^{n_\alpha} \right\rangle \\ &= \prod_{\alpha \in \Phi_+^{\text{red}}} \delta_{n_\alpha, m_\alpha} \Psi_n^{((-1)^{p(\alpha)} q^{(\alpha, \alpha)}) \langle E_\alpha, E_\alpha \rangle^{n_\alpha}}. \end{aligned}$$

(See 5.3 for the notation $\prod_{\alpha \in \Phi_+^{\text{red}}} \langle \cdot, \cdot \rangle$.)

Proof. Note that $\langle \cdot, \cdot \rangle$ is symmetric. Let $\gamma \in \Phi_+^{\text{red}}$ be such that $m_\gamma + n_\gamma \neq 0$ and $m_{\gamma'} = n_{\gamma'} = 0$ for all $\gamma' > \gamma$. Assume $m_\gamma \geq n_\gamma$. By Lemma 5.2.1 and Lemma 10.1.1, we have:

$$\left\langle \prod_{\alpha \in \Phi_+^{\text{red}}} E_\alpha^{m_\alpha}, \prod_{\alpha \in \Phi_+^{\text{red}}} E_\alpha^{n_\alpha} \right\rangle$$

$$\begin{aligned}
&= \langle \left(\prod_{\alpha \in \Phi_+^{\text{red}}(\gamma)} E_\alpha^{m_\alpha} \right) E_\gamma^{m_\gamma - 1} \otimes E_\gamma, \left(\prod_{\alpha \in \Phi_+^{\text{red}}} \Delta'(E_\alpha)^{n_\alpha} \right) \rangle \\
&= \langle \left(\prod_{\alpha \in \Phi_+^{\text{red}}(\gamma)} E_\alpha^{m_\alpha} \right) E_\gamma^{m_\gamma - 1} \otimes E_\gamma, \\
&\quad \left(\prod_{\alpha \in \Phi_+^{\text{red}}(\gamma)} (E_\alpha \otimes 1)^{n_\alpha} \right) (E_\gamma \otimes 1 + \exp(\sqrt{h} H_\gamma) \sigma^{p(\gamma)} \otimes E_\gamma)^{n_\gamma} \rangle \\
&= \langle \left(\prod_{\alpha \in \Phi_+^{\text{red}}(\gamma)} E_\alpha^{m_\alpha} \right) E_\gamma^{m_\gamma - 1} \otimes E_\gamma, \\
&\quad \left(\prod_{\alpha \in \Phi_+^{\text{red}}(\gamma)} (E_\alpha \otimes 1)^{n_\alpha} \right) \\
&\quad \cdot (\Psi_{n_\gamma}((-1)^{p(\gamma)} q(\gamma, \gamma)) E_\gamma^{n_\gamma - 1} \exp(\sqrt{h} H_\gamma) \sigma^{p(\gamma)} \otimes E_\gamma) \rangle \\
&= (\Psi_{n_\gamma}((-1)^{p(\gamma)} q(\gamma, \gamma))) \\
&\langle \left(\prod_{\alpha \in \Phi_+^{\text{red}}(\gamma)} E_\alpha^{m_\alpha} \right) E_\gamma^{m_\gamma - 1}, \left(\prod_{\alpha \in \Phi_+^{\text{red}}(\gamma)} E_\alpha^{n_\alpha} \right) E_\gamma^{n_\gamma - 1} \rangle
\end{aligned}$$

where we regarded E_γ^{-1} as 0.

Iterating this procedure, we can prove the lemma.

10.3. Here we determine the values $\langle E_\alpha, E_\alpha \rangle$ ($\alpha \in \Phi_+^{\text{red}}$).

Define $d_\alpha \in \mathbb{Z}$ ($\alpha \in \Phi_+^{\text{red}}$) by

$$d_\alpha = \begin{cases} 1 & \text{if } (\alpha, \alpha) = 0, \\ 2 & \text{if } \Phi_+^{\text{red}} \text{ is of type } G_3 \text{ and } \alpha = \alpha_1 + 2\alpha_2 + \alpha_1, \\ \frac{|(\alpha, \alpha)|}{2} & \text{otherwise.} \end{cases}$$

For $\alpha = c_1 \alpha_1 + \dots + c_n \alpha_n \in \Phi_+^{\text{red}}$, put

$$b(\alpha) = (q^{d_\alpha} - q^{-d_\alpha}) \langle E_\alpha, E_\alpha \rangle / \prod_{i=1}^n (q^{d_i} - q^{-d_i}) c_i \in K.$$

Lemma 10.3.1. For any $\alpha \in \Phi_+^{\text{red}}$, $b(\alpha)$ can be written as

$$b(\alpha) = (-1)^a q^b$$

for some $a, b \in \mathbb{Z}$. More precisely, for each type of Φ_+^{red} , $b(\alpha)$ ($\alpha \in \Phi_+^{\text{red}}$) are given by:

(i) (Type A_{N-1})

$$b(\bar{\varepsilon}_i - \bar{\varepsilon}_j) = \prod_{i < \ell < j} \bar{d}_\ell q^{\bar{d}_\ell} \quad (i < j).$$

(ii) (Type B_N)

$$b(\bar{\varepsilon}_i - \bar{\varepsilon}_j) = \prod_{i < \ell < j} \bar{d}_\ell q^{\bar{d}_\ell} \quad (i < j), \quad b(\bar{\varepsilon}_i) = \prod_{i < \ell \leq N} \bar{d}_\ell q^{\bar{d}_\ell},$$

$$b(\bar{\varepsilon}_i + \bar{\varepsilon}_j) = (-1)^{p(\alpha_N)} \bar{d}_N \left(\prod_{i < \ell \leq j} \bar{d}_\ell q^{\bar{d}_\ell} \right) \left(\prod_{j < \ell \leq N} \bar{d}_\ell q^{2\bar{d}_\ell} \right) \quad (i < j).$$

(iii) (Type C_N)

$$b(\bar{\varepsilon}_i - \bar{\varepsilon}_j) = \prod_{i < \ell < j} \bar{d}_\ell q^{\bar{d}_\ell} \quad (i < j), \quad b(2\bar{\varepsilon}_i) = \prod_{i < \ell \leq N} q^{2\bar{d}_\ell} \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 0),$$

$$b(\bar{\varepsilon}_i + \bar{\varepsilon}_j) = q \bar{d}_N \left(\prod_{i < \ell \leq j} \bar{d}_\ell q^{\bar{d}_\ell} \right) \left(\prod_{j < \ell \leq N} \bar{d}_\ell q^{2\bar{d}_\ell} \right) \quad (i < j).$$

(iv) (Type D_N)

$$b(\bar{\varepsilon}_i - \bar{\varepsilon}_j) = \prod_{i < l < j} \bar{d}_l q^{\frac{d}{2}l} \quad (i < j), \quad b(2\bar{\varepsilon}_i) = \bar{d}_N \prod_{i < l \leq N} q^{2\frac{d}{2}l} \quad (p(\bar{\varepsilon}_i - \bar{\varepsilon}_N) = 0),$$

$$b(\bar{\varepsilon}_i + \bar{\varepsilon}_j) = \bar{d}_N q^{\frac{d}{2}N} (\prod_{i < l \leq j} \bar{d}_l q^{\frac{d}{2}l}) (\prod_{j < l \leq N-1} \bar{d}_l q^{2\frac{d}{2}l}) \quad (i < j).$$

(v) (*Type F₄*) Here b_{abcd} denotes $b(a\alpha_1 + b\alpha_4 + c\alpha_3 + d\alpha_2)$.

$$\begin{aligned} b_{1000} &= 1, \quad b_{1100} = -q^{-2}, \quad b_{1110} = q^{-4}, \quad b_{1120} = -q^{-4}, \\ b_{1111} &= -q^{-5}, \quad b_{1220} = q^{-6}, \quad b_{1121} = q^{-6}, \quad b_{1221} = -q^{-8}, \\ b_{1231} &= q^{-9}, \quad b_{1232} = -q^{-12}. \\ b_{0001} &= 1, \quad b_{0011} = -q^{-1}, \quad b_{0111} = q^{-3}, \quad b_{0121} = -q^{-4}, \\ b_{0010} &= 1, \quad b_{0110} = -q^{-2}, \quad b_{0120} = q^{-2}, \quad b_{0100} = 1. \end{aligned}$$

(vi) (*Type G₃*) Here b_{abc} denotes $b(a\alpha_1 + b\alpha_3 + c\alpha_2)$.

$$\begin{aligned} b_{100} &= 1, \quad b_{110} = q, \quad b_{111} = q^4, \quad b_{121} = q^6, \\ b_{131} &= q^6, \quad b_{132} = q^9, \quad b_{142} = q^{10}, \\ b_{001} &= 1, \quad b_{011} = q^3, \quad b_{021} = q^4, \quad b_{032} = q^6, \\ b_{031} &= q^3, \quad b_{010} = 1. \end{aligned}$$

Proof. Here we sketch how to calculate $\langle E_\alpha, E_\alpha \rangle$ ($\alpha \in \Phi_+^{\text{red}}$). Put $L_\alpha = \exp(\sqrt{h}H_\alpha) \sigma^{p(\alpha)}$ for $\alpha \in \Phi_+^{\text{red}}$. We are in one of the cases (1) $c_\alpha = 1$ and (2) $c_\alpha = 2$. Firstly assume that we are in case (1). Suppose $\text{ht}(\alpha) \geq 2$ and $\alpha \in \Phi_{+,i}^{\text{red}}$. In this case, there exists $\alpha_j \in \Pi$ such that $\alpha - \alpha_j \in \Phi_{+,i}^{\text{red}}$. Let $r \in \mathbb{Z}_+$ be such that $\alpha - u\alpha_j \in \Phi_{+,i}^{\text{red}}$ ($0 \leq u \leq r$) and $\alpha - (r+1)\alpha_j \notin \Phi_{+,i}^{\text{red}}$.

Put $\beta = \alpha - \alpha_j$ and $\gamma = \alpha - r\alpha_j$. By the definition of

E_α (see Definition 5.1.1), $E_\beta = y[[E_\gamma, E_j], E_j] \dots, E_j]$ and $E_\beta = x[E_\gamma, E_j]$ for some $x, y \in R^\times$. By (10.1.3), we have:

$$\begin{aligned} \langle E_\alpha, E_\alpha \rangle &= x \langle E_\beta \otimes E_j - (-1)^{\frac{p(\beta)p(\alpha_j) - (\beta, \alpha_j)}{q}} E_j \otimes E_\beta, \Delta'(E_\alpha) \rangle \\ x^2 y \langle E_\beta \otimes E_j - (-1)^{\frac{p(\beta)p(\alpha_j) - (\beta, \alpha_j)}{q}} E_j \otimes E_\beta, \\ &\quad [[\dots [[E_\gamma \otimes 1 + L_\gamma \otimes E_\gamma, \Delta'(E_j)], \Delta'(E_j)], \dots, \Delta'(E_j)] \dots] \rangle. \end{aligned}$$

By direct computations, we see that this equals

$$\begin{aligned} &x^2 \langle -(-1)^{\frac{p(\beta)p(\alpha_j) - (\beta, \alpha_j)}{q}} E_j \otimes E_\beta, \\ &(-1)^{\frac{p(\gamma)p(\alpha_j) - (\gamma, \alpha_j)}{q}} (1 - (-1)^{\frac{(r-1)p(\alpha_j) - (2\gamma + (r-1)\alpha_j, \alpha_j)}{q}})} \\ &\frac{1 - (-1)^{\frac{rp(\alpha_j) - r(\alpha_j, \alpha_j)}{q}}}{1 - (-1)^{\frac{p(\alpha_j) - (\alpha_j, \alpha_j)}{q}}} E_j L_\beta \otimes E_\beta \rangle \\ &= -x^2 (-1)^{\frac{(r-1)p(\alpha_j) - (r-1)(\alpha_j, \alpha_j)}{q}} \\ &(1 - (-1)^{\frac{(r-1)p(\alpha_j) - (2\gamma + (r-1)\alpha_j, \alpha_j)}{q}}) \frac{1 - (-1)^{\frac{rp(\alpha_j) - r(\alpha_j, \alpha_j)}{q}}}{1 - (-1)^{\frac{p(\alpha_j) - (\alpha_j, \alpha_j)}{q}}} \\ &\langle E_\beta, E_\beta \rangle. \end{aligned}$$

Hence we can calculate $\langle E_\alpha, E_\alpha \rangle$ by the induction on $ht(\alpha)$.

Next assume that we are in case (2). Suppose $\alpha \in \Phi_+^{red}$. By the definition of E_α (see Definition 5.1.1), there exist $\beta, \gamma \in \Phi_+^{red}$ such that $\alpha = \beta + \gamma$, $ht(\gamma) - ht(\beta) \leq 1$ and $E_\alpha = z[E_\beta, E_\gamma]$ for some $z \in R^\times$. If $ht(\gamma) - ht(\beta) = 1$, then $E_\gamma = w[E_\beta, E_{\gamma-\beta}]$ for some $w \in R^\times$. In this case, since $c_\gamma = 1$, similarly to the proof in (1), we have:

$$\langle E_{\gamma-\beta} \otimes E_\beta, \Delta'(E_\gamma) \rangle = -w^{-1}(-1)^{p(\gamma-\beta)p(\beta)} q^{(\gamma-\beta, \beta)} \langle E_\gamma, E_\gamma \rangle .$$

By (10.1.3), we have:

$$\begin{aligned} \langle E_\alpha, E_\alpha \rangle &= z \langle E_\beta \otimes E_\gamma - (-1)^{p(\beta)p(\gamma)} q^{-(\beta, \gamma)} E_\gamma \otimes E_\beta, \Delta'(E_\alpha) \rangle \\ &= z^2 \langle E_\beta \otimes E_\gamma - (-1)^{p(\beta)p(\gamma)} q^{-(\beta, \gamma)} E_\gamma \otimes E_\beta, \\ &\quad [\{ E_\beta \otimes 1 + L_\beta \otimes E_\beta \}, \{ E_\gamma \otimes 1 \\ &\quad - \delta_{ht(\gamma), ht(\beta)+1} w^{-1}(-1)^{p(\beta)p(\gamma-\beta)} q^{(\beta, \gamma-\beta)} \langle E_\gamma, E_\gamma \rangle \langle E_\beta, E_\beta \rangle^{-1} \\ &\quad \cdot E_{\gamma-\beta} L_{\gamma-\beta} \otimes E_\beta + L_\gamma \otimes E_\gamma \}] \rangle \\ &= -z^2 (-1)^{p(\beta)p(\gamma)} q^{-(\beta, \gamma)} ((-1)^{p(\beta)p(\gamma)} (q^{(\beta, \gamma)} - q^{-(\beta, \gamma)}) \\ &\quad - \delta_{ht(\gamma), ht(\beta)+1} w^{-2} (-1)^{p(\beta)p(\gamma-\beta)} q^{(\beta, \gamma-\beta)} \langle E_\gamma, E_\gamma \rangle \langle E_\beta, E_\beta \rangle^{-1}) \\ &\quad \langle E_\gamma \otimes E_\beta, E_\gamma L_\gamma \otimes E_\beta \rangle \\ &= -z^2 ((1-q^{-2}(\beta, \gamma)) \langle E_\gamma, E_\gamma \rangle \langle E_\beta, E_\beta \rangle \\ &\quad - \delta_{ht(\gamma), ht(\beta)+1} w^{-2} (-1)^{p(\beta)} q^{-(\beta, \beta)} \langle E_\gamma, E_\gamma \rangle^2) . \end{aligned}$$

Since $c_\gamma = c_\beta = 1$, using results in case (1), we can get

$$\langle E_\alpha, E_\alpha \rangle .$$

10.4.

*Proposition 10.4.1. (The Ponincaré-Birkhoff-Witt theorem
for N_+ and N_+)*

(i) The R -module N_+ is a free module with a basis

$$\{ \underset{\alpha \in \Phi_+^{\text{red}}}{\prod} \underset{n_\alpha \in \mathbb{Z}_+}{E_\alpha^{n_\alpha}} \mid (n_\alpha \in \mathbb{Z}_+ \text{ if } (\alpha, \alpha) \neq 0, n_\alpha = 0, 1 \text{ if } (\alpha, \alpha) = 0) \} .$$

(ii) Let N_+ and I_+ (resp. \mathcal{N}_+ and \mathcal{I}_+) be the R -algebra and the ideal defined in 2.9 (resp. 4.2) respectively. Then $N_+ = \mathcal{N}_+$ and $I_+ = \mathcal{I}_+$.

Proof. By Lemma 10.3.1, $\langle E_\alpha, E_\alpha \rangle \neq 0$. Therefore, from Proposition 5.3.1 and Lemma 10.2.1, the proposition follows.

10.5. As an immediate consequence of Theorem 2.9.4 and Proposition 10.4.1, we have:

Theorem 10.5.1. Let $(\mathcal{E}, \Pi = \{\alpha_1, \dots, \alpha_n\}, p)$ and D be the triple system and the diagonal matrix of type X_N described in 3.1. Let \mathcal{G} be the complex simple Lie superalgebra of type X_N . Let $U(\mathcal{G})$ be the enveloping superalgebra of \mathcal{G} . Let U_h be the R -Hopf superalgebra defined for the R -Hopf algebra $U_h^\sigma((\mathcal{E}, \Pi, p), D)$ (see 1.9). Let \hat{c} be an ideal of U_h h -adically generated by the elements $\{H \in \mathcal{E} \mid \alpha_i(H) = 0 \quad (1 \leq i \leq n)\}$. Then an R -Hopf superalgebra $\hat{U}_h(\mathcal{G})$ defined by $\hat{U}_h(\mathcal{G}) = U_h / \hat{c}$ is topologically free. Moreover, as \mathbb{C} -Hopf superalgebras, $U(\mathcal{G}) \cong \hat{U}_h(\mathcal{G}) / h\hat{U}_h(\mathcal{G})$.

10.6. Here we give the main theorem. For $\alpha \in \Phi_+^{\text{red}}$, put $F_\alpha = E_\alpha^\circ \sigma^{p(\alpha)} \in U_h^\sigma$. Let ε_i 's be basis elements of \mathcal{E} such that $(\varepsilon_i, \varepsilon_j) = \delta_{ij}$. Put $t_0 = \sum_{i=1}^n H_{\varepsilon_i} \otimes H_{\varepsilon_i} \in \mathcal{E} \otimes \mathcal{E}$. Let

$e(u; t) = \sum_n (u^n / \Psi_n(t))$ be the formal power series called the "q-exponential". Put $u(\alpha) = (-1)^{ht(\alpha)} b(\alpha)^{-1}$.

Theorem 10.6.1. (*Universal R-matrix of U_h^σ*) Let R be an element of $U_h^\sigma \otimes U_h^\sigma$ defined by

$$R = \left\{ \prod_{\alpha \in \Phi_+^{\text{red}}} e((q^{d_\alpha} - q^{-d_\alpha}) u(\alpha) E_\alpha \otimes F_\alpha \sigma^{p(\alpha)}) ; (-1)^{p(\alpha)} q^{(\alpha, \alpha)} \right\} \\ \cdot \left\{ \frac{1}{2} \sum_{c, d \in \{0, 1\}} (-1)^{cd} \sigma^c \otimes \sigma^d \right\} \cdot \exp(-ht_0).$$

Then (U_h^σ, Δ, R) is a quasi-triangular Hopf algebra.

Proof. Use Lemma 2.9.10, Lemma 10.2.1, Lemma 10.3.1 and Proposition 10.4.1. Here we note the facts $\Omega'(E_\alpha) = E_\alpha$, $\Omega'(F_\alpha) = \prod_{i=1}^n (q^{-d_i} - q^{d_i})^{c_i} F_\alpha$ for $\alpha = c_1 \alpha_1 + \dots + c_n \alpha_n \in \Phi_+^{\text{red}}$.

Appendix

A.1. We use the notation in 2.1. Let \mathbb{I}_+ be an ideal of $\tilde{\mathbb{N}}_+$ generated by the elements of Definition 4.2.1 (i) - (iv). Put $\mathbb{N}_+ = \tilde{\mathbb{N}}_+ / \mathbb{I}_+$. We define an ideal \mathbb{I}_- of $\tilde{\mathbb{N}}_-$ in a similar way. Let \mathbb{L} be the ideal of \tilde{U}_h^σ h-adically generated by the elements in $\mathbb{I}_+ \cup \mathbb{I}_-$. Put $U_h^\sigma = \tilde{U}_h^\sigma / \mathbb{L}$.

Lemma A.1.1. Let $i_+ : N_+ \rightarrow U_h^\sigma$ be an R -algebra map defined by $x + I_+ \mapsto x + L$. Then i_+ is injective.

Proof. By direct computations, we see that

$F_i \cdot I_+ \subset I_+ \cdot F_i + I_+$ ($1 \leq i \leq n$). Hence $L_+ = \overline{I_+ S[\mathbb{R}]R<\sigma>N_-}$ is an ideal of U_h^σ . Similarly, we see that $L_- = \overline{N_+ S[\mathbb{R}]R<\sigma>I_-}$ is also an ideal of U_h^σ . Hence $L = L_+ + L_-$. In particular, we see that $L \cap N_+ = I_+$. This completes the proof.

A.2. For $v \in P_+$, let $N_{+,v} = \tilde{N}_{+,v} + I_+ \subset N_+$ and

$I_{+,v} = \tilde{N}_{+,v} \cap I_+$. Then $N_{+,v} = \tilde{N}_{+,v}/I_{+,v}$.

If $\begin{array}{c} \times \\ \diagup \\ \otimes \\ \diagdown \end{array}^{N-2} \circ^{N-1} \circ_N$, then $I_{+,v}^{\alpha_{N-2} + \alpha_{N-1} + \alpha_N} = \{0\}$.

In particular, $\text{rank } N_{+,v}^{\alpha_{i-1} + 2\alpha_i + \alpha_{i+1}} = 6$. Hence

Poincaré-Birkhoff-Witt type theorem can not hold for U_h^σ .

A.3. Let U_h^* be the Hopf superalgebra called the "quantized Kac-Moody superalgebra" in [7]. Here we understand that U_h^* is defined as an h -adic R -Hopf superalgebra. Even if we take the *Note added in proof* in [7] into account, we can show that there exists a natural epimorphism $(U_h^*)^\sigma \rightarrow U_h^\sigma$ of Hopf algebras. Hence, for U_h^* , a P.B.W. type theorem can not hold contrary to their assertion (Proposition 3.3 and Remark under it) in [7].

Acknowledgements. I would like to express my heartfelt thanks to Professor M. Noumi, Professor T. Tanisaki, Professor J. Murakami and Professor M. Okado for valuable communication. Specially I thank Professor T. Tanisaki for showing me a proof of the result on the "quantum duoble construction". I also thank Professor M. Rosso for sending me his preprint of [13] and Professor E. Date for informing me about the existence of the preprint [7] of Khoroshkin and Tolstoy.

References

- [1] E. Abe. Hopf algebras. Cambridge Univ. Press, Cambridge-New York, 1980.
- [2] V.G. Drinfeld: Quantum groups, Proc. Int. Congr. Math., Berkeley, vol. 1 (1986); Amer. Math. Soc.(1988), 798-820.
- [3] M. Jimbo: A q-difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation. Lett. Math. Phys., 10 (1985), 63-69.
- [4] V.G. Kac: Infinite dimensional Lie algebras, Progress in Mathematics 44 Birkhäuser, Boston/Basel/Stuttgart, 1983.
- [5] V.G. Kac: Lie superalgebras, Adv. Math. 26 (1977), 8-96.
- [6] A.N. Kirillov and N.Yu. Reshetikhin: q-Weyl group and a multiplicative formula for universal R-matrix, Commun. Math. Phys. 134 (1990), 421-431.

- [7] S.M. Khoroshkin and V.N. Tolstoy: Universal R-matrix for quantized (super)algebras (preprint).
Commun. Math. Phys. 141 (1991), 599-617.
- [8] S. Levendorskii and Ya. Soibelman: Some applications of quantum Weyl group (preprint).
- [9] G. Lusztig: Finite dimensional Hopf algebras arising from quantized universal enveloping algebras, Jour. Amer. Math. Soc. 3 (1990), 257-296.
- [10] G. Lusztig: Quantum groups at roots of 1, Geom. Dedicata 35 (1990), 89-144.
- [12] H. Matsumura: Commutative Algebra, Second ed. London: Benjamin 1980.
- [13] M. Rosso: An analogue of P.B.W. Theorem and the Universal R-matrix for $U_h\mathfrak{sl}(N+1)$, Commun. Math. Phys. 124 (1989), 307-318.
- [14] M. Scheunert: The theory of Lie superalgebras, Lecture note in Math. 716, Springer, Berlin 1979.
- [15] T. Tanisaki: Harish-Chandra isomorphism for quantum algebras, Commun. Math. Phys. 127 (1990), 555-571.
- [16] T. Tanisaki: Killing forms, Harish-Chandra isomorphisms, and universal R-matrices for quantum algebras, to appear in Proc. of Project 91, RIMS.
- [17] H. Yamane: Universal R-matrices for quantum groups associated to simple Lie superalgebras, Proc. Japan Acad. Ser. A, No. 4 (1991), 108-112.

Department of mathematics

Osaka University

Toyonaka 560, Japan