

Title	Clinical Survey on Adverse Reaction of Iodinated Contrast Media-Interim Report-
Author(s)	片山, 仁; 小塚, 隆弘; 高島, 力 他
Citation	日本医学放射線学会雑誌. 1988, 48(2), p. 214-216
Version Type	VoR
URL	https://hdl.handle.net/11094/15737
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Clinical Survey on Adverse Reactions of Iodinated Contrast Media —Interim Report—

Hitoshi Katayama¹⁾, Takahiro Kozuka²⁾, Tsutomu Takashima³⁾,
Keiichi Matsuura⁴⁾ and Koichi Yamaguchi⁵⁾

1) Department of Radiology, Juntendo University, School of Medicine, Tokyo, Japan

2) Department of Radiology, Osaka University, School of Medicine, Osaka

3) Department of Radiology, Kanazawa University, School of Medicine, Kanazawa

4) Saga Medical College, Saga

5) Department of Radiology, Yamagata University, School of Medicine, Yamagata

Research Code No. : 502

Key Words : Adverse reactions, Contrast media, Side-effects,
Iodinated contrast media

ヨード造影剤の副作用に関する臨床調査

（中間報告）

—123,060例の分析—

片山 仁¹⁾ 小塚 隆弘²⁾ 高島 力³⁾
松浦 啓一⁴⁾ 山口 昂一⁵⁾

1) 順天堂大学医学部放射線医学教室

2) 大阪大学医学部放射線医学教室

3) 金沢大学医学部放射線医学教室

4) 佐賀医科大学

5) 山形大学医学部放射線医学教室

（昭和62年11月20日受付）

（昭和62年12月17日最終原稿受付）

Introduction

This study consisted of a comprehensive survey to evaluate adverse reactions to iodinated contrast media. It included comparisons of ionic and non-ionic low osmolar contrast media at a time of changes from a generation of the former to a generation of the latter.

Material and Method

From September 1986 to April 1987 we collected data concerning 123,060 cases examined in 196 hospitals throughout Japan. For several reasons, 3,439 cases were excluded from analyses. Only procedures involving the intravenous administration of contrast media were studied.

Severity of reactions	ADR No. (%)		Odds rate of non-ionic to ionic (95% confidence limit)
	Ionic CM 77,040 cases	Non-ionic CM 42,581 cases	
generalized	10,429 (13.54)	1,789 (4.2)	0.28 ** (0.27~0.29)
severe a)	344 (0.45)	41 (0.10)	0.21 ** (0.16~0.29)
very severe b)	38 (0.05)	3 (0.01)	0.14 ** (0.05~0.39)
death	*(1) (0.001)	*(1) (0.002)	1.81 NS (0.12~27.79)

a) Severe adverse reactions included : facial edema, rigor/tremor, dyspnea, sudden drop in blood pressure, cardiac arrest and loss of consciousness.
 b) Very severe adverse reactions : severe adverse reactions requiring anesthesiologist's help or hospitalization.
 * causal relation unknown under investigation. (**: $\alpha < 0.01$)
 (Odds rate and 95% confidence limit according to generalized Mantel-Haenszel's method) (NS: $\alpha > 0.05$)

Fig. 1 Over-all frequency and severity of adverse reactions (ADR) to contrast media.

Reactions to pretesting	Ionic CM			Non-ionic CM		
	Total No.	ADR No. (%)	Severe ADR No. (%)	Total No.	ADR No. (%)	Severe ADR No. (%)
Positive	487	238 (48.87)	11 (2.26)	244	47 (19.26)	0 (0)
Negative	60,907	8,061 (13.23)	262 (0.43)	30,339	1,263 (4.16)	30 (0.099)
Unknown	67	16 (23.88)	1 (1.49)	39	3 (7.69)	0 (0)
No response	1,836	—	—	1,049	—	—

Fig. 2 Frequency of adverse reactions (ADR) to contrast media related to whether pretest was positive.

History of previous reactions	Ionic CM			Non-ionic CM		
	Total No.	ADR No. (%)	Severe ADR No. (%)	Total No.	ADR No. (%)	Severe ADR No. (%)
Yes	2,962	1,335 (45.07)	44 (1.49)	3,166	390 (12.32)	9 (0.28)
No	32,620	3,105 (9.52)	79 (0.24)	17,445	533 (3.06)	13 (0.07)
Unknown	571	123 (21.54)	6 (1.05)	382	21 (5.50)	0 (0)
No response	1,623	—	—	974	—	—

Fig. 3 Frequency of adverse reactions (ADR); correlation of history of previous adverse reactions and current reactions to contrast media.

History of allergy	Ionic CM			Non-ionic CM		
	Total No.	ADR No. (%)	Severe ADR No. (%)	Total No.	ADR No. (%)	Severe ADR No. (%)
Yes	6,096	1,474 (24.18)	72 (1.18)	4,048	354 (8.75)	8 (0.20)
No	66,708	8,356 (12.53)	252 (0.38)	36,219	1,338 (3.69)	31 (0.09)
Unknown	3,848	533 (13.85)	16 (0.42)	2,094	88 (4.20)	2 (0.10)
No response	388	—	—	220	—	—

Fig. 4 Frequency of adverse reactions (ADR) to contrast media; effect of history of allergy.

Results

There was an over-all rate of adverse reactions of 13.5% for the ionic contrast media and a rate of 4.2% for non-ionic low osmolar contrast media (Fig. 1).

One death occurred in each group related to the media, but the causal relationships are unknown and still under investigation.

Among those who had positive reactions to test dose, there were tendencies for higher rates of adverse reactions.

Among positive test individuals adverse reactions were observed in 234 (48.9%) of 487 persons who received the ionic contrast media and in 47 (19.3%) of 244 who received nonionic contrast media (Fig. 2).

The effect of histories of previous reactions was demonstrated. For both ionic and non-ionic contrast media, previous reactors were liable to have other kinds of adverse reactions and hypersensitivities (Fig. 3).

Histories of allergy had a definite correlations with higher rates of adverse reactions, even when non-ionic low osmolar contrast media were used. The allergies investigated in this study included atopy, asthma, pollenosis, drugs and foods.

Discussion

The frequency of such adverse reactions varies with reported series¹⁾²⁾. In the present study, the rate was 13.5% for ionic and 4.2% for non-ionic contrast media. The rate may range from 5 to 15%.

The introduction of low osmolar iodine contrast media obviously minimized the rate of adverse reactions.

From a clinical standpoint it is generally agreed that alleviation of pain at the time of injection of the contrast media greatly facilitates the radiological examinations. Sogn et al (1987)³⁾ reported that comparison of iohexol and ionic monomeric contrast media disclosed the rate of adverse reactions to be about 3.5 times less using iohexol than ionic monomers.

We are also of this opinion.

Conclusion

The over-all rate of adverse reactions were 13.5% among 77,040 persons who received ionic contrast media and 4.2% among 42,581 who received non-ionic low osmolar contrast media. Crude data are currently available and investigations are now in progress.

References

- 1) Witten DH, Hirsch FD and Hartman GW: Acute reaction to Urographic contrast medium. Incidence to clinical characteristics and relationships to history of hypersensitive states. *AJR* 119: 832—840, 1973
- 2) Shehadi WH and Toniolo G: Adverse reaction to contrast media. *Radiology* 137: 299—302, 1980
- 3) Sogn EE, Ødegård T, Haider, et al: Adverse reactions following two separate intravascular injections of contrast media in the same patient —A comparison between iohexol and monomeric ionic media. *Acta Radiol.* 28(1): 93—98, 1987

*Request for reprints: Hitoshi Katayama, M.D. Department of Radiology, Juntendo University, School of Medicine, Hongo 3-1-3, Bunkyo-ku Tokyo 113 Japan
