

Title	Multicommodity Flows in Graphs II
Author(s)	岡村, 治子
Citation	大阪大学, 1984, 博士論文
Version Type	VoR
URL	<a href="https://hdl.handle.net/11094/1691">https://hdl.handle.net/11094/1691</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

---

Multicommodity Flows in Graphs II

---

Haruko OKAMURA

---

---

## 1. Introduction

Let  $G=(V,E)$  be a graph (finite undirected, possibly with multiple edges but without loops), and let  $V=V(G)$ ,  $E=E(G)$  be the sets of vertices and edges of  $G$  respectively. In this paper a path has no repeated edges, and we permit paths with one vertex and no edges. For two distinct vertices  $x,y$ , let  $\lambda(x,y)=\lambda_G(x,y)$  be the maximum number of edge-disjoint paths between  $x$  and  $y$ , and let  $\lambda(x,x)=\infty$ .

We first consider the following problem.

Let  $(s_1,t_1),\dots,(s_k,t_k)$  be pairs (not necessarily distinct) of vertices of  $G$ . When is the following true ?

(1.1) There exist edge-disjoint paths  $P_1,\dots,P_k$  such that  $P_i$  has ends  $s_i,t_i$  ( $1 \leq i \leq k$ ).

Seymour [10] and Thomassen [12] characterized such graphs when  $k=2$ , and Seymour [10] when  $s_1,\dots,s_k,t_1,\dots,t_k$  take only three distinct values.

Our result is the following

Theorem 1. Suppose that  $s_1,s_2,s_3,t_1,t_2,t_3$  are vertices of a graph  $G$ . If for each  $i=1,2,3$ ,

$$\lambda(s_i,t_i) \geq 3,$$

then there exist edge-disjoint paths  $P_1,P_2,P_3$  of  $G$ , such that  $P_i$  has ends  $s_i$  and  $t_i$  ( $i=1,2,3$ ).

If  $\lambda(s_i, t_i) \leq 2$  for some  $i$ , then the conclusion does not always hold. Figure 1 gives a counterexample.




Figure 1.

For a positive integer  $k$ , let  $g(k)$  be the smallest integer such that for every  $g(k)$ -edge-connected graph and for every vertices  $s_1, \dots, s_k, t_1, \dots, t_k$  of the graph, (1.1) holds. Thomassen [12] conjectured the following.

Conjecture. For each odd integer  $k \geq 1$ ,  $g(k)=k$ , and for each even integer  $k \geq 2$ ,  $g(k)=k+1$ .

If  $k$  is even then  $g(k) > k$  (see [12]). It follows easily from Menger's theorem that  $g(k) \leq 2k-1$ , thus  $g(1)=1$ ,  $g(2)=3$ ; and Cypher [1] proved  $g(4) \leq 6$  and  $g(5) \leq 7$ . As a corollary of Theorem 1 we have the following.

Corollary.  $g(3)=3$ .

The second problem we consider is the multicommodity flow problem.

Suppose that each edge  $e \in E$  has a real-valued capacity  $w(e) \geq 0$ , and each path has a positive value. We assume that  $w \equiv 1$  and each path has value 1 when there is no explanation. For a positive number  $\alpha$ , paths  $\alpha P$ ,  $P$  denote paths of value  $\alpha, 1$  respectively. We say that a set of paths  $\alpha_1 P_1, \dots, \alpha_n P_n$  is feasible if for each edge  $e \in E$ ,

$$\sum_{i \in \{i | e \in E(P_i)\}} \alpha_i \leq w(e),$$

where  $E(P_i)$  is the set of edges of  $P_i$ .

For two vertices  $x, y$  and a real number  $q > 0$ , a flow  $F$  of value  $q$  between  $x$  and  $y$  is a set of paths  $\alpha_1 P_1, \dots, \alpha_n P_n$  between  $x$  and  $y$  such that  $\alpha_1 + \dots + \alpha_n = q$ . When  $\alpha_1, \dots, \alpha_n$  are all integers (half-integers),  $F$  is called an integer (half-integer) flow. We say that a set of flows  $F_1, \dots, F_k$  is feasible if the set of paths of  $F_1, \dots, F_k$  is feasible.

Now the multicommodity flow problem is as follows.

Let  $(s_i, t_i), \dots, (s_k, t_k)$  be pairs of vertices of  $G$ , as before, and suppose that  $q_i \geq 0$  ( $1 \leq i \leq k$ ) are real-valued demands. When is the following true?

(1.2) There exist feasible flows  $F_1, \dots, F_k$ , such that  $F_i$  has ends  $s_i$  and  $t_i$  and value  $q_i$  ( $1 \leq i \leq k$ ).

Remark. When  $k=3$ ,  $w \equiv 1$ , and  $q_i = 1$  ( $1 \leq i \leq 3$ ), Theorem 1

implies that (1.2) is true if  $\lambda(s_i, t_i) \geq 3$  ( $1 \leq i \leq 3$ ), and then the flows may be chosen as integer flows.

For a set  $X \subseteq V$ , let  $\partial(X) = \partial_G(X) \subseteq E$  be the set of edges with one end in  $X$  and the other in  $V-X$ , and let  $D(X) = D_G(X) \subseteq \{1, 2, \dots, k\}$  be

$$\{i \mid 1 \leq i \leq k, X \cap (s_i, t_i) \neq \emptyset \neq (V-X) \cap (s_i, t_i)\}.$$

It is clear that if (1.2) is true, then the following holds.

(1.3) For each  $X \subseteq V$ ,

$$\sum_{e \in \partial(X)} w(e) \geq \sum_{i \in D(X)} q_i.$$

Note that  $\sum_{e \in \partial(X)} w(e) = |\partial(X)|$  if  $w \equiv 1$ , and  $\sum_{i \in D(X)} q_i = |D(X)|$

if  $q_i = 1$  for any  $i$ .

Our second result is the following

**Theorem 2.** Suppose that  $G$  is a graph and  $w$  is integer-valued, and that  $k=3$ ,  $q_1=q_2=q_3=1$ . Then (1.2) and (1.3) are equivalent.

Moreover if (1.3) holds, then the flows  $F_i$  in (1.2) may be chosen as half-integer flows.

(1.4) In general (1.2) and (1.3) are not equivalent, but in the following cases they are equivalent.

(1.4.1)  $k=1$  (Ford and Fulkerson [2]).

(1.4.2)  $k=2$  (Hu [3] and Seymour [8])

(1.4.3)  $k=5$ ,  $t_i=s_{i+1}$  ( $i=1,2,3,4$ ) and  $t_5=s_1$  (Papernov [7]).

(1.4.4)  $k=6$ , and  $(s_1, t_1), \dots, (s_6, t_6)$  correspond to the six pairs of a set of four vertices (Papernov [7] and Seymour [9]).

(1.4.5)  $s_1=s_2=\dots=s_j$  and  $s_{j+1}=\dots=s_k$  (obvious extension of (1.4.2)).

(1.4.6) The graph  $(V, E \cup \{e_1, \dots, e_k\})$  is planar, where the edge  $e_i$  has ends  $s_i$  and  $t_i$  ( $1 \leq i \leq k$ ) (Seymour [11]).

(1.4.7)  $G$  is planar and can be drawn in the plane so that  $s_1, \dots, s_k, t_1, \dots, t_k$  are all on the boundary of the infinite face (Okamura and Seymour [5]).

(1.4.8)  $G$  is planar and can be drawn in the plane so that  $s_1, \dots, s_j, t_1, \dots, t_j$  are all on the boundary of a face and  $s_{j+1}, \dots, s_k, t_{j+1}, \dots, t_k$  are all on the boundary of the infinite face (Okamura [6]).

(1.4.9)  $G$  is planar and can be drawn in the plane so that  $s_{j+1}, \dots, s_k, t_1, t_2, \dots, t_k$  are all on the boundary of the infinite face, and  $t_1 = \dots = t_j$  (Okamura [6]).

Moreover if (1.3) and the following (1.5) hold in each case except (1.4.3), or if (1.3) holds and  $w, q_j$  are even-integer valued in the case (1.4.3), then the flows  $F_i$  of (1.2) may be chosen as integer flows.

(1.5)  $w$  and  $q_i$  are integer-valued, and for each vertex  $x \in V$ ,

$$\sum_{e \in \partial(x)} w(e) - \sum_{i \in D(x)} q_i$$

is even.

(1.4.1), ..., (1.4.5) are all the configurations of  $(s_i, t_i)$  for which (1.2) and (1.3) are equivalent for all graphs  $G$  and all  $w, q_i$  (see [9]). When  $q_i > 0$  ( $1 \leq i \leq 3$ ), the case of Theorem 2 is the only case for which (1.2) and (1.3) are equivalent for all graphs  $G$  and all  $w, (s_i, t_i)$ . Figure 1 gives a counterexample with  $q_1=2, q_2=q_3=1$ .

Notations and definitions. We call  $S \subseteq E$  an  $n$ -cut if  $|S|=n$  and  $S=\partial(X)$  for some  $X \subseteq V$  such that  $\langle X \rangle$  (which is the subgraph induced by  $X$ ) and  $\langle V-X \rangle$  are both connected; and an  $n$ -cut  $\partial(X)$  is called nontrivial if  $|X| \geq 2$  and  $|V-X| \geq 2$ , trivial otherwise. For two vertices  $x, y$  a path  $P[x, y]$  or a path  $[x, y]$  denotes a path between  $x$  and  $y$ , and let  $xy$  be an edge with ends  $x, y$ , and let  $d(x, y) = d_G(x, y)$  be the distance between  $x$  and  $y$ . If vertices  $x, y$  belong to a path  $P$ , then  $P(x, y)$  denotes the subpath of  $P$  between  $x$  and  $y$ . For a vertex  $x$   $\deg(x) = \deg_G(x)$  denotes the degree of  $x$ , and we let  $N(x) = N_G(x)$  be  $\{y \in V \mid xy \in E\}$ . For a set  $X \subseteq V$  and an edge  $e$ , we denote graphs  $\langle V-X \rangle$ ,  $(V, E-e)$  by  $G-X$ ,  $G-e$  respectively. For a set  $X \subseteq V$  ( $S \subseteq E$ ) and an element  $x \in V$  ( $e \in E$ ), we denote  $X \cup \{x\}$  ( $S \cup \{e\}$ ) by  $X \cup x$  ( $S \cup e$ ).



## 2. Proof of Theorem 1.

In this section disjoint means edge-disjoint. We require the following lemmas.

Lemma 2.1. Suppose that  $s_1, s_2, t_1, t_2$  are vertices of a graph  $G$ . If  $\lambda(s_1, t_1) \geq 3$  and  $\lambda(s_2, t_2) \geq 1$ , then  $G$  contains disjoint paths  $[s_1, t_1]$  and  $[s_2, t_2]$ .

Proof. Since  $\lambda(s_1, t_1) \geq 3$ ,  $G$  contains disjoint paths  $P_1[s_1, t_1], P_2[s_1, t_1]$  and  $P_3[s_1, t_1]$ .  $G$  contains a path  $P_4[s_2, t_2]$ . There exist vertices  $x, y \in V(P_4)$  such that  $P_4(s_2, x)$  and  $P_4(t_2, y)$  are disjoint from  $P_1, P_2, P_3$ . Choose  $x, y$  with this property such that  $P_4(s_2, x), P_4(t_2, y)$  have the maximum length respectively. If  $x$  or  $y \notin V(P_1) \cup V(P_2) \cup V(P_3)$ , then  $x=t_2$  or  $y=s_2$ , and so the result follows. We may therefore assume that  $x \in V(P_2)$  and  $y \in V(P_i)$  ( $i=2$  or  $3$ ). When  $i=2$  ( $i=3$ ), let  $P_5$  be the path obtained by combining  $P_4(s_2, x), P_2(x, y)$  and  $P_4(y, t_2)$  ( $P_4(s_2, x), P_2(x, s_1), P_3(s_1, y)$  and  $P_4(y, t_2)$ ). Now  $P_1$  and  $P_5$  are required paths of  $G$ .

Lemma 2.2. If  $G$  is 3-regular 3-edge-connected graph with no nontrivial 3-cut and with  $4 \leq |V| \leq 8$ , then  $G$  is  $K_4, K_{3,3}$ , a cube or the graph in Figure 2.




Figure 2.

Proof. Since  $G$  is 3-regular 3-edge-connected,  $G$  has no multiple edges. Thus if  $|V|=4$ , then  $G$  is  $K_4$ . If  $|V|>4$ , then  $G$  has no cycle of length three. If  $|V|=6$ , then let  $V=\{x_1, \dots, x_6\}$ . We may let  $N(x_1)=\{x_2, x_3, x_4\}$ . Since  $x_i x_j \notin E$  ( $2 \leq i < j \leq 4$ ), we have  $x_i x_j \in E$  ( $i=2,3,4; j=5,6$ ). Thus  $G$  is  $K_{3,3}$ . If  $|V|=8$ , then it easily follows that  $G$  is a cube or the graph in Figure 2.

Lemma 2.3. Suppose that  $G$  is a 3-regular 3-edge-connected graph, and that  $a, x_1, x_2, x_3, x_4$  are vertices such that  $a \neq x_i$  ( $1 \leq i \leq 4$ ). Then  $G-a$  contains disjoint paths  $[x_1, x_2]$  and  $[x_3, x_4]$ .

Proof. We proceed by induction on  $|V|$ . If  $|V|=2$ , then  $G$  is the graph of triple edges, and the result holds. Therefore we assume  $|V| \geq 4$ .

First we assume that  $G$  contains a nontrivial 3-cut  $\{e_1, e_2, e_3\} = \partial(X)$  ( $X \subseteq V$ ). Let  $b_i \in X$ ,  $c_i \in V-X$ ,  $e_i = b_i c_i$  ( $i=1,2,3$ ),

then  $b_i \neq b_j, c_i \neq c_j$  if  $i \neq j$ , since  $G$  is 3-edge-connected. Let  $H, K$  be the graphs obtained from  $G$  by contracting  $V-X, X$  to one vertex respectively. Let  $V(H)=X \cup v, V(K)=(V-X) \cup u$ . Then  $H, K$  are 3-regular 3-edge-connected graphs and  $|V(H)| < |V|, |V(K)| < |V|$ . We may assume  $a \in V-X$ . It suffices to prove the lemma in the following cases.

Case 1.  $\{x_1, x_2, x_3, x_4\} \subseteq V-X$ . By induction the result holds in  $K$ , and so in  $G$ .

Case 2.  $x_1 \in X$  and  $\{x_2, x_3, x_4\} \subseteq V-X$ . By induction the result holds in  $K$  (note that  $x_1 = u$  in  $K$ ). Thus the result holds in  $G$ , since  $G$  contains a subgraph  $G_1$  homeomorphic to  $K$ , such that  $x_1$  corresponds to  $u$  and each vertex of  $V-X$  to itself.

Case 3.  $\{x_1, x_2, x_3, x_4\} \subseteq X$ .  $G$  contains a subgraph  $G_2$  homeomorphic to  $H$ , such that  $a$  corresponds to  $v$  and each vertex of  $X$  to itself, and so the result holds in  $G$ .

Case 4.  $\{x_1, x_2\} \subseteq X$  and  $\{x_3, x_4\} \subseteq V-X$ . Since  $K-(a, u)$  is connected, this contains a path  $[x_3, x_4]$ ; and  $H-v$  contains a path  $[x_1, x_2]$ .

Case 5.  $\{x_1, x_3\} \subseteq X$  and  $\{x_2, x_4\} \subseteq V-X$ . By induction  $K-a$  contains disjoint paths  $P_1 [u, x_2]$  and  $P_2 [u, x_4]$ . We may let  $c_i \in V(P_i)$  ( $i=1,2$ ), and  $H-v$  contains disjoint paths  $[x_1, b_1]$  and  $[x_3, b_2]$ . Thus the result follows.

Case 6.  $\{x_1, x_2, x_3\} \subseteq X$  and  $x_4 \in V-X$ .  $K-a$  contains a path  $P [u, x_4]$ , and we may let  $c_1 \in V(P)$ .  $H-v$  contains disjoint paths  $[x_1, x_2]$  and  $[x_3, b_1]$ . Thus the result follows.

Next we assume that  $G$  does not contain a nontrivial 3-cut. If  $G$  contains an edge  $e$  which is not incident to any of  $a, x_1, x_2, x_3, x_4$ , then let  $\widetilde{G-e}$  be the 3-regular graph homeomorphic to the graph  $G-e$ . Then  $\widetilde{G-e}$  is 3-edge-connected. By induction the result holds in  $\widetilde{G-e}$ , and so in  $G$ . Thus we assume that any edge is incident to one of  $a, x_1, x_2, x_3, x_4$ . Then  $|E| \leq 15$  and  $|V| \leq 10$ . We put  $T = \{a, x_1, x_2, x_3, x_4\}$ . We may assume that  $x_1, x_2, x_3$  and  $x_4$  are all distinct. For if not, then the result follows, since  $G-a$  is 2-edge-connected. Thus  $|V| \geq 5$ . If  $|V| = 10$ , then  $N(x_i) \subseteq V-T$  ( $1 \leq i \leq 4$ ) and  $|V-T| = 5$ . Thus for some  $y \in V-T$ ,  $y \in N(x_1) \cap N(x_2)$ .  $G-\{a, y\}$  is connected, and so the result follows. If  $|V| = 6$  or  $8$ , then by Lemma 2.2  $G$  is  $K_{3,3}$ , a cube, or the graph in Figure 2. We omit the proofs for them.

Lemma 2.4. Suppose that  $G$  is a 3-regular 3-edge-connected graph, and that  $a, a_1, a_2, a_3, x_1, x_2, x_3$  are vertices such that  $N(a) = \{a_1, a_2, a_3\}$  and  $a \neq x_i$  ( $1 \leq i \leq 3$ ). Then

$$|I_G| \geq 4.$$

Here  $I_G = I_G(a, a_1, a_2, a_3, x_1, x_2, x_3)$  is

$$\left\{ \begin{array}{l} (i, j, k) \\ \left. \begin{array}{l} (i, j, k) = (1, 2, 3). \text{ } G-a \text{ contains disjoint paths} \\ [x_1, a_i], [x_2, a_j] \text{ and } [x_3, a_k]. \end{array} \right\} \end{array} \right\}.$$

Proof. We proceed by induction on  $|V|$ . We assume  $|V| \geq 4$ . First we assume that  $G$  contains a nontrivial 3-cut

$(e_1, e_2, e_3) = \partial(X)$  ( $X \subseteq V$ ). We define  $b_i, c_i$  ( $i=1,2,3$ ),  $H, K, v$  and  $u$  similarly as in the proof of Lemma 2.3. We may assume  $a \in V-X$ . Then  $|X \cap N(a)| \leq 1$ . If  $a_i \in X$  for some  $i$ , then  $a_i = u$  in  $K$ . It suffices to prove the lemma in the following cases.

Case 1.  $(x_1, x_2, x_3) \subseteq V-X$ ;  $x_1 \in X$  and  $(x_2, x_3) \subseteq V-X$ ; or  $(x_1, x_2, x_3) \subseteq X$ . Similar as Case 1, 2 or 3 in the proof of Lemma 2.3.

Case 2.  $(x_1, x_2) \subseteq X$  and  $x_3 \in V-X$ . By induction  $|I_K(a, a_1, a_2, a_3, u, u, x_3)| \geq 4$ . For each  $(i, j, k)$  of  $I_K$ ,  $K-a$  contains disjoint paths  $P_1 [u, a_i]$ ,  $P_2 [u, a_j]$  and  $P_3 [x_3, a_k]$ . If  $u \notin N_K(a)$ , then we may let  $c_i \in V(P_i)$  ( $i=1,2$ ). By Induction  $H-v$  contains disjoint paths  $[x_1, b_1]$  and  $[x_2, b_2]$ . Thus  $(i, j, k) \in I_G(a, a_1, a_2, a_3, x_1, x_2, x_3)$ , and so  $|I_G| \geq 4$ . If  $u \in N_K(a)$ , then we may let  $a_1 = u, a_2 = c_1$ . Now  $k \neq 1$  and we may let  $i=1, j=2, k=3, c_2 \in V(P_2)$ . Since  $H-v$  contains disjoint paths  $[x_1, b_1]$  and  $[x_2, b_2]$ ,  $|I_G| \geq 4$ .

Next we assume that  $G$  does not contain a nontrivial 3-cut. We may assume that any edge is incident to one of  $a, x_1, x_2, x_3$  (see the proof of Lemma 2.3). Thus  $|E| \leq 12$  and  $|V| \leq 8$ . By Lemma 2.2  $G$  is  $K_4, K_{3,3}$ , a cube or the graph in Figure 2, but in the last graph any four vertices do not cover all edges of the graph. Thus  $G$  is one of the first three graphs. If  $G$  is a cube, then in Figure 3 it suffices to check the case  $y_1 = a, y_3 = x_1, y_6 = x_2, y_8 = x_3$ . We omit the proofs for  $K_4, K_{3,3}$ .




Figure 3.

Lemma 2.5. Suppose that  $s_1, s_2, s_3, t_1, t_2, t_3$  are vertices of a graph  $G$ . If  $G$  is 3-regular 3-edge-connected, then  $G$  contains disjoint paths  $[s_1, t_1], [s_2, t_2]$  and  $[s_3, t_3]$ .

Proof. We proceed by induction on  $|V|$ . We put  $T = \{s_1, s_2, s_3, t_1, t_2, t_3\}$ . If  $s_i = t_i$  for some  $i$ , then the result follows by Lemma 2.1, and if  $s_1 = s_2 = s_3$ , then the result follows from Menger's theorem. Thus we may assume that these are not the cases.

First we assume that  $G$  contains a nontrivial 3-cut  $(e_1, e_2, e_3) = \partial(X)$  ( $X \subseteq V$ ). We define  $b_i, c_i$  ( $i=1,2,3$ ),  $H, K, v$  and  $u$  similarly as in the proof of Lemma 2.3. It suffices to prove the lemma in the following cases.

Case 1.  $T \cap X = \emptyset$ . By induction the result holds in  $K$ , and so in  $G$ .

Case 2.  $s_1 \in X$  and  $\{s_2, s_3, t_1, t_2, t_3\} \subseteq V - X$ .  $G$  contains a subgraph  $G_1$  homeomorphic to  $K$ , such that  $s_1$  corresponds to  $u$

and each vertex of  $V-X$  to itself.

Case 3.  $\{s_1, t_1\} \subseteq X$  and  $\{s_2, s_3, t_2, t_3\} \subseteq V-X$ . By Lemma 2.3  $K-u$  contains disjoint paths  $[s_2, t_2]$  and  $[s_3, t_3]$ , and  $H-v$  contains a path  $[s_1, t_1]$ .

Case 4.  $\{s_1, s_2\} \subseteq X$  and  $\{s_3, t_1, t_2, t_3\} \subseteq V-X$ . By induction  $K$  contains disjoint paths  $P_1 [u, t_1], P_2 [u, t_2]$  and  $[s_3, t_3]$ . Let  $c_i \in V(P_i)$  ( $i=1,2$ ). By Lemma 2.3  $H-v$  contains disjoint paths  $[s_1, b_1]$  and  $[s_2, b_2]$ . Now the result follows.

Case 5.  $\{s_1, s_2, t_1\} \subseteq X$  and  $\{s_3, t_2, t_3\} \subseteq V-X$ . We can get the result by applying Lemma 2.3 on  $H$  and  $K$ .

Case 6.  $\{s_1, s_2, s_3\} \subseteq X$  and  $\{t_1, t_2, t_3\} \subseteq V-X$ . By Lemma 2.4  $I_H(v, b_1, b_2, b_3, s_1, s_2, s_3) \cap I_K(u, c_1, c_2, c_3, t_1, t_2, t_3) \neq \emptyset$ , and so the result follows.

Next we assume that  $G$  does not contain a nontrivial 3-cut. We may assume that every edge of  $G$  is incident to a vertex of  $T$  (see the proof of Lemma 2.3). Thus  $|E| \leq 18$  and  $|V| \leq 12$ . We require the following.

(2.1) We may assume that  $d(s_i, t_i) \geq 2$  ( $i=1,2,3$ ). If  $d(s_i, t_i)=2$  for some  $i$  and  $s_i, t_i$  are adjacent to a common vertex  $x$ , say for  $i=1$ , then we may assume that

$$x \in (s_2, t_2) \cap (s_3, t_3).$$

Proof. Let  $d(s_1, t_1)=1$ . If  $(s_i, t_i) \cap (s_1, t_1) = \emptyset$ , for  $i=2$  or 3, say  $i=2$ , then  $\lambda_{G-s_1 t_1}(s_2, t_2)=3$  and by Lemma 2.1

$G - s_1, t_1$  contains disjoint paths  $[s_2, t_2]$  and  $[s_3, t_3]$ , and so the result of Lemma 2.5 follows; if not, then we may let  $s_2 = s_1, s_3 = t_1$  and  $s_i \neq t_i$  ( $i=2,3$ ). Let  $y \in N(s_1) - t_1$ . By Lemma 2.3  $G - s_1$  contains disjoint paths  $[s_3, t_3]$  and  $[t_2, y]$ . Thus the result of Lemma 2.5 follows. Hence we may assume that  $d(s_i, t_i) \geq 2$  ( $i=1,2,3$ ). Assume that  $s_1$  and  $t_1$  are adjacent to a vertex  $x$ . Let  $y \in N(x) - \{s_1, t_1\}$ . If  $x \notin T$ , then by Lemma 2.3  $G - x$  contains disjoint paths  $[s_2, t_2]$  and  $[s_3, t_3]$ . If  $x \in T$  and  $x \notin \{s_2, t_2\} \cap \{s_3, t_3\}$ , then we may let  $x = s_2$  and  $s_3 \neq x \neq t_3$ . By Lemma 2.3  $G - x$  contains disjoint paths  $[s_3, t_3]$  and  $[t_2, y]$ , hence Lemma 2.5 holds. Thus (2.1) is proved.

Now we return to the proof of Lemma 2.5. If  $G = K_4$ , then  $d(s_1, t_1) = 1$ , and if  $G = K_{3,3}$ , then  $s_1$  and  $t_1$  are adjacent to common three vertices, contrary to (2.1). If  $G$  is the graph in Figure 2, then we may let  $s_1 = y_1$  without loss of generality. Then  $t_i \neq y_i$  ( $i=4,5,6$ ) by (2.1). If  $t_i = y_i$  ( $i=2$  or  $8$ ), say for  $i=8$ , then  $\{y_4, y_5\} \subseteq \{s_2, t_2\} \cap \{s_3, t_3\}$  by (2.1). So we may let  $y_4 = s_2 = s_3$  and  $y_5 = t_2 = t_3$ , contrary to (2.1). If  $t_i = y_i$  ( $i=3$  or  $7$ ), say for  $i=3$ , then we may let  $y_4 = s_2 = s_3$  by (2.1). Now we can not choose  $\{t_2, t_3\}$  such that  $T$  covers  $E$ , a contradiction. When  $G$  is a cube, in Figure 3 we may let  $s_1 = y_1$  and  $t_i \neq y_i$  ( $i=2,4,5$ ). If  $t_i = y_i$  ( $i=3,6$  or  $8$ ), say for  $i=3$ , then we may let  $y_2 = s_1 = s_3$  and  $y_4 = t_2 = t_3$ , and the result



follows. Thus we may let  $t_1 = y_7$ . Since  $T$  covers all edges, we may let  $(s_1, t_2) = (y_2, y_8)$  and  $(s_3, t_3) = (y_3, y_5)$ , then the result easily follows.

By Lemma 2.2 we may let  $|V| = 10$  or  $12$ . Thus  $|T| \geq 5$ . Note that for each distinct vertices  $x, y \in V$ ,  $N(x) \neq N(y)$ , because  $G$  has no nontrivial 3-cut. We distinguish three cases.

Case 1.  $|T| = 5$ . Let  $s_1 = s_2$ . Now  $|V| = 10$ , and  $G$  is a bipartite graph and the partition of  $V$  is  $(T, V-T)$ . The number of vertices which have distance two from  $s_1 = s_2$  is at least three, and so  $d(s_i, t_i) = 2$  for  $i = 1$  or  $2$ , contrary to (2.1).

Case 2.  $|T| = 6$  and  $|V| = 12$ . Now  $G$  is a bipartite graph and the partition of  $V$  is  $(T, V-T)$ . If the number of vertices which have distance two from  $s_1$  is at least five, then one of such vertices is  $t_1$ , a contradiction; if not, then the number is four, since  $G$  does not contain a nontrivial 3-cut. Thus  $G$  contains a subgraph as illustrated in Figure 4, where  $T = (s_1, x_1, x_2, x_3, x_4, x_5)$ . By (2.1)  $t_1 \neq x_i$  ( $i = 1, 2, 3, 4$ ) and  $(s_j, t_j)$  is not  $(x_1, x_2)$ ,  $(x_1, x_4)$  nor  $(x_2, x_3)$  ( $j = 2, 3$ ), and so we may let  $(x_1, x_3) = (s_2, t_2)$ ,  $(x_2, x_4) = (s_3, t_3)$  and  $x_5 = t_1$ . Now  $(x_5 y_1, x_5 y_2, x_5 y_3) \subseteq E$ . If  $x_i y_i \in E$  ( $i = 1$  or  $2$ ), say for  $i = 1$ , then  $(x_3 y_2, x_3 y_3) \subseteq E$  and  $x_2 y_3 \in E$ . Now the result follows. If  $x_1 y_3 \in E$ , then  $x_3 y_3 \notin E$ , and so  $(x_3 y_1, x_3 y_2) \subseteq E$ , contrary to  $N(y_1) \neq N(y_2)$ .




Figure 4.

Case 3.  $|T|=6$  and  $|V|=10$ . Now both ends of just three edges are in  $T$ , and by (2.1)  $d(s_i, t_i) \geq 3$  ( $i=1,2,3$ ). Thus  $G$  contains a subgraph as illustrated in Figures 5a, 5b, 5c or 5d, where  $T=(x_1, \dots, x_6)$  and  $V-T=(y_1, \dots, y_4)$ .





Figure 5a.





Figure 5b.




Figure 5c.




Figure 5d.

In Figure 5a, we may let  $\{x_1, x_3\} = \{s_1, t_1\}$ ,  $\{x_2, x_5\} = \{s_2, t_2\}$ ,  $\{x_4, x_6\} = \{s_3, t_3\}$  and  $\{x_1, y_1, x_1, y_2\} \subseteq E$ . Then  $x_i y_j \in E$  ( $i=2,3$ ;  $j=1,2$ ). Since  $N(y_1) \neq N(y_2)$ , one of them contains  $\{x_5, x_6\}$  or  $\{x_4, x_6\}$ , a contradiction. In Figure 5b, we may let  $\{x_6, y_1, x_6, y_2, x_6, y_3\} \subseteq E$ . If for some  $i=1,3,4,5$   $\{x_i, x_6\} = \{s_1, t_1\}$ , then  $d(s_1, t_1) = 2$ , a contradiction. Thus we may let  $\{x_2, x_6\} = \{s_1, t_1\}$ ,  $\{x_1, x_4\} = \{s_2, t_2\}$  and  $\{x_3, x_5\} = \{s_3, t_3\}$ . Thus  $x_2 y_4 \in E$ . We may let  $\{x_1, y_1, x_1, y_2\} \subseteq E$ , and so  $\{x_4 y_3, x_4 y_4, x_5 y_1, x_5 y_2\} \subseteq E$ , contrary to  $N(y_1) \neq N(y_2)$ . In Figure 5c, for some  $i=1,2,3$   $d(s_i, t_i) \leq 2$ , a contradiction. In Figure 5d, we may let  $\{x_2, x_5\} = \{s_1, t_1\}$ ,  $\{x_3, x_6\} = \{s_2, t_2\}$ ,  $\{x_1, x_4\} = \{s_3, t_3\}$  and  $\{x_1, y_1, x_1, y_2, x_4 y_3, x_4 y_4\} \subseteq E$ .

Now  $x_2 y_i \in E$  ( $i=3$  or  $4$ ), say for  $i=3$ , then  $\{x_5 y_1, x_5 y_2, x_5 y_4\} \subseteq E$ .  $x_3 y_i \in E$  ( $i=1$  or  $2$ ), say for  $i=1$ , then  $\{x_6 y_2, x_6 y_3, x_6 y_4\} \subseteq E$ . Now the result easily follows.

Proof of Theorem 1. We proceed by induction on  $|V|$ . If  $G$  is not 2-connected, then we can deduce the result by using induction on blocks. Thus we may assume that  $G$  is 2-connected. If  $G$  contains a vertex of degree  $k$  ( $\geq 4$ ), then we replace this vertex by a  $k$ -gon with  $k$  vertices of degree 3. (Figure 6 gives an example with  $k=5$ .) If this vertex of  $G$  is  $s_i(t_i)$  for some  $i$ , then we assign  $s_i(t_i)$  on any vertex of this  $k$ -gon, producing a 3-regular graph  $G'$  such that  $\lambda_{G'}(s_i, t_i) \geq 3$  for each  $i$ . If the result holds in  $G'$ , then the result clearly holds in  $G$ , and so we may assume that  $G$




Figure 6.

is 3-regular. By Lemma 2.5 we may assume that  $G$  contains a 2-cut  $(e_1, e_2) = \partial(X)$  ( $X \subseteq V$ ). Let  $b_i \in X$ ,  $c_i \in V-X$  and  $e_i = b_i c_i$  ( $i=1,2$ ). We define new graphs  $H, K$  as follows.

$$H=(X, E(\langle X \rangle) \cup f),$$

$$K=(V-X, E(\langle V-X \rangle) \cup g),$$

where  $f, g$  are new edges with ends  $b_1, b_2$  and  $c_1, c_2$  respectively. Then  $H$  and  $K$  are 2-edge-connected. Since  $\lambda_G(s_i, t_i) \geq 3$ ,  $\{s_i, t_i\} \subseteq X$  or  $\{s_i, t_i\} \subseteq V-X$  for each  $i$ .

Thus it suffices to consider the following cases.

Case 1.  $\{s_1, s_2, s_3, t_1, t_2, t_3\} \subseteq X$ . By induction the result holds in  $H$ .

Case 2.  $\{s_1, s_2, t_1, t_2\} \subseteq X$  and  $\{s_3, t_3\} \subseteq V-X$ . By Lemma 2.1  $H$  contains disjoint paths  $P_1 [s_1, t_1]$  and  $P_2 [s_2, t_2]$ . Let  $P_3, P_4, P_5$  be disjoint paths of  $K$  between  $s_3$  and  $t_3$ , and let  $c_1, c_2 \notin E(P_3) \cup E(P_4)$ . If  $b_1, b_2 \notin E(P_1) \cup E(P_2)$ , then  $P_1, P_2, P_3$  are required paths of  $G$ . Thus let  $b_1, b_2 \in E(P_1)$ . If  $c_1, c_2 \notin E(P_5)$ , then by Lemma 2.1  $K - c_1, c_2$  contains disjoint paths  $[s_3, t_3]$  and  $[c_1, c_2]$ ; and if  $c_1, c_2 \in E(P_5)$ , then let  $P_6 [c_1, c_2]$  be the path obtained by combining  $P_5 - c_1, c_2$  and  $P_4$ . In each case we can construct required paths of  $G$ .

### 3. Proof of Theorem 2.

For an integer  $n \geq 3$  and vertices  $x_1, x_2, \dots, x_n$ , we denote feasible paths  $\frac{1}{2}[x_1, x_2], \frac{1}{2}[x_2, x_3], \dots, \frac{1}{2}[x_{n-1}, x_n]$ , and  $\frac{1}{2}[x_n, x_1]$  by  $\frac{1}{2}[x_1, \dots, x_n, x_1]$ . For a vertex  $x \in V$  and  $a, b \in N(x)$ , let  $G_x^{a,b}$  be the graph  $(V, E \cup e_1 - \{e_2, e_3\})$ , where  $e_1$  is a new edge with ends  $a, b$  and  $e_2, e_3$  are edges of  $E$  with ends  $a, x$  and  $b, x$  respectively.

Lemma 3.1 (Mader [4]). Suppose that  $G$  is a graph and  $x$  is a non-separating vertex of  $G$  with  $\deg x \geq 4$  and with  $|N(x)| \geq 2$ . Then there exist two vertices  $a, b \in N(x)$ , such that for each two vertices  $y, z \in V - x$ ,

$$\lambda_{G_x^{a,b}}(y, z) = \lambda_G(y, z).$$

Lemma 3.2. Suppose that  $x_1, \dots, x_5$  are vertices of a graph  $G$ . If for each  $1 \leq i < j \leq 5$ ,

$$\lambda(x_i, x_j) \geq 4,$$

and each vertex of  $G$  has even degree, then  $G$  contains edge-disjoint paths  $[x_1, x_2], [x_2, x_3], [x_3, x_4], [x_4, x_5]$ , and  $[x_5, x_1]$ .

Proof. We proceed by induction on  $|E|$ . We put  $T = \{x_1, \dots, x_5\}$ . If  $|T| \leq 4$ , then the result follows from (1.4.4), and so we may let  $|T| = 5$ . We may assume that  $G$  is 2-connected, and that for each vertex  $x$  of  $G$   $\deg x \geq 4$ . If there exists a vertex  $x$  in  $V - T$ , then by Lemma 3.1 there

exist two vertices  $a, b \in N(x)$  such that  $\lambda_{G-x}^{a,b}(x_i, x_j) = \lambda_G(x_i, x_j)$  ( $1 \leq i < j \leq 5$ ).  $|E(G_x^{a,b})| < |E|$  and each vertex of  $G_x^{a,b}$  has even degree, thus by induction the result holds in  $G_x^{a,b}$ , and so in  $G$ . Let  $V=T$ . If  $x_5 x_i \in E$ , then we can apply (1.4.5) for the graph  $G - x_5 x_i$ , and for pairs  $(s_i, t_i) = (x_i, x_{i+1})$  and  $q_i = 1$  ( $1 \leq i \leq 4$ ). Thus we may let  $x_5 x_i \notin E$  and  $x_i x_{i+1} \notin E$  ( $1 \leq i \leq 4$ ). Now  $G$  contains a subgraph as illustrated in Figure 7a or 7b, and the result holds.




Figure 7a.




Figure 7b.

Lemma 3.3. Suppose that  $G$  is a 2-edge-connected graph and  $a, b, c, d, x, y$  are vertices such that  $\deg a = 3$ ,  $N(a) = \{b, c, d\}$ ,  $\deg b \geq 3$ , and  $a, x, y$  are all distinct, and that for each

2-cut  $\partial(X)$  ( $X \subseteq V$ ,  $|X| \leq |V-X|$ ),

$X=\{x\}$ ,  $X=\{y\}$  or  $X=\{x,y\}$  and  $|E(\langle X \rangle)|=1$ .

Then  $G-a$  contains  $\frac{1}{2}[b,c,x,d,y,b]$ , if it is not the cases that

$\deg c=2$ ,  $c=x$ ,  $\deg c_1=2$  ( $N(c)=\{a,c_1\}$ ) or,

$\deg c=2$ ,  $c=y$ .

Proof. We distinguish four cases.

Case 1.  $\deg c \geq 3$  and  $\deg d \geq 3$ . Now  $G-a$  is 2-edge-connect. Let  $G'$  be the graph obtained by replacing each edge of  $G$  by double edges. Then  $G'-a$  is 4-edge-connected, and so by applying Lemma 3.2 on  $G'-a$  we can deduce the result.

Case 2.  $\deg c=2$  and  $\deg d \geq 3$ . Let  $N(c)=\{a,c_1\}$ . By the hypothesis  $c \neq y$ , and so  $c=x$  and  $\deg c_1 \geq 3$ .  $G-\{a,c\}$  is 2-edge-connected, and so this contains  $\frac{1}{2}[b,c_1,d,y,b]$  by Lemma 3.2.

Case 3.  $\deg c \geq 3$  and  $\deg d=2$ . Let  $d=x$  and  $N(d)=\{a,d_1\}$ . If  $\deg d_1 \geq 3$ , then  $G-\{a,d\}$  is 2-edge-connected, and so this contains  $\frac{1}{2}[b,c,d_1,y,b]$ . If  $\deg d_1=2$ , then  $d_1=y$ . By (1.4.4)  $G-\{a,d\}$  contains  $\frac{1}{2}[b,c,d_1,b]$ , thus  $G$  contains  $\frac{1}{2}[b,c,x,d,y,b]$ . When  $d=y$ , the proof is similar.

Case 4.  $\deg c=\deg d=2$ . Now  $c \neq d$  and  $c \neq y$ , thus  $c=x, d=y$ , and  $G-\{a,c,d\}$  is 2-edge-connected. By (1.4.4)  $G-a$  contains  $\frac{1}{2}[b,c,d,b]$ .

If we prove following Lemma 3.4, Theorem 2 follows.

Lemma 3.4. Suppose that  $G$  is a graph with  $w \equiv 1$ ,  $(s_1, t_1)$ ,  $(s_2, t_2)$ ,  $(s_3, t_3)$  are pairs of vertices of  $G$ , and  $q_1=q_2=q_3=1$ .



If (1.3) holds, then  $G$  contains feasible paths  $\frac{1}{2}P_1[s_1, t_1]$ ,  $\frac{1}{2}P_2[s_1, t_1]$ ,  $\frac{1}{2}P_3[s_2, t_2]$ ,  $\frac{1}{2}P_4[s_2, t_2]$ ,  $\frac{1}{2}P_5[s_3, t_3]$ , and  $\frac{1}{2}P_6[s_3, t_3]$ .

Proof. We proceed by induction on  $|E|$ . We put  $T = \{s_1, s_2, s_3, t_1, t_2, t_3\}$ . We require the following

(3.1) We may assume the following.

(3.1.1)  $G$  is 2-connected, and  $|T| = 6$ .

(3.1.2) For each 2-cut  $\partial(X)$  ( $X \subseteq V$ ),  $|X| = |X \cap T| = 1$  or  $|X \cap T| \geq 2$ .

(3.1.3) For each edge  $e \in E$ , there exists  $X \subseteq V$  such that  $|\partial(X)| = |D(X)|$  and  $e \in \partial(X)$ .

(3.1.4) For each  $1 \leq i \leq 3$ ,  $s_i$  and  $t_i$  are not adjacent.

(3.1.5) If for vertices  $x_1, x_2, x_3, x_4$  of  $G$   $\deg x_2 = \deg x_3 = 2$  and  $\{x_1 x_2, x_2 x_3, x_3 x_4\} \subseteq E$ , then  $\deg x_1 \geq 3$  and  $\deg x_4 \geq 3$ .

Proof. (1) If  $|T| \leq 5$ , then Lemma 3.4 follows from (1.4.5).

(2) Let  $\{e_1, e_2\} = \partial(X)$  be a 2-cut, and let  $a_i \in X$ ,  $b_i \in V - X$  and  $a_i b_i = e_i$  ( $i = 1, 2$ ). We define new graphs  $H, K$  as follows.

$$H = (X, E(\langle X \rangle) \cup f),$$

$$K = (V - X, E(\langle V - X \rangle) \cup g),$$

where  $f, g$  are new edges with ends  $a_1, a_2$  and  $b_1, b_2$  respectively. If  $X \cap T = \emptyset$ , then by induction the result of Lemma 3.4 holds in  $K$ , and so in  $G$ . If  $|X \cap T| = 1$  (say  $s_i \in X$ ) and  $|X| \geq 2$ , then we assign  $s_i$  on the midpoint of  $g$  in  $K$ ,

producing a new graph  $K'$ . Now by induction the result of Lemma 3.4 holds in  $K'$ , and so in  $G$ .

(3) If there exists  $e \in E$  such that for each  $X \subseteq V$  with  $e \in \partial(X)$  and  $|\partial(X)| > |D(X)|$ , then the hypothesis of Lemma 3.4 holds in  $G-e$ , and so we can apply induction on  $G-e$ .

(4) If  $s_3 t_3 \in E$ , then we can apply (1.4.2) for the graph  $G-s_3 t_3$ , and for two pairs  $(s_1, t_1), (s_2, t_2)$  and  $q_1 = q_2 = 1$ .

(5) If  $\deg x_i = 2$ , then  $x_i \in T$  ( $1 \leq i \leq 3$ ) by (3.1.2), and so we may let  $x_1 = s_2$ ,  $x_2 = s_1$  and  $x_3 = t_2$  by (3.1.4) and (1.3). Let  $x_0 \in N(x_1) - x$ . Let  $G'$  be the graph obtained by contracting the edge  $x_0 x_1$ . By induction  $G'$  contains feasible paths  $\frac{1}{2}P_1[s_1, t_1]$ ,  $\frac{1}{2}P_2[s_1, t_1]$ ,  $\frac{1}{2}P_3[s_2, t_2]$ ,  $\frac{1}{2}P_4[s_2, t_2]$ ,  $\frac{1}{2}P_5[s_3, t_3]$  and  $\frac{1}{2}P_6[s_3, t_3]$ . Let  $Q_1, \dots, Q_6$  be the corresponding paths of  $G$ . We may let  $x_1 x_2 \in E(Q_1) \cap E(Q_2)$  or  $x_1 x_2 \in E(Q_1) \cap E(Q_3)$ . In the former case, let  $Q_7$  be the path of  $G$  such that  $E(Q_7) = (x_1 x_2, x_2 x_3)$  and let  $Q_8$  be the path of  $G$  obtained by combining  $x_2 x_3$ ,  $Q_3(t_2, x_0)$  and  $Q_2(x_0, t_1)$ . Then  $\frac{1}{2}Q_1, \frac{1}{2}Q_8, \frac{1}{2}Q_7, \frac{1}{2}Q_4, \frac{1}{2}Q_5, \frac{1}{2}Q_6$  are required paths of  $G$ . In the latter case  $\frac{1}{2}Q_1, \dots, \frac{1}{2}Q_6$  are required paths of  $G$ .

Now we come to the proof of Lemma 3.4. We distinguish three cases.

Case 1.  $G$  contains a nontrivial 2-cut  $(e_1, e_2) = \partial(X)$  ( $X \subseteq V$ ). We define  $H, K, a_i, b_i, f$  and  $g$  similarly as in the proof of (3.1.2). Then  $H$  and  $K$  are 2-edge-connected. It suffices to consider the following cases by (3.1.2).

Case 1a.  $\{s_1, t_1\} \subseteq X$  and  $\{s_2, s_3, t_2, t_3\} \subseteq V-X$ . Assume that  $K-g$  contains feasible paths  $\frac{1}{2}P_1[s_2, t_2]$ ,  $\frac{1}{2}P_2[s_2, t_2]$ ,  $\frac{1}{2}P_3[s_3, t_3]$  and  $\frac{1}{2}P_4[s_3, t_3]$ . Then  $H-f$  contains a path  $P_5[s_1, t_1]$ , and  $P_5, \frac{1}{2}P_1, \frac{1}{2}P_2, \frac{1}{2}P_3, \frac{1}{2}P_4$  are required paths of  $G$ . If this is not the case, then by (1.4.2) for the graph  $K-g$ , and for two pairs  $(s_2, t_2), (s_3, t_3)$  and  $q_2=q_3=1$ , (1.3) does not hold. Thus for some  $Y \subseteq V-X$  with  $b_1 \in Y$ ,

$$D_{K-g}(Y) = \{2, 3\} \text{ and } |\partial_{K-g}(Y)| = 1.$$

For each  $Z \subseteq X$  such that  $a_1 \in Z$ ,  $f \in \partial_H(Z)$  and  $D_H(Z) = \{1\}$ ,

$$|D_G(Y \cup Z)| = 3,$$

$$\text{and so } |\partial_G(Y \cup Z)| = |\partial_{H-f}(Z)| + |\partial_{K-g}(Y)| \geq 3,$$

$$\text{thus } |\partial_{H-f}(Z)| \geq 2.$$

Hence by (1.4.2)  $H-f$  contains feasible paths  $\frac{1}{2}P_1[s_1, t_1]$ ,  $\frac{1}{2}P_2[s_1, t_1]$ ,  $\frac{1}{2}P_3[a_1, a_2]$  and  $\frac{1}{2}P_4[a_1, a_2]$ , and  $K$  contains feasible paths  $\frac{1}{2}P_5[s_2, t_2]$ ,  $\frac{1}{2}P_6[s_2, t_2]$ ,  $\frac{1}{2}P_7[s_3, t_3]$  and  $\frac{1}{2}P_8[s_3, t_3]$ . Now we can construct required paths of  $G$ .

Case 1b.  $\{s_1, s_2\} \subseteq X$  and  $\{s_3, t_1, t_2, t_3\} \subseteq V-X$ . If  $H-f$  is 2-edge-connected, then we assign a new vertex  $u$  on the midpoint of  $g$ , producing a new graph  $K'$ . By induction  $K'$  contains feasible paths  $\frac{1}{2}P_1[u, t_1]$ ,  $\frac{1}{2}P_2[u, t_1]$ ,  $\frac{1}{2}P_3[u, t_2]$ ,  $\frac{1}{2}P_4[u, t_2]$ ,  $\frac{1}{2}P_5[s_3, t_3]$  and  $\frac{1}{2}P_6[s_3, t_3]$ . We may let  $ub_1 \in E(P_1) \cap E(P_2)$  or  $ub_1 \in E(P_1) \cap E(P_3)$ . In each case we can construct required paths of  $G$ , since by (1.4.5)  $H-f$  contains feasible paths  $\frac{1}{2}P_7[s_1, a_1]$ ,  $\frac{1}{2}P_8[s_1, a_1]$ ,  $\frac{1}{2}P_9[s_2, a_2]$  and  $\frac{1}{2}P_{10}[s_2, a_2]$  and contains  $\frac{1}{2}[s_1, a_1, s_2, a_2, s_1]$ . Thus we may assume that  $H-f$  is not 2-edge-connected, and so  $H$  contains a 2-cut

$\{f, f'\}$ . Then  $\{f', e_1\}$  and  $\{f', e_2\}$  are 2-cuts of  $G$ . By (3.1.2)  $H = (\{a_1, a_2\}, \{f, f'\})$ , and so we may let  $a_1 = s_1$  and  $a_2 = s_2$ . By (3.1.5)  $\deg b_i \geq 3$  ( $i=1,2$ ). If  $K$  contains a 2-cut  $\{g, g'\} = \partial_K(Y)$  ( $Y \subseteq V-X$ ), then  $\{g', e_1\}$  and  $\{g', e_2\}$  are 2-cuts of  $G$ . Since  $\deg b_i \geq 3$  ( $i=1,2$ ), by (3.1.2)  $|Y \cap T| = 2$  and  $|(V-X-Y) \cap T| = 2$ . By Case 1a we may let  $Y \cap T \neq \{s_3, t_3\}$ , and so we may let  $\langle Y \rangle$  is an edge, contrary to (3.1.5). Thus assume that  $K-g$  is 2-edge-connected. By (3.1.3), there exists  $X \subseteq V$  such that  $|\partial(X)| = |D(X)|$  and  $s_1, s_2 \in \partial(X)$ . Thus we may assume that  $G$  is the graph as illustrated in Figure 8. Let  $Y_1, Y_2$  be the subsets




Figure 8.

of  $V$  such that  $b_i \in Y_i$  and  $\partial(Y_i) = \{s_i, b_i, c_1, c_2, d_1, d_2\}$  ( $i=1,2$ ). We construct new graphs  $K_1, K_2$  as follows.

$$K_i = (Y_i \cup v_i, E(\langle Y_i \rangle) \cup \{b_i, v_i, c_i, v_i, d_i, v_i\}), \quad i=1,2,$$

where  $v_1, v_2$  are new vertices. If for  $i=1$  or  $2$ ,  $K_i$  contains a 2-cut  $\partial_{K_i}(Z_i)$  ( $Z_i \subseteq Y_i$ ) such that  $|Z_i| \geq 2$  and  $|V(K_i) - Z_i| \geq 2$ , say  $i=1$ , then  $\partial_G(Z_1)$  is a 2-cut of  $G$ , and by (3.1.2)  $Z_1 \cap T = \{s_3, t_2\}$ . Thus we may assume that  $Z_1 = \{s_3, t_2\}$  and  $\deg s_3 = \deg t_2 = 2$ . This allows that we can apply Lemma 3.3 on

$K_1$  and  $K_2$ .

Assume that  $\deg c_1 \geq 3$ ,  $\deg d_2 \geq 3$  or  $\deg c_2 \geq 3$ ,  $\deg d_1 \geq 3$ , say the former. By Lemma 3.3  $K_1 - v_1$  contains  $\frac{1}{2}[b_1, c_1, s_3, d_1, t_2, b_1]$  and  $K_2 - v_2$  contains  $\frac{1}{2}[b_2, d_2, t_3, c_2, t_1, b_2]$ . Now we can construct required paths of  $G$ . Assume that for  $i=1$  or  $2$ ,  $\deg c_i \geq 3$  and  $\deg d_i \geq 3$ , say for  $i=1$ . Now we may assume that  $\deg c_2 = \deg d_2 = 2$ .  $c_2 = t_1$ ,  $d_2 = t_3$  or  $c_2 = t_3$ ,  $d_2 = t_1$ , say the former, then by Lemma 3.3  $K_2 - v_2$  contains  $\frac{1}{2}[b_2, d_2, t_3, c_2, t_1, b_2]$  and  $K_1 - v_1$  contains  $\frac{1}{2}[b_1, c_1, s_3, d_1, t_2, b_1]$ . Assume that  $\deg c_i = 2$  ( $i=1,2$ ) or  $\deg d_j = 2$  ( $j=1,2$ ), say the former. Let  $y_1 \in N(c_1) - c_2$ , and let  $y_2 \in N(c_2) - c_1$ , then by (3.1.5)  $\deg y_i \geq 3$  ( $i=1,2$ ). By (3.1.4) we may let  $c_1 = t_2$ ,  $c_2 = t_1$  or  $c_1 = s_3$ ,  $c_2 = t_1$ . If  $c_1 = t_2$ , then by (1.4.2)  $K_1 - (v_1, c_1)$  contains feasible paths  $\frac{1}{2}P_1[s_3, d_1]$ ,  $\frac{1}{2}P_2[s_3, d_1]$ ,  $\frac{1}{2}P_3[b_1, y_1]$  and  $\frac{1}{2}P_4[b_1, y_1]$ , and  $K_2 - (v_2, c_2)$  contains feasible paths  $\frac{1}{2}P_5[t_3, d_2]$ ,  $\frac{1}{2}P_6[t_3, d_2]$ ,  $\frac{1}{2}P_7[b_2, y_2]$  and  $\frac{1}{2}P_8[b_2, y_2]$ , and so the result follows. If  $c_1 = s_3$ , then by Lemma 3.3  $K_1 - (v_1, c_1)$  contains  $\frac{1}{2}[b_1, y_1, d_1, t_2, b_1]$  and  $K_2 - (v_2, c_2)$  contains  $\frac{1}{2}[b_2, d_2, t_3, y_2, b_2]$ , and so the result follows.

Case 1c.  $\{s_1, s_2, t_1\} \subseteq X$  and  $\{s_3, t_2, t_3\} \subseteq V - X$ . We may assume that neither Case 1a nor Case 1b occurs. If  $\deg a_1 = 2$ , then  $\partial(X - a_1)$  is a 2-cut of  $G$  and  $|(X - a_1) \cap T| = 2$ , a contradiction. Thus  $\deg a_i \geq 3$  and  $\deg b_i \geq 3$  ( $i=1,2$ ). We assign new vertices  $v_1, u_2$  on the midpoints of  $f, g$  respectively, producing new graphs  $H', K'$ . For the graph  $H'$ , and for two pairs  $(s_1, t_1), (s_2, v_2)$  and  $q_1 = 1, q_2 = 2$ , if (1.3) does not hold, then

there exists  $Z \subseteq V(H')$  such that  $D_{H'}(Z) = \{1, 2\}$  and  $|\partial_{H'}(Z)| = 2$ . Now Case 1b occurs in  $G$ , thus (1.3) holds, and so (1.2) holds. Hence  $H' - v_2$  contains feasible paths  $\frac{1}{2}P_1[s_1, t_1]$ ,  $\frac{1}{2}P_2[s_1, t_1]$ ,  $\frac{1}{2}P_3[s_2, a_1]$  and  $\frac{1}{2}P_4[s_2, a_2]$ . Similarly  $K' - u_2$  contains feasible paths  $\frac{1}{2}P_5[s_3, t_3]$ ,  $\frac{1}{2}P_6[s_3, t_3]$ ,  $\frac{1}{2}P_7[t_2, b_1]$  and  $\frac{1}{2}P_8[t_2, b_2]$ , and so the result follows.

Case 2. Every 2-cut of  $G$  is trivial, and  $G$  contains a 2-cut. Now we may let  $\deg s_1 = 2$ , and let  $e_1, e_2$  be the edges incident to  $s_1$ . By (3.1.3), for  $i=1, 2$  there exists  $X_i \subseteq V$  such that  $s_1 \in X_i$ ,  $|\partial(X_i)| = |D(X_i)|$  and  $e_i \in \partial(X_i)$ . For  $i=1, 2$ , since  $|\partial(X_i)| = 3$ , let  $\partial(X_i) = \{e_i, f_i, g_i\}$ . We put  $X_3 = V - (X_1 \cup X_2)$ , then  $t_1 \in X_3$ . By simple counting we have

$$(3.2) \quad |\partial(X_1 \cup X_2)| = |\partial(X_1)| + |\partial(X_2)| - |\partial(X_1 \cap X_2)| \\ - 2|\partial(X_1 - X_2) \cap \partial(X_2 - X_1)|.$$

If  $|\partial(X_1 \cap X_2)| \geq 4$ , then by (3.2)

$$|\partial(X_3)| = |\partial(X_1 \cup X_2)| \leq 3 + 3 - 4 = 2.$$

Thus  $|\partial(X_3)| = 2$  and  $|\partial(X_1 \cap X_2)| = 4$ . Then  $|X_3| = 1$  and  $X_1 \cap X_2 = \{s_1, x\}$  for some  $x \in V$  with  $\deg x = 2$ . We may let  $x = s_2$ , then  $t_2 \in X_3$ , and so  $t_1 = t_2$ , a contradiction. Thus  $|\partial(X_1 \cap X_2)| = 2$  and  $X_1 \cap X_2 = \{s_1\}$ .

Case 2a.  $f_1, f_2, g_1, g_2$  are not all distinct. We may let  $f_1 = f_2$ . Since  $f_1 \notin \partial(X_1 \cap X_2) = \{e_1, e_2\}$ ,  $f_1 \in \partial(X_1 - X_2) \cap \partial(X_2 - X_1)$ . By (3.2)

$$|\partial(X_3)| = |\partial(X_1 \cup X_2)| \leq 3 + 3 - 2 - 2 = 2.$$

Thus  $X_3 = \{t_1\}$ , and we may assume that  $G$  is the graph as illustrated in Figure 9.




Figure 9.

Since every 2-cut is trivial,  $\deg b_i \geq 3$  and  $\deg c_i \geq 3$  ( $i=1,2$ ). By Lemma 3.3  $\langle X_1 \rangle$  contains  $\frac{1}{2}[b_1, s_3, d_1, s_2, c_1, b_1]$  and  $\langle X_2 \rangle$  contains  $\frac{1}{2}[b_2, t_3, d_2, t_2, c_2, b_2]$ , and so the result follows.

Case 2b.  $f_1, f_2, g_1, g_2$  are all distinct. Now  $\partial(X_1 - X_2) \cap \partial(X_2 - X_1) = \emptyset$ . From (3.2) we have

$$|\partial(X_3)| = 3 + 3 - |\partial(X_1 \cap X_2)| = 4.$$

Thus we may assume that  $G$  is the graph as illustrated in Figure 10.




Figure 10.

$\langle X_3 \rangle$  is connected. For if not, then there exist  $Y_1, Y_2 \subseteq X_3$  such that  $X_3 = Y_1 \cup Y_2$ ,  $Y_1 \cap Y_2 = \emptyset$ , and  $|\partial(Y_1)| = |\partial(Y_2)| = 2$ . Then  $|\partial(X_3 \cap T)| = 2$ , a contradiction.  $\deg c_i \geq 3$  and  $\deg d_i \geq 3$  ( $i=3,4$ ), for if not, then  $\deg t_i = 2$  and one of  $f_1, f_2, g_1, g_2$  is incident to

$t_1$ , say  $g_1$ , and Case 2a occurs for  $X_1, X_2 \cup X_3 - t_1$  instead of  $X_1, X_2$ . If  $\langle X_3 \rangle$  contains a 1-cut  $(h) = \partial_{\langle X_3 \rangle}(Y_1)$  ( $Y_1 \subseteq X_3$ ), then  $Y_1$  contains just two of  $c_3, c_4, d_3, d_4$ . Put  $Y_2 = X_3 - Y_1$ .  $(c_3, d_3) \not\subseteq Y_1$ , thus we may let  $(c_3, c_4) \subseteq Y_1, (d_3, d_4) \subseteq Y_2$  and  $t_1 \in Y_1$ . Let  $v_1, v_2$  be the vertices such that  $v_i \in Y_i$  ( $i=1,2$ ) and  $v_1 v_2 = h$ .  $\langle Y_1 \rangle$  contains  $\frac{1}{2}[c_3, c_4, v_1, t_1, c_3]$ , for if  $\langle Y_1 \rangle$  is not 2-edge-connected, then  $\deg v_1 = 2$  and  $v_1 = t_1$ .  $\langle Y_2 \rangle$  contains  $\frac{1}{2}[d_3, d_4, v_2, d_3]$ . Thus  $\langle X_3 \rangle$  contains  $\frac{1}{2}[c_3, t_1, d_4, d_3, c_4, c_3]$  and feasible paths  $\frac{1}{2}[c_3, c_4], \frac{1}{2}[c_3, t_1], \frac{1}{2}[t_1, c_4]$  and  $[d_3, d_4]$ . If  $\langle X_3 \rangle$  is 2-edge-connected, then  $\langle X_3 \rangle$  contains  $\frac{1}{2}[c_3, t_1, d_4, d_3, c_4, c_3]$  by Lemma 3.2. Assume that  $\deg c_1 = 2$ . We may let  $c_1 = s_2$ .  $\langle X_1 \rangle$  contains  $\frac{1}{2}[b_1, s_2, d_1, s_3, b_1]$  and  $\langle X_3 \rangle$  contains  $\frac{1}{2}[c_3, t_1, d_4, d_3, c_4, c_3]$ . If  $\deg d_2 \geq 3$  or  $\deg d_2 = 2, d_2 = t_2$ , then by Lemma 3.3  $\langle X_2 \rangle$  contains  $\frac{1}{2}[b_2, t_3, c_2, t_2, d_2, b_2]$ . If  $\deg d_2 = 2$  and  $d_2 = t_3$ , then  $\langle X_2 \rangle$  contains feasible paths  $\frac{1}{2}P_1[b_2, t_3], \frac{1}{2}P_2[b_2, t_3], \frac{1}{2}P_3[c_2, t_2]$  and  $\frac{1}{2}P_4[c_2, t_2]$ . Now we can deduce the result. Thus we may assume that  $\deg c_i \geq 3$  ( $i=1,2$ ). By Lemma 3.3  $\langle X_1 \rangle$  contains  $\frac{1}{2}[b_1, c_1, s_2, d_1, s_3, b_1]$  and  $\langle X_2 \rangle$  contains  $\frac{1}{2}[b_2, c_2, t_2, d_2, t_3, b_2]$ . If  $\langle X_3 \rangle$  is 2-edge-connected, then by Lemma 3.2  $\langle X_3 \rangle$  contains  $\frac{1}{2}[c_3, t_1, c_4, d_3, d_4, c_3]$ ; and if not, then  $\langle X_3 \rangle$  contains feasible paths  $\frac{1}{2}[c_3, c_4], \frac{1}{2}[c_3, t_1], \frac{1}{2}[t_1, c_4]$  and  $[d_3, d_4]$ . Now we can deduce the result.

Case 3.  $G$  is 3-edge-connected. By Theorem 1 the result follows.



## References

- [1] A. Cypher: An approach to the k paths problem, Proc. 12th Annual ACM Symposium on Theory of Computing (1980) 211-217.
- [2] L. R. Ford, Jr. and D. R. Fulkerson: Maximal flow through a network, Canad. J. Math. 8 (1956), 399-404.
- [3] T. C. Hu: Multi-commodity network flows, Operations Res. 11 (1963), 344-360.
- [4] W. Mader: A reduction method for edge-connectivity in graphs, Annals of Discrete Math. 3 (1978), 145-164.
- [5] H. Okamura and P. D. Seymour: Multicommodity flows in planar graphs, J. Combinatorial Theory Ser. B 31 (1981), 75-81.
- [6] H. Okamura: Multicommodity flows in graphs, Discrete Applied Math. 6 (1983), 55-62.
- [7] B. A. Papernov: Realizability of multi-product flows, (in Russian), Compendium Investigations in Discrete Optimization, Nauka, Moscow, (1976).
- [8] P. D. Seymour: A short proof of two-commodity flow theorem, J. Combinatorial Theory Ser. B 26 (1979), 370-371.
- [9] P. D. Seymour: Four terminus flows, Networks 10 (1980), 79-86.

- [10] P. D. Seymour: Disjoint paths in graphs, *Discrete Math.* 29 (1980), 293-309.
- [11] P. D. Seymour: On odd cuts and plane multicommodity flows, *Proc. London Math. Soc.* (3) 42 (1981), 178-192.
- [12] C. Thomassen: 2-linked graphs, *Europ. J. Combinatorics* 1 (1980), 371-378.