

Title	Nash G manifold structures of compact or compactifiable $C^\infty$ G manifolds
Author(s)	川上, 智博
Citation	大阪大学, 1996, 博士論文
Version Type	VoR
URL	<a href="https://doi.org/10.11501/3110235">https://doi.org/10.11501/3110235</a>
rights	
Note	


*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

NASH  $G$  MANIFOLD STRUCTURES OF COMPACT  
OR COMPACTIFIABLE  $C^\infty G$  MANIFOLDS

TOMOHIRO KAWAKAMI



# NASH $G$ MANIFOLD STRUCTURES OF COMPACT OR COMPACTIFIABLE $C^\infty G$ MANIFOLDS

TOMOHIRO KAWAKAMI

Department of Liberal Arts, Osaka Prefectural  
College of Technology, Neyagawa Osaka 572, Japan

## 1. Introduction.

Let  $G$  be a compact affine Nash group. We say that a  $C^\infty G$  manifold  $X$  admits a (resp. an *affine*, a *nonaffine*) Nash  $G$  manifold structure if there exists a (resp. an affine, a nonaffine) Nash  $G$  manifold  $Y$  such that  $X$  is  $C^\infty G$  diffeomorphic to  $Y$ . In the present paper we consider Nash  $G$  manifold structures of compact or compactifiable  $C^\infty G$  manifolds.

We have the following when  $X$  is compact.

**Theorem 1.** *Let  $G$  be a compact affine Nash group and let  $X$  be a compact  $C^\infty G$  manifold with  $\dim X \geq 1$ .*

(1)  *$X$  admits exactly one affine Nash  $G$  manifold structure up to Nash  $G$  diffeomorphism.*

(2) *If  $G$  acts on  $X$  transitively then a Nash  $G$  manifold structure of  $X$  is unique up to Nash  $G$  diffeomorphism.*

(3) *If  $X$  is connected and the action on  $X$  is not transitive, then  $X$  admits a continuum number of nonaffine Nash  $G$  manifold structures.*

In the non-equivariant category, M. Shiota in [4] proved that any compactifiable  $C^\infty$  manifold  $X$  admits a continuum number of nonaffine Nash manifold structures. When  $X$  is not compact but compactifiable, an affine Nash compactification of  $X$  is not unique, and the number of affine ones can be investigated by the cardinality of the Whitehead torsion of  $X$  [6]. Here an affine Nash compactification of  $X$  means an affine Nash manifold  $Y$  with boundary so that  $X$  is  $C^\infty$  diffeomorphic to the interior of  $Y$ .

We say that a  $C^\infty G$  manifold  $X$  is *compactifiable as a  $C^\infty G$  manifold* if there exists a compact  $C^\infty G$  manifold  $Y$  with boundary so that  $X$  is  $C^\infty G$  diffeomorphic to the interior of  $Y$ . We obtain the following.

**Theorem 2.** *Let  $G$  be a compact affine Nash group and let  $X$  be a non-compact compactifiable  $C^\infty G$  manifold with  $\dim X \geq 1$ .*

(1)  *$X$  admits an affine Nash  $G$  manifold structure.*

(2)  *$X$  admits a continuum number of nonaffine Nash  $G$  manifold structures.*

This paper consists of two parts. The first half is to investigate Nash  $G$  manifold structures of compact  $C^\infty G$  manifolds. We consider Nash  $G$  manifold structures of compactifiable (not compact)  $C^\infty G$  manifolds in the latter half.

In this paper all Nash  $G$  manifolds and all Nash  $G$  maps are of class  $C^\omega$  unless otherwise stated.

**Acknowledgement.** I would like to thank Professor M. Shiota for many useful conversations and suggestions. Theorem 1 (2) is due to the cooperation of Professor M. Shiota.

## 2. Nash $G$ manifolds.

First of all we recall the definition of Nash groups.

*Definition 2.1.* A group is called a (resp. an *affine*) *Nash group* if it is a (resp. an affine) Nash manifold and that the multiplication  $G \times G \rightarrow G$ , the inversion  $G \rightarrow G$  are Nash maps.

We remark that one-dimensional Nash groups are classified by J.J. Madden and C.M. Stanton [2].

Let  $G$  be an affine Nash group. In this paper, a *representation* of  $G$  means a Nash group homomorphism  $G \rightarrow GL(\mathbb{R}^n)$  for some  $\mathbb{R}^n$ . Here a Nash group homomorphism means a group homomorphism which is a Nash map. We use a representation as a representation space.

*Definition 2.2.* Let  $G$  be an affine Nash group

- (1) An affine Nash submanifold in some representation of  $G$  is called an *affine Nash  $G$  submanifold* if it is  $G$  invariant. A Nash manifold  $X$  with  $G$  action is said to be a *Nash  $G$  manifold* if the action map  $G \times X \rightarrow X$  is a Nash map.
- (2) Let  $X$  and  $Y$  be Nash  $G$  manifolds. A Nash map  $f : X \rightarrow Y$  is called a *Nash  $G$  map* if it is a  $G$  map. We say that  $X$  is *Nash  $G$  diffeomorphic to  $Y$*  if there exist Nash  $G$  maps  $f : X \rightarrow Y, h : Y \rightarrow X$  so that  $f \circ h = id, h \circ f = id$ .
- (3) A Nash  $G$  manifold  $X$  is said to be *affine* if there exists an affine Nash  $G$  submanifold  $Y$  so that  $X$  is Nash  $G$  diffeomorphic to  $Y$ .

Tubular neighborhood theorem and collaring theorem are well known in the smooth equivariant category. They are proved in the Nash category by M. Shiota (Lemma 1.3.2 [7], Lemma 6.1.6 [7]). Since M. Shiota's proofs work in the equivariant Nash category, the following two propositions are obtained.

**Proposition 2.3.** *Let  $G$  be a compact affine Nash group and let  $X$  be an affine Nash  $G$  submanifold in a representation  $\Omega$  of  $G$ . Then there exists a Nash  $G$  tubular neighborhood  $(U, p)$  of  $X$  in  $\Omega$ , namely,  $U$  is an affine Nash  $G$  submanifold in  $\Omega$  and the orthogonal projection  $p : U \rightarrow X$  is a Nash  $G$  map.  $\square$*

**Proposition 2.4.** *Let  $G$  be a compact affine Nash group. Any compact affine Nash  $G$  manifold  $X$  with boundary  $\partial X$  admits a Nash  $G$  collar, that is, there exists a Nash  $G$  imbedding  $\phi : \partial X \times [0, 1] \rightarrow X$  so that  $\phi|_{\partial X \times 0} = id_{\partial X}$ , where the action on the closed unit interval  $[0, 1]$  is trivial.  $\square$*

## 3. Compact $C^\infty G$ manifolds.

Recall a theorem proved by K.H. Dovermann, M. Masuda, and T. Petrie [1], which is a partial solution of the equivariant Nash conjecture.

**Theorem 3.1 [1].** *Let  $G$  be a compact affine Nash group and let  $X$  be a compact  $C^\infty G$  manifold so that  $X$  is  $G$  cobordant to a nonsingular algebraic  $G$  set. Then  $X$  is  $C^\infty G$  diffeomorphic to a nonsingular algebraic  $G$  set. Here an algebraic  $G$  set means a  $G$  invariant algebraic subset of some representation of  $G$ .  $\square$*

*Proof of Theorem 1.* The disjoint union  $X \amalg X$  is null cobordant. By Theorem 3.1,  $X \amalg X$  is  $C^\infty G$  diffeomorphic to a nonsingular algebraic  $G$  set in some representation  $\Omega$  of  $G$ . Since a  $G$  invariant collection of connected components of a nonsingular algebraic  $G$  set is an affine Nash  $G$  submanifold in  $\Omega$ ,  $X$  admits an affine Nash  $G$  manifold structure  $Y \subset \Omega$ . Let  $Z$  be another affine Nash  $G$  manifold structure of  $X$  in  $\Omega'$ . We have to prove  $Y$  is Nash  $G$  diffeomorphic to  $Z$ . Let  $f$  be a  $C^\infty G$  diffeomorphism from  $Y$  to  $Z$ . Let  $F$  denote the composition of  $f$  with the inclusion  $Z \rightarrow \Omega'$ . By [1]  $F$  can be approximated by a polynomial  $G$  map  $q : Y \rightarrow \Omega'$ . By Proposition 2.3, we have a Nash  $G$  tubular neighborhood  $(U, p)$  of  $Z$  in  $\Omega'$ . Since  $Y$  is compact, if the approximation is close then the image of  $q$  lies in  $U$ . Thus  $k := p \circ q$  is an approximation of  $f$ . If the approximation is close then a Nash  $G$  map  $k : Y \rightarrow Z$  is a Nash  $G$  diffeomorphism. Therefore (1) is proved.

Next we prove (2). Let  $X_1, X_2$  be two Nash  $G$  manifold structures (may not be affine) of  $X$  and let  $k$  be a  $C^\infty G$  diffeomorphism from  $X_1$  to  $X_2$ . Fix  $x_1 \in X_1$ , and let  $x_2 = k(x_1)$ . Then the map  $f_i : G \rightarrow X_i : f_i(g) = gx_i$  ( $i = 1, 2$ ) is a surjective Nash  $G$  map because  $G$  acts on  $X_i$  ( $i = 1, 2$ ) transitively, and  $f_2 = k \circ f_1$ .

To prove  $k$  is a Nash map, it is enough to show  $k$  is a  $C^0$  Nash map. By [4] we can find a  $C^0$  Nash imbedding  $I_i$  from  $X_i$  to some Euclidean space  $\mathbb{R}^s$  ( $i = 1, 2$ ). Let  $X'_i = I_i(X_i)$  ( $i = 1, 2$ ),  $f'_i = I_i \circ f_i$  ( $i = 1, 2$ ) and  $k' = I_2 \circ k \circ I_1^{-1}$ . Then  $f'_i : G \rightarrow X'_i$  ( $i = 1, 2$ ) is a  $C^0$  Nash map. Since  $G$  and  $X_i$  ( $i = 1, 2$ ) are affine, there exists a finite semialgebraic open covering  $\{O_t\}_t$  of  $G$  such that each  $f'_i|_{O_t}$  is semialgebraic. Therefore  $f'_i$  ( $i = 1, 2$ ) is semialgebraic. Since  $k'$  is  $C^0$  Nash if and only if  $k$  is  $C^0$  Nash, we have only to show that  $k'$  is  $C^0$  Nash.

Since  $f'_i$  ( $i = 1, 2$ ) is a  $C^0$  Nash map, there exist finite systems of coordinate neighborhoods  $\{\phi_i : W_i \rightarrow \mathbb{R}^m\}$  of  $G$ ,  $\{\psi_j : U_j \rightarrow \mathbb{R}^n\}$  of  $X_1$ , and  $\{\varphi_l : V_l \rightarrow \mathbb{R}^n\}$  of  $X_2$  such that, for any  $i, j$  and  $l$ ,  $\phi_i((f'_1)^{-1}(U_j) \cap W_i)$ ,  $\phi_i((f'_2)^{-1}(V_l) \cap W_i)$  are semialgebraic, and that  $\psi_j \circ f'_1 \circ \phi_i^{-1} : \phi_i((f'_1)^{-1}(U_j) \cap W_i) \rightarrow \mathbb{R}^n$ ,  $\varphi_l \circ f'_2 \circ \phi_i^{-1} : \phi_i((f'_2)^{-1}(V_l) \cap W_i) \rightarrow \mathbb{R}^n$  are  $C^0$  Nash maps, where  $m$  (resp.  $n$ ) denotes the dimension of  $G$  (resp.  $X_1$ ). We have only to show that each  $\varphi_l \circ k' \circ \psi_j^{-1}$  is semialgebraic. For a map  $h$ , let  $\text{graph}(h)$  denote the graph of  $h$ . For  $j$  and  $l$ , let

$$K = \bigcup_i \text{graph}(\psi_j \circ f'_1 \circ \phi_i^{-1}) \times \text{graph}(\varphi_l \circ f'_2 \circ \phi_i^{-1}).$$

Then  $K$  is semialgebraic in  $(\mathbb{R}^m \times \mathbb{R}^n) \times (\mathbb{R}^m \times \mathbb{R}^n)$ , hence the image  $K'$  of  $K$  by the projection  $(\mathbb{R}^m \times \mathbb{R}^n) \times (\mathbb{R}^m \times \mathbb{R}^n) \rightarrow \mathbb{R}^n \times \mathbb{R}^n$  is semialgebraic in  $\mathbb{R}^n \times \mathbb{R}^n$ . Since  $f'_i$  ( $i = 1, 2$ ) is surjective and  $f'_2 = k' \circ f'_1$ ,  $\text{graph}(\varphi_l \circ k' \circ \psi_j^{-1}) = K'$ . Thus each  $\varphi_l \circ k' \circ \psi_j^{-1}$  is semialgebraic. Hence  $k'$  is a  $C^0$  Nash. Therefore  $k$  is a Nash  $G$  diffeomorphism.

Now we prove (3). By (1) we can assume that  $X$  is an affine Nash  $G$  submanifold of a representation  $\Omega$  of  $G$ . For any  $x \in X$ , the orbit  $G(x)$  of  $x$  is a  $C^\infty G$  submanifold of  $\Omega$  because  $G$  is compact. Moreover  $G(x)$  is a semialgebraic set. Hence  $G(x)$

is an affine Nash  $G$  submanifold in  $\Omega$ . Since the action on  $G$  is not transitive and by Proposition 2.3, there exists some Nash  $G$  tubular neighborhood  $(U', p)$  of some orbit  $G(x)$  in  $\Omega$  with  $X \neq U := U' \cap X$ .

For  $0 < c < 1$ , set

$$\begin{aligned} a &= 2^{2.5}(1+c)^2/(1-c)^2, \\ d &= 2 + 2^{0.5}3a + a^2 - (a + \sqrt{2})\sqrt{a^2 + 2^{2.5}a}. \end{aligned}$$

Then  $a > 2^{2.5}$ ,  $1 < d < 2$ . Suppose  $k$  is a Nash function satisfying

$$\sqrt{2}(x + k(x)) = (x - k(x))^2/a.$$

The graph of  $k$  comes to a rotation of the graph of  $y = x^2/a$  with center at the origin. It follows from this and  $a > 2^{2.5}$  that  $k$  and its Nash extension  $k'$  to

$$[1 - 2^{-0.25}\sqrt{a}, 1 + 2^{-0.25}\sqrt{a}] (\supset (-1, 3))$$

is well-defined, and that  $k'$  satisfies

$$k'[1 - 2^{-0.25}\sqrt{a}, 1 + 2^{-0.25}\sqrt{a}] = [1 - 2^{-0.25}\sqrt{a}, 1 + 2^{-0.25}\sqrt{a}],$$

the derivative of  $k'$  is negative,  $k' \circ k' = id$ .

Let

$$N_1 = (-\infty, d), N_2 = (0, \infty), N_3 = (0, 1).$$

Define the Nash maps  $h_1 : N_3 \rightarrow N_1, h_2 : N_3 \rightarrow N_2$  by

$$h_1(t) = t^2 + k(t)^2 \text{ and } h_2(t) = 2t - t^2.$$

Then  $h_1$  and  $h_2$  are Nash imbeddings so that  $h_1(N_3) = (0, d), h_2(N_3) = (0, 1)$ . We can extend  $h_1$  to

$$h'_1 : [1 - 2^{-0.25}\sqrt{a}, 1 + 2^{-0.25}\sqrt{a}] \rightarrow \mathbb{R}$$

as a Nash function such that the derivative vanishes at only 0 and that  $h'_1 = h'_1 \circ k'$  because the derivative of  $k'$  is negative and  $k' \circ k' = id$ .

Applying Proposition 2.3 to the boundary  $\partial\bar{U}$  of the closure  $\bar{U}$  of  $U$  in  $X$ , there exists a Nash  $G$  collar  $\phi : \partial\bar{U} \times [0, 1] \rightarrow \bar{U}$ . Let  $D(\varepsilon)$  ( $0 < \varepsilon < 1$ ) denote  $\phi(\partial\bar{U} \times (0, \varepsilon))$ . Take a Nash diffeomorphism  $f : \mathbb{R} \rightarrow (0, 1)$  (e.g. the inverse map of the composition of  $f : (0, 1) \rightarrow (-1, 1) : f(x) = 2x - 1$  with  $h : (-1, 1) \rightarrow \mathbb{R} : h(x) = x/(1 - x^2)$ ). Set

$$U_1 = D(f(d)), U_2 = X - \overline{D(f(0))}, U_3 = D(f(1)) - \overline{D(f(0))}.$$

Then each  $U_i$  is an open affine Nash  $G$  submanifold of  $X$ . Let

$$H_1 = \phi \circ (id \times (f \circ h_1 \circ f^{-1})) \circ \phi^{-1} : U_3 \rightarrow U_1,$$

$$H_2 = \phi \circ (id \times (f \circ h_2 \circ f^{-1})) \circ \phi^{-1} : U_3 \rightarrow U_2.$$

We define  $X_c$  by the quotient topological space of the disjoint union  $\coprod_{i=1}^3 U_i$ , and the equivalence relation  $x \sim H_1(x) \sim H_2(x)$  for  $x \in U_3$  on the union. Then one can check that  $X_c$  is a Nash  $G$  manifold which is  $C^\infty G$  diffeomorphic to  $X$ . Next we prove  $X_c$  is nonaffine. To prove this, we use the following lemma.

**Lemma 3.2 (c.f. Remark 1.2.2.15 [7]).** *Let  $f$  be a locally semialgebraic  $C^\infty$  map from a Nash manifold  $M$  to a Nash manifold  $N$ . If  $N$  is affine then  $f$  is a Nash map.  $\square$*

Fix  $0 < c < 1$  and  $z \in S(f(1))$ , where  $S(f(1))$  denotes  $\phi(\partial\bar{U} \times \{f(1)\})$ . Let  $\psi_c : (f(0), f(1)) \rightarrow X_c$  be the composition

$$(f(0), f(1)) \rightarrow S(f(1)) \times (f(0), f(1)) \rightarrow U_3 \rightarrow X_c,$$

where the first map is  $x \rightarrow (z, x)$ , the second is the natural Nash  $G$  diffeomorphism from  $S(f(1)) \times (f(0), f(1))$  to  $U_3$ , and the third is the natural imbedding from  $U_3$  into  $X_c$ . Then  $\psi_c$  is an imbedding. We extend  $\psi_c$  as follows. Let  $l_{ci}$  ( $i = 1, 2, 3$ ) be the natural imbedding  $U_i \rightarrow X_c$  and let  $V_{ci}$  ( $i = 1, 2, 3$ ) denote its image. Then

$$p \circ k_1^{-1} \circ l_1^{-1} \circ \psi_c = f \circ h_1 \circ f^{-1}, p \circ k_2^{-1} \circ l_2^{-1} \circ \psi_c = f \circ h_2 \circ f^{-1} \text{ on } (f(0), f(1)),$$

where  $p$  denotes the projection  $\partial\bar{U}_3 \times (f(0), f(d)) \rightarrow (f(0), f(d))$  and  $k_i$  ( $i = 1, 2$ ) stands for the natural imbedding  $\partial\bar{U}_3 \times (f(0), f(d)) \rightarrow U_i$ . We extend  $\psi_c$  to  $(f(0), f(1 + \varepsilon))$  for small positive  $\varepsilon$ . It suffices to consider  $p \circ k_2^{-1} \circ l_2^{-1} \circ \psi_c$  because the image of  $\psi_c$  lies in  $V_{c2}$  and  $\lim_{t \rightarrow f(1)} \psi_c(t) \in V_{c2}$ . Now  $p \circ k_2^{-1} \circ l_2^{-1} \circ \psi_c = f(2f^{-1}(t) - (f^{-1}(t))^2)$  on  $(f(0), f(1))$ . Thus  $p \circ k_2^{-1} \circ l_2^{-1} \circ \psi_c$  and  $\psi_c$  are extensible to  $(f(0), f(2))$  and

$$p \circ k_2^{-1} \circ l_2^{-1} \circ \psi_c(t) = f(2f^{-1}(t) - (f^{-1}(t))^2) \text{ on } [f(1), f(2)).$$

Clearly we can extend  $\psi_c$  to  $[f(0), f(1)]$ , and  $\psi_c((f(0), f(2)) \subset \psi_c([f(0), f(1)])$ . Hence

$$\psi_{c0}^{-1} \circ \psi_c(t) = f(2 - f^{-1}(t)) \text{ on } [f(1), f(2)),$$

$f(1)$  is the only and nondegenerate critical point, where  $\psi_{c0}$  denotes the homeomorphism  $\psi_c : [f(0), f(1)] \rightarrow \psi_c([f(0), f(1)])$ . In the same way,  $\psi_c$  can be defined on  $(f(k'(1)), f(0)]$  satisfying

$$\psi_{c0}^{-1} \circ \psi_c(t) = f(k'(f^{-1}(t))) \text{ for } t \in (f(k'(1)), f(0)],$$

and the critical point is only  $f(0)$  and nondegenerated. Repeating this argument,  $\psi_c$  is extensible on

$$(f(1 - 2^{-0.25}\sqrt{a}), f(1 + 2^{-0.25}\sqrt{a})),$$

and  $\psi_c$  is locally semialgebraic, the image of  $\psi_c$  is  $\psi_c([f(0), f(1)])$ , and that for any  $e \in (f(0), f(1))$ ,  $(\psi_{c0}^{-1} \circ \psi_c)^{-1}(e)$  is discrete and consists of infinitely many elements. The set of critical points of  $\psi_c$  is  $(\psi_{c0}^{-1} \circ \psi_c)^{-1}(f(0)) \cup (\psi_{c0}^{-1} \circ \psi_c)^{-1}(f(1))$ , and they are nondegenerate ones. Since  $\psi_c$  is locally semialgebraic and not semialgebraic and by Lemma 3.2,  $X_c$  is not affine.

Finally we prove that  $X_c$  is not Nash  $G$  diffeomorphic to  $X_{c'}$  if  $0 < c, c' < 1$ ,  $\alpha = \log f(c')/\log f(c)$  is irrational. Assume that there exists a Nash  $G$  diffeomorphism  $u : X_c \rightarrow X_{c'}$ . Then we have to prove  $\log f(c')/\log f(c)$  is rational. Set

$$a = 2^{2.5}(1+c)^2/(1-c)^2, a' = 2^{2.5}(1+c')^2/(1-c')^2,$$

$$\psi_c : (f(1 - 2^{-0.25}\sqrt{a}), f(1 + 2^{-0.25}\sqrt{a})) \rightarrow X_c,$$

$$\psi_{c'} : (f(1 - 2^{-0.25}\sqrt{a'}), f(1 + 2^{-0.25}\sqrt{a'})) \rightarrow X_{c'}.$$

We also write

$$\psi_{c0} = \psi_c|[f(0), f(1)] : [f(0), f(1)] \longrightarrow \psi_c([f(0), f(1)]),$$

$$\psi_{c'0} = \psi_{c'}|[f(0), f(1)] : [f(0), f(1)] \longrightarrow \psi_{c'}([f(0), f(1)]),$$

Let  $L_1$  be the composition of the diffeomorphism  $S(f(1)) \times (f(-10d), f(10d)) \longrightarrow D(f(10d)) - \overline{D(f(-10d))}$  with the projection  $D(f(10d)) \longrightarrow X_{c'}$ , and let  $L_2$  be the projection  $S(f(1)) \times (f(-10d), f(-10d)) \longrightarrow S(f(1))$ . By Lemma 3.2 and the infinite vibration of  $\psi_c$ ,  $L_2 \circ L_1^{-1} \circ u \circ \psi_c$  is constant. Let  $z'$  denote this constant. Clearly the images of  $\phi(z' \times f(N_2))$  and  $\phi(z' \times (f(1), f(d)))$  via  $\pi_{c'}$  in  $X_{c'}$  are affine Nash  $G$  submanifolds. Let  $k_c$  be the natural homeomorphism from  $\overline{U_3}$  into  $X_c$ . Thus  $u \circ k_c \circ \phi(z \times [f(0), f(1)])$  is not contained in these affine Nash  $G$  submanifolds because the image of  $\psi_c$  is not affine. This implies that

$$u \circ k_c \circ \phi(z \times [f(0), f(1)]) \subset k_{c'} \circ \phi(z' \times [f(0), f(1)]).$$

Applying the same argument to  $u^{-1}$ , we have

$$u^{-1} \circ k_{c'} \circ \phi(z' \times [f(0), f(1)]) \subset k_c \circ \phi(z \times [f(0), f(1)]).$$

Therefore

$$u \circ k_c \circ \phi(z \times [f(0), f(1)]) = k_{c'} \circ \phi(z' \times [f(0), f(1)]).$$

For any  $e \in (f(0), f(1))$ , let  $(\psi_{c0}^{-1} \circ \psi_c)^{-1}(e) = \{e_i\}_{i \in \mathbb{Z}}$ ,  $(\psi_{c'0}^{-1} \circ \psi_{c'})^{-1}(e) = \{e'_i\}_{i \in \mathbb{Z}}$ . Then

$$[3.1] \quad \lim_{i \rightarrow \infty} (f(1 + 2^{-0.25} \sqrt{a}) - e_{-i-2}) / (f(1 + 2^{-0.25} \sqrt{a}) - e_{-i}) = f(c),$$

$$[3.2] \quad \lim_{i \rightarrow \infty} (f(1 + 2^{-0.25} \sqrt{a'}) - e'_{-i-2}) / (f(1 + 2^{-0.25} \sqrt{a'}) - e'_{-i}) = f(c'),$$

are obtained as follows. The map  $t \longrightarrow f(k'(2 - f^{-1}(t)))$  has fixed points only at the end of the interval, it repels from  $f(1 + 2^{-0.25} \sqrt{a})$ , attracts to  $f(1 - 2^{-0.25} \sqrt{a})$  and its derivatives at the latter point is  $f(c)$ . Thus  $(f(1 + 2^{-0.25} \sqrt{a}) - e_{-i-2}) / (f(1 + 2^{-0.25} \sqrt{a}) - e_{-i})$  converges  $f(c)$  because  $e_{-i-2} = f(k'(2 - f^{-1}(e_{-i})))$ . Hence we have [3.1]. A similar argument shows [3.2]. Since

$$u \circ k_c \circ \phi(z \times [f(0), f(1)]) = k_{c'} \circ \phi(z' \times [f(0), f(1)]),$$

for a pair

$$\begin{aligned} e_0 &\in (f(1 - 2^{-0.25} \sqrt{a}), f(1 + 2^{-0.25} \sqrt{a})) \text{ and} \\ e'_0 &\in (f(1 - 2^{-0.25} \sqrt{a'}), f(1 + 2^{-0.25} \sqrt{a'})) \text{ with} \\ &\psi_{c'}(e'_0) = u \circ \psi_c(e_0) \end{aligned}$$

there exists a homeomorphism

$$\tau : (f(1 - 2^{-0.25} \sqrt{a}), f(1 + 2^{-0.25} \sqrt{a})) \longrightarrow (f(1 - 2^{-0.25} \sqrt{a'}), f(1 + 2^{-0.25} \sqrt{a'}))$$



so that  $\tau(e_0) = e'_0$  and  $\psi_{c'} \circ \tau = u \circ \psi_c$  on  $(f(1 - 2^{-0.25}\sqrt{a}), f(1 + 2^{-0.25}\sqrt{a}))$ . Remember that all critical points of  $\psi_c, \psi_{c'}$  are nondegenerate. This shows that  $\tau$  is of class  $C^\omega$ . Therefore, by Lemma 3.2,  $\tau$  is a Nash diffeomorphism. Set

$$\begin{aligned}\psi_{c_0}^{-1} \circ \psi_c(e_0) &= e, \psi_{c'_0}^{-1} \circ \psi_{c'}(e_0) = e', \\ (\psi_{c_0}^{-1} \circ \psi_c)^{-1}(e) &= \{e_i\}_{i \in \mathbb{Z}}, \\ (\psi_{c'_0}^{-1} \circ \psi_{c'})^{-1}(e') &= \{e'_i\}_{i \in \mathbb{Z}}.\end{aligned}$$

Then  $\tau$  satisfies

$$\begin{aligned}\tau(e_i) &= e'_i \text{ for any } i \in \mathbb{Z} \text{ or } , \\ \tau(e_i) &= e'_{-i} \text{ for any } i \in \mathbb{Z}.\end{aligned}$$

A map  $f \circ (\text{translation}) \circ f^{-1}$  takes  $(f(1 - 2^{-0.25}\sqrt{a}), f(1 + 2^{-0.25}\sqrt{a}))$  to  $(f(0), f(2^{0.75}\sqrt{a}))$ , and a similar map  $f \circ (\text{translation}) \circ f^{-1}$  takes  $(f(1 - 2^{-0.25}\sqrt{a'}), f(1 + 2^{-0.25}\sqrt{a'}))$  to  $(f(0), f(2^{0.75}\sqrt{a'}))$ , we may suppose that  $e_i$  and  $e'_i$  are converge to 0 as  $i \rightarrow \infty$ . Assume that  $e$  and  $e'$  lie in  $(f(0), f(1))$ . Then it follows from [3.1] and [3.2] that

$$[3.3] \quad \lim_{i \rightarrow \infty} e_{-i-2}/e_{-i} = f(c),$$

$$[3.4] \quad \lim_{i \rightarrow \infty} e'_{-i-2}/e'_{-i} = f(c').$$

Let  $Z$  denote the Zariski closure of  $\text{graph}(\tau)$ . This is of dimension 1 because  $\tau$  is semialgebraic. It is clear that  $Z$  contains all  $(e_i, e'_i)$ . Let  $P(x, y) = \sum_{j=1}^s \delta_j x^{\beta_j} y^{\gamma_j}$  ( $\delta_j \in \mathbb{R}, \beta_j, \gamma_j \in \mathbb{N}$ ) be a defining polynomial of  $Z$ . Then

$$P(e_i, e'_i) = 0 \text{ for any } i \in \mathbb{Z}.$$

Since  $\alpha$  is irrational,

$$[3.5] \quad \beta_i + \alpha\gamma_i \neq \beta_j + \alpha\gamma_j \text{ for } i \neq j.$$

Set

$$P_i(x, y) = x^{\beta_i} y^{\gamma_i}.$$

For each  $n \in \mathbb{Z}$ , let  $E(n)$  denote the  $s \times s$ -matrix whose  $(i, j)$  entry is

$$P_i(e_{-n-2j+1}, e'_{-n-2j+1}).$$

Then

$$(\delta_1, \dots, \delta_s)E(n) = (P(e_{-n-1}, e'_{-n-1}), \dots, P(e_{-n-2s+1}, e'_{-n-2s+1})) = 0.$$

In particular  $\det E(n) = 0$ . On the other hand, we have

$$\det E(n) = \left( \prod_{i=1}^s P_i(e_{-n-1}, e'_{-n-1}) \right) \det F(n),$$

where  $F(n)$  is the  $s \times s$ -matrix whose  $(i, j)$  entry is

$$P_i(e_{-n-2j+1}, e'_{-n-2j+1})/P_i(e_{-n-1}, e'_{-n-1}).$$

Now [3.3] and [3.4] mean that each entry of  $F(n)$  converges to

$$(c^{\beta_i}(c')^{\gamma_i})^{j-1} = c^{(\beta_i+\alpha\gamma_i)(j-1)}$$

as  $n \rightarrow \infty$ . Thus  $\det F(n)$  converges to a Vandermonde's determinant equals

$$\prod_{i < j} (c^{\beta_j + \alpha\gamma_j} - c^{\beta_i + \alpha\gamma_i}) \neq 0,$$

by [3.5]. Therefore  $\det E(n) \neq 0$  for large  $n$ . This proves the result.  $\square$

#### 4. Compactifiable $C^\infty G$ manifolds.

The same argument of the proof of Theorem 1 (3) proves Theorem 2 (2). To prove Theorem 2 (1), we show a relative version of Theorem 3.1. After proving Theorem 4.2, we give a proof of Theorem 2 (1).

*Definition 4.1.* (1) An algebraic subset of a representation of  $G$  is said to be an *algebraic  $G$  set* if it is  $G$  invariant. Moreover we call it a *nonsingular algebraic  $G$  set* if it is nonsingular.

(2) Let  $X$  be a  $C^\infty G$  manifold and let  $X'$  be a  $C^\infty G$  submanifold of  $X$ . A pair  $(X, X')$  is called *algebraically  $G$  cobordant* if there exist a nonsingular algebraic  $G$  set  $Y$ , a nonsingular algebraic  $G$  subset  $Y'$  of  $Y$ , a  $G$  cobordism  $N$  between  $X$  and  $Y$ , and a  $G$  cobordism  $N'$  between  $X'$  and  $Y'$  such that  $N'$  is a  $C^\infty G$  submanifold of  $N$ .

**Theorem 4.2.** *Let  $G$  be a compact affine Nash group,  $X$  a compact  $C^\infty G$  manifold, and  $X'$  a compact  $C^\infty G$  submanifold of  $X$ . If the pair  $(X, X')$  is algebraically  $G$  cobordant then there exist a nonsingular algebraic  $G$  set  $Z$  in  $X \times \Omega$  for some representation  $\Omega$  of  $G$ , a nonsingular algebraic subset  $Z'$  of  $Z$ , and a  $C^\infty G$  diffeomorphism  $\phi : X \rightarrow Z$  with  $\phi(X') = Z'$ .*

For any  $C^\infty G$  manifold  $X$  and  $C^\infty G$  submanifold  $X'$  of  $X$ , the pair  $(X \amalg X, X' \amalg X')$  is algebraically  $G$  cobordant. Therefore we have the next corollary because a  $G$  invariant collection of connected components of a nonsingular algebraic  $G$  set is an affine Nash  $G$  submanifold in some representation of  $G$ .

**Corollary 4.3.** *Let  $G$  be a compact affine Nash group,  $X$  a compact  $C^\infty G$  manifold, and  $X'$  a compact  $C^\infty G$  submanifold of  $X$ . Then there exist an affine Nash  $G$  manifold  $Y$ , an affine Nash  $G$  submanifold  $Y'$  of  $Y$ , and a  $C^\infty G$  diffeomorphism  $\phi : X \rightarrow Y$  so that  $\phi(X') = Y'$ .  $\square$*

*Proof of Theorem 4.2.* By the proof of Theorem 1.3 [1],  $X'$  is  $G$  isotopic to a nonsingular algebraic  $G$  subset  $Z'$  of  $X \times \Omega$  by an arbitrarily small isotopy, for some representation of  $G$ . Extending this isotopy, we may assume that it maps  $X \times 0$  to some  $C^\infty G$  manifold  $M$  in  $X \times \Omega$  so that  $M - X \times 0$  has compact closure and that the composition of the inclusion  $M \rightarrow X \times \Omega$  with the projection

$X \times \Omega \longrightarrow X$  is a  $C^\infty G$  diffeomorphism. In particular  $Z' \subset M$ . Since  $Z'$  is compact and by Lemma 4.7 [1], one can find a proper  $G$  invariant polynomial  $\rho$  such that  $\rho^{-1}(0) = Z'$ . Let  $\alpha : X \longrightarrow \Omega$  be a  $C^\infty G$  map with compact support so that

$$M = \{(x, y) \in X \times \Omega \mid y = \alpha(x)\}.$$

Take a  $G$  invariant  $C^\infty$  function  $\beta : X \times \Omega \longrightarrow [0, 1]$  with compact support with  $\beta(x, y) = 1$  when  $|y| < 2|\alpha(x)|$ . Let  $\gamma : X \times \Omega \longrightarrow \Omega$  be

$$\gamma(x, y) = \beta(x, y)(y - \alpha(x)) + (1 - \beta(x, y))\rho^2(x, y)y.$$

Then 0 is a regular value of  $\gamma$ ,  $\gamma^{-1}(0) = M$ , and  $\gamma$  is equal to the polynomial  $\rho^2(x, y)y$  outside of a  $G$  invariant compact set. By Lemma 5.1 [1], one can  $C^1$  approximate  $\gamma(x, y) - \rho^2(x, y)y$  by an equivariant entire rational map  $u : (X \times \Omega, Z') \longrightarrow (\Omega, 0)$ . Here an entire rational map means a fraction of polynomial maps with nowhere vanishing denominator. This approximation is close on all  $X \times \Omega$ . Thus

$$w(x, y) = u(x, y) + \rho^2(x, y)y$$

is  $C^1$  approximation of  $\gamma$  on  $X \times \Omega$ . Since  $\rho$  is proper and by equivariant Morse theory, there exists a  $C^\infty G$  diffeomorphism from  $Z := w^{-1}(0)$  to  $M = \gamma^{-1}(0)$  fixing  $Z'$ .  $\square$

*Proof of Theorem 2 (1).* Since  $X$  is compactifiable, there exists a  $C^\infty G$  manifold  $X'$  with boundary  $\partial X$  so that  $X$  is  $C^\infty G$  diffeomorphic to the interior of  $X'$ . Let  $Y$  be the double of  $X'$ . Applying Corollary 4.3 to the pair  $(Y, \partial X')$ , one can find a representation  $\Omega$  of  $G$  and a  $C^\infty G$  imbedding  $F : Y \longrightarrow \Omega$  such that  $F(Y)$  and  $F(\partial X')$  are affine Nash  $G$  manifolds. Hence  $F(X)$  is an affine Nash  $G$  manifold. Therefore  $X$  admits an affine Nash  $G$  manifold structure.  $\square$

On the other hand, T. Petrie [3] proved that any nonsingular algebraic  $G$  set is compactifiable as a  $C^\infty G$  manifold when  $G$  is an algebraic group. A similar proof shows the next theorem, because the number of connected components of the zeros of a Nash map is finite.

**Theorem 4.4.** *Let  $G$  be a compact affine Nash group. Then every affine Nash  $G$  manifold is compactifiable as a  $C^\infty G$  manifold.  $\square$*

M. Shiota studied compactifications of Nash manifolds as either  $C^\infty$  manifolds [4] or Nash manifolds [5].

By Theorem 2 (1) and Theorem 4.4, we have the following.

**Theorem 4.5.** *Let  $G$  be a compact affine Nash group. Then a  $C^\infty G$  manifold is compactifiable if and only if it admits an affine Nash  $G$  manifold structure.  $\square$*

## REFERENCES

1. K.H. Dovermann, M. Masuda and T. Petrie, *Fixed point free algebraic actions on varieties diffeomorphic to  $\mathbf{R}^n$* , Progress in Math. 80, Birkhäuser (1990), 49-80.
2. J.J. Madden and C.M. Stanton, *One-dimensional Nash group*, Pacific J. Math. **154**, (1992), 331-344.

3. T. Petrie, *Smooth conjugacy of algebraic actions on affine varieties*, *Topology* **27** (1988), 473-477.
4. M. Shiota, *Abstract Nash manifolds*, *Proceedings of Amer. Math. Soc.* **96** (1986), 155-162.
5. M. Shiota, *a book Chap 2. Nash functions and manifolds*.
6. M. Shiota, *Classification of Nash manifolds*, *Ann. Inst. Fourier, Grenoble* **33** (1983), 209-232.
7. M. Shiota, *Nash manifolds*, *Lecture Note in Mathematics* vol 1269, Springer-Verlag, 1987.