

Title	The Thermal Sensitivity of Sarcoplasmic Reticulum Ca <sup>2+</sup> -ATPase
Author(s)	永田, 善明
Citation	大阪大学, 1999, 博士論文
Version Type	VoR
URL	<a href="https://doi.org/10.11501/3155163">https://doi.org/10.11501/3155163</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

*The Thermal Sensitivity  
of Sarcoplasmic Reticulum Ca<sup>2+</sup>-ATPase*

*Yoshiaki Nagata*

*Departement of Biology  
Graduate School of Science  
Osaka University  
February 1999*

The thermal sensitivity of sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase

Yoshiaki Nagata  
Department of Biology  
Faculty of Science  
Osaka University  
1999

## CONTENTS

ABBREVIATIONS	2
GENERAL INTRODUCTION	3
PART I	12
PART II	33
PART III	50
PART IV	75
BIBLIOGRAPHY	95
ACKNOWLEDGEMENTS	96

## ABBREVIATIONS

AMP-PNP	adenyl-5'-imidodiphosphate
C <sub>12</sub> E <sub>9</sub>	polyoxyethylene-9-laurylether
DACM	N-(7-Dimethylamino-4-methyl-3-coumarinyl)-maleimide
DMSO	dimethyl sulfoxide
DOC	deoxycholate
DTNB	5,5'-dithiobis-(2-nitrobenzoic acid)
EDTA	Ethylenediaminetetraacetic acid
EGTA	ethylene glycol bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid
EP	phosphoenzyme
FITC	fluoresceine isothiocyanate
GSH	glutathione
HPLC	high-performance liquid chromatography
IAA	iodoacetamide
IAEDANS	5-(2-iodoacetamidoethyl)aminonaphthalene-1-sulfonate
NCD-4	N-cyclohexyl-N'-(4-dimethylamino-1-naphthyl)carbodiimide
PAGE	polyacrylamide gel electrophoresis
PEP	phosphoenolpyrvate
PMI	N-(3-pyrene)maleimide
SDS	sodium dodecyl sulfate
SR	sarcoplasmic reticulum
TCA	trichloroacetic acid
TES	N-tris(hydroxymethyl)methyl-2-aminoethane sulfonate
TFA	trifluoroacetic acid

## GENERAL INTRODUCTION


### **Physiological functions of sarcoplasmic reticulum**

In skeletal muscle, intracellular  $\text{Ca}^{2+}$  ions play a critical role in the regulation of contraction and relaxation of muscle; When muscle cell is relaxed, concentration of  $\text{Ca}^{2+}$  is kept below  $0.1 \mu\text{M}$ . When muscle cell is excited by electric current from a nerve end, the depolarization of plasma membrane is transferred to interior of the cell *via* transverse tubules, which are vertically running against the muscle fiber, then the signal stimulates the  $\text{Ca}^{2+}$ -burst from the sarcoplasmic reticulum (SR) vesicles which are surrounding myofibrils in the cell. Upon the  $\text{Ca}^{2+}$ -release from SR, intracellular  $\text{Ca}^{2+}$  level increases above several  $\mu\text{M}$  which suffices to trigger the muscle contraction. The entire process from excitation to contraction of muscle is called "E-C coupling". After the excitation, cytoplasmic  $\text{Ca}^{2+}$  ions are removed by SR to a level lower than  $0.1 \mu\text{M}$ , which results in the muscle relaxation ( see Ref. 1 for a review ).

SR vesicles can be easily isolated with retention of the full functional activities by differential centrifugation of a muscle homogenate. The isolated SR vesicles have been proven suitable for the study of the molecular mechanisms of active cation transport since this system possesses several features distinct from other transport membranes as follow; (i) A high content of the  $\text{Ca}^{2+}$ -pump ATPase, (ii) a clear distinction between the inside and outside of the membrane, (iii) tightly coupled ATP hydrolysis to  $\text{Ca}^{2+}$ -transport, and (iv) readily measured reaction intermediates, EP, which facilitates the kinetic analysis and characterization of the elementary steps of the reaction ( see Ref. 2 for review ).

### **Reaction scheme of $\text{Ca}^{2+}$ -ATPase**

Figure 1 shows the coupling mechanism of ATP hydrolysis and  $\text{Ca}^{2+}$ -transport by SR. This scheme has been proposed on the basis of a number of transient and steady state kinetic analysis ( see Ref. 3 for review ). In the first step of the  $\text{Ca}^{2+}$ -pump reaction,  $\text{Ca}^{2+}$ -ATPase forms  $\text{E}_1\text{Ca}$  complex by the binding of 2 mol  $\text{Ca}^{2+}$  to 1 mol of enzyme. In the intact SR membrane  $\text{Ca}^{2+}$ -ATPase exists in an equilibrium between two different configurations of  $\text{E}_1$  and  $\text{E}_2$ . In the absence of  $\text{Ca}^{2+}$ , the equilibrium shifts to  $\text{E}_2$ . When  $\text{Ca}^{2+}$  is added to the cytoplasmic side, the enzyme in  $\text{E}_1$  form, but not in  $\text{E}_2$  form, can sequentially bind 2 mol of Ca to form  $\text{E}_1\text{Ca}$  complex. In the second step,  $\text{E}_1\text{Ca}$  is reacted with ATP at the outer surface of the SR membrane to form phosphoprotein intermediate



**Fig. 1 Coupling mechanism of ATP hydrolysis and Ca transport in SR.** The scheme is based on the assumption of two major conformational states, E1 and E2, which are characterized by the ability to react with ATP and Pi, respectively. In the formation of E1P from E1Ca and ATP, 2 mol Ca<sup>2+</sup> ions are occluded in the enzyme. After conversion of E1P to E2P, 2 mol Ca<sup>2+</sup> ions are deoccluded. This is followed by dephosphorylation of E2P to E2 and Pi. For further details see text.



(E<sub>1</sub>P) by transferring a  $\gamma$ -phosphate of ATP to the specific Asp residue of the catalytic site. E<sub>1</sub>P is highly sensitive to ADP and it can form ATP by the reversal reaction with ADP. In this step, Ca<sup>2+</sup> is occluded in the enzyme and inaccessible to both of cytoplasmic and luminal sides of the SR membrane. The third step is a conversion of the Ca<sup>2+</sup>-ATPase from E<sub>1</sub>P to E<sub>2</sub>P in which Ca<sup>2+</sup> ions move into luminal side. The apparent affinity of E<sub>2</sub>P for Ca<sup>2+</sup> becomes 3 to 4 order of magnitude less than that of E<sub>1</sub>P and 2 mol of Ca<sup>2+</sup> are randomly released from the enzyme into the SR lumen. In the final step, E<sub>2</sub>P is hydrolyzed by H<sub>2</sub>O into E<sub>2</sub> and inorganic phosphate, Pi at the outer surface of SR membrane and the reaction cycle is accomplished. This catalytic cycle of the Ca<sup>2+</sup>-ATPase is completely reversible. When EGTA is added to remove outer medium Ca<sup>2+</sup> and, Pi and ADP are simultaneously added to the medium which contains SR vesicles that have been loaded with Ca<sup>2+</sup>, 2 mol of Ca<sup>2+</sup> are released from the vesicles which is accompanied by the synthesis of 1 mol of ATP *via* the steps of E<sub>2</sub>P formation which is followed by the conversion from E<sub>2</sub>P to E<sub>1</sub>P. This reverse reaction indicates that free energy required for ATP synthesis is supplied by the dissipation of Ca<sup>2+</sup> gradient across the SR membrane.

### Structure of Ca<sup>2+</sup>-ATPase

Although the kinetics has been extensively studied, the exact mechanism of this chemiosmotic energy coupling can only be obtained by detailed studies on the structure-function relationships. The importance of the structural studies and characterization of the Ca<sup>2+</sup>-ATPase has been increasingly recognized recently. Ca<sup>2+</sup>-ATPase is about 110,000 Dalton membrane spanning protein. From amino acid sequence and hydropathy profile, the second structure of Ca<sup>2+</sup>-ATPase is expected to be constructed by 10 transmembrane helices and large cytoplasmic domain ( Fig. 2 ). Analysis of the amino acid sequence (4, 5) along with extensive site-directed mutagenesis studies (6) together with the studies on the two-dimensional membrane crystals (7) have concurred in a structural model for the Ca<sup>2+</sup>-ATPase in which the Ca<sup>2+</sup>-binding region is located within a cluster of putative transmembrane  $\alpha$ -herices, M4, M5, and M6, (8) while the ATP binding site, Lys<sup>515</sup>, and phosphorylation site, Asp<sup>351</sup> (6) are located in the cytoplasmic domains. Therefore energy coupling between ATP hydrolysis and Ca<sup>2+</sup> transport seems to require long distance communication through conformational changes. Recent studies of mutational analysis have suggested that several kinds of residues play a key role in




Fig. 2. **Secondary structural model of the Ca<sup>2+</sup>-ATPase.** This model is based on the structure proposed by MacLennan et al (4) and Andersen, J.P. (5). Mutated residues by site-directed mutagenesis are shown by colored circles.

the coupling of ATP hydrolysis to  $\text{Ca}^{2+}$  transport across the SR membrane (9). There are number of investigations by using chemical reagents and mutagenetic methods in analyzing various aspects of structure function relationship in the  $\text{Ca}^{2+}$ -ATPase protein ( see Ref. 10 for review )

### **Functional movements of the ATPase in the SR membrane**

Employing conformational probes such as spin labels and fluorescent dyes, many investigators attempted to detect a conformational changes associated with the transport of  $\text{Ca}^{2+}$  across the SR membrane ( see Ref. 3 for review ). However, none of these studies provide solid evidence that the ATPase molecules undergo specific movements or conformational changes that are directly coupled with the translocation of  $\text{Ca}^{2+}$  across the membrane. One of the major problems in understanding the functional movements of the enzyme molecule is to determine whether the catalytic function is performed by the  $\text{Ca}^{2+}$ -ATPase as a structural unit of a monomer or an oligomer. The oligomericity of the  $\text{Ca}^{2+}$ -ATPase in the SR membrane has been suggested, following the experimental results from a variety of techniques such as analytical centrifugation on the detergent solubilized protein, electron microscopy, fluorescence energy transfer, fluorescence anisotropy measurements as laser flash induced photodichroism ( see Ref. 11 for review ). However, it has not been determined whether or not the molecular interaction of the  $\text{Ca}^{2+}$ -ATPase in the SR membrane plays an essential role in the  $\text{Ca}^{2+}$  transport.

### **Purposes of studies on the structure function relationship of scallop $\text{Ca}^{2+}$ -ATPase**

For the SR isolated from poikirothermal animal muscle, temperature has been considered to be one of a crucial factors influencing its catalytic functions. Abe *et al.* (12) were successful in preparing SR membrane from scallop (*Patinopecten yessoensis*) adductor muscle with retention of full  $\text{Ca}^{2+}$ -transport activity. They found that the isolated scallop SR has characteristics in the temperature sensitivity that pronouncedly differs from those of homiothermal animals such as a rabbit. Both activities of ATP hydrolysis and  $\text{Ca}^{2+}$  transport by scallop SR were irreversibly lost above  $37^{\circ}\text{C}$ , while almost activities of rabbit SR retained even at a temperature as high as  $50^{\circ}\text{C}$ .

The first purpose of this work is to elucidate and characterize the reaction step in the catalytic cycle of scallop  $\text{Ca}^{2+}$ -ATPase upon which thermal disruption drastically

affects. We found that the basic aspects of ATP hydrolysis by the scallop SR are similar for that of rabbit SR and that when the scallop enzyme exist in  $E_1$  state it can be easily inactivated by heat, whereas the enzyme was completely protected from the heat inactivation when the enzyme was fixed in  $E_2$  state (13). However, no protection was observed for the  $Ca^{2+}$ -transport across the membrane even when the enzyme was fixed in the  $E_2$  state. This uncoupling between the  $Ca^{2+}$  transport and the ATP hydrolysis did not lead to a rise in the  $Ca^{2+}$  permeability of SR membrane ( Part I ).

The second purpose of this work is to characterize the primary structure which contribute to the stabilization of the rabbit  $Ca^{2+}$ -ATPase. For this purpose we have cloned cDNA of the  $Ca^{2+}$ -ATPase (SERCA1) from scallop adductor muscle, and predicted entire amino acid sequence of this enzyme (14). Based on detailed comparison of the amino acid sequence with that of rabbit  $Ca^{2+}$ -ATPases, sequence analysis by using fluorescent SH reagent, and site-directed mutagenesis, it is suggested that Cys<sup>675</sup> may binds to Cys<sup>349</sup> through disulfide bond on the rabbit  $Ca^{2+}$ -ATPase, which can contribute to the thermal stabilization of the enzyme, while that lacking in the disulfide bond in the scallop  $Ca^{2+}$ -ATPase may cause the thermal destabilization of the enzyme ( Part II and Part III).

The third purpose of this work is to provide direct evidence suggesting that the catalytic function of  $Ca^{2+}$  transport is performed by the  $Ca^{2+}$ -ATPase in the oligomeric form. In this study, rabbit and scallop  $Ca^{2+}$ -ATPases at various protein ratios were cross-reconstituted into proteoliposomes. When proteoliposomes containing rabbit and scallop SR at a protein ratio of 1:1 were pre-incubated at 39°C for 10 min, the Ca-transport activity was almost completely lost (15). These results indicate that molecular interaction in the SR membrane is essential for  $Ca^{2+}$ -transport ( part IV ).

## REFERENCES

- 1) Melzer, W., Herrmann-Frank, A., Lüttgau, H.Ch. (1995) The role of  $\text{Ca}^{2+}$  ions in excitation-contraction coupling of skeletal muscle fibres. *Biochim. Biophys. Acta* 1241, 59-116
- 2) Tada, M., Yamamoto, T., and Tonomura, Y. (1978) Molecular mechanism of active calcium transport by sarcoplasmic reticulum. *Physiol. Rev.* 58, 1-79
- 3) Suzuki, H. (1998) Calcium pump of plasma membrane and sarco(endo)plasmic reticulum. *Protein, Nucleic acid, Enzyme* 43, 1610-1621
- 4) MacLennan, D.H., Brandl, C.J., Korezak, B., and Green, N.M. (1985) Amino-acid sequence of a  $\text{Ca}^{2+} + \text{Mg}^{2+}$ -dependent ATPase from rabbit muscle sarcoplasmic reticulum, deduced from its complementary DNA sequence. *Nature* 316, 696-700
- 5) Andersen, J.P. (1995) Dissection of the function domains of the sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase by site-directed mutagenesis. *Bioscience Reports.* 15, 243-261
- 6) MacLennan, D.H., Clarke, D.M., Loo, T.W., and Skerjanc, I.S. (1992) Site-directed mutagenesis of the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum. *Acta Physiol. Scand.* 146, 141-150
- 7) Zhang, P., Toyoshima, C., Yonekura, K., Green, N.M., and Stokes, D. (1998) Structure of the calcium pump from sarcoplasmic reticulum at 8-Å resolution *Nature* 392, 835-839
- 8) Clarke, D.M., Loo, T.W., and MacLennan, D.H. (1989) Location of high affinity  $\text{Ca}^{2+}$ -binding sites within the predicted transmembrane domain of the sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase. *Nature* 339, 476-478
- 9) Andersen, J.P. and Sorensen, T. (1996) Site-directed mutagenesis studies of energy coupling in the sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase. *Biochim. Biophys. Acta* 1275, 118-122
- 10) Møller, J.V., Juul, B., and leMaire, M. (1996) Structural organization, ion transport, and energy transduction of P-type ATPases. *Biochim. Biophys. Acta* 1286, 1-51
- 11) Inesi, G. (1985) Mechanism of calcium transport. *Ann. Rev. Physiol.* 47, 573-601
- 12) Abe, M., Shirakata, Y., Sato, D., Konishi, K., Watanabe, T., and Nakamura, J. (1992) Isolation and characteristics of scallop sarcoplasmic reticulum with calcium transport activity. *J. Biochem.* 112, 822-827

- 13) Nagata, Y., Nakamura, J., and Yamamoto, T. (1996) Protection of scallop sarcoplasmic reticulum ATPase from thermal inactivation by removal of calcium from high-affinity binding sites on the enzyme. *J. Biochem.* 119, 1100-1105
- 14) Nagata, Y., Yamamoto, T., Ema, M., Mimura, j., Fujii-Kuriyama, Y., Suzuki, T., Furukohri, T., Konishi, K., Sato, D., Tajima, G., and Nakamura, J. (1998) cDNA cloning and predicted structure of scallop sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase. *Comp. Biochem. Physiol. Part B* 119, 777-785
- 15) Nagata, Y., Nakamura, J., and Yamamoto, T. (1997) Temperature sensitivity of proteoliposomes reconstituted from a mixture of scallop and rabbit sarcoplasmic reticulum Ca<sup>2+</sup>-ATPases. *J. Biochem.* 121, 648-653

## Part I

Protection of Scallop Sarcoplasmic Reticulum ATPase from Thermal Inactivation by  
Removal of Calcium from High-Affinity Binding Sites on the Enzyme

## SUMMARY

Sarcoplasmic reticulum (SR) vesicles were isolated from scallop muscle by the method of Abe *et al.* (J.Biochem.112,822-827,1992) and their thermolability was examined in the presence and absence of  $\text{Ca}^{2+}$ . When SR was preincubated at  $38^{\circ}\text{C}$  in the presence of  $0.1\text{ mM Ca}^{2+}$ ,  $\text{Ca}^{2+}$ -transport activity decreased as a function of time with a half-inhibition time of about 5 min. Activities of the  $\text{Ca}^{2+}$ -dependent ATPase, phosphoenzyme (EP) formation and  $\text{E}_2$  to  $\text{E}_1$  transition were decreased by the heat treatment in parallel with the  $\text{Ca}^{2+}$ -transport activity. In contrast, when SR was preincubated at  $38^{\circ}\text{C}$  in the presence of  $2\text{-}5\text{ mM EGTA}$ , all of these activities, except for the  $\text{Ca}^{2+}$ -transport, were markedly protected from the heat inactivation. The uncoupling between  $\text{Ca}^{2+}$ -transport and the ATPase reaction did not lead to a rise in the  $\text{Ca}^{2+}$  permeability of SR membrane. Plots of the ATPase activity or steady-state level of EP against  $\text{pCa}$  in the thermal incubation medium revealed a typical sigmoidal curve with a half-inhibition concentration and Hill number of about  $0.5\ \mu\text{M}$  and 1.80, respectively. These results suggest that 2 mol of  $\text{Ca}^{2+}$  must be removed from the high-affinity  $\text{Ca}^{2+}$  binding sites on the ATPase to stabilize the  $\text{Ca}^{2+}$ -ATPase against heat inactivation.

The protection from heat inactivation disappeared if SR was preincubated at  $38^{\circ}\text{C}$  after having been solubilized with a nonionic detergent, but returned when the detergent was removed to reconstitute the SR membrane. These results suggest that the protection of ATPase from thermal inactivation in EGTA may require a membrane structure in which the ATPase molecules exist in an appropriate arrangement.



## INTRODUCTION

$\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum (SR) couples the energy derived from ATP hydrolysis with the transport of  $\text{Ca}^{2+}$  across the SR membrane against a concentration gradient (1). In the presence of  $\text{Ca}^{2+}$ , ATP is hydrolyzed into ADP and Pi through two kinds of phosphorylated intermediates ( $\text{E}_1\text{P}$  and  $\text{E}_2\text{P}$ ), followed by a transition of the enzyme state from  $\text{E}_2$  to  $\text{E}_1$  (2). Two  $\text{Ca}^{2+}$  externally bound to high-affinity binding sites of the ATPase are occluded by the enzyme when  $\text{E}_1\text{P}$  is formed, and they are released into the SR lumen when  $\text{E}_1\text{P}$  is converted into  $\text{E}_2\text{P}$ .

Much work has been done to characterize the molecular movements of ATPase which could be associated with the  $\text{Ca}^{2+}$ -transport across the SR membrane by means of variety of methods, such as chemical cross linking (3-5), fluorometric studies (6,7), saturation transfer EPR methods (8-11), and electron microscopic observations of two-dimensional crystallization of the ATPase on SR membrane (12-14), but the movements remain poorly understood.

In the present study, I investigated the effects of thermal treatment on the catalytic functions of scallop SR under various conditions. In the preceding study, Abe *et al* (15) found that scallop SR is highly sensitive to heat; both  $\text{Ca}^{2+}$ -transport and ATP hydrolysis activities are rapidly lost at temperatures higher than  $37^\circ\text{C}$ . In the present work, I found that susceptibility of ATP hydrolysis to heat could be prevented by removing 2 mol of  $\text{Ca}^{2+}$  from the high-affinity binding sites on the enzyme or by adding unhydrolyzable nucleotide. This protection disappeared when SR membrane was destroyed by a nonionic detergent, while it was restored by removing the detergent to reconstitute the membrane. These results raise the possibility that the thermal stability of  $\text{Ca}^{2+}$ -ATPase might be related to the formation of an oligomeric ATPase in the scallop SR membrane.

## EXPERIMENTAL PROCEDURES

**Materials** --- SR was isolated from the striated portion of scallop (*Patinopecten yessoensis*) adductor muscle according to a method described previously (15). The SR sample was divided into small pieces, quickly frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$ . Pyruvate kinase, lactate dehydrogenase, NADH, PEP were purchased from Boehringer Mannheim.  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  was purchased from Amersham,  $^{45}\text{CaCl}_2$  was from Japan RI Association, and Antipyrilazo III was from Nacalai.

**Heat treatment of SR** --- A small portion of the SR suspension was added at 0.5-2 mg/ml to the preincubation medium, which contained 0.1 M KCl, 10% glycerol, 5 mM  $\text{MgCl}_2$ , and 20 mM TES (pH 7.2) usually at  $38^{\circ}\text{C}$ . At the indicated times, 0.1 ml of the SR suspension was transferred into 5-20 volumes of assay medium which had been kept cool on ice. In some experiments, the preincubation medium was frozen in liquid nitrogen to stop the thermal treatment.

**Measurements of activities** ---  $\text{Ca}^{2+}$ -transport by SR was measured in a medium containing 0.025-0.05 mg/ml SR protein, 0.1 M KCl, 5 mM  $\text{MgCl}_2$ , 5-10% glycerol, 40  $\mu\text{M}$   $\text{CaCl}_2$ , 0.2 mM Antipyrilazo III, and 20 mM TES at pH 7.2. The reaction was started at  $23^{\circ}\text{C}$  by the addition of 0.2-1 mM ATP.  $\text{Ca}^{2+}$  uptake into SR was determined by measuring the absorption at 700 nm in the reaction mixture (16).

The permeability of the SR membrane to  $\text{Ca}^{2+}$  was measured by essentially the same method as that described previously (16) except that  $^{45}\text{Ca}^{2+}$  loading and  $\text{Ca}^{2+}$  efflux assay were both carried out at 0 and  $25^{\circ}\text{C}$ , and that  $^{45}\text{Ca}^{2+}$ -loaded SR vesicles were incubated in the 5 mM EGTA medium for 5 s.

ATP hydrolysis was measured under similar conditions to those of  $\text{Ca}^{2+}$ -transport assay except that the  $\text{Ca}^{2+}$  indicator was omitted and 2-5  $\mu\text{M}$  A23187 was added to the reaction mixture. ATP hydrolysis was measured in the presence of an ATP-regenerating system (0.5 mM NADH, and 1.5 mM PEP, 0.1-1 unit of lactate dehydrogenase and 0.2-2 unit of pyruvate kinase), and the amount of ADP liberated from ATP was determined by measuring the decrease in absorbance of NADH at 340 nm.

**Phosphorylation of  $\text{Ca}^{2+}$ -ATPase** --- SR, 0.05-0.1 mg/ml, was phosphorylated with 10-100  $\mu\text{M}$   $\text{AT}^{32}\text{P}$  at  $0^{\circ}\text{C}$ . At the indicated time, 5% TCA with 2 mM ATP and 0.5 mM phosphate were added to stop the reaction. The amount of EP was determined as

described previously (17).

$E_2$ - $E_1$  transition was measured as a fluorescence change of tryptophan in  $Ca^{2+}$ -ATPase after addition of 2 mM  $CaCl_2$  to the SR suspension containing 2 mM EGTA. For these measurements, a Shimadzu 1000 fluorometer was used.


**Reconstitution of SR membrane** --- SR (2.5 mg/ml) was solubilized with 30 mg/ml of  $C_{12}E_9$ . The suspension was centrifuged at  $540,000 \times g$  for 20 min to remove insoluble debris. Soybean asolectin (25 mg/ml) was added to the supernatant and the suspension was incubated at  $23^\circ C$  for 1 h with 0.2 g/ml of Bio-beads SM-2 to remove the detergent. The Bio-beads were removed by filtration, and the filtrate was centrifuged at  $540,000 \times g$  for 30 min to precipitate reconstituted SR vesicles.

## RESULTS

**Inactivation of Ca<sup>2+</sup>-transport and ATP hydrolysis by heat treatment of SR in the presence of Ca<sup>2+</sup>** --- 2 mg/ml SR was preincubated at 38°C for various time periods in 0.1 ml of reaction medium containing 0.1 mM CaCl<sub>2</sub>, 10% glycerol, 0.1 M KCl, 20 mM TES (pH 7.2) and 5 mM MgCl<sub>2</sub>, then rapidly transferred into liquid nitrogen to stop the heat treatment. Ca<sup>2+</sup>-transport assay was carried out at 23°C under the conditions described in "EXPERIMENTAL PROCEDURES". The Ca<sup>2+</sup>-transport showed a slow decrease in rate immediately after initiating the preincubation. This was followed by a rapid decrease, and the activity almost disappeared within 8 min (Fig. 1). The rate of inactivation increased as the preincubation temperature increased (data not shown). In the following experiments, scallop SR was mainly preincubated at 38°C, because the catalytic activities of SR decreased at a rate slow enough to allow accurate analysis of the inactivation process. The activity of Ca<sup>2+</sup>-dependent ATPase decreased essentially in parallel to that of Ca<sup>2+</sup>-transport. These complicated time courses of heat inactivation might suggest the existence of different kinds of thermally sensitive regions on the ATPase protein. The loss of ATPase activity may be attributed to the lack of EP formation but not to EP decomposition, because the steady-state level of EP decreased at a similar rate to that of the Ca<sup>2+</sup>-ATPase (Fig. 1).

Figure 2 compares the Ca<sup>2+</sup> concentration dependence of EP formation after preincubation of SR for 6 min at 38 and 23°C. In both cases, plots of the EP level against pCa revealed a typical sigmoidal curve with a K<sub>d</sub> and Hill number of about 0.2 μM and 1.8, respectively. Therefore, the inhibition of EP formation by heat treatment of SR does not seem to be due to a decrease in the affinity of this enzyme for Ca<sup>2+</sup>.

**Protection of ATPase from thermal inactivation in the presence of EGTA** -- The Ca<sup>2+</sup>-ATPase of scallop SR became very stable at high temperature if EGTA was added to the preincubation mixture. Under this condition, more than 90, 70, and 80% of the activities of ATP hydrolysis, EP formation and E<sub>2</sub>-E<sub>1</sub> transition, respectively, remained at 12 min after initiation of the thermal treatment (Fig. 3). Figure 4 shows the dependence of susceptibility to thermal inactivation of ATPase and E<sub>2</sub>-E<sub>1</sub> transition on the Ca<sup>2+</sup> concentration of the preincubation medium. Scallop SR was preincubated at 38°C for 10 min at different concentrations of Ca<sup>2+</sup> from pCa 9 to 5. Significant



**Fig. 1. Time courses of heat inactivation of Ca<sup>2+</sup> -transport, ATP hydrolysis and EP formation.** SR was added at 38°C to the preincubation medium containing 0.1 mM CaCl<sub>2</sub> to a final concentration of 2 mg/ml. At the indicated times, 0.1 ml of the SR suspension was immediately frozen in liquid nitrogen to terminate the thermal treatment. Measurements of Ca<sup>2+</sup>-transport ( O ), ATP hydrolysis( □ ), and EP formation( Δ ) were carried out at 23°C as described in "EXPERIMENTAL PROCEDURES".





Fig. 2. Effect of heat treatment of SR on the  $\text{Ca}^{2+}$  concentration dependence of EP formation. SR was preincubated at 38°C (○) or 23°C (□) for 6 min in the presence of 0.1 mM  $\text{CaCl}_2$ , then centrifuged to wash the SR membrane. Phosphorylation of ATPase was initiated at 23°C by adding 1 mM  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  in the presence of 0.1 mM  $\text{CaCl}_2$ . 5 sec later, the reaction was terminated by addition of 5% TCA, and the amount of EP was determined as described in "EXPERIMENTAL PROCEDURES".



**Fig. 3. Protection of ATP hydrolysis, EP formation, and E<sub>2</sub>-E<sub>1</sub> transition against thermal treatment of SR in EGTA.** SR was preincubated at 38°C in the presence of 5 mM EGTA (open symbols) or 0.1 mM CaCl<sub>2</sub> (closed symbols) for the indicated times. These samples were washed by centrifugation, and resuspended in the assay medium. ATP hydrolysis(○, ●), EP formation (△, ▲), and E<sub>1</sub>-E<sub>2</sub> transition (□, ■) were measured as described in "EXPERIMENTAL PROCEDURES". All the activities were presented as a percentage of those obtained with the untreated SR.

enhancement of the thermal inactivation was observed by increasing the concentration of  $\text{Ca}^{2+}$  above pCa 6, while maximal activities of both ATP hydrolysis and  $\text{E}_2\text{-E}_1$  transition were retained even after 12 min of thermal treatment below pCa 8. The plots of the remaining activities of ATP hydrolysis and  $\text{E}_2\text{-E}_1$  transition against pCa of the preincubation medium revealed typical sigmoidal curves with a dissociation constant and Hill number of about 0.5  $\mu\text{M}$  and 1.6-1.8, respectively. Essentially the same sigmoidal curve of  $\text{Ca}^{2+}$  dependence was observed for EP formation (data not shown). These observations indicate that removal of 2 mol of  $\text{Ca}^{2+}$  from the high-affinity sites on the  $\text{Ca}^{2+}$ -ATPase or fixing the enzyme in the  $\text{E}_2$  state might be required to make this enzyme resistant to heat inactivation.

**Uncoupling of  $\text{Ca}^{2+}$ -transport from ATP hydrolysis after thermal treatment of SR in EGTA** --- As shown in Fig. 5, the thermal treatment of SR in the presence of EGTA unexpectedly failed to protect the  $\text{Ca}^{2+}$ -transport activity. These results support the possibility that the thermally sensitive region for the  $\text{Ca}^{2+}$ -transport is different from that for ATP hydrolysis. Uncoupling of the  $\text{Ca}^{2+}$ -transport from the ATPase reaction may lead to an increase in the  $\text{Ca}^{2+}$  permeability of the SR membrane. In order to test this possibility, I examined the leakiness of the SR membrane to  $\text{Ca}^{2+}$  as shown in Fig. 6. SR vesicles were preincubated in the presence of 2 mM EGTA at 38°C for the indicated times. The treated SR vesicles were passively loaded with  $^{45}\text{Ca}^{2+}$  by incubation with 10 mM  $^{45}\text{Ca}^{2+}$  for 5 h at 0°C, or for 2 h at 23°C. After the  $\text{Ca}^{2+}$  concentrations inside and outside the SR membrane had reached equilibrium, a small amount of SR suspension was transferred into a solution containing 5 mM EGTA and incubated for 5 s. The amount of  $^{45}\text{Ca}^{2+}$  remaining in the SR vesicles was measured by the filtration method (16). Heat treatment of SR at 38°C for up to at least 15 min did not result in any rise in the  $\text{Ca}^{2+}$  permeability of the SR membrane. Therefore the lack of  $\text{Ca}^{2+}$ -transport in the SR is due to some facts other than  $\text{Ca}^{2+}$  leakiness of the membrane.

Lepock *et al.*(18,19) observed that incubation of rabbit SR at 37°C in the presence of EGTA resulted in a rapid loss of  $\text{Ca}^{2+}$ -transport with no effect on ATPase activity. With rabbit SR, however, neither ATP hydrolysis nor  $\text{Ca}^{2+}$ -transport was destroyed by incubation with  $\text{Ca}^{2+}$  below 49°C. These observations are in contrast with our data for scallop SR, indicating that the mechanism of thermal inactivation in scallop SR may differ from that in rabbit SR.








Fig. 4. Ca<sup>2+</sup> concentration dependence of heat sensitivity of Ca<sup>2+</sup>-ATPase and E2-E1 transition. SR was preincubated at 38°C for 10 min in the presence of various concentrations of Ca<sup>2+</sup>. Ca<sup>2+</sup>-ATPase (□) and E2-E1 transition (○) activities were measured as described in "EXPERIMENTAL PROCEDURES".




**Fig. 5. Uncoupling of ATP hydrolysis from Ca<sup>2+</sup> transport after thermal treatment of SR in the presence of EGTA.** SR was preincubated at 38°C in the presence of 2 mM EGTA for the indicated times. ATP hydrolysis (○) and Ca<sup>2+</sup> transport (●) were measured as described in "EXPERIMENTAL PROCEDURES".




**Fig. 6. Effects of heat treatment on the  $\text{Ca}^{2+}$  permeability of SR membrane.** SR was treated at  $38^\circ\text{C}$  for the indicated times as presented in Fig. 5, then the SR vesicles were loaded with  $10\text{ mM }^{45}\text{Ca}^{2+}$  at  $0^\circ\text{C}$  for 5 h (□) or at  $25^\circ\text{C}$  for 2 h (○). The SR suspension was transferred into 20 volumes of assay medium containing 5 mM EGTA. After 5 s, the reaction medium was passed through a membrane filter, and the amount of  $\text{Ca}^{2+}$  retained on the filter was determined as described in "EXPERIMENTAL PROCEDURES".

**Role of membrane structure in thermal stabilization of Ca<sup>2+</sup>-ATPase in EGTA** --- To test whether or not EGTA protection of Ca<sup>2+</sup>-ATPase from heat inactivation requires some membrane structure of SR, I compared the thermal lability of the ATPase solubilized in C<sub>12</sub>E<sub>9</sub> and the reconstituted membrane-bound ATPase (Fig. 7). These samples were preincubated at 38°C in the presence and absence of 2 mM EGTA for various time periods. As shown in Fig. 7a, the Ca<sup>2+</sup>-dependent ATPase activity of the solubilized ATPase was completely lost during the preincubation even in the presence of EGTA. When solubilized ATPase was reconstituted into the proteoliposome, EGTA protection of the enzyme from thermal lability was fully restored (Fig. 7b).

**Other factors affecting thermal inactivation of SR functions** --- Figure 8 shows the effect of nucleotide on the thermal inactivation of SR function. Preincubation of SR at 38°C in the presence of 1 mM AMP-PNP induced rapid inactivation of Ca<sup>2+</sup>-transport with complete retention of ATP hydrolytic activity. The Ca<sup>2+</sup>-ATPase activity shows strong resistance to the heat treatment in spite of the presence of 0.1 mM Ca<sup>2+</sup>. Other nucleotides (ADP and AMP) up to 10 mM showed no effect on the SR functions (data not shown). I examined the effects of other reagents such as β-mercaptoethanol, dithiothreitol, vanadate, and polyclonal antibody against the Ca<sup>2+</sup>-ATPase purified from scallop SR, but they failed to offer protection against heat inactivation of SR catalytic functions.



**Fig. 7. a: Inactivation of the solubilized ATPase by thermal treatment in the presence and absence of  $\text{Ca}^{2+}$ .** SR at 2 mg/ml was solubilized with 30 mg/ml C12E9, then preincubated in the presence of 0.1 mM  $\text{CaCl}_2$  (●) or 2 mM EGTA (○) at 38°C for the indicated times. The SR suspension was added to 20 volumes of assay medium. ATPase activity was measured at the final  $\text{Ca}^{2+}$  concentration of 0.1 mM as described in "EXPERIMENTAL PROCEDURES". **b: Restoration of thermal resistance of the  $\text{Ca}^{2+}$  ATPase in EGTA on removal of detergent from solubilized ATPase.** SR membrane was reconstituted as described in "EXPERIMENTAL PROCEDURES", and the reconstituted membrane was preincubated at 38°C in the presence of 0.1 mM  $\text{CaCl}_2$  (●) or 5 mM EGTA (○) for the indicated times. ATPase activity of the reconstituted SR was measured as described in (a).



**Fig. 8. Effects of thermal treatment of SR in the presence of AMP-PNP on the ATP hydrolysis and Ca<sup>2+</sup> transport.** SR (5 mg/ml) was preincubated at 38°C in the presence of 0.1 mM CaCl<sub>2</sub>, and 1 mM AMP-PNP for the indicated time, then washed by centrifugation to remove the nucleotide. ATP hydrolysis (O) and Ca<sup>2+</sup> transport (□) were measured as presented in Fig. 5.

## DISCUSSION

SR membranes isolated from cold water fish, including scallop SR, have striking characteristics of thermolability (15,20,21). As shown in Fig. 1, most of the  $\text{Ca}^{2+}$ -transport and ATPase activities of scallop SR were almost completely lost within 10 min of incubation at  $38^\circ\text{C}$  in the presence of  $\text{Ca}^{2+}$ . Based on preliminary tests, it was considered that the thermal treatment of SR directly affects the  $\text{Ca}^{2+}$ -ATPase in the SR membrane, because the thermal inactivation was still observed when  $\text{Ca}^{2+}$ -ATPase was purified from scallop SR by HPLC in  $\text{C}_{12}\text{E}_9$  and reconstituted into soybean liposome (data not shown).

The data in Figs. 3 and 4 suggest that removal of 2 mol of  $\text{Ca}^{2+}$  from the high-affinity sites on the  $\text{Ca}^{2+}$ -ATPase makes the enzyme thermally resistant. One of the simplest explanations is that the enzyme becomes heat stable if it exists in the SR membrane in the  $\text{E}_2$  form. However, this is ruled out by the fact that the  $\text{Ca}^{2+}$ -ATPase of scallop SR was easily inactivated by heat treatment at low pH without KCl, where the enzyme is believed to exist in the  $\text{E}_2$  state (22-25). In addition, the heat treatment of SR with 1 mM AMP-PNP, which stimulates the conversion of the enzyme from the  $\text{E}_2$  to the  $\text{E}_1$  state (26), completely protected  $\text{Ca}^{2+}$ -ATPase from thermal lability (Fig. 8).

As shown in Fig. 7a, when SR membrane was destroyed by  $\text{C}_{12}\text{E}_9$ , the  $\text{Ca}^{2+}$ -ATPase became thermolabile in EGTA. On the other hand, the thermal resistance of the ATPase was fully restored when the detergent was removed to reconstitute the SR membrane (Fig. 7b). These results suggest that the thermal stability of  $\text{Ca}^{2+}$ -ATPase requires not only removal of  $\text{Ca}^{2+}$  from the enzyme but also correct location in the SR membrane.

Based on these considerations, I propose the following mechanism for the thermal inactivation of  $\text{Ca}^{2+}$ -ATPase in scallop SR membrane. The susceptibility of  $\text{Ca}^{2+}$ -ATPase to heat inactivation may vary depending upon the arrangement of the enzyme molecules in the SR membrane. When the enzyme molecules exist in an oligomeric form, they will be heat resistant, while in a monomeric form, they are easily inactivated by heat treatment. Using fluorometric techniques (6,7), EPR spectrometric analysis (8-11), and electron microscopic observations of unstained SR membrane (12-14,27), many studies have suggested that arrangement of the ATPase molecules in the SR membrane can be varied by adjusting the concentrations of divalent cations and

nucleotides. Stokes and Lacapere (27) demonstrated that simultaneous addition of vanadate and EGTA induced two-dimensional crystallization of ATPase on the surface membrane of rabbit SR, and that the crystal formation was prevented by  $\text{Ca}^{2+}$  at physiological concentrations. They observed that non-hydrolyzable nucleotides affected the crystal formation. These observations are consistent with our present results. Moreover, Yamasaki and co-worker (3,5) suggested that when the SR membrane is allowed to react with a cross linking reagent, N, N'-(1,4-phenylene)bismaleimide (PBM), in the presence of 1 mM AMP-PNP, the  $\text{Ca}^{2+}$ -ATPase molecules predominantly form the dimer in the SR membrane. This corresponds well with the finding that the nucleotide prevented heat inactivation of the ATPase (Fig. 8).

Thus, I might be able to determine whether the  $\text{Ca}^{2+}$ -ATPase molecules exist in the scallop SR membrane in a monomeric form or an oligomeric form simply by examining the susceptibility of this enzyme to thermal inactivation.



## REFERENCES

- 1) Yamamoto, T., Takisawa, H., and Tonomura, Y. (1979) Reaction mechanisms for ATP hydrolysis and synthesis in the sarcoplasmic reticulum. *Curr. Top. Bioenerg.* 9, 179-235
- 2) deMeis, L. and Vianna, A. (1979) Energy interconversion by the  $\text{Ca}^{2+}$ -dependent ATPase of the sarcoplasmic reticulum. *Annu. Rev. Biochem.* 48, 275-292
- 3) Yamasaki, K. and Yamamoto, T. (1989) Effects of adenylyl-5'-imidodiphosphate and vanadate ion on the intermolecular cross-linking of  $\text{Ca}^{2+}$ -ATPase in the sarcoplasmic reticulum membrane with N,N'-(1,4-phenylene)bismaleimide. *J. Biochem.* 106, 1114-1120
- 4) Keresztes, T., Jona, I., Pikula, S. Vegh, M., Mullner, N., Papp, S., and Martonosi, A. (1989) Effect of calcium on the interactions between  $\text{Ca}^{2+}$ -ATPase molecules in sarcoplasmic reticulum. *Biochim. Biophys. Acta* 984, 326-338
- 5) Yamasaki, K., Sano, N., Ohe, M., and Yamamoto, T. (1990) Determination of the primary structure of intermolecular cross-linking sites on the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum using  $^{14}\text{C}$ -labeled N,N'-(1,4-phenylene)bismaleimide or N-ethylmaleimide. *J. Biochem.* 108, 918-925
- 6) Kutchai, H., Mahaney, J.E., Geddis, L.M., and Thomas, D.D. (1994) Hexanol and lidocaine affect the oligomeric state of the Ca-ATPase of sarcoplasmic reticulum. *Biochemistry* 33, 13208-13212
- 7) Karon, B.S., Mahaney, J.E., and Thomas, D.D. (1994) Halothane and cyclopiazonic acid modulate Ca-ATPase oligomeric state and function in sarcoplasmic reticulum. *Biochemistry* 33, 13928-13937
- 8) Bigelow, D.J. and Thomas, D.D. (1987) Rotational dynamics of lipid and the Ca-ATPase in sarcoplasmic reticulum: The molecular basis of activation by diethyl ether. *J. Biol. Chem.* 262, 13449-13456
- 9) Squier, T.C., Bigelow, D.J., and Thomas, D.D. (1988) Lipid fluidity directly modulates the overall protein rotational mobility of the Ca-ATPase in sarcoplasmic reticulum. *J. Biol. Chem.* 263, 9178-9186
- 10) Birmachu, W. and Thomas, D.D. (1990) Rotational dynamics of the Ca-ATPase in sarcoplasmic reticulum studied by time-resolved phosphorescence anisotropy. *Biochemistry* 29, 3904-3914

- 11) Karon, B.S. and Thomas, D.D. (1993) Molecular mechanism of Ca-ATPase activated by halothane in sarcoplasmic reticulum. *Biochemistry* 32, 7503-7511
- 12) Buhle, E.L., Knox, B.E., Serpersu, E., and Aebi, U. (1983) The structure of the Ca<sup>2+</sup> ATPase as revealed by electron microscopy and image processing of ordered arrays. *J. Ultrastruct. Res.* 85, 186-203
- 13) Tayler, K.A., Dux, L., and Martonosi, A. (1986) Three-dimensional reconstruction of negative stained crystals of the Ca<sup>2+</sup>-ATPase from muscle sarcoplasmic reticulum. *J. Mol. Biol.* 187, 417-427
- 14) Dux, L., Taylor, K.A., Ting-Beall, H.P., and Martonosi, A. (1985) Crystallization of the Ca<sup>2+</sup>-ATPase of sarcoplasmic reticulum by calcium and lanthanide ions. *J. Biol. Chem.* 260, 11730-11743
- 15) Abe, M., Shirakata, Y., Sato, D., Konishi, K., Watanabe, T., and Nakamura, J. (1992) Isolation and characteristics of scallop sarcoplasmic reticulum with calcium transport activity. *J. Biochem.* 112, 822-827
- 16) Hirose, T., Yamasaki, K. and Yamamoto, T. (1995) Irradiation with ultraviolet light in the presence of vanadate increases Ca<sup>2+</sup> permeability of the sarcoplasmic reticulum membrane *via* Ca<sup>2+</sup>-ATPase. *J. Biochem.* 117, 324-330
- 17) Kanazawa, T., Yamada, S., Yamamoto, T., and Tonomura, Y. (1971) Reaction mechanism of the Ca<sup>2+</sup>-dependent ATPase of sarcoplasmic reticulum from skeletal muscle. V. Vectorial requirements for calcium and magnesium ions of three partial reactions of ATPase: formation and decomposition of a phosphorylated intermediate and ATP formation from ADP and the intermediate. *J. Biochem.* 70, 95-123
- 18) Lepock, J.R., Rodahl, A.M., Zhang, C., Heynen, M.L., Waters, B., and Cheng, K.-H. (1990) Thermal denaturation of the Ca<sup>2+</sup>-ATPase of sarcoplasmic reticulum reveals two thermodynamically independent domains. *Biochemistry* 29, 681-689
- 19) Cheng, K.H. and Lepock, J.R. (1992) Inactivation of calcium uptake by EGTA is due to an irreversible thermotropic conformational change in the calcium binding domain of the Ca<sup>2+</sup>-ATPase. *Biochemistry* 31, 4074-4080
- 20) Chini, E.N., de Toledo, F.G.S., Albuquerque, M.C., and deMeis, L. (1993) The Ca<sup>2+</sup>-transporting ATPases of rabbit and trout exhibit different pH- and temperature-dependences. *Biochem. J.* 293, 469-473
- 21) de Toledo, F.G.S., Albuquerque, M.C., Goulart, B.H., and Chini, E.N. (1995) Different thermostabilities of sarcoplasmic reticulum (Ca<sup>2+</sup>, Mg<sup>2+</sup>)-ATPase from

rabbit and trout muscles. *Comp. Biochem. Physiol.* 111C, 93-98

- 22) Pick, U. (1982) The interaction of vanadate ions with the Ca-ATPase from sarcoplasmic reticulum. *J. Biol. Chem.* 257, 6111-6126
- 23) Pick, U. and Karlish, J.D. (1982) Regulation of the conformational transition in the Ca-ATPase from sarcoplasmic reticulum by pH, temperature, and calcium ions. *J. Biol. Chem.* 257, 6120-6126
- 24) Henderson, I.M.J., Khan, Y.M., and Lee, A.G. (1994) Binding of  $\text{Ca}^{2+}$  to the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum. *Biochem. J.* 305, 615-624
- 25) Lee, A.G., Baker, K., Khan, Y.M., and East, J.M. (1995) Effect of  $\text{K}^+$  on the binding of  $\text{Ca}^{2+}$  to the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum. *Biochem. J.* 315, 225-231
- 26) deMeis, L. (1981) *The Sarcoplasmic Reticulum: Transport and Energy Transduction.* pp. 69-81, John Wiley & Sons, Inc., NY
- 27) Stokes, D.L. and Lacapere, J.J. (1994) Conformation of  $\text{Ca}^{2+}$ -ATPase in two crystal forms: Effects of  $\text{Ca}^{2+}$ , thapsigargin, adenosine 5'-(methylene)triphosphate, and chromium(III)-ATP on crystallization. *J. Biol. Chem.* 269, 11606-11613

## Part II

cDNA cloning and predicted primary structure of scallop sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase

## SUMMARY

Sarcoplasmic reticulum (SR)  $\text{Ca}^{2+}$ -ATPase of the scallop cross-striated adductor muscle was purified with deoxycholate and digested with lysyl endopeptidase for sequencing of the digested fragments. Overlapping cDNA clones of the ATPase were isolated by screening the cDNA library with a RT-PCR product, as a hybridization probe, which encodes the partial amino acid sequence of the ATPase. The predicted amino acid sequence of the ATPase contained all the partial sequences determined with the proteolytic fragments and consisted of the 993 residues with about 70% overall sequence similarity to those of the SR ATPases from rabbit fast-twitch and slow-twitch muscles. An outline of the structure of the scallop ATPase molecule is predicted to mainly consist of 10 transmembrane and 5 stalk domains with two large cytoplasmic regions as observed with the rabbit SR. The sequence relationship between scallop and other sarco/endoplasmic reticulum-type  $\text{Ca}^{2+}$ -ATPases is discussed.

## INTRODUCTION

Intracellular calcium concentrations in eucaryotic cells are regulated by the plasma membrane and by intracellular organelles. In cardiac and skeletal muscle cells, the sarcoplasmic reticulum (SR) plays a primary role in regulating cytoplasmic calcium concentrations (1). The  $\text{Ca}^{2+}$ -ATPase of the SR is an integral membrane protein that helps to relax the striated muscles contraction by pumping calcium from the cytoplasm into the SR by hydrolytic coupling with ATP. The transport reaction is performed with a cyclic change of  $E_1$  (high affinity conformation of the enzyme for calcium) and  $E_2$  (low affinity conformation of the enzyme for calcium) states of the ATPase, accompanied by the monomer-dimer transition of the ATPase molecules on the SR membrane (2). By using the SR of rabbit fast-twitch skeletal muscle, the mechanism of calcium transport in the SR has been extensively studied from physiological, biochemical, and molecular biological points of view (3, 4). Almost all such studies on the invertebrate SR have been carried out with the SR from cross-striated adductor muscle of adult scallop (5-11). Although the scallop SR shares many important properties with the rabbit SR (9), distinct differences between them have been found as follows: (i) Scallop ATPase activity was found to be much more labile to heat than that of rabbit ATPase (9). (ii) When these ATPase molecules were monomerized in the absence of the calcium ion, the scallop molecules became inactivated, while the rabbit counterparts maintained their activities (10). (iii) The dimer state of the scallop molecules was stable without vanadate or phosphate which is required for stabilization of the rabbit molecules in the dimer state (5). The relationship between stability of the activity and the organized state of the scallop ATPase is currently being investigated (10-12). For this purpose, it is necessary to know the primary structure of the scallop ATPase. Although the primary structures of several invertebrate SR  $\text{Ca}^{2+}$ -ATPases have been deduced from the nucleotide sequences of their cDNAs (13-15), the structure of the scallop  $\text{Ca}^{2+}$ -ATPase is not known. Here, we analyzed proteolytic fragments of the ATPase protein from the cross-striated muscle cells of adult scallop, isolated and sequenced its cDNA, and predicted its amino acid sequence. The relationship of amino acid sequences between scallop and other sarco/endoplasmic reticulum-type  $\text{Ca}^{2+}$ -ATPases (SERCA) is discussed.

## EXPERIMENTAL PROCEDURES

**Sequencing of fragments of the ATPase protein** --- The SR was prepared from the cross-striated adductor muscle of adult scallop (*Patinopecten yessoensis*) as reported previously (9). The Ca<sup>2+</sup>-ATPase was purified by washing the SR with sodium deoxycholate in a 2 : 5 ratio of sodium deoxycholate to the reticulum protein according to the methods previously used for purification of Ca<sup>2+</sup>-ATPase from the SR of rabbit fast-twitch skeletal muscle (16). The preparations of the SR and the purified ATPase were analyzed by SDS-PAGE by using gels containing 7.5 and 4% acrylamide as separating and stacking gels, respectively. The gel was stained with Coomassie blue. After delipidation of the purified ATPase preparation with chloroform and methanol (2 : 1), the protein was reduced and carboxymethylated according to the method of Suzuki *et al.* (17), and then digested with lysyl endopeptidase (18). The digested products were first isolated by reverse-phase chromatography on a column of Cosmosil 5C<sub>18</sub>-300 (4.6 × 150 mm) with a linear gradient of 0-90% acetonitrile in 0.1% trifluoroacetic acid. Peptides for sequence analysis were purified further by re-chromatography on the same column with a linear gradient of the acetonitrile in 10 mM ammonium acetate (18). By the re-chromatography, 12 preparations of the purified peptide were obtained. The amino acid sequences of the peptides were determined by an automated protein sequencer (Applied BioSystems 476A).

**Cloning and sequencing of cDNA** --- Total RNA was extracted from the cross-striated adductor muscle of a one-year-old adult scallop by the method (19) of guanidium thiocyanate/CsCl ultracentrifugation using cesium TFA instead of CsCl. Poly(A)<sup>+</sup> RNA was isolated from the total RNA by oligo(dT)-cellulose chromatography. cDNA with average length of 1.5-kb was synthesized from the isolated poly(A)<sup>+</sup> RNA by using Superscript<sup>TM</sup> reverse transcriptase (GIBCO BRL) with random primers. A part of the cDNA was inserted into λgt10 for construction of a cDNA library. The remaining cDNA was used for RT-PCR using a pair of degenerated primers, i.e., 5'-AA(A/G)AA(A/G)GAGTT(T/C)AC(A/T/C/G)CT(T/C/G)GA(A/G)TT-3' as an upstream primer and 5'-AT(A/T/C/G)GCCAT(A/G)TA(A/G)CG(A/G/T/C)AA(A/G)AA-3' as a downstream primer. These primers were designed based on the two conserved amino acid sequences of SERCA-type Ca<sup>2+</sup>-ATPases, SKEFILEF and FFRYMAI, which are close to the ATP-binding region (19) and in one of the

transmembrane regions (19), respectively (see text for details). After 3 cycles of PCR (94°C for 1 min, 45°C for 1 min and then 72°C for 1 min), 27 cycles of PCR (94°C for 1 min, 55°C for 1 min and then 72°C for 1 min) were performed. The PCR products (about 1,000 bp) were subcloned into a phagemid vector pBluescript SK(+). The nucleotide sequences of the products were determined by the dideoxy chain-termination method using the Dye Terminator Cycle sequencing kit and an automated DNA sequencer (Applied Biosystems, model 373A18). One of the cDNA clones was found to include one of the partial amino acid sequences (see text for details) of the scallop ATPase protein. This was used as a probe for screening the cDNA library by the method of plaque hybridization. The hybridization was carried out under the standard protocol of hybridization (20), and then, the filter was washed with  $0.5 \times$  SSC (0.15M NaCl, 0.015M sodium citrate) containing 0.1% SDS at 65°C. 25,000 independent plaques of the library were screened and 11 clones were obtained. By sequencing cDNA of the obtained 11 clones, the longest four clones were found to overlap with the screening probe which was mentioned above. The overlapping clones were sequenced with the automated DNA sequencer, described above. The plasmid pBluescript SK(+) was utilized for subcloning the different DNA fragments of the overlapping clones to be sequenced.



## RESULTS AND DISCUSSION

In rabbit cross-striated muscle cells, two types of  $\text{Ca}^{2+}$ -ATPase genes have been found to be expressed, i.e., they are fast-twitch form (SERCA1) gene and slow-twitch-form (SERCA2) gene, respectively (19). The difference between the predicted amino acid sequences of these ATPases have been shown to be about 17% and be scattered over the whole sequences rather than concentrated in some region of the sequences (19). In order to obtain enough data to examine the homogeneity of the ATPase gene in cross-striated muscle cells of adult scallop, we thoroughly analyzed the amino acid sequence of the scallop  $\text{Ca}^{2+}$ -ATPase (Fig. 1). The ATPase was purified with deoxycholate from the SR preparation in which the ATPase protein comprised about 40 % of the total SR proteins (Fig. 1A). The content of the ATPase protein in the purified ATPase preparation was estimated to be more than 90% by gel electrophoresis. From lysyl endopeptidase digests of the purified ATPase, 12 peptides, L1-L12, were isolated for sequencing (Fig. 1B). The determined amino acid sequences of the peptides are as follows: IRASLLVPGDIVEISV (L1), IRDE (L2), MFVF (L3), TCDY (L4), MNFFNTGRSGLNLREQGTVCNHVIQ (L5), GAPE (L6), NCRK (L7), ATAEAICRRIGVFGENESTEGMSFT (L8), SRLF (L9), AEIGIAMGSGTAVAK (L10), SASEMVLADDNFATIVSAVEEGRAIYNN (L11) and ISMPVILIDETLK (L12).

4 overlapping cDNA clones were obtained by screening 25,000 plaques of the cDNA library by using a RT-PCR product as a probe; the product, which was used, included one (AEIGIAMGSGTAVAK) of the partial amino acid sequences which were mentioned above. The complete nucleotide sequence of these clones is shown in Fig. 2A. The sequence contains the longest open reading frame of 2,979 bp, which encodes a polypeptide of 993 amino acids with a calculated  $M_r$  of 109,581 and contains a 5'-untranslated leader sequence of 172 nucleotides and a 3'-untranslated 732-nucleotide sequence. The putative initiator methionine was assigned, because the nucleotide sequence around the methionine fit the Kozak's consensus sequence (GCCA/GCCATGG) for translational initiation and a stop codon (TAA) was present at 5' codons upstream of the methionine. The conserved amino acid sequences (SKEFTLEF and FFRYMAI) of SERCA-type  $\text{Ca}^{2+}$ -ATPases, which were used to design the RT-PCR primers, were in the sequences of 479-486 and 833-839, respectively. All of the partial amino acid sequences determined with the purified ATPase were found in

the deduced sequences of 139-154, 234-237, 365-389, 403-406, 451-475, 515-518, 612-615, 629-653, 672-675, 714-727, 728-755 and 972-984. That is, no sequence, which does not correspond to the deduced sequences, was obtained. This suggests that the cDNA encodes the scallop ATPase. In other word, one type of SERCA gene seems to be expressed in the scallop striated muscle cells. In some of vertebrates and invertebrates, SERCA genes have been found to undergo alternative splicing of their primary transcripts, i.e., In rabbit (21), human (22), rat (23) and bird (24), it is shown that the position of amino acid, 993, in their translated amino acid sequences is involved in such pre-mRNA processing. As mentioned above, the position of 993 is that of the last amino acid in scallop amino acid sequence. The scallop sequence size, 993, is smallest in a SERCA family of 20 species as ever known. In Crustacean *Artemia* (25), amino acid residue, 997, was recently shown to be in a position involving such processing. It is, therefore, interesting to know the splicing in the scallop gene.

A secondary structure predicted according to the methods of Kyte and Doolittle (26) and of Brandl *et al.* (19) shows an outline of scallop ATPase molecule which mainly consists of 10 transmembrane domains (M1-M10), 5 amphipathic  $\alpha$ -helical regions (stalk regions) (S1-S5) continuous to the transmembrane domains, and 2 large cytoplasmic domains (Fig. 2B), as observed with the ATPase (19) from rabbit fast-twitch skeletal muscles. This supports the report of their similar three-dimensional structures determined from their electron micrographies (2, 5). Based on the similarity of the scallop sequence to that of the rabbit (19), Asp350 and Lys514, respectively, in one (amino acid residues 327-727) of the cytoplasmic domains seem to be a phosphorylation site and a fluorescein 5-isothiocyanate binding site which forms a part of the ATP-binding sites. Although the present results of the primary structure do not provide any direct information about the difference (5, 9, 10, 11) of the scallop ATPase from the rabbit ATPase with regard to the relationship between stability of the ATPase activity and the aggregation state of the ATPase molecules, it provides a molecular basis for examining the differences of those ATPases.

The overall amino acid sequence of the scallop  $\text{Ca}^{2+}$ -ATPase exhibited about 70 (70.1-75.7)% similarity to those of the SERCA-type ATPases of *Schistosoma*, *Artemia*, *Drosophila*, rabbit, etc., which belong to Animalia, while it exhibited about 30 (28.8-34.3) and 50 (47.1-48.5)% similarity to those of yeasts, and *Trypanosoma* and *Plasmodium*, which belong to Fungi and Protista, respectively (Fig. 3). Sequence




Fig. 1. **Isolation of lysyl endopeptidase peptides of scallop  $\text{Ca}^{2+}$ -ATPase.** (A) Sodium dodecyl sulfate-polyacrylamide gel electrophoresis profile of the SR (a) and the purified ATPase (b). The protein band indicated by an arrow was shown in a previous paper (9) to have almost the same molecular weight as that (100kDa) of rabbit SR  $\text{Ca}^{2+}$ -ATPase of the fast-twitch muscle. The protein was also shown to be  $\text{Ca}^{2+}$ -ATPase protein based on the observations of calcium-dependent phosphorylation of the protein with ATP, hydroxylamine-sensitivity of the phosphoprotein and time-dependent liberation of the phosphate incorporated into the protein (9). (B) First reverse-phase chromatography of the lysyl endopeptidase digest of scallop ATPase. Twelve peptides, L1-L12, were sequenced (see "Materials and Methods" for details).



Figure2-A (Continued)

1740 1770 1800  
CGCCGCGAGGACATGGACCTGGAAGATTCCCGCAAGTTCATCCAGTATGAGACCAATATGACATTCGTAGGAGTCGTAGGAATGTTGGAC  
R R E D M D L E D S R K F I Q Y E T N M T F V G V V G M L D

1830 1860 1890  
CCCCACGTATGGAAGTGTTCGACTCCATCAAGAACTGCCGCAAGGCTGGAATCCGTGTCAATTGTCATCACCGGCGACAACAAGGCCACA  
P P R M E V F D S I K N C R K A G I R V I V I T G D N K A T  
L7 L8

1920 1950 1980  
GCTGAGGCTATCTGTCTAGGATCGGTGTGTTCCGGCGAAAACGAGTCTACAGAAGGCATGTCTTCACTGGCCGCGAGTTTGTGACCTG  
A E A I C R R I G V F G E N E S T E G M S F T G R E F D D L

2010 2040 2070  
TCACATGAAGAGCAGCGATTGGCGGTAAACAAAGAGTCGTCTGTTTCCCGTGTAGAGCCTGCTCACAAGAGTAAGATCGTAGAGTACCTG  
S H E E Q R L A V T K S R L F A R V E P A H K S K I V E Y L  
L9

2100 2130 2160  
CAAGGAGAAGGAGAAATCTCGGCCATGACAGGAGATGGAGTAAACGACGCCCTGCTCTGAAGAAAGCCGAAATTGGTATCGCCATGGGA  
Q G E G E I S A M T G D G V N D A P A L K K A E I G I A M G  
L10

2190 2220 2250  
TCTGGAAGTGTGTGGCCAAGTCTGCCTCTGAGATGGTGTGGCTGATGACAACCTCGCAACAATTGTGTCCGCTGTGGAAGAGGGCCCGC  
S G T A V A K | S A S E M V L A D D N F A T I V S A V E E G R

2280 2310 2340  
GCATCTACAACAACATGAAACAATTTCATCAGATATCTCATCTCCTCCAACATTTGGAGAGGTCTGATGTATCTTCTTACTGCTGCTCTT  
A I Y N N M K Q F I R Y L I S S N I G E V V C I F L T A A L  
S5 M5

2370 2400 2430  
GGCATCCCTGAGGCTCTCATCCCCGTACAGCTTCTGTGGGTGAACCTGGTAACTGATGGTCTTCCAGCCACTGCCCTCGGCTTCAACCCC  
G I P E A L I P V Q L L W V N L V T D G L P A T A L G F N P  
M6

2460 2490 2520  
CCAGATATGGACATCATGAAGAACTCCAGGAATGCAAAGGAAGTCTCATCACGGCTGGTGTCTTTCAGATATATGGCTATGGGA  
P D M D I M K K P P R N A K E G L I T G W L F F R Y M A I G  
M7

2550 2580 2610  
GGCTACGTTGGATGTGCGACTGTTGGTGTGCTGCCGCTGGTGGTTCATGGTCTATGACAAAGGGCCCAACTCAACTACTACCAGCTGACC  
G Y V G C A T V G A A A W W F M V Y D K G P Q L N Y Y Q L T

2640 2670 2700  
CATCACTCACAGTGTCTGGCTCAAGATGAGCGTTTCTCGGTGTGACTGCAAGGTTTTTGACCACCCTGCACCTATGACAATGGCTCTG  
H H S Q C L A Q D E R F L G V D C K V F D H P A P M T M A L  
M8

2730 2760 2790  
TCCGCTCGGTCGTCAATGAAATGCTTAACGCTCTCAACAGCTTGTCTGAGAACCAGTCCCTGCTGGTGTATGCCCCCTTGGTGTAAACAAG  
S V L V V I E M L N A L N S L S E N Q S L L V M P P W C N K  
M9

2820 2850 2880  
TGGCTGTAGGAGCCATGGCTCTGTCTATGGGACTCCACTTTTGCATCCTCTACATTGACGTCATGTCGACCATCTTCAAATCAGCCCT  
W L L G A M A L S M G L H F C I L Y I D V M S T I F Q I T P  
M10

2910 2940 2970  
CTTGGATTGAAGAATGGTTCCGCGTACTCAAGATCTCAATGCCTGTGATACTTATAGACGAAACATTGAAGTTCTGTGCTCGCAAATTC  
L G F E E W F A V L K I S M P V I L I D E T L K F C A R K F  
L12

3000 3030 3060  
ACTGACGCGTAAATATTTGACACATTATACAAGACACAGACAAGGACAAGTTGTGAGGTCCAGTGAAGAAGGCAGCCATTTTCGTACAAC  
T D A \*

3090 3120 3150  
CTGCTGGACTCGAGTCACACAAGAAACATCGGTAAGGGAAGGACGCTGGTGTATCCCGACTCGTGTGAATTAGTGATTTTAAACCAATCAGT

3180 3210 3240  
GTGGATGCTATAGGACCATTCGGCTTCTCAGTGAATAACAACAGTTGACGATGATTGCCTTATATGGTGTGTACTAGTTTCCTT

3270 3300 3330  
CAACCTGAACTACCAGAGTGGATGTTATATTTAGATAGAAGCTTTTCAGCGTCTACCAACCCAGGAAGCACATAAGAAGCTCATGTGTG

3360 3390 3420  
ATTGTCCCTCACTTTGCTCCAAATTTTACTCCTTACCCAAAATTTCTCCAGGAATGAGGAAATTCAAAATAGGTGAATTATCAATCTTCAA


3450 3480 3510  
AACTACTCCGTTTAGCAAAAATGATTGTAACCCCGCGGAAGTGTGGTACCGGAAACGCTTCCCTCGTGTGACGTATTGTACCAGATACCCG

3540 3570 3600  
GACTATTTTACGTGTATATTTATGCAAATGTAAGCTAATTTATGACAAGTCATCGAATTTATTTATGTGACGTGGTGA AAAACGTATAATT

3630 3660 3690  
GTGGCATTTTCTAAAGAATATTAGTTCATGAATATTGTAGATATATATTTCTGTGATTGGACTCTTATGCACCTCCTGACCTTTGACCT

CCTTCAATGATAAACAGGGGG


Figure2-B (Continued)



**Fig. 2. Predicted primary and secondary structures of the scallop  $\text{Ca}^{2+}$ -ATPase.** (A) Nucleotide and deduced amino-acid sequence of the  $\text{Ca}^{2+}$ -ATPase. Nucleotides are numbered with '1' being A of the ATG starting codon. The translation stop codon is denoted by an asterisk. Amino acids are numbered below the nucleotide sequence. The amino acid sequences which were used to design the primers for RT-PCR are marked in broken lines. The sequences (L1-L12) of the lysyl endopeptidase peptides of the ATPase are boxed. Solid lines and dotted lines represent the  $\alpha$ -helical stalk regions (S1-S5) and the transmembrane segments (M1-M10), respectively. (B) Scheme of the predicted ATPase topography with 10 transmembrane domains (M1-M10), 5 stalk regions (S1-S5) and two large cytoplasmic domains. The numbers above and below the membrane indicate position of the amino acid residues at each end of the transmembrane helices. Asp<sup>350</sup> and Lys<sup>514</sup> are a phosphorylation site and a fluorescein 5-isothiocyanate binding site, respectively.

similarity between the two yeasts, which belong to the same phylum of Ascomycota, was about 40%, while the similarity between *Trypanosoma* and *Plasmodium*, which belong to the different phyla of Mastigophora and Apicomplexa, was about 30%. These results suggest the existence of differences in the phylogenetic classification among the ATPases of these kingdoms and in the diversity of the ATPases within each kingdom. As to the ATPases in Animalia, the following relations among them could be recognized: The ATPases in Chordata are in three branches of SERCA1, 2 and 3 in a group, as shown by Nolan *et al.*, (27) and Song and Fambrough (28). The ATPases of scallop, *Artemia* and *Drosophila* are in two branches of Mollusca and Arthropoda in a group. *Schistosoma* is in a branch of common origin for these Animalia ATPases. It is interesting to know an evolutionary relation of amino acid sequences between the SERCA-type ATPases and the plasma membrane type  $\text{Ca}^{2+}$ -ATPases (PMCA) which exhibit the differences in calmodulin-regulation of the ATPase activity (29) and in a ratio of calcium transported to ATP hydrolyzed (30,31). At present time, however, it is difficult to discuss such relation, because that the variety of the species, of which PMCA-type ATPases have been sequenced, is poor (4 for Chordata and one for Fungi).

The nucleotide sequence reported in this paper has been submitted to the DDBJ Data Bank with accession number AB002112.



**Fig. 3. Schematic representation of the relationship of overall amino acid sequences between scallop ATPase and other SERCA-type ATPases.** The amino acid sequence of scallop ATPase, which was deduced from the nucleotide sequence of its cDNA, was analyzed for similarity to those from the following organisms, by using the program GENE-TYX-Mace with UPGMA (unweighted pair-group method with arithmetic mean) and arranged in the form of a ladder to illustrate common origins: fission yeast (Ref.32), budding yeast (Ref.33), *Trypanosoma brucei* (Ref.34), *Plasmodium falciparum* (Ref.35), *Schistosoma mansoni* (Ref.13), *Crustacean Artemia* (Ref. 14), *Drosophila melanogaster* (Ref. 15), frog -1 (Ref. 36), chicken -1 (Ref. 37), rabbit -1 (Ref.19), cat -2 (Ref. 38) human -2 (Ref. 39), rat -2 (Ref.40), rat -3 (Ref.41) and Human-3 (Ref.42). The appended numbers, 1, 2 and 3, represent the fast-twitch skeletal muscle cell form (SERCA1), the slow twitch skeletal/cardiac form (SERCA2), and the non-muscle form (SERCA3), respectively.



## REFERENCES

- 1) Tada, M., Yamamoto, T., Tonomura, Y. (1978) Molecular mechanism of active calcium transport by sarcoplasmic reticulum. *Physiol. Rev.* 56, 1-79
- 2) Taylor, K.A., Dux, L., Martonosi, A. (1986) Three-dimensional reconstitution of negatively stained crystals of the  $\text{Ca}^{2+}$ -ATPase from muscle sarcoplasmic reticulum. *J. Mol. Biol.* 187, 417-427
- 3) Carafoli, E. (1987) Intercellular calcium homeostasis. *Annu. Rev. Biochem.* 56, 395-433
- 4) Inesi, G., Sumbilla, C., Kirtley, M.E. (1990) Relationships of molecular structure and function in  $\text{Ca}^{2+}$ -transport ATPase. *Physiol. Rev.* 70, 749-760
- 5) Castellani, L., Hardwicke, P.M.D., Vibert, P. (1985) Dimer ribbons in the three-dimensional structure of sarcoplasmic reticulum. *J. Mol. Biol.* 185, 579-594
- 6) Kalabokis, V., Hardwicke, P. (1988) Variation of scallop sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase activity with temperature. *J. Biol. Chem.* 263, 15184-15188
- 7) Castellani, L., Hardwicke, P.M.D., Franzini-Armstrong, C. (1989) Effect of  $\text{Ca}^{2+}$  on the dimeric structure of scallop sarcoplasmic reticulum. *J. Cell Biol.* 108, 551-520
- 8) Kalabokis, V.N., Bozzola, J.J., Castellani, L., Hardwicke, P.M.D. (1991) A possible role for the dimer ribbon state of scallop sarcoplasmic reticulum: Dimer ribbons are associated with stabilization of the  $\text{Ca}^{2+}$ -free Ca-ATPase. *J. Biol. Chem.* 266, 22044-22050
- 9) Abe, M., Shirakata, Y., Sato, D., Konishi, K., Watanabe, T., Nakamura, J. (1992) Isolation and characteristics of scallop sarcoplasmic reticulum with calcium transport activity. *J. Biochem.* 112, 822- 827
- 10) Kalabokis, V.N., Santoro, M.M., Hardwicke, P.M.D. (1993) Effect of  $\text{Na}^+$  and nucleotide on the stability of solubilized  $\text{Ca}^{2+}$ - free Ca-ATPase from sarcoplasmic reticulum. *Biochemistry* 32, 4389-4396
- 11) Nagata, Y., Nakamura, J., Yamamoto, T. (1996) Protection of scallop sarcoplasmic reticulum ATPase from thermal inactivation by removal of calcium from high-affinity binding sites on the enzyme. *J. Biochem.* 119, 1100-1105
- 12) Nagata, Y., Nakamura, J., Yamamoto, T. (1997) Temperature sensitivity of proteoliposomes reconstituted from a mixture of scallop and rabbit sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPases. *J. Biochem.* 121: 648-653

- 13) de Mendonca, R.L., Beck, E., Rumjanek, F.D., Goffeau, A. (1995) Cloning and characterization of a putative calcium-transporting ATPase gene from *Schistosoma mansoni*. *Mol. Biochem. Parasitol.* 72, 129-139
- 14) Palmero, I., Sastre, L. (1989) Complementary DNA cloning of a protein highly homologous to mammalian sarcoplasmic reticulum Ca-ATPase from the *Crustacean artemia*. *J. Mol. Biol.* 210, 737-748
- 15) Magyar, A., Varadi, A. (1990) Molecular cloning and chromosomal location of a sarco/endoplasmic reticulum Ca<sup>2+</sup>-ATPase of *Drosophila melanogaster*. *Biochem. Biophys. Res. Commun.* 173, 872-877
- 16) Nakamura, J. (1983) The ADP- and Mg<sup>2+</sup>-reactive calcium complex of the phosphoenzyme in skeletal sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase. *Biochim. Biophys. Acta* 723, 182-190
- 17) Suzuki, T., Furukohri, T., Gotoh, T. (1985) Subunit structure of extracellular hemoglobin from the polychaete *Tylorrhynchus heterochaetus* and amino acid sequence of the constituent polypeptide chain (IIC). *J. Biol. Chem.* 260, 3145-3154
- 18) Suzuki, T., J. (1986) Amino acid sequence of myoglobin from the mollusc *Dolabella auricularia*. *Biol. Chem.* 261, 3692-3699
- 19) Brandl, C. J., Green, N. M., Korczak, B., MacLennan, D.H. (1986) Two Ca<sup>2+</sup> ATPase genes, Homologies and mechanistic implications of deduced amino acid sequences. *Cell* 44, 597-607
- 20) Sambrook, J., Fritsch, E.F., Maniatis, T. (1989) *Molecular Cloning. In, A Laboratory Manual*, second edition. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press
- 21) Branbl, C.J., deLeon, S., Martin, D.R., MacLennan, D.H. (1987) Adult forms of the Ca<sup>2+</sup>-ATPase of sarcoplasmic reticulum: Expression in developing skeletal muscle. *J. Biol. Chem.* 262, 3768-3774
- 22) Lytton, J., MacLennan, D. H. (1988) Molecular cloning of cDNAs from human kidney coding for two alternatively spliced products of the cardiac Ca<sup>2+</sup>-ATPase gene. *J. Biol. Chem.* 263, 15024-15031
- 23) Genteski-Hamblin, A.M., Greeb, J., Shull, G.E. (1988) A novel Ca<sup>2+</sup> pump expressed in brain, kidney, and stomach is encoded by an alternative transcript of the slow-twitch muscle sarcoplasmic reticulum Ca-ATPase gene: Identification of cDNAs encoding Ca<sup>2+</sup> and other cation-transporting ATPase using an

- oligonucleotide probe derived from the ATP-binding site. *J. Biol. Chem.* 263, 15032-15040
- 24) Campbell, A.M., Kessler, P.D., Sagara, Y., Inesi, G., Fambrough, D.M. (1991) Nucleotide sequences of avian cardiac and brain SR/ER Ca<sup>2+</sup>-ATPases and functional comparisons with fast twitch Ca<sup>2+</sup>-ATPase: Calcium affinities and inhibitor effects. *J. Biol. Chem.* 266, 16050-16055
- 25) Escalante, R., Sastre, L. (1993) Similar alternative splicing events generate two sarcoplasmic or endoplasmic reticulum Ca-ATPase isoforms in the Crustacean *Artemia franciscana* and in vertebrates. *J. Biol. Chem.* 268, 14090-14095
- 26) Kyte, J., Doolittle, R.F. (1982) A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* 157, 105-132
- 27) Nolan, D.P., Revelard, P., Pays, E. (1994) Overexpression and characterization of a gene for a Ca<sup>2+</sup>-ATPase of the endoplasmic reticulum in *Typososoma brucei*. *J. Biol. Chem.* 269, 26045-26051
- 28) Song, I, Fambrough, D. (1994) Molecular evolution of the calcium-transporting ATPases analyzed by the maximum parsimony method. In: Fambrough D.M., editor. *Molecular evolution of physiological processes*. The Rockefeller University Press Press, 271-283
- 29) Carafoli, E. (1992) The Ca<sup>2+</sup> pump of the plasma membrane. *J. Biol. Chem.* 267, 2115-2118
- 30) Niggli, J, Adunyah, E.S., Penniston, J. T., Carafoli, E. (1981) Purified (Ca<sup>2+</sup>- Mg<sup>2+</sup>)-ATPase of the erythrocyte membrane: Reconstitution and effect of calmodulin and phospholipids. *J. Biol. Chem.* 256, 395-401
- 31) Hao, L., Rigaud, J.L., Inesi, G. (1994) Ca<sup>2+</sup>/H<sup>+</sup> countertransport and electrogenicity in proteoliposomes containing erythrocyte plasma membrane Ca-ATPase and exogenous lipids. *J. Biol. Chem.* 269, 14268-14275
- 32) Ghislain, M., Goffeau, A., Halachmi, D., Eilam, Y. (1990) Calcium homeostasis and transport are affected by disruption of *cta3*, a novel gene encoding Ca<sup>2+</sup>-ATPase in *Schizosaccharomyces pombe*. *J. Biol. chem.* 265, 18400-18407
- 33) Rudolph, H.K., Antebi, A., Fink, G.R., Buckley, C.M., Dorman, T.E., Levitre, J., Davidow, L.S., Mao, J.I., Moir, D.T. (1989) The yeast secretory pathway is perturbed by mutations in *PMRI*, a member of a Ca<sup>2+</sup> ATPase family. *Cell* 58, 133-145
- 34) Revelard, P., Pays, E. (1991) Structure and transcription of a P-ATPase gene from

*Typanosoma brucei*. Mol. Biochem. Parasitol. 46, 241-252

- 35) Kimura, M., Yamaguchi, Y., Takada, S., Tanabe, K. (1993) Cloning of a  $\text{Ca}^{2+}$ -ATPase gene of *Plasmodium falciparum* and comparison with vertebrate  $\text{Ca}^{2+}$ -ATPases. J. Cell. Sci. 104, 1129-1136
- 36) Vilsen, B., Andersen, J.P. (1992) Deduced amino acid sequence and  $E_1$ - $E_2$  equilibrium of the sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase of frog skeletal muscle: Comparison with the  $\text{Ca}^{2+}$ -ATPase of rabbit twitch muscle. FEBS Lett. 306, 213-218
- 37) Karin, N.J., Kaprielian, Z., Fambrough, D.M. (1989) Expression of avian  $\text{Ca}^{2+}$ -ATPase in cultured mouse myogenic cells. Mol. Cell. Biol. 9, 1978-1986
- 38) Gambel, A.M., Gallien, T.N., Dantzler-Whitworth, T., Bowes, M., Menick, D.R. (1992) Sequence of the feline cardiac sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase. Biochim. Biophys. Acta 1131, 203-206
- 39) Lytton, J., MacLennan, D.H., (1988) Molecular cloning of cDNAs from human kidney coding for two alternatively spliced products of the cardiac  $\text{Ca}^{2+}$ -ATPase. J. Biol. Chem. 263, 15024-15031
- 40) Guntjeski-Hamblin, A.M., Greeb, J., Shull, G.E. (1988) A novel  $\text{Ca}^{2+}$  pump expressed in brain, kidney, and stomach is encoded by an alternative transcript of the slow-twitch muscle sarcoplasmic reticulum Ca-ATPase gene: Identification of cDNAs encoding  $\text{Ca}^{2+}$  and other cation-transporting ATPases using an oligonucleotide probe derived from the ATP-binding site. J. Biol. Chem. 263, 15032-15040
- 41) Burk, S.E., Lytton, J., MacLennan, D.H., Shull, G.E. (1989) cDNA cloning, functional expression, and mRNA tissue distribution of a third organellar  $\text{Ca}^{2+}$  pump. J. Biol. Chem. 264, 18561-18568
- 42) Dole, L., Wuytack, F., Kools, P.F.J., Baba-Aissa, F., Raeymaekers, L., Brik, F., Van De Ven, W.J.N., Casteels, R. (1996) cDNA cloning, expression and chromosomal location of the human sarco/endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase 3 genes. Biochem. J. 318, 689-699

### Part III

**Change in the thermal sensitivity of sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase by chemical modifications and mutagenesis of specific Cys residues**

## SUMMARY

In the proceeding study I have cloned cDNA of the sarcoplasmic reticulum (SR)  $\text{Ca}^{2+}$ -ATPase (SERCA) from scallop adductor muscle, and predicted entire amino acid sequence of this enzyme (1). In comparison of the amino acid sequence with that of rabbit SR  $\text{Ca}^{2+}$ -ATPases, I have further explored the role of Cys residues on thermal stability of this enzyme. When rabbit SR was incubated above  $36^{\circ}\text{C}$  in the presence of 6.4 mM glutathione,  $\text{Ca}^{2+}$ -transport activity was lost within 5 min, while in the absence of glutathione, full activity was retained up to  $45^{\circ}\text{C}$ . This is in contrast that  $\text{Ca}^{2+}$  pump activity of scallop SR was lost above  $36^{\circ}\text{C}$  irrespective of the presence or absence of . These results raise a possibility that the pronounced difference in thermal stability between rabbit and scallop SR is caused by the formation of a disulfide bond among Cys residues on their  $\text{Ca}^{2+}$ -ATPase.

I have attempted to identify the Cys residues linking through a disulfide bond by examining accessibility of SH-groups to N-(7-Dimethylamino-4-methyl-3-coumarinyl)-maleimide (DACM). Sequence analysis of the DACM-labelled peptides isolated from rabbit  $\text{Ca}^{2+}$ -ATPase revealed that almost all of the cytoplasmically exposed Cys residues were readily accessible to DACM except for Cys12, Cys377, Cys349, Cys674, and Cys675. By contrast Cys349 in the Scallop  $\text{Ca}^{2+}$ -ATPase was found to be easily modified by DACM.

Site-directed mutagenesis was performed on Cys12, Cys377, Cys674, and Cys675 in the rabbit  $\text{Ca}^{2+}$ -ATPase to respective amino acids at the same position of sequence in the scallop  $\text{Ca}^{2+}$ -ATPase and the effects of mutations are compared for heat sensitivity of the mutants expressed in Sf21 cells. It was demonstrated that the activity of phosphoenzyme (EP) formation was pronouncedly inhibited when Cys675 $\rightarrow$ Leu mutant was incubated at  $41^{\circ}\text{C}$  for 10 min. While other mutants, Cys12 $\rightarrow$ Val, C377 $\rightarrow$ Ile, C674 $\rightarrow$ Arg, as well as the wild type of rabbit  $\text{Ca}^{2+}$ -ATPase, were virtually unaffected by the heat treatment. Single substitution of Cys349 of the rabbit  $\text{Ca}^{2+}$ -ATPase with Ala resulted in destabilization of the enzyme by the thermal treatment. These results suggest that on the rabbit  $\text{Ca}^{2+}$ -ATPase, Cys675 may bind to Cys349 through disulfid bond which can contribute to the thermal stabilization of the enzyme. while that lacking in the disulfide bond in the scallop  $\text{Ca}^{2+}$ -ATPase may cause the thermal destabilization of the enzyme.

## INTRODUCTION

The active transport of  $\text{Ca}^{2+}$  into SR is mediated by an enzyme  $\text{Ca}^{2+}$ -ATPase which catalyzes the hydrolysis of ATP. In the reaction cycle,  $\gamma$ -phosphate from ATP is transferred to a specific Asp residue to form a phosphorylated intermediate (EP). Its formation and decomposition strictly couple to the  $\text{Ca}^{2+}$  transport across the membrane (2, 3). Although considerable information has been obtained regarding kinetic function of the enzyme, the structural basis for coupling of ATP hydrolysis to  $\text{Ca}^{2+}$  translocation across the membrane remains unsolved as major issues. Recent protein chemical and molecular biology techniques have been providing important information about structural features of the  $\text{Ca}^{2+}$ -ATPase.

In the proceeding paper I have studied the effects of temperature on the catalytic functions of the scallop SR (1, 4-6). The specific activity of the scallop  $\text{Ca}^{2+}$ -ATPase was lost after several minutes incubation of SR at temperatures as high as  $39^{\circ}\text{C}$ . whereas the activity of the rabbit  $\text{Ca}^{2+}$ -ATPase decreased only marginally with almost 80% of the activity retained even after the incubation above  $50^{\circ}\text{C}$ . In order to provide answers to the question what structural difference between rabbit and scallop  $\text{Ca}^{2+}$ -ATPases is related to the pronounced difference in the temperature sensitivity, we have cloned cDNA of the scallop  $\text{Ca}^{2+}$ -ATPase and predicted its primary structure (1). The detailed comparison of the amino acid sequence with that of rabbit SR  $\text{Ca}^{2+}$ -ATPases revealed interesting differences in the distribution of Cys residues on the cytoplasmically exposed region of these enzymes.

In the present study, I have identified and characterized the Cys residues on the rabbit  $\text{Ca}^{2+}$ -ATPase which could contribute to the structural stabilization of the enzyme. It was demonstrated that mutation of Cys675 to Leu or Cys349 to Ala led to expression of  $\text{Ca}^{2+}$ -pump activity which was sharply decreased by incubation at temperatures similar to those required for the inactivation of scallop  $\text{Ca}^{2+}$ -ATPase.

## EXPERIMENTAL PROCEDURES

**Materials** --- Scallop SR was isolated from the striated portion of scallop (*Patinopecten yessoensis*) adductor muscle according to the method described previously (4). Rabbit SR was isolated from the skeletal muscle of rabbit dorsal and hind leg as described previously (5). These SR preparations were divided into small pieces, quickly frozen using liquid nitrogen, and stored at  $-80^{\circ}\text{C}$ .  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  was purchased from Amersham, Antipyrilazo III was from Nacalai, glutathione and DACM were from Wako. Other Chemicals were analytical grade.

**Modification of  $\text{Ca}^{2+}$ -ATPase with DACM** --- SR vesicles at 2-2.5 mg/ml protein were exposed to 0.2-0.3 mM DACM for 30 min at  $23^{\circ}\text{C}$  in the standard medium which contained 0.1 M NaCl, 20 mM TES (pH7.2), 5 mM  $\text{MgCl}_2$ , 5% glycerol. At the end of the incubation period, the mixture was taken into 25 ml of the standard solution. SR was washed twice by centrifugation in the same solution. The amount of probe incorporated was determined spectrophotometrically using molar extinction coefficients of 19,800 at 398 nm (7).

**Purification of DACM labeled peptides** --- The  $\text{Ca}^{2+}$ -ATPase was purified according to Meissner *et al.* (8) by solubilizing SR with DOC to remove extrinsic proteins. The purified  $\text{Ca}^{2+}$ -ATPase at 10 mg/ml protein was allowed to react with 0.1 mg/ml thermolysin at  $52^{\circ}\text{C}$  for 90 min as described previously (8). Separation and purification of DACM labeled peptides from the thermolysin digest was carried out by using HPLC with ODS-120T reversed-phase column essentially as described by our previous work (9). The final purified DACM-labeled peptides were subjected to Edman degradation on a model LF3600 protein sequencer (Applied BioSystems), and the amino acid sequences were determined.

**Measurements of Activities** ---  $\text{Ca}^{2+}$ -transport by SR was measured in a reaction medium containing 0.025 -0.05 mg/ml SR protein, 0.1 M KCl, 5 mM  $\text{MgCl}_2$ , 5 mM oxalate, 5% glycerol, 50  $\mu\text{M}$   $\text{CaCl}_2$ , 0.2 mM Antipyrilazo III, and 20 mM TES at pH 7.2. The reaction was started at  $23^{\circ}\text{C}$  by addition of 100  $\mu\text{M}$  ATP.  $\text{Ca}^{2+}$  uptake into SR was determined by monitoring the absorption at 700 nm in the reaction mixture (6). Phosphorylation of SR was measured in 20  $\mu\text{l}$  of reaction mixture which contained 0.5 mg/ml protein of microsome membrane, 0.1 M KCl, 5 mM  $\text{MgCl}_2$ , 5% glycerol, and 50 mM TES at pH 7.2. The reaction was initiated at  $23^{\circ}\text{C}$  by the addition of 1 mM  $[\gamma\text{-}$




$^{32}\text{P}$ ]ATP and quenched after 10 sec with an ice cold solution containing 5% TCA, 10 mM ATP and 0.1 M Pi. The amount of  $[\gamma\text{-}^{32}\text{P}]$ phosphoprotein formed was determined as described elsewhere (10).

**Construction and expression of mutant cDNAs** --- Full-length rabbit SERCA1a gene and monoclonal antibody A52 were provided from MacLennan, D.H. Full-length scallop  $\text{Ca}^{2+}$ -ATPase gene (Scal-SERCA) was described previously (1). Each genes were subcloned into pBluescript SK+. Site-directed mutagenesis was used to insert a Spe I site at the initiator methionine codon of SERCA1a and a Hin dIII site immediately after the stop codon. In the same way, a Bam HI site was inserted into the scallop  $\text{Ca}^{2+}$ -ATPase gene (Scal-SERCA) before first ATG. Mutant genes of SERCA1a and Scal-SERCA were constructed used PCR with LA-Taq (TAKARA). Mutants and mutagenic primers are as follows; C12V; CACAGAGGAAGTTCTGGCCTATTTTGG. C377I; GACGGAGACTTCATTTGCTGAACGAG, C674R; CTGCCGCCGCGCCC-GCTGCTTCGCGCG, C675L; CGCCGCGCCTGCCTCTTCGCGCGCGTG. The PCR reaction was carried under the standard condition suggested by TAKARA. After PCR reaction, template DNA was digested by restriction enzyme Dpn I which digests only methylated DNA and then PCR products were separated by agarose gel electrophoresis and purified by Prep-A-Gene DNA purification systems (BIO-RAD). Then self-ligated mutant cDNAs were transfected in E.coli DH5a competent cells. The presence of the desired mutations and the absence of unexpected ones were verified by DNA sequencing. 4 desired mutation, 1683C  $\rightarrow$  T, 2202C  $\rightarrow$  G, 2253G  $\rightarrow$  C and 2367C  $\rightarrow$  T, were available in SERCA1a gene used in this study, but these mutations do not change the native amino acid.


Recombinant baculoviruses which contain wild type or mutant genes of  $\text{Ca}^{2+}$ -ATPase were constructed by Bac-to-Bac baculovirus expression systems (GIBCO). Preparation of ER fraction from Sf21 cells was as follows; 60 hour passed from virus infection, Sf21 cells were harvested and homogenized by polytron homogenizer after washing with PBS twice. Microsomal fraction was purified by step-wise centrifugation by the method described previously (5) and finally suspended in 0.1 M KCl, 5 mM Tris-maleate, and 10% glycerol and stocked at  $-80^{\circ}\text{C}$ . An anti-peptidase was not used through preparation.

## RESULTS

We have demonstrated a striking difference in temperature sensitivity between the catalytic functions of rabbit and scallop SR (1, 4-6). As illustrated in Fig. 1, incubation of scallop SR for 5 min at temperature at 38°C suffices to produce complete inactivation of ATP hydrolysis, whereas rabbit SR ATPase is not inhibited by the incubation at temperatures up to at least 50°C. The difference can be attributed to a difference in the amino acid sequences of these enzymes. In the proceeding study I have cloned cDNA of SR Ca<sup>2+</sup>-ATPase from scallop adductor muscle, and predicted entire amino acid sequence of this enzyme (1). Based on the detailed comparison between the entire amino acid sequences of the rabbit and scallop Ca<sup>2+</sup>-ATPases, it was revealed that there are several candidate amino acids which could contribute to the stabilization of the enzyme structure. One of the most striking difference relates to the distribution of Cys residues on the cytoplasmically exposed domain in the Ca<sup>2+</sup>-ATPase molecule. In the experiment shown in Fig. 2, I studied the effects of glutathione on the rabbit and scallop Ca<sup>2+</sup>-ATPases, as expressed by Ca<sup>2+</sup>-pump activity. When rabbit SR was incubated above 36°C in the presence of 6.4 mM glutathione, Ca<sup>2+</sup>-transport activity was lost within 5 min, while in the absence of glutathione, full activity was retained up to 50°C (Fig. 2A, open and closed circles). Fig. 2B shows the time courses of decrease in Ca<sup>2+</sup>-pump activity of rabbit SR by the incubation at 43°C in the presence of different concentration of glutathione. The half time of inactivation is 10 min for 3.2 mM, 5 min for 6.4 mM and 30 sec for 10 mM. Ca<sup>2+</sup>-pump activity of rabbit SR was decreased in the presence of DTT at high temperature in fashion similar to that observed with glutathione, although half concentration of DTT for heat inactivation is much higher than that for glutathione (Data not shown). These results suggest that disulfide bonds of the Ca<sup>2+</sup>-ATPase are involved in thermal stabilization of the structure which is essential to the enzymatic function. On the other hand, Ca<sup>2+</sup>-pump activity of scallop SR was lost after 5 min incubation above 35°C irrespective of the presence or absence of glutathione (Fig. 2A, open and closed triangles). It may be concluded from these observations that the pronounced difference in thermal stability between rabbit and scallop SR is presumably caused by formation of a disulfide bond among Cys residues on the enzyme. This conclusion seems to be in accord with experimental observations demonstrated in Fig. 3. We compared between reactivities of SH groups with DTNB in the rabbit and



**Fig. 1. Temperature sensitivities of the  $\text{Ca}^{2+}$ -ATPase activities in rabbit and scallop SR.** 2 mg/ml protein of rabbit (○) or scallop (□) SR was incubated at indicated temperatures in the medium containing 0.1 mM  $\text{CaCl}_2$ , 0.1 M KCl, 20 mM TES (pH 7.2), 10% glycerol, 5 mM  $\text{MgCl}_2$ . After 5 min incubation, 0.1 ml of SR suspension was transferred into 20 volumes of assay medium which had been kept cool on ice to stop the thermal treatment. ATP hydrolysis was measured in a medium which contained 0.1 mg/ml SR, 40  $\mu\text{M}$   $\text{CaCl}_2$ , 2  $\mu\text{M}$  A23187, 0.1 M KCl, 20 mM TES (pH 7.2), 5% glycerol, 5 mM  $\text{MgCl}_2$ , and ATP regenerating system (0.5 mM NADH, 1.5 mM Phosphoenolpyruvate, 1 unit of lactate dehydrogenase, 1 unit of pyruvate kinase). Reaction was initiated by the addition of 0.2 mM ATP at 23°C, and the amount of ADP liberated from ATP was determined by monitoring the decrease in absorbance of NADH at 340 nm.



**Fig. 2. A: Temperature sensitivity of Ca<sup>2+</sup>-transport activity in the presence and absence of glutathion.** 3.5 mg/ml protein of rabbit (O,●) and scallop (Δ,▲) SR were incubated with (O,Δ) or without (●,▲) 6.4 mM GSH at the indicated temperatures in the solution containing 0.1 MKCl, 10 mM Tris-maleate (pH 6.5), and 10% glycerol. After 5 min incubation was terminated by cooling the medium in ice. Ca<sup>2+</sup>-transport activity remaining after the incubation was monitored at 23°C as described in "EXPERIMENTAL PROCEDURE"

**B: Time courses of thermal inactivation of rabbit SR Ca<sup>2+</sup>-transport at various concentrations of GSH.** 3.5 mg/ml protein of rabbit SR was incubated with 0 (□), 4.8 (Δ), and 6.4 (O) mM GSH at 42°C under the conditions described above. Ca<sup>2+</sup>-transport activities remaining after various incubation times were measured as in "EXPERIMENTAL PROCEDURE"

scallop  $\text{Ca}^{2+}$ -ATPase. For the case of rabbit SR, the number of SH group modified with DTNB increased very slowly. It reached a steady level of 9 -10 mol per mol of ATPase after 30 min of incubation, while it increased to about 22 mol immediately after the addition of 0.2% SDS. For the case of scallop SR, the number of reacted SH increased to 12 mol per mol of ATPase within 5 min. It reached a maximum level of as high as 23 mol upon the addition of SDS. Since rabbit and scallop  $\text{Ca}^{2+}$ -ATPases share a very high sequence homology in transmembrane segments but not in cytoplasmically exposed regions (1), candidate Cys residues which contribute to formation of disulfide bond can be presumably located in the latter regions. In an attempt to determine the Cys residues which are linked through disulfide bond, intact rabbit SR vesicles were allowed to react with DACM at a concentration enough to label the accessible SH groups under similar conditions to that used for glutathione or DTNB modification.

For purification of  $\text{Ca}^{2+}$ -ATPase, the SR vesicles were washed by centrifugation and solubilized with 0.5% DOC and processed as described in "EXPERIMENTAL PROCEDURES". After the purification step, about 30% of the original SR protein was recovered as  $\text{Ca}^{2+}$ -ATPase protein. As estimated from the absorption at 398 nm, as high as 11 mol of DACM was incorporated per mole of ATPase. This value is in agreement with observations using DTNB as in Fig.3. The DACM-labeled  $\text{Ca}^{2+}$ -ATPase of rabbit SR was then extensively digested with thermolysine for 90 min at 52°C. More than 75% of the amount of DACM were recovered in the supernatant when thermolysine digest was centrifuged to removed insoluble fractions. These results indicate that SH groups in the extra cellular portion of the  $\text{Ca}^{2+}$ -ATPase were mainly reacted with DACM under the conditions used.

In the experiment shown in Fig. 4, DACM-labeled peptides were applied to an ODS-120T column and eluted with 10-50%  $\text{CH}_3\text{CN}$  gradient. By repeating the chromatography under various  $\text{CH}_3\text{CN}$  concentration gradients in the presence or absence of salt, more than 15 fluorescent peaks have been obtained as homogeneous peptides. These peptide were subjected to Edman degradation on a protein sequencer and the position of Cys residues on the primary structure of  $\text{Ca}^{2+}$ -ATPase was estimated (Table I).

Comparison of these motifs with the primary structure of the rabbit  $\text{Ca}^{2+}$ -ATPase revealed that Cys residues at positions of 344\* 364\* 417\* 420\* 471\* 498\* 525\* 561\* 614\* 636\* and 670 were readily modified with DACM (\* represents common residue to






Fig. 3. Comparison of the DTNB accessibilities between rabbit and scallop SR vesicles. 0.34 mg/ml rabbit SR vesicles ( O , ● ) and 0.19 mg/ml scallop SR vesicles ( Δ , ▲ ) were allowed to react at 23°C with 0.2 mM DTNB in the solution containing 0.1 M NaCl, 50 mM TES (pH 7.0), and 5% glycerol. At the indicated times, 0.5% SDS was added to determine total amount of DTNB reacted with SR. Sulfhydryl content was calculated from the absorbance at 412 nm using molar extinction coefficients of 14,100 for the SR suspension ( ● , ▲ ), and 13,000 for the suspension in the presence of SDS ( O , Δ ). These coefficients were obtained by referring to the calibration curve for the absorption which was obtained with 2-mercaptoethanol of known concentration. In addition, purity of the Ca<sup>2+</sup>-ATPase in rabbit and scallop SR assumed to be 60 and 65%, respectively.



**Fig. 4. Elution profile of thermolytic digest of the rabbit SR Ca<sup>2+</sup>-ATPase on reversed-phase HPLC.** Rabbit SR vesicles were allowed to react with DACM, and the Ca<sup>2+</sup>-ATPase was extracted with DOC and digested with thermolysin as described in "EXPERIMENTAL PROCEDURES". The digest was applied to an ODS-120T column and eluted with a 10-50%CH<sub>3</sub>CN gradient in 0.1% trifluoroacetic acid. The absorption of the eluted peptides (black solid line) was monitored at 214 nm and DACM fluorescence in the fraction (red solid line) was monitored for excitation and emission at 398 and 465nm, respectively. The fluorescent fraction was collected, evaporated and redissolved in 10% CH<sub>3</sub>CN and subjected to further purification step.

[Table-I] Separation and purification of SR-DACM-peptides from rabbit Ca<sup>2+</sup>-ATPase

Peak	DACM (nmol)	Protein (nmol AA)	AA/DACM	Sequence
3b	0.26	1.9	7-8	SKVER--- 471C*
				FVKGA--- 525C*
3c	0.38	2.3	6-7	EQREA--- 670C*
4	3.2	22.0	7-8	KVER----- 471C*
5b	0.8	13.4	17	EQREA--- 670C*
				LSKVI---- 471C*
5d	1.4	4.9	3-5	
5c	1.6	6.7	4-5	VY----- 498C*
5e	2.5	10.0	4-5	VRXLP--- 344C*
6a	1.7	5.2	4-5	
6c	2.7	21.0	7-8	VY----- 498C*
7a	0.66	57.0	86	NA----- 471C*
7b	0.56	3.8	7-8	
8	0.5	23.0	45	TGRDT--- 561C*
9a	4.5	23.0	4-5	LVEL----- 417C*, 420C*
				LSKV----- 471C*
9b-1	2.5	19.0	7-8	EQREA--- 670C*
				AI----- 636C*
				VIKEW--- 561C*
9b-2	1.8	28.5	16	AIAI----- 636C*
				FVKG----- 525C*
10	3.0	11.0	3-4	FVKGAP- 525C*
11a	2.1	15.2	7-8	LTTN----- 364C*
11b	1.2	9.7	8-9	VMGS----- 614C*

\*represents common residue to both Ca<sup>2+</sup>-ATPases of rabbit and scallop SR



both Ca<sup>2+</sup>-ATPases), while Cys residues at positions of 12, 349\*, 377, 674, and 675 seemed to be inaccessible to the reagent irrespective of their existence on the cytoplasmically exposed regions of the enzyme (Fig. 5). Therefore the five inaccessible Cys residues are considered to be candidate amino acids which could contribute to formation of disulfide bonds. Accessibility of Cys residues on the scallop Ca<sup>2+</sup>-ATPase were also tested under similar conditions to those for rabbit Ca<sup>2+</sup>-ATPase, and it was found that Cys349 was readily reacted to DACM. This is in contrary to the result obtained with rabbit Ca<sup>2+</sup>-ATPase.

In order to determine which Cys residues are cross linked through disulfide bond, site-directed mutagenesis was performed on the Cys residues at positions of 12, 377, 674, or 675 in rabbit Ca<sup>2+</sup>-ATPase to the respective amino acid, Val, Ile, Arg, or Leu each of which locates at the same position of sequence in scallop one, and Cys349, which is common to that of scallop Ca<sup>2+</sup>-ATPase, was substituted with Ala. Recombinant baculovirus that contain wild type or mutant genes of Ca<sup>2+</sup>-ATPase were transfected to Sf21 cells. To avoid thermal inactivation of expressed Ca<sup>2+</sup>-ATPase, the infected cells were cultured below 27°C. After 60 hours, cells were harvested and the microsomal fraction was obtained by differential centrifugation of homogenized cell as described in "EXPERIMENTAL PROCEDURES".

Figure 6 shows SDS-PAGE and immunoblotting of the microsomes isolated from the mutant cells which were infected by wild type, C12V, C377I, C674R, and C675L baculovirus. All of these mutant ATPases are expressed in the Sf21 cells as well as that of wild type. Functional consequences of the site-directed mutation were summarized in Table II. The activity of EP formation in the presence of Ca<sup>2+</sup> significantly varied from one preparation to another presumably depending upon the amount of SERCA1 expressed in the cell. In the control experiment using no-transfected cell the phosphorylation activity was negligibly low.

In the experiments in Table II, thermostabilities of the mutants were examined by monitoring the phosphorylation activity remaining after 5 min incubation at 43°C. It was found that single substitution of Cys residue at the position of 675 in the rabbit Ca<sup>2+</sup>-ATPase with Leu residue which locates at the same position as that of the scallop ATPase resulted in reduction of the phosphorylating activity to about 30% of original one, while other mutants as well as wild type ATPase maintained the activities at almost the same level after the incubation. As illustrated in Fig. 7, incubation of C675L mutant




Fig. 5. Schematic representation of the Cys residue positions on the cytoplasmically exposed regions of rabbit and scallop  $\text{Ca}^{2+}$ -ATPases.





Fig. 6. **SDS-PAGE and immunoblotting of recombinant  $\text{Ca}^{2+}$ -ATPases expressed in Sf21 cell.** Sf21 microsomes from wild type, C12V, C337I, C674R, C675L baculovirus-infected cells as well as rabbit SR were subjected to SDS-PAGE and immunoblotting. Panel A shows a Coomassie Blue-stained gel, and panel B shows a corresponding immunoblot developed with SERCA1a monoclonal antibody. Arrow indicates the position of  $\text{Ca}^{2+}$ -ATPase with a molecular weight of 115,000. Long bars also indicate positions of molecular weight markers.

**[Table II] Effect of various mutations on the EP formation activity of SERCA-1 enzyme expressed in Sf21 cells.**

Sample	Amount of EP (pmol/mg)		Remained activity (%)
	No treatment	Heat treated	
Wild Type	6.33	-	-
	5.86	-	-
	16.29 (2)	14.16	87
	17.09	12.66	74
	43.72	38.49	88
	47.16	38.81	82
Control	2.51	1.55	-
	0.93 (2)	1.34	-
C12V	11.67	12.60	108
	22.80	21.58	95
	24.24	21.53	89
C377I	18.27	12.54	69
	15.45	10.80	70
	29.52 (2)	29.18	99
C674R	13.82	15.05	107
	18.03	17.77	99
	7.23 (2)	7.71	107
	31.64 (3)	40.21	112
C675L	5.76	1.88	32
	8.01	1.10	60
	15.84 (2)	9.55	32


Number in ( ) represents number of experiments. The amount of EP was measured as described in "EXPERIMENTAL PROCEDURES". Control means ER fraction from no transfected cells.



**Fig. 7. Effects of single substitution of Cys675 to Leu on the thermal stability of the Ca<sup>2+</sup>-ATPase.** Microsomes from C675L (□) mutant cells as well as that of wild type (O) were incubated for 10 min at indicated temperatures in the presence of 0.1 mM CaCl<sub>2</sub>. Phosphorylation of the microsomes by AT<sup>32</sup>P was measured as described in "EXPERIMENTAL PROCEDURES".

for 5 min at 42°C lowered EP formation activity by 30%, whereas more than 80% of the activity of wild type ATPase remained even after the incubation at 50°C. From these results, it can be concluded that Cys675 presumably binds with another Cys through disulfide bond and contribute to structural stability of the rabbit Ca<sup>2+</sup>-ATPase.

To obtain further support for this conclusion I determined the effect of single substitution of Cys349 with Ala on the thermostability of rabbit Ca<sup>2+</sup>-ATPase. Autoradiogram in Fig.8 shows that EP formation activity of the C349A mutant ATPase decreased with increase in the temperature in the preincubation medium, and no activity was seen when the mutant was incubated for 10 min at temperatures higher than 40°C. Our conclusion to be drawn from these results combined with those of Table II and Fig. 7 is that Cys residues 675 and 349 may form a disulfide bond which can contribute to the thermal stabilization of the rabbit Ca<sup>2+</sup>-ATPase, while that lacking in the disulfide bond in the scallop Ca<sup>2+</sup>-ATPase may cause the thermal destabilization of the enzyme.



**Fig. 8. Autoradiogram of the phosphorylated Cys349->Ala mutant ATPase by  $AT^{32}P$  after incubation at various temperature.** Microsomes containing C349A mutant ATPase were incubated for 10 min at indicated temperatures in the presence of 0.1 mM  $CaCl_2$ .  $Ca^{2+}$ -ATPase was phosphorylated in the standard medium with 0.1 mM  $AT^{32}P$  at 0 °C and quenched after 15 sec by 5% TCA containing 5 mM Pi. Incorporation of  $^{32}P$  was determined by separating the proteins in acidic NaDodSo<sub>4</sub>/polyacrylamide gels for autoradiography essentially according to Maruyama and MacLennan (25).

## DISCUSSION

It is generally accepted that disulfide bond plays a key role in stabilizing enzyme structure. The number of related reports is increasing in the cation transport ATPase such as  $F_1$ -ATPase of mitochondria (11, 12), Na/K-pump ATPase and  $Ca^{2+}$ -pump ATPase of SR. For the Na/K-pump ATPase, reduction of disulfide bond (13, 14) or mutation (15) of Cys residues have been shown to result in a loss of ATPase activity. Further evidence was provided that structural disruption of the extra cellular portion of  $\beta$ -subunit drastically affects catalytic activity of  $\alpha$ -subunit of Na/K-ATPase (16, 17). It has been shown that the SR  $Ca^{2+}$ -ATPase contains 2-3 disulfide bonds (18-20), and that the ATPase is inhibited by the treatment with  $\beta$ -mercaptoethanol (20, 21), indicating that disulfide bonds can be involved in stabilization of the structure essential to the enzymatic function. Recently evidence has been presented that reduction of disulfide bonds on the  $Ca^{2+}$ -ATPase by DTT in the presence of nucleotide and  $Ca^{2+}$  diminish its catalytic function (22), although no evidence has been obtained for identification of Cys residues which are linking through disulfide bridge.

In this report, we have further explored the role of Cys residues on thermal stability of the  $Ca^{2+}$ -ATPase in comparison between primary sequences of scallop and rabbit  $Ca^{2+}$ -ATPase. The detailed comparison revealed a pronounced difference between the distribution of Cys residues on the cytoplasmically exposed region of rabbit and scallop  $Ca^{2+}$ -ATPases; Cys 12, 377, 670, 674, and 675 in rabbit  $Ca^{2+}$ -ATPase are replaced with Val, Ile, Val, Arg, and Leu, respectively, in scallop  $Ca^{2+}$ -ATPase. Cys349 of rabbit  $Ca^{2+}$ -ATPase is inaccessible to DACM, whereas Cys349 of scallop ATPase is readily reacted with the reagent under the same condition described above. In addition, Thr9, Ala219, Gly405, and Val449 of rabbit ATPase are replaced with Cys residues in scallop one (1). If we assume that the Cys residues linking through disulfide bond can contribute to thermal stabilization of the  $Ca^{2+}$ -ATPase, at least one of these Cys residues must be located on the rabbit ATPase, but not on scallop one. Furthermore, these residues should be inaccessible to SH reagents even they are exposed to outer medium. Among these candidates, Cys674 has already been reported to be accessible to IAF (23) and IAEDANS or IAA (24) under physiological conditions.

More definitive evidence has come from mutagenesis studies, in which single substitution of Cys349 to Ala or Cys675 to Leu were shown to drastically diminish



catalytic activity of the rabbit  $\text{Ca}^{2+}$ -ATPase when the enzyme was incubated above  $40^{\circ}\text{C}$  (Figs. 7 and 8). As a reasonable explanation for the lower sensitivity of the C675L mutant to thermal treatment, we postulate that substitution of any residues other than Cys675 may be also required for complete heat inactivation of the enzyme. These results suggested that Cys349 and Cys675 involve in the thermal stabilization of the structure essential for the enzymatic functions. However, it remains to be elucidated whether Cys349 and Cys675 are actually linking through disulfide bond on the rabbit  $\text{Ca}^{2+}$ -ATPase to contribute to thermal stabilization of the enzyme, or the local changes of environment around these residues might in some manner cause a pronounced increase in thermal stability of the enzyme. Experiments are presently under way to determine these possibilities.

In Fig.9, we propose a simplest model to explain a role of Cys residues 349 and 675 in thermal stabilization of rabbit  $\text{Ca}^{2+}$ -ATPase. According to the mobile pore model (26), a portion of the ATPase molecule could be mobile, permitting to act as a carrier through which  $\text{Ca}^{2+}$  ions can pass. In our model,  $\alpha$ -helices of Stalk-4 and at the top of Stalk-5, whose transmembrane segments M4 and M5 form a passage of  $\text{Ca}^{2+}$  ions together with M6, are connected through disulfide bond between Cys349 and Cys675. This linkage presumably serve as a stopper which can restrict to the minimal motion for the  $\text{Ca}^{2+}$ -passage. It is possible that since the motional restriction by the stopper is lacking in the scallop  $\text{Ca}^{2+}$ -ATPase, this enzyme is readily destructed by thermal treatment. This possibility may be more critical in E1 state of the enzyme where binding of  $\text{Ca}^{2+}$  to the enzyme causes a conformational change allowing the  $\text{Ca}^{2+}$ -entrance to open, than E2 state where the entrance is closed. This speculation is in good agreement with our previous studies (4) suggesting that removal of  $\text{Ca}^{2+}$  from the high affinity sites can protect the enzyme from thermal inactivation.




Fig. 8. Schematic model of heat-inactivation of rabbit  $\text{Ca}^{2+}$ -ATPase which lacks the Cys residue at the site of 349 or 675. Cys residues which were mutated in this study is shown by spheres. Cys349 and Cys675 are painted in orange, and others are in blue. Phosphorylation site and nucleotide binding site are also indicated in green and red, respectively (see text for details).

## REFERENCES

- 1) Nagata, Y., Yamamoto, T., Ema, M., Mimura, J., Fujii-Kuriyama, Y., Suzuki, T., Furukohri, T., Konishi, K., Sato, D., Tajima, G., and Nakamura, J. (1998) cDNA cloning and predicted primary structure of scallop sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase. *Comp. Biochem. Physiol. Part B* 119, 777-785
- 2) Tada, M., Yamamoto, T., and Tonomura, Y. (1978) molecular mechanism of active calcium transport by sarcoplasmic reticulum. *Physiol. Rev.* 56, 1-79
- 3) Andersen JP. (1995) Dissection of the functional domains of the sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase by site-directed mutagenesis. *Bioscience Reports* 15, 243-261
- 4) Abe, M., Shirakata, Y., Sato, D. Konishi, K., Watanabe, T., and Nakamura, J. (1992) Isolation and characteristics of scallop sarcoplasmic reticulum with calcium transport activity. *J. Biochem.* 112, 822-827
- 5) Nagata, Y., Nakamura, J., and Yamamoto, T. (1996) Protection of scallop sarcoplasmic reticulum ATPase from thermal inactivation by removal of calcium from high-affinity binding sites on the enzyme. *J. Biochem.* 119, 1100-1105
- 6) Nagata, Y., Nakamura, J., and Yamamoto, T. (1997) Temperature sensitivity of proteoliposomes reconstituted from a mixture of scallop and rabbit sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPases. *J. Biochem.* 121, 648-653
- 7) Yantorno, R.E., Yamamoto, T., and Tonomura, Y. (1983) Energy transfer between fluorescent dyes attached to  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ -ATPase in the sarcoplasmic reticulum. *J. Biochem.* 94, 1137-1145
- 8) Meissner, G., Conner, G.E., and Fleischer, S. (1973) Isolation of sarcoplasmic reticulum by zonal centrifugation, and purification of  $\text{Ca}^{2+}$ -pump and  $\text{Ca}^{2+}$ -binding proteins. *Biochim. Biophys. Acta* 298, 246-269
- 9) Yamasaki, K., Sano, N., Ohe, M., and Yamamoto, T. (1990) Determination of the primary structure of intermolecular cross-linking sites on the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum using  $^{14}\text{C}$ -labeled N,N'-(1,4-phenylene) bismaleimide or N-ethyl-maleimide. *J. Biochem.*, 108,918-925.
- 10) Yamasaki, K. and Yamamoto, T. (1992) Inhibition of phosphoenzyme formation from phosphate and sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase by vanadate binding to high- or low-affinity site on the enzyme. *J. Biochem.*,112,658-664.

- 11) Hisabori, T., Motohashi, K., Kroth, P., Strotmann, H., and Amano, T. (1998) The formation or the reduction of a disulfide bridge on the  $\beta$ -subunit of chloroplast ATP synthase affects the inhibitory effect of the  $\alpha$ -subunit. *J. Biol. Chem.* 273, 15901-15905
- 12) Nalin, C.M. and McCarty, R.E. (1984) Role of a disulfide bond in the gamma subunit in activation of the ATPase of chloroplast coupling factor 1. *J. Biol. Chem.* 259, 7275-7280
- 13) Kawamura, M. and Nagano, K. (1984) Evidence for essential disulfide bonds in the beta-subunit of  $(\text{Na}^{++}\text{K}^{+})$ -ATPase. *Biochim. Biophys. Acta* 774, 188-192
- 14) Kirley, T.L. (1990) Inactivation of  $(\text{Na}^{+},\text{K}^{+})$ -ATPase by  $\beta$ -mercapto ethanol. Differential sensitivity to reduction of the three beta subunit disulfide bonds. *J. Biol. Chem.* 265, 4227-4232
- 15) Kawamura, M. and Noguchi, S.H. (1992) In *The Sodium Pump: Structure, Mechanism, and Regulation* (Kaplan, J.H., and De Weer, P., Eds.) pp 45-61, Rockefeller University Press, New York.
- 16) Lutsenko, S. and Kaplan, J.H. (1993) An essential role for the extra cellular domain of the Na, K-ATPase  $\alpha$ -subunit in cation occlusion. *Biochemistry* 32, 6737-6743
- 17) Lutsenko, S. and Kaplan, J.H. (1992) Evidence of a role for the Na,K-ATPase beta-subunit in active cation transport. *Ann. N.Y. Acad. Sci.* 671, 147-155
- 18) Thorley-Lawson, D.A. and Green, N.M. (1977) The reactivity of the thiol groups of the adenosine triphosphatase of sarcoplasmic reticulum and their location on tryptic fragments of the molecule. *Biochem. J.* 167, 739-748
- 19) Abramson, J.J. and Shamoo, A.E. (1978) Purification and characterization of the 45,000-Dalton fragment from tryptic digestion of  $(\text{Ca}^{2+}+\text{Mg}^{2+})$ -adenosine triphosphatase of sarcoplasmic reticulum. *J. Membr. Biol.* 44, 233-257
- 20) Mutoh, Y., Sugino, A., Higashi, K., Takasugi, M., Kuroiwa, A., and Kawamura, M. (1992) Reduction and inactivation of the sarcoplasmic  $\text{Ca}(2+)$ -ATPase by 2-mercaptoethanol--contrast to the  $(\text{Na}^{+},\text{K}^{+})$  ATPase. *J. Univ. Occup. Environ. Health* 14, 253-260
- 21) Georgoussi, Z. and Sotiroidis, T.G. (1985) Inhibition of sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase by 2-mercaptoethanol. *Biochem. Biophys. Res. Commun.* 126, 1196-1200

- 22) Daiho, T. and Kanazawa, T. (1994) Reduction of disulfide bond in sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase by dithiothreitol causes inhibition of phosphoenzyme isomerization in catalytic cycle: The reduction requires binding of both purine nucleotide and  $\text{Ca}^{2+}$  to enzyme. *J. Biol. Chem.* 269, 11060-11064
- 23) Bishop, J.E., Squier, T.C., Bigelow, D.J., and Inesi, G. (1988) (Iodoacetamido)-fluorescein labels a pair of proximal cysteines on the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum. *Biochemistry* 27, 5233-5240
- 24) Yamashita, T. and Kawakita, M. (1987) Reactive sulfhydryl groups of sarcoplasmic reticulum ATPase. II Site of labeling with iodacetamide and its fluorescent derivative. *J. Biochem.* 101, 377-385
- 25) Maruyama, K. and MacLennan, D.H. (1988) Mutation of aspartic acid-351, lysine-352, and lysine-515 alters the  $\text{Ca}^{2+}$  transport activity of the  $\text{Ca}^{2+}$ -ATPase expressed in COS-1 cells. *Proc. Natl. Acad. Sci. USA* 85, 3314-3318
- 26) Singer, S.J. (1974) The molecular organization of membranes. *Ann. Rev. Biochem.* 43, 805-833

## Part IV

Temperature Sensitivity of Proteoliposomes Reconstituted from A Mixture of Scallop and Rabbit Sarcoplasmic Reticulum  $\text{Ca}^{2+}$ -ATPases

## SUMMARY

I reconstituted proteoliposomes by mixing scallop sarcoplasmic reticulum, SR with the rabbit one at different protein weight ratios, and investigated the effects of temperature on their  $\text{Ca}^{2+}$ -transport activity. When proteoliposomes containing scallop and rabbit SR at a protein ratio of 1:1 were pre-incubated in the presence of  $\text{Ca}^{2+}$  at  $39^\circ\text{C}$  for 10 min, the  $\text{Ca}^{2+}$ -transport activity was almost completely lost, whereas the activity of proteoliposomes containing rabbit SR alone decreased only slightly. Essentially the same results were obtained for proteoliposomes reconstituted with  $\text{Ca}^{2+}$ -ATPases partially purified from scallop and rabbit SR. The susceptibility of the reconstituted proteoliposomes to heat inactivation increased with increase of the protein weight ratio of the scallop to rabbit SR, with the maximum being approached at a ratio higher than 1. When the scallop SR was thermally treated before reconstitution, the resulting vesicles showed  $\text{Ca}^{2+}$ -transport activity as high as that of control vesicles reconstituted from rabbit SR alone. The former vesicles were not inactivated by further treatment at high temperature. In contrast, when scallop SR was heated in EGTA before, followed by reconstitution of vesicles with rabbit SR, their  $\text{Ca}^{2+}$ -transport activity was strongly inhibited by subsequent treatment at high temperature in the presence of  $\text{Ca}^{2+}$ . These results can be easily explained if I assume that  $\text{Ca}^{2+}$ -transport of the reconstituted vesicles can be catalyzed through dimeric interaction between the scallop and rabbit  $\text{Ca}^{2+}$ -ATPases in the membrane. Pre-incubation of these vesicles at  $39^\circ\text{C}$  for 10 min in the presence of  $\text{Ca}^{2+}$  may destroy the dimeric interaction due to denaturation of scallop  $\text{Ca}^{2+}$ -ATPase.

## INTRODUCTION

The  $\text{Ca}^{2+}$ -ATPase bound to SR membrane is responsible for the active transport of  $\text{Ca}^{2+}$  across the membrane. The coupling mechanism of ATP hydrolysis with  $\text{Ca}^{2+}$ -transport has been investigated extensively (1, 2). However, the movements of the ATPase molecules during the transport cycle have remained unknown. A number of investigators have suggested that  $\text{Ca}^{2+}$ -ATPase exists in the SR membrane in an oligomeric form (3-6), but the physiological role of the molecular interaction has been poorly understood.

In the preceding study, I investigated heat inactivation of scallop  $\text{Ca}^{2+}$ -ATPase under a variety of conditions, and concluded that the thermal lability of the catalytic function may closely correlated with the arrangement of the ATPase molecules in the SR membrane (7). As an extension of our previous studies on the ATPase-ATPase interaction, I reconstituted proteoliposomes by mixing the  $\text{Ca}^{2+}$ -ATPase of heat-labile scallop SR and heat-resistant rabbit SR at various protein weight ratios, and investigated the effect of temperature on their catalytic function. If there was no interaction between the two ATPases in the membrane, I could expect that the  $\text{Ca}^{2+}$ -transport activity of the reconstituted vesicle would additively change depending on their respective heat sensitivities. However, incubation of the vesicles at  $39^{\circ}\text{C}$  for 10 min almost completely eliminated their  $\text{Ca}^{2+}$ -transport activity, whereas heat treatment of the control vesicles reconstituted with rabbit SR alone only slightly decreased it.

These results suggest that the  $\text{Ca}^{2+}$ -ATPases of scallop and rabbit SR form an oligomer in the reconstituted membrane and that this molecular interaction is essential for  $\text{Ca}^{2+}$ -transport.



## EXPERIMENTAL PROCEDURES

**Materials** --- Scallop SR was isolated from the striated portion of scallop adductor muscle as described previously (7, 8). Rabbit SR was prepared from rabbit dorsal and leg muscles as described previously (9). These SR were suspended in medium containing 500 mM KCl, 10% glycerol, and 20 mM TES (pH 7.3), and divided into small pieces. They were quickly frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$ . The  $\text{Ca}^{2+}$ -ATPases were partially purified from scallop and rabbit SR according to the method described by Meissner *et al* (10).

**Chemical reagents** ---  $\text{C}_{12}\text{E}_9$  was purchased from Nikko Chemicals. Bio-beads SM-2 were obtained from Bio Rad Laboratories and repeatedly washed with methanol and water. Soybean asolectin was purchased from Sigma, A23187 from Calbiochem-Behring, and Antipyrylazo III from Nacalai. All other chemicals were of reagent grade.

**Reconstitution of proteoliposomes** --- Proteoliposomes were prepared by the method of Ando and Yamamoto (11) with the following modifications. SR was added to the 1 ml of reconstitution medium containing 5 mM  $\text{MgCl}_2$ , 0.1 M oxalate, 0.1 M KCl, 5-10% glycerol, and 0.1 M TES (pH 6.8) to a final concentration of 1-3 mg/ml protein. When proteoliposomes were reconstituted from a mixture of rabbit and scallop SR at indicated weight ratios, protein concentration of rabbit SR was kept at 1 mg/ml, and that of scallop SR was varied from 0 to 2 mg/ml. SR was solubilized by addition of 45 mg/ml  $\text{C}_{12}\text{E}_9$ . After several minutes, 40 mg/ml soybean asolectin solubilized in 45 mg/ml  $\text{C}_{12}\text{E}_9$  was added to the medium. The SR mixture was further incubated at  $20^{\circ}\text{C}$  for 15-20 min under continuous supply of  $\text{N}_2$  gas, then centrifuged to remove insoluble materials. Next, Bio-beads SM-2, 0.2-0.4 g dry weight, were added to the supernatant, and the mixture was vigorously stirred for 60-90 min under  $\text{N}_2$  gas. The Bio-beads were removed by passing the mixture through a cotton layer, and the filtrate was centrifuged as described above to remove debris. The supernatant was again incubated with 0.2 g/ml of Bio-beads for 45 min, then centrifuged at  $530,000 \times g$  for 30 min to collect the proteoliposomes. The pellet was suspended in 0.2-0.3 ml of the reconstitution medium from which oxalate had been omitted. The final sample was quickly frozen in liquid nitrogen and stored at  $-20^{\circ}\text{C}$ .

**Heat treatment of SR** --- SR suspension (0.1 ml) was transferred into incubation medium containing 5 mM  $\text{MgCl}_2$ , 0.1 M KCl, 5% glycerol, and 50 mM TES

(pH 6.8) which had been warmed in a glass test tube at the indicated temperature. At the indicated time, heat treatment was stopped by transferring the mixture into ice cold assay medium.

**Measurement of Ca<sup>2+</sup>-transport** --- Ca<sup>2+</sup>-transport was measured at 23°C in assay medium containing SR or reconstituted vesicles at 0.02-0.1 mg/ml protein, 20-50 μM CaCl<sub>2</sub>, and 0.2 mM Antipyrylazo III, as the Ca<sup>2+</sup> indicator. The reaction was initiated by addition of 0.2 mM ATP. The change in the Ca<sup>2+</sup> concentration in the reaction mixture was continuously monitored by measuring the absorption of the Ca<sup>2+</sup> indicator at 700 nm.

Ca<sup>2+</sup> efflux from the reconstituted vesicles was measured under conditions similar to those for the Ca<sup>2+</sup>-transport except for the absence of oxalate. The reaction was started by the addition of 5 μM A23187 to the reaction mixture.

## RESULTS

Sarcoplasmic reticulum isolated from cold water fish muscles including scallop adductor muscle has higher sensitivity to temperature than those from mammal muscles (8,12,13). Figure 1 compares the time courses of  $\text{Ca}^{2+}$ -transport by SR isolated from rabbit and scallop muscles after their pre-incubation at 39°C for 0, 5, and 10 min in medium containing 5 mM  $\text{MgCl}_2$ , 0.1 M KCl, 5% glycerol, and 50 mM TES (pH 6.8). Noted that the pre-incubation medium is usually contaminated with  $\mu\text{M}$   $\text{Ca}^{2+}$  under the conditions used, except when  $\text{Ca}^{2+}$  is removed by addition of 2-4 mM EGTA. The  $\text{Ca}^{2+}$ -transport activity of rabbit SR was unaffected by the heat treatment for up to 10 min, while the activity of scallop SR completely disappeared within 5 min.

Such a big difference in the thermolability between scallop and rabbit SR offers me the means to directly examine molecular interaction of  $\text{Ca}^{2+}$ -ATPase within the SR membrane. In the following experiments, I reconstituted proteoliposomes by mixing heat-labile scallop SR and heat-resistant rabbit SR to investigate the effects of thermal treatment upon  $\text{Ca}^{2+}$ -transport by the proteoliposomes. Figure 2 presents the time courses of  $\text{Ca}^{2+}$ -transport by the reconstituted vesicles before and after their incubation at 39°C for 10 min. The velocity of the  $\text{Ca}^{2+}$ -transport to decrease with increasing reaction time, probably due to the formation of some membrane potential accompanied by  $\text{Ca}^{2+}$  movements across the reconstituted membrane. The transport activity of the proteoliposomes reconstituted with rabbit SR alone (panel a) was only slightly decreased by pre-incubation at 39°C for 10 min, while the activity of the proteoliposomes reconstituted with scallop SR completely disappeared with the treatment (panel c). For the panel b data, proteoliposomes were reconstituted with a 1:1 mixture of scallop and rabbit SR by protein weight. Surprisingly, when they were incubated for 10 min at 39°C,  $\text{Ca}^{2+}$ -transport activity decreased to almost the same level as that observed for the control vesicles reconstituted with scallop SR alone (panel c). If I assume that the  $\text{Ca}^{2+}$ -transport can be independently catalyzed by the scallop and rabbit  $\text{Ca}^{2+}$ -ATPases in the reconstituted membrane, almost the same activity as observed for the heated vesicles which contain rabbit SR alone should remain after the heat treatment (compare between upper traces in panel a and b). Furthermore, it does not seem that  $\text{Ca}^{2+}$ -transport activity of the untreated vesicles was additively changed by mixing rabbit and scallop SR. These results suggest that scallop and rabbit  $\text{Ca}^{2+}$ -ATPase





Fig. 1. **Effects of heat treatment on  $\text{Ca}^{2+}$  transport by rabbit and scallop SR.** SR was added at  $39^{\circ}\text{C}$  to preincubation medium containing 5 mM  $\text{MgCl}_2$ , 0.1 M KCl, 5% glycerol, and 50 mM TES (pH 6.8) to a final concentration of 1 mg/ml. At 0 (a), 5 (b), and 10 min (c), the SR suspension was rapidly cooled in ice water to stop the heat treatment.  $\text{Ca}^{2+}$ -transport was initiated at  $23^{\circ}\text{C}$  by addition of 0.2 mM ATP ( $\downarrow$ ) to the assay solution containing 50  $\mu\text{M}$   $\text{CaCl}_2$ , 5 mM oxalate, 5 mM  $\text{MgCl}_2$ , 0.1 M KCl, 5% glycerol, and 50 mM TES (pH 6.8) in the presence of 0.2 mM Antipyrilazo III and 20  $\mu\text{g}/\text{ml}$  of SR.



**Fig. 2. Effect of heat treatment on  $Ca^{2+}$  transport by proteoliposomes reconstituted from scallop and rabbit SR.** Proteoliposomes were reconstituted as described in "EXPERIMENTAL PROCEDURES" by mixing scallop and rabbit SR at the protein weight ratio of 0:1 (a), 1:1 (b), and 1:0 (c). The proteoliposomes were suspended in the preincubation medium and thermally treated as given in Fig. 1, for 0 (lower traces) and 10 min (upper traces).  $Ca^{2+}$  transport by the proteoliposomes was measured at 23°C under the conditions described in Fig. 1. ( $\downarrow$ ), addition of ATP.

molecules exist in the reconstituted membrane mainly as an oligomer, and that interaction between them is essential for the enzyme to catalyze  $\text{Ca}^{2+}$ -transport. Heat denaturation of the scallop  $\text{Ca}^{2+}$ -ATPase may result in destruction of the oligomeric interaction.

Another possible explanation for the susceptibility of the reconstituted vesicles to heat inactivation is that the thermal treatment may specifically increase  $\text{Ca}^{2+}$  permeability of the membrane *via* the scallop  $\text{Ca}^{2+}$ -ATPase. In the experiments presented in Fig. 3, I measured the  $\text{Ca}^{2+}$  efflux from proteoliposomes reconstituted with scallop and rabbit SR at different protein weight ratios. These vesicles were loaded with  $\text{Ca}^{2+}$  during the reconstitution in the presence of 10 mM  $\text{CaCl}_2$ . The  $\text{Ca}^{2+}$ -loaded proteoliposomes were further incubated at 39°C for 10 min. Panel a, b, and c of Fig. 3 show the time courses of  $\text{Ca}^{2+}$  release from the respective proteoliposomes reconstituted with rabbit SR alone, a 1:1 mixture of scallop and rabbit SR, and scallop SR alone, after addition of 5  $\mu\text{M}$  A23187. Upon addition of the ionophore,  $\text{Ca}^{2+}$  was rapidly released from the proteoliposomes and reached the steady level within 30 sec. From the calibration curve of the  $\text{Ca}^{2+}$  concentration in panel d, about 15  $\mu\text{M}$   $\text{Ca}^{2+}$  was released from each sample of reconstituted vesicles.

The amount of  $\text{Ca}^{2+}$  released from the thermally treated vesicles was about 20% lower than that from untreated vesicles possibly due to loss of internal  $\text{Ca}^{2+}$  during the treatment. The reduction in the amount of  $\text{Ca}^{2+}$  release after the heat treatment was seen in the vesicles reconstituted with rabbit SR alone (panel a) as well as in those containing scallop SR (panel b and c). These results do not support the possibility of heat inactivation of  $\text{Ca}^{2+}$ -transport of proteoliposomes being caused by a rise in  $\text{Ca}^{2+}$  leakage of the membrane *via* the scallop  $\text{Ca}^{2+}$ -ATPase.

As shown in Fig. 4, the susceptibility of the reconstituted proteoliposomes to heat inactivation increased as a function of the protein weight ratio of scallop SR to rabbit one, and appeared to reach the maximal level at a ratio higher than 1 (Fig. 4, inset). These observations support the possibilities that the scallop and rabbit  $\text{Ca}^{2+}$ -ATPases form a dimer in the reconstituted membrane at the molar ratio of 1:1 and that the molecular interaction between them plays an important role in the active transport of  $\text{Ca}^{2+}$  across the membrane. I decided to test whether or not the scallop  $\text{Ca}^{2+}$ -ATPase could still form an oligomer in the reconstituted membrane even after the SR was thermally treated. Figure 5 shows the results for proteoliposomes reconstituted by





Fig. 3.  $Ca^{2+}$  release from the  $Ca^{2+}$ -loaded proteoliposomes after addition of  $Ca^{2+}$  ionophore. Proteoliposomes were reconstituted by mixing scallop and rabbit SR at a ratio of 0:1 (a), 1:1 (b), and 1:0 (c) in the solution described in "EXPERIMENTAL PROCEDURES" except for substitution of 10 mM  $CaCl_2$  for oxalate. The  $Ca^{2+}$ -loaded proteoliposomes were incubated for 10 min at 23°C (upper traces) and 39°C (lower traces). They were diluted with 50-fold of the assay medium containing 0.2 mM Antipyryl azo III.  $Ca^{2+}$  release was measured at 23°C after addition of 5  $\mu M$  A23187 (+) using the calibration curve as shown in panel d.



**Fig. 4. Dependence of the heat sensitivity of proteoliposomes on the protein weight ratio of scallop to rabbit SR in the reconstituted membrane.** A. Proteoliposomes were reconstituted as described in "EXPERIMENTAL PROCEDURES" by mixing scallop and rabbit SR at a protein weight ratio of 0:1 (1), 0.4:1 (2), 0.6:1 (3), 1:1 (4), and 2:1 (5), respectively. Each sample was incubated at 39°C for 0 (lower traces) and 10 min (upper traces), and the time courses of Ca<sup>2+</sup> transport were measured as given in Fig. 1. B. Heat inactivation was represented as a percentage and plotted against the protein weight ratio of scallop to rabbit SR. They were determined from the Ca<sup>2+</sup> concentration difference at 1 min after addition of ATP (↓) between the time courses of Ca<sup>2+</sup> transport in the control (lower traces) and heat-treated samples (upper traces).



mixing pre-heated scallop SR with native rabbit SR at the protein weight ratios of 0:1 (panel a), 1:1 (panel b), and 2:1 (panel c). In each case, the control vesicles were reconstituted with native scallop and rabbit SR at the respective ratios, then these vesicles were incubated for 10 min at 39°C. It is clear from panel b or c of Fig. 5 that heat treatment of scallop SR before reconstitution did not inhibit the Ca<sup>2+</sup>-transport activity, whereas heat treatment after reconstitution strongly inhibited it. The former vesicles possess transport activity as high as those vesicles reconstituted with rabbit SR alone (panel a). On the other hand, heat treatment of rabbit SR either before or after reconstitution had virtually no effect on the Ca<sup>2+</sup>-transport as shown in panel a. These observations indicate that heat treatment of scallop SR before reconstitution resulted in failure of molecular interaction among the Ca<sup>2+</sup>-ATPases. In such a case, native Ca<sup>2+</sup>-ATPase of rabbit SR forms a functional dimer by itself in the reconstituted membrane.

In the preceding study (7), I found that when scallop SR was heated in the presence of 2 to 5 mM EGTA, the Ca<sup>2+</sup>-dependent ATP hydrolysis, as well as EP formation was almost completely protected against heat inactivation, although Ca<sup>2+</sup>-transport activity was completely lost. Since the protection was not seen when SR was heated in the presence of a free Ca<sup>2+</sup> concentration higher than 0.5 μM, I concluded that removal of Ca<sup>2+</sup> from the high-affinity binding sites of the scallop ATPase resulted in complete protection against heat inactivation. The difference in thermolability of scallop ATPase between in the presence and absence of Ca<sup>2+</sup> may be attributed to differences in protein structure. It would be interesting to know if the scallop Ca<sup>2+</sup>-ATPase can still interact with rabbit Ca<sup>2+</sup>-ATPase to form a dimer in the membrane when scallop SR is thermally treated in the presence of EGTA, then reconstituted it into proteoliposomes.

As shown in panel b of Fig. 6, the resulting proteoliposomes exhibited Ca<sup>2+</sup>-transport activity as high as that of the control vesicles which were reconstituted from native scallop and rabbit SR (panel a). The transport activity of the proteoliposomes was inhibited, in the similar manner to the control vesicles, by subsequent incubation at 39°C in the presence of Ca<sup>2+</sup> (upper curve in panel a, b). For the experiments in panel c, proteoliposomes were reconstituted by mixing the native rabbit SR with scallop SR which had been heated in Ca<sup>2+</sup> instead of EGTA. These vesicles similarly showed high Ca<sup>2+</sup>-transport activity, but did not show any inhibition of the activity by further treatment at high temperature in the presence of Ca<sup>2+</sup> (panel c). These findings suggest that the heat treatment of scallop SR in the presence of Ca<sup>2+</sup> may prevent not only the

activity of  $\text{Ca}^{2+}$ -transport but also the interaction between the ATPase molecules. On the other hand, heat treatment of SR in the absence of  $\text{Ca}^{2+}$  may fully protect the molecular interaction among the  $\text{Ca}^{2+}$ -ATPases in the reconstituted membrane.





Fig. 5. **Reconstitution of heat-resistant proteoliposomes from pre-heated scallop SR and native rabbit SR.** Lower traces: SR was heated in the presence of  $\text{Ca}^{2+}$  for 10 min before reconstitution. Proteoliposomes were reconstituted by mixing heat-inactivated scallop SR with intact rabbit SR at a ratio of 0:1 (a), 1:1 (b), and 2:1 (c). Upper traces: Proteoliposomes were reconstituted from native scallop and rabbit SR at the respective ratios given above, then incubated at  $39^\circ\text{C}$  for 10 min. The time courses of  $\text{Ca}^{2+}$  transport were measured under the conditions described in Fig. 1. ( $\downarrow$ ), addition of ATP.



**Fig. 6. Reconstitution of heat-labile proteoliposomes from scallop SR pre-heated in EGTA and native rabbit SR.** Proteoliposomes were reconstituted by mixing native rabbit SR with native scallop SR (a), with SR pre-heated in the presence (b) or absence of 2 mM EGTA (c), at a protein weight ratio of 1:1.  $Ca^{2+}$ -transport activity was measured after heat treatment of these samples in the presence of 0.1 mM  $CaCl_2$  for 0 (lower traces) and 10 min (upper traces). ( $\downarrow$ ), addition of ATP.

## DISCUSSION

Much work has been done on the molecular interaction among the  $\text{Ca}^{2+}$ -ATPases in the SR membrane by adopting a variety of methods and techniques such as electron microscopic observations (14-16), saturation transfer EPR (17-20), fluorescence spectroscopy (21, 22), chemical cross linking methods (23-25), and kinetic analysis (5, 27-31). However, the precise interactions remain poorly understood due to critical restrictions on the investigation conditions.

In the present study, I examined the temperature effects on the  $\text{Ca}^{2+}$ -transport by reconstituted proteoliposomes consisting of heat-labile scallop SR and heat-stable rabbit one. If there is no interaction between their  $\text{Ca}^{2+}$ -ATPase molecules in the reconstituted membrane, incubation of the vesicles at  $39^{\circ}\text{C}$  for 10 min in the presence of  $\text{Ca}^{2+}$  will inhibit only the  $\text{Ca}^{2+}$ -transport catalyzed by the scallop  $\text{Ca}^{2+}$ -ATPase, but not that by the rabbit one. In fact, however, heat treatment of the proteoliposomes resulted in almost complete attenuation of  $\text{Ca}^{2+}$ -transport, indicating that it inactivated not only the scallop but also the rabbit SR  $\text{Ca}^{2+}$ -ATPase (Fig. 2). Similar phenomenon was observed with the proteoliposomes reconstituted from the mixture of purified  $\text{Ca}^{2+}$ -ATPases of scallop and rabbit SR (Data not shown). This observation supports our previous work (7) which suggests that the thermal treatment directly affects scallop  $\text{Ca}^{2+}$ -ATPase.

It is unlikely that the heat denaturation of scallop  $\text{Ca}^{2+}$ -ATPase would increase the  $\text{Ca}^{2+}$  permeability of the reconstituted membrane to reduce the  $\text{Ca}^{2+}$ -transport activity, since the amount of  $\text{Ca}^{2+}$  released from the vesicles upon addition of a  $\text{Ca}^{2+}$  ionophore was slightly smaller in the thermally treated vesicles than the untreated ones (Fig. 3). In addition, the difference of the amount of  $\text{Ca}^{2+}$  release from the vesicles containing scallop and rabbit  $\text{Ca}^{2+}$ -ATPase was almost the same as that from the control vesicle containing rabbit  $\text{Ca}^{2+}$ -ATPase alone. These results well agreed with our previous observation that heat treatment of scallop SR in the presence of EGTA strongly inhibits  $\text{Ca}^{2+}$ -transport activity without a rise in  $\text{Ca}^{2+}$  permeability of the SR membrane (7).

The susceptibility of the proteoliposomes to heat inactivation increased as a function of the protein weight ratio of scallop to rabbit SR in the reconstituted membrane, and appeared to reach maximum at a ratio higher than 1 (Fig. 4). These

results suggest that, in the reconstituted membrane, each mol of  $\text{Ca}^{2+}$ -ATPases of scallop and rabbit SR may dominantly form a dimer as a minimum functional unit, and that the dimeric interaction between them may be involved in the catalytic cycle of  $\text{Ca}^{2+}$ -transport. Heat treatment of reconstituted vesicles at  $39^{\circ}\text{C}$  for 10 min may easily destroy the dimeric interaction in the functional unit *via* denaturation of scallop  $\text{Ca}^{2+}$ -ATPase.

As shown in Fig. 5, when the scallop SR was heated in the presence of  $\text{Ca}^{2+}$  before reconstituting vesicles, the resulting vesicles still possessed  $\text{Ca}^{2+}$ -transport activity as high as those reconstituted from rabbit SR alone. This finding can be interpreted that, since heat-denatured scallop ATPase does not form a dimer in the membrane, the remaining rabbit ATPase molecules will form a dimer by themselves to serve the  $\text{Ca}^{2+}$  pump. In the preceding study (7), I found that removal of  $\text{Ca}^{2+}$  by EGTA from the high-affinity sites on the scallop ATPase fully protected ATPase reaction, but not  $\text{Ca}^{2+}$ -transport activity, against heat inactivation. In addition, several authors including as have suggested that removal of  $\text{Ca}^{2+}$  from the enzyme may stabilize it in an oligomeric form (7, 16, 23, 26). As shown in Fig. 6b, heat sensitive proteoliposomes were obtained when scallop SR was previously heated in EGTA, then reconstituted it into vesicle together with the same amount of rabbit intact SR. Since catalytic activity of scallop  $\text{Ca}^{2+}$ -ATPase had been lost by the treatment (7), the  $\text{Ca}^{2+}$ -transport into a vesicle would be catalyzed by rabbit ATPase in the reconstituted membrane. This phenomenon is interpreted that, scallop  $\text{Ca}^{2+}$ -ATPase may be still capable to interact with rabbit one, and that the interaction was easily destroyed by the subsequent treatment of the vesicles at high temperature in the presence of  $\text{Ca}^{2+}$ . Work is underway to identify and characterize the heat-sensitive domain as well as the molecular interaction domain on scallop  $\text{Ca}^{2+}$ -ATPase by adopting genetic method.

## REFERENCES

- 1) Yamamoto, T., Takisawa, H., and Tonomura, Y. (1979) Reaction mechanisms of ATP hydrolysis and synthesis in the sarcoplasmic reticulum. *Curr. Top. Bioenerg.* 9, 179-236
- 2) Inesi, G., Lewis, D., Hussain, A., and Kirtley, M.E. (1992) Long-range intramolecular linked functions in the calcium transport ATPase. *Adv. Enzymol.* 65, 185-215
- 3) Scales, D., and Inesi, G. (1976) Assembly of ATPase protein in sarcoplasmic reticulum membrane. *Biophys. J.* 16, 735-75
- 4) Andersen, J.P., Fellmann, P., Moller, J.V., and Devaux, P.L. (1981) Immobilization of a spin-labeled fatty acid chain covalently attached to  $\text{Ca}^{2+}$ -ATPase from sarcoplasmic reticulum suggests an oligomeric structure. *Biochemistry* 20, 4928-4936
- 5) Ikemoto, N. (1982) Structure and function of the calcium pump protein of sarcoplasmic reticulum. *Annu. Rev. Physiol.* 44, 297-317
- 6) Birmachou, W., and Thomas, D.D. (1990) Rotational dynamics of the Ca-ATPase in sarcoplasmic reticulum studied by time-resolved phosphorescence anisotropy. *Biochemistry* 29, 3904-3914
- 7) Nagata, Y., Nakamura, J., and Yamamoto, T. (1996) Protection of scallop sarcoplasmic reticulum ATPase from thermal inactivation by removal of calcium from high-affinity binding sites on the enzyme. *J. Biochem.* 119, 1100-1105
- 8) Abe, M., Shirakata, Y., Sato, D., Konishi, K., Watanabe, T., and Nakamura, J. (1992) Isolation and characteristics of scallop sarcoplasmic reticulum with calcium transport activity. *J. Biochem.* 112, 822-827
- 9) Yamada, S., Yamamoto, T., and Tonomura, Y. (1970) Reaction mechanism of the  $\text{Ca}^{2+}$ -dependent ATPase of sarcoplasmic reticulum from skeletal muscle. *J. Biochem.* 67, 789-794
- 10) Meissner, G., Conner, G.E., and Fleischer, S. (1973) Isolation of sarcoplasmic reticulum by zonal centrifugation and purification of  $\text{Ca}^{2+}$ -pump and  $\text{Ca}^{2+}$ -binding proteins. *Biochim. Biophys. Acta* 298, 246-269
- 11) Ando, R. and Yamamoto, T. (1985) Restoration of  $\text{Ca}^{2+}$ -transport in sarcoplasmic reticulum ATPase solubilized with octaethyleneglycol mono n-dodecyl ether. *J.*

Biochem. 97, 877-882

- 12) Chini, E.N., de Toledo, F.G.S., Albuquerque, M.C., and de Meis, L. (1993) The  $\text{Ca}^{2+}$ -transporting ATPases of rabbit and trout exhibit different pH- and temperature-dependences. *Biochem. J.* 293, 469-473
- 13) de Toledo, F.G.S., Albuquerque, M.C., Goulart, B.H., and Chini, E.N. (1995) Different thermostabilities of sarcoplasmic reticulum ( $\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ )-ATPase from rabbit and trout muscles. *Comp. Biochem. Physiol.* 111C, 93-98
- 14) Buhle, E.L., Knox, B.E., Serpersu, E., and Aebi, U. (1983) The structure of the  $\text{Ca}^{2+}$ -ATPase as revealed by electron microscopy and image processing of ordered arrays. *J. Ultrastruct. Res.* 85, 186-203
- 15) Tayler, K.A., Dux, L., and Martonosi, A. (1986) Three-dimensional reconstruction of negative stained crystals of the  $\text{Ca}^{2+}$ -ATPase from muscle sarcoplasmic reticulum. *J. Mol. Biol.* 187, 417-427
- 16) Dux, L., Taylor, K.A., Ting-Beall, H.P., and Martonosi, A. (1985) Crystallization of the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum by calcium and lanthanide ions. *J. Biol. Chem.* 260, 11730-11743
- 17) Bigelow, D.J. and Thomas, D.D. (1987) Rotational dynamics of lipid and the  $\text{Ca}$ -ATPase in sarcoplasmic reticulum: The molecular basis of activation by diethyl ether. *J. Biol. Chem.* 262, 13449-13456
- 18) Squier, T.C., Bigelow, D.J., and Thomas, D.D. (1988) Lipid fluidity directly modulates the overall protein rotational mobility of the  $\text{Ca}$ -ATPase in sarcoplasmic reticulum. *J. Biol. Chem.* 263, 9178-9186
- 19) Birmachou, W. and Thomas, D.D. (1990) Rotational dynamics of the  $\text{Ca}$ -ATPase in sarcoplasmic reticulum studied by time-resolved phosphorescence anisotropy. *Biochemistry* 29, 3904-3914
- 20) Karon, B.S. and Thomas, D.D. (1993) Molecular mechanism of  $\text{Ca}$ -ATPase activated by halothane in sarcoplasmic reticulum. *Biochemistry* 32, 7503-7511
- 21) Kutchai, H., Mahaney, J.E., Geddis, L.M., and Thomas, D.D. (1994) Hexanol and lidocaine effect the oligomeric state of The  $\text{Ca}$ -ATPase of sarcoplasmic reticulum. *Biochemistry* 33, 13208-13212
- 22) Karon, B.S., Mahaney, J.E., and Thomas, D.D. (1994) Halothane and cyclopiazonic acid modulate  $\text{Ca}$ -ATPase oligomeric state and function in sarcoplasmic reticulum. *Biochemistry* 33, 13928-1393



- 23) Yamasaki, K. and Yamamoto, T. (1989) Effects of adenylyl-5'-imidodiphosphate and vanadate ion on the intermolecular cross-linking of  $\text{Ca}^{2+}$ -ATPase in the sarcoplasmic reticulum membrane with N,N'-(1,4-phenylene)bismaleimide. *J. Biochem.* 106, 1114-1120
- 24) Keresztes, T., Jona, I., Pikula, S. Vegh, M., Mullner, N., Papp, S., and Martonosi, A. (1989) Effect of calcium on the interactions between  $\text{Ca}^{2+}$ -ATPase molecules in sarcoplasmic reticulum. *Biochim. Biophys. Acta* 984, 326-338
- 25) Yamasaki, K., Sano, N., Ohe, M., and Yamamoto, T. (1990) Determination of the primary structure of intermolecular cross-linking sites on the  $\text{Ca}^{2+}$ -ATPase of sarcoplasmic reticulum using  $^{14}\text{C}$ -labeled N,N'-(1,4-phenylene)bismaleimide or N-ethylmaleimide. *J. Biochem.* 108, 918-925
- 26) Stokes, D.L. and Lacapere, J.J. (1994) Conformation of  $\text{Ca}^{2+}$ -ATPase in two crystal forms: Effects of  $\text{Ca}^{2+}$ , thapsigargin, Adenosine 5'-(-methylene)triphosphate, and chromium(III)-ATP on crystallization. *J. Biol. Chem.* 269, 11606-11611
- 27) Froehlich, J.P. and Taylor, E. (1975) Transient state kinetic studies of sarcoplasmic reticulum adenosine triphosphatase. *J. Biol. Chem.* 250, 2013-2021
- 28) Dupont, Y., Pougeois, R., Ronjat, M., and Verjovsky-Almeida, S. (1985) Two distinct classes of nucleotide binding sites in sarcoplasmic reticulum Ca-ATPase by 2',3'-o-(2,4,6-trinitrocyclohexadienylidene)-ATP. *J. Biol. Chem.* 260, 7241-7249
- 29) Ferreira, S.T. and Verjovski-Almeida, S. (1988) Stoichiometry and mapping of the nucleotide sites in sarcoplasmic reticulum ATPase. *J. Biol. Chem.* 263, 9973-9980
- 30) Yamasaki, K. and Yamamoto, T. (1991) Existence of high- and low-affinity vanadate-binding sites on  $\text{Ca}^{2+}$ -ATPase of the sarcoplasmic reticulum. *J. Biochem.* 110, 915-921
- 31) Yamasaki, K. and Yamamoto, T. (1992) Inhibition of phosphoenzyme formation from phosphate and sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase by vanadate binding to high- or low-affinity site on the enzyme. *J. Biochem.* 112, 658-664

## BIBLIOGRAPHY

1. Yoshiaki Nagata, Jun Nakamura, and Taibo Yamamoto  
Protection of scallop sarcoplasmic reticulum ATPase from thermal inactivation by removal of calcium from high-affinity binding sites on the enzyme  
J. Biochem. 119, 1100-1105 (1996)
2. Yoshiaki Nagata, Jun Nakamura, and Taibo Yamamoto  
Temperature sensitivity of proteoliposomes reconstituted from a mixture of scallop and rabbit sarcoplasmic reticulum Ca<sup>2+</sup>-ATPases  
J. Biochem. 121, 648-653 (1997)
3. Yoshiaki Nagata, Taibo Yamamoto, Masatsugu Ema, Junsei Mimura, Yoshiaki Fujii-Kuriyama, Tomohiko Suzuki, Takahiro Furukohri, Kazuhiko Konishi, Dai Sato, Genichi Tajima, Jun Nakamura  
cDNA cloning and predicted primary structure of scallop sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase  
Comp. Biochem. Physiol. B 119, 777-785 (1998)
4. Yoshiaki Nagata, Jun Nakamura, and Taibo Yamamoto  
Change in the thermal sensitivity of sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase by chemical modifications and mutagenesis of specific Cys residues  
In preparation

## ACKNOWLEDGEMENTS

I wish to express my sincere thanks to Professor Akihiko Ogura and Associate Professor Taibo Yamamoto of Osaka University for their helpful guidance and valuable advices through this work. I am greatly indebted to Associate Professor Jun Nakamura, Dr. Genichi Tajima, and Mr. Dai Satoh of Tohoku University for their assistance, support, and efforts during my work in Tohoku University.

I am indebted to Professor Yoshiaki Fujii-Kuriyama, Dr Masatsugu Ema, and all the member of their laboratory for their gracious guidance of the techniques of molecular biology. I am also grateful to Associate Professor Haruhiko Takisawa for his excellent advices and valuable suggestions. I would like to thank Dr. Akio Inoue, Dr. Toshiaki Arata, and Dr. Keiko Yoshino-Tominaga for their helpful advices.

I wish to thank Professor Takahiro Furukohri and Associate Professor Tomohiko Suzuki of Kochi University for analysis of amino acid composition. I am also grateful to Professor David H. MacLennan of Toronto University for providing SERCA1a gene and monoclonal antibody A52.

I would like to thank Professor Tohru Kanazawa, Associate Professor Hiroshi Suzuki, Dr. Takashi Daiho, and Dr. Kazuo Yamasaki of Asahikawa medical college for their valuable suggestion and discussion. Thanks also go to Dr. Makoto Ushimaru of Kyorin University for his useful advices.

I wish to thank Professor Seiki Kuramitsu and Yuji Goto of Osaka University for his useful advice about protein stability. I am also grateful to Professor Masao Kawakita of The Tokyo Metropolitan Institute of Medical Science for his useful advices about amino acid modification on  $\text{Ca}^{2+}$ -ATPase.

I am also grateful to the member of Professor Ogura's laboratory for their kind help during this study.

I wish to thank Akiko Harada for her warmhearted encouragement.

I am deeply grateful to my parents, sister, and grandfather for their continuous encouragement and support for my study in University.