



Title	起立負荷 $99mTc$ -HMPAO脳SPECTによる脳灌流圧の変化の観察
Author(s)	林田, 孝平; 西村, 恒彦; 植原, 敏勇 他
Citation	日本医学放射線学会雑誌. 1992, 52(7), p. 1036-1038
Version Type	VoR
URL	https://hdl.handle.net/11094/19667
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

研究速報

起立負荷^{99m}Tc-HMPAO脳SPECTによる脳灌流圧の観察

国立循環器病センター放射線診療部¹⁾

大阪大学医学部トレーサー解析²⁾

林田 孝平¹⁾ 西村 恒彦^{1,2)} 植原 敏勇¹⁾

神長 達郎¹⁾ 広瀬 義晃¹⁾

（平成4年4月16日受付）

（平成4年5月15日最終原稿受付）

Raise up ^{99m}Tc-HMPAO Brain SPECT for Detecting the Change in Cerebral Perfusion Pressure

Kohei Hayashida¹⁾, Tsunehiko Nishimura^{1,2)}, Toshiisa Uehara¹⁾,
Tatsuro Kaminaga¹⁾ and Yoshiaki Hirose¹⁾

¹⁾Department of Diagnostic Radiology, National Cardiovascular Center

²⁾Department of Tracer Data Analysis, Medical School, Osaka University

Research Code No. : 721

Key Words : Cerebral vascular disease, Raise up stress test,
^{99m}Tc-HMPAO, Brain SPECT

^{99m}Tc-HMPAO crossed the blood brain barrier instantly in proportional to cerebral blood flow. Raise up stress and supine resting ^{99m}Tc-HMPAO with brain SPECT was devised for detecting abnormal response in the changes of cerebral perfusion pressure. Asymmetric ratio in the middle cerebral artery area in patients with internal carotid artery occlusion (n=5) was changed significantly from 0.82 ± 0.06 to 0.90 ± 0.06 and that of normal (n=5) was not changed from 0.94 ± 0.03 to 0.95 ± 0.01 . Raise up ^{99m}Tc-HMPAO brain SPECT enhanced the detectability in abnormality of regional cerebral blood flow and visualized the dys-autoregulated area during blood pressure falls.

1. はじめに

^{99m}Tc-Hexamethylpropyleneamine oxime (= HMPAO)d, 1体は¹⁾、脳血液閥門を通過し、瞬時に低脂溶性に変換されることにより脳血流分布を観察できる²⁾。さらに、Acetazolamide 静注法³⁾、CO₂吸入法⁴⁾などの脳血流増加、Matas test⁵⁾、Balloon occ lusion test⁶⁾、Wada test⁷⁾など脳血流減少の負荷法に応用してきた。今回、起立負荷時、臥位安静時にそれぞれ^{99m}Tc-HMPAOを投与し、脳血流の変化を捉える方法を開発したので報告する。

2. 方法と対象

ベッドサイド^{99m}Tc-HMPAO標識法 (Fig. 1)：起立直前に三方活栓をつけた19Gエラスタ針をセレック・バイアル（アマシャム薬品）に注入し、注射器で^{99m}TcO₄⁻ 1,300MBqを注入、吸引を繰り返すことにより標識し、初回投与分500MBqと減衰を見込んだ二回目投与分630MBqの^{99m}Tc-HMPAOを調整した。

起立負荷法 (Fig. 2)：臥位にて両足を挙上し15分後、起立させ直ちに^{99m}Tc-HMPAO 500MBqを静注する。初回SPECT撮像終了後^{99m}Tc-HMPAO 600MBqを再投与し二回目のSPECT




Fig. 1 Labeling method of HMPAO with Tc-99m at bed-side. (HMPAO = Hexamethyl-propyleneamine oxime)




Fig. 2 Procedure of raise up stress and supine rest testing with 99m Tc-HMPAO brain SPECT.

の撮像を行う。SPECT像において両側の中大脳動脈領域の皮質に関心領域を設定し、初回と二回目の対側比を比較した。有意差検定は paired t-test を用いた。また起立前後に血圧測定を行った。対象は、片側内頸動脈閉塞(=ICO) 7例(年齢65±9歳、男：女=6:1)で、このうち5例は、X線CTにて低吸収域を認めなかった。

尚、対照は脳血管に異常を認めない正常5例(=NR)(年齢62±15歳、男：女=3:2)である。使用機種は、GAPコリメータを装着したGeneral Electric社製STARCAM 400AC/Tである。収集マトリックス 64×64 、初回は15秒間、二回目20秒間の64stepのデータを収集した。吸収補正(0.12 cm^{-1})、Butter-worth フィルターによる前処理後、Ramp フィルターにて再構成した。また、初回と二回目の差し引像にて臥位の脳血流分布も観察した。

3. 結 果

収縮期の血圧変動は、X線CTにて低吸収域を認めないICO群で $-12 \pm 4 \text{ mmHg}$ 、NR群で $-10 \pm 5 \text{ mmHg}$ と有意差がなかった。また、中大脳動脈領域の対側比は、ICO群の起立負荷では 0.82 ± 0.06 、臥位では 0.90 ± 0.06 と有意差($p < 0.01$)があったが、NR群の起立負荷では 0.94 ± 0.03 、臥位では 0.95 ± 0.01 と有意差がなかった。X線CTにて低吸収域を認めたICO 2例を示す。(症例1: Fig. 3)63歳男性、梅毒性大動脈瘤の置換手術時に左総頸、左鎖骨下動脈を結紮した。X線CTでは左放射冠にのみ低吸収域があり、中大脳動脈域では、脳萎縮を認めた。起立時(血圧変動 -10 mmHg)に前頭葉から頭頂葉に血流低下があり、臥位にて前頭葉に血流増加を認めた。(症例2: Fig. 4)77歳、男性。血管造影では左内頸動脈が閉塞し、X線CTにて左前頭葉に低吸収域を認めた。発症15日目の 99m Tc-HMPAO脳SPECTにて、座位の起立負荷(血圧変動 -10 mmHg)で左




Fig. 3 Case of postoperative thoracic aneurysm with ligation of left common carotid artery. Comparing raise up with supine resting 99m Tc-HMPAO brain SPECT, a perfusion defect in the left frontal area diminishes in size in supine resting indicating an abnormal responded area in cerebral blood flow during blood pressure fall in the vicinity of the infarcted area (\rightarrow). CR=Coronal view. TX=Transaxial view. UP=Raise up 99m Tc-HMPAO brain SPECT. Down=Supine resting 99m Tc-HMPAO brain SPECT. SUB=Subtracted 99m Tc-HMPAO brain SPECT.




Fig. 4 Case of subacute cerebral infarction. Comparing raise up with supine resting ^{99m}Tc-HMPAO brain SPECT, a hyperemic area in the left frontal area being interpreted as luxury perfusion shows an excessive decrease in vascularity during blood pressure fall indicating the dys-autoregulated area (→). Abbreviations as in Fig. 3.

前頭葉に hyperemia があり、この領域は臥位にて異常な血流増加を認めた。さらに、発症後22日のポジトロン検査にて hyperemia の領域は酸素代謝が低下しており“Luxury perfusion”と判定できた。

4. 考 察

起立負荷時、臥位安静時に^{99m}Tc-HMPAO を投与し脳血流の変化を観察した。起立負荷法でICO群では、対側比の低下がみられ、脳血流異常の領域の検出能を向上させた。また、亜急性期あるいは術後脳梗塞例では、血圧の下降に伴い^{99m}Tc-HMPAO の不均等な分布がみられた。これらの領域は、脳灌流圧の低下に伴う血流の異常な反応、Dysautoregulation の領域と解される⁸⁾。このようにベッドサイド^{99m}Tc-HMPAO 標識法にて脂溶性分画を有効に利用し⁹⁾、^{99m}Tc-HMPAO 脳

SPECT に起立負荷法を併用することで脳血流異常の検出能が向上し、脳灌流圧の変化をも観察しうる。

文 献

- 1) Neirinckx RD, Canning LR, Piper IM, et al: Technetium-99m d,l-HM-PAO: A new pharmaceutical for SPECT imaging of regional cerebral blood perfusion. J Nucl Med 28: 191-202, 1987
- 2) Sharp PF, Smith FW, Gemmell HG, et al: Technetium-99m HM-PAO stereo-isomers as potential agents for imaging regional cerebral blood flow: Human volunteer studies. J Nucl Med 27: 171-177, 1986
- 3) Sullivan HG, Kingsbury TB, Morgan ME, et al: The rCBF response to diamox in normal subjects and cerebrovascular disease patients. J Neurosurg 67: 525-534, 1987
- 4) Herold S, Brown MM, Frackowiak RSL, et al: Assessment of cerebral haemodynamic reserve: Correlation between PET parameters and CO₂ reactivity measured by the intravenous ¹³³Xenon technique. J Neurol Neurosur Psychiatry 51: 1045-1050, 1988
- 5) Matsuda H, Higashi S, Terada H, et al: Evaluation of cerebral collateral circulation by technetium-99m HM-PAO brain SPECT during Natas test: Report on three cases. J Nucl Med 29: 1724-1729, 1988
- 6) Monsein LH, Jeffrey PJ, Van Heerden BB, et al: Assessing adequacy of collateral circulation during balloon test occlusion of the internal carotid artery with ^{99m}Tc-HMPAO SPECT. AJNR 12: 1045-1051, 1991
- 7) Biersack HJ, Linke D, Brassel F, et al: Tc-99m HM-PAO brain SPECT in epileptic patients before and during unilateral hemisphere anesthesia (Wada Test): Report of three cases. J Nucl Med 28: 1763-1767, 1987
- 8) Power WJ, Grubb RL, Raichile ME: Physiological responses to focal cerebral ischemia in human. Ann Neurol 16: 546-552, 1984
- 9) 林田孝平, 西村恒彦, 植原敏勇, 他: ^{99m}Tc-HMPAO の脳血流シンチグラフィーとしての臨床的有用性の検討, 核医学, 25: 231-241, 1988