



Title	A LARGE DEVIATION PRINCIPLE FOR SYMMETRIC MARKOV PROCESSES NORMALIZED BY FEYNMAN-KAC FUNCTIONALS
Author(s)	Takeda, Masayoshi; Tawara, Yoshihiro
Citation	Osaka Journal of Mathematics. 2013, 50(2), p. 287-307
Version Type	VoR
URL	https://doi.org/10.18910/25087
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

A LARGE DEVIATION PRINCIPLE FOR SYMMETRIC MARKOV PROCESSES NORMALIZED BY FEYNMAN–KAC FUNCTIONALS

MASAYOSHI TAKEDA and YOSHIHIRO TAWARA

(Received January 24, 2011, revised July 12, 2011)

Abstract

We establish a large deviation principle for the occupation distribution of a symmetric Markov process normalized by Feynman–Kac functional. The obtained theorem means a large deviation from a ground state, not from an invariant measure.

1. Introduction

Let $\mathbf{M} = (\Omega, X_t, \mathbb{P}_x, \zeta)$ be an m -symmetric irreducible Markov process on a locally compact separable metric space X . Here ζ is the lifetime and m is a positive Radon measure with full support. Let $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ be the Dirichlet form on $L^2(X; m)$ generated by \mathbf{M} (for the definition, see (2.1)). We denote by \mathcal{P} the set of probability measures with the weak topology, and for a positive Green-tight Kato measure μ (Definition 2.1) define the function I^μ on the set \mathcal{P} by

$$(1.1) \quad I^\mu(v) = \begin{cases} \mathcal{E}^\mu(\sqrt{f}, \sqrt{f}) & \text{if } v = f \cdot m, \sqrt{f} \in \mathcal{D}(\mathcal{E}), \\ \infty & \text{otherwise,} \end{cases}$$

where $\mathcal{E}^\mu = \mathcal{E} - (\cdot, \cdot)_\mu$. Given $\omega \in \Omega$ with $0 < t < \zeta(\omega)$, let $L_t(\omega) \in \mathcal{P}$ be the normalized occupation distribution: for a Borel set A of X

$$L_t(\omega)(A) = \frac{1}{t} \int_0^t 1_A(X_s(\omega)) ds,$$

where 1_A is the indicator function of the set A . We denote by A_t^μ the positive continuous additive functional with Revuz measure μ . One of authors proved Donsker–Varadhan type large deviation principle with rate function I^μ .

Theorem 1.1 ([24]). *Assume that the Markov process \mathbf{M} possesses the strong Feller property and the tightness property (see (III) in Section 2).*

(i) For each open set $G \subset \mathcal{P}$

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_x(e^{A_t^\mu}; L_t \in G, t < \zeta) \geq -\inf_{v \in G} I^\mu(v).$$

(ii) For each closed set $K \subset \mathcal{P}$

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \sup_{x \in X} \mathbb{E}_x(e^{A_t^\mu}; L_t \in K, t < \zeta) \leq -\inf_{v \in K} I^\mu(v).$$

Varadhan [29] gave an abstract formulation for the large deviation principle. The statement in Theorem 1.1 is slightly different from his formulation. In fact, the rate function I^μ is not always non-negative because it is defined by the Schrödinger form \mathcal{E}^μ , not by the Dirichlet form \mathcal{E} . Furthermore, Theorem 1.1 does not represent a large deviation from a invariant measure because the Markov process is allowed to be explosive. By this reason, we consider the normalized probability measure $Q_{x,t}$ on \mathcal{P} defined by, for a Borel set $B \subset \mathcal{P}$,

$$Q_{x,t}(B) = \frac{\mathbb{E}_x(e^{A_t^\mu}; L_t \in B, t < \zeta)}{\mathbb{E}_x(e^{A_t^\mu}; t < \zeta)},$$

and prove that the family of probability measures $\{Q_{x,t}\}_{t>0}$ obeys the large deviation principle as $t \rightarrow \infty$ in the sense of Varadhan's formulation. In other words, $\{Q_{x,t}\}_{t>0}$ satisfies the *full* large deviation principle with a *good* rate function in the sense of [11, Section 2.1]. This is the main theorem of this paper (Theorem 4.1). The rate function is given by

$$(1.2) \quad J(v) := I^\mu(v) - \lambda_2(\mu), \quad v \in \mathcal{P}.$$

Here $\lambda_2(\mu)$ is the bottom of the spectrum of the Schrödinger type operator $\mathcal{L} + \mu$, where \mathcal{L} is the generator of the Markov process:

$$\lambda_2(\mu) = \inf\{\mathcal{E}^\mu(u, u) : u \in \mathcal{D}(\mathcal{E}), \|u\|_2 = 1\}.$$

To obtain the main theorem, we need to show that the rate function J is good, that is, enjoys the properties (i)–(iv) in Lemma 4.1. In particular, we must show that J has a unique zero point, that is, the existence of a ground state ϕ_0 of the operator $\mathcal{L} + \mu$. In order to show the existence of a ground state, we usually use the L^2 -weak compactness of the set $\{u \in \mathcal{D}(\mathcal{E}) : \mathcal{E}^\mu(u, u) \leq l\}$ ($l \in \mathbb{R}$) and the lower semi-continuity of the Schrödinger form \mathcal{E}^μ with respect to the L^2 -weak topology (e.g. [17]); however we can not derive these properties from our general setting. Hence we here use the following properties instead, the tightness of the level set $\{v \in \mathcal{P} : I^\mu(v) \leq l\}$ and the lower semi-continuity of the function I^μ with respect to the weak topology. This is a key to the proof of the

goodness of the rate function J . We would like to emphasize that the tightness follows from the condition (III) and the Green-tightness of μ , and the lower semi-continuity of I^μ follows from a variational formula for the Schrödinger form (Proposition 2.1), that is, the identification of the Schrödinger form with the modified I -function defined in (2.8). The latter is an extension of a well-known fact due to Donsker and Varadhan that for a symmetric Markov process, the I -function is identical with the Dirichlet form. On account of Lemma 4.1, we can regard the main theorem as a large deviation from the ground state of the Schrödinger operator.

In [10], [25], [28], L^p -independence of growth bounds of non-local Feynman–Kac semigroups have been considered. In this paper we also deal with non-local Feynman–Kac transforms and extended Theorem 1.1 to symmetric Markov processes with non-local Feynman–Kac functional (Theorem 2.1). The existence of ground states implies the existence of a quasi-stationary distribution, $\eta(B) := \int_B \phi_0(x) dm(x) / \int_X \phi_0(x) dm(x)$. In [16], they prove that if a Markov semigroup is intrinsically ultracontractive, then the measure η is the so-called Yaglom limit and a unique quasi-stationary distribution. In the last section, we will give an extension of this fact to generalized Feynman–Kac semigroups by employing Fukushima’s ergodic theorem.

2. Symmetric Markov processes with non-local Feynman–Kac functionals

Let X be a locally compact separable metric space and $\mathcal{B}(X)$ the Borel σ -field. Adjoining an extra point ∞ to the measurable set $(X, \mathcal{B}(X))$, we set $X_\infty = X \cup \{\infty\}$ and $\mathcal{B}(X_\infty) = \mathcal{B}(X) \cup \{B \cup \{\infty\}: B \in \mathcal{B}(X)\}$. Let $\mathbf{M} = (\Omega, X_t, \mathbb{P}_x, \zeta)$ be a right Markov process on X with lifetime $\zeta := \inf\{t > 0: X_t = \infty\}$. We define the semigroup and the resolvent by

$$p_t f(x) = \mathbb{E}_x(f(X_t); t < \zeta), \quad R_\beta f(x) = \int_0^\infty e^{-\beta t} p_t f(x) dt$$

for a bounded Borel function f on X . We assume that the Markov process \mathbf{M} is m -symmetric, $(p_t f, g)_m = (f, p_t g)_m$, where m is a positive Radon measure with full support. Let $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ be the Dirichlet form on $L^2(X; m)$ generated by \mathbf{M} :

$$(2.1) \quad \begin{cases} \mathcal{D}(\mathcal{E}) = \left\{ u \in L^2(X; m): \lim_{t \rightarrow 0} \frac{1}{t} (u - p_t u, u)_m < \infty \right\}, \\ \mathcal{E}(u, v) = \lim_{t \rightarrow 0} \frac{1}{t} (u - p_t u, v)_m. \end{cases}$$

For basic materials on right processes and associated Dirichlet forms (quasi-regular Dirichlet forms), we refer to [7], [18].

We impose three assumptions on \mathbf{M} .

(I) (*Irreducibility*) If a Borel set A is p_t -invariant, i.e., $p_t(1_A f)(x) = 1_A p_t f(x)$ m -a.e. for any $f \in L^2(X; m) \cap \mathcal{B}_b(X)$ and $t > 0$, then A satisfies either $m(A) = 0$ or $m(X \setminus A) = 0$. Here $\mathcal{B}_b(X)$ is the space of bounded Borel functions on X .

(II) (*Strong Feller property*) For each t , $p_t(\mathcal{B}_b(X)) \subset C_b(X)$, where $C_b(X)$ is the space of bounded continuous functions on X .

(III) (*Tightness*) For any $\epsilon > 0$, there exists a compact set K such that

$$\sup_{x \in X} R_1 1_{K^c}(x) \leq \epsilon.$$

Here 1_{K^c} is the indicator function of the complement of the compact set K .

The assumption (II) implies that \mathbf{M} satisfies the *absolute continuity condition*, that is, its transition probability $p_t(x, \cdot)$ is absolutely continuous with respect to m for each $t > 0$ and $x \in X$. As a result, the resolvent kernel is also absolutely continuous with respect to m , $R_\beta(x, dy) = R_\beta(x, y)m(dy)$. By [14, Lemma 4.2.4] the density $R_\beta(x, y)$ is assumed to be a non-negative Borel function such that $R_\beta(x, y)$ is symmetric and β -excessive in x and in y . Under the absolute continuity condition, “quasi everywhere” statements are strengthened to “everywhere” ones. Moreover, we can define notions without exceptional set, for example, *smooth measures in the strict sense* or *positive continuous additive functional in the strict sense* (cf. [14, Section 5.1]). Here we only treat the notions in the strict sense and omit the phrase “in the strict sense”.

We denote S_{00} the set of positive Borel measures μ such that $\mu(X) < \infty$ and $R_1\mu(x) (= \int_X R_1(x, y)\mu(dy))$ is uniformly bounded in $x \in X$. A positive Borel measure μ on X is said to be *smooth* if there exists a sequence $\{E_n\}_{n=1}^\infty$ of Borel sets increasing to X such that $1_{E_n} \cdot \mu \in S_{00}$ for each n and

$$\mathbb{P}_x \left(\lim_{n \rightarrow \infty} \sigma_{X \setminus E_n} \geq \zeta \right) = 1, \quad \forall x \in X,$$

where $\sigma_{X \setminus E_n}$ is the first hitting time of $X \setminus E_n$. The totality of smooth measures is denoted by S_1 .

If an additive functional $\{A_t\}_{t \geq 0}$ is positive and continuous with respect to t for each $\omega \in \Lambda$, it is said to be a *positive continuous additive functional* (PCAF in abbreviation). By [14, Theorem 5.1.7], there exists a one-to-one correspondence between positive smooth measures and PCAF's (*Revuz correspondence*): for each smooth measure μ , there exists a unique PCAF $\{A_t\}_{t \geq 0}$ such that for any positive Borel function f on X and γ -excessive function h ($\gamma \geq 0$), that is, $e^{-\gamma t} p_t h \leq h$,

$$(2.2) \quad \lim_{t \rightarrow 0} \frac{1}{t} \mathbb{E}_{h \cdot m} \left(\int_0^t f(X_s) dA_s \right) = \int_X f(x) h(x) \mu(dx).$$

Here $\mathbb{E}_{h \cdot m}(\cdot) = \int_X \mathbb{E}_x(\cdot) h(x) m(dx)$. We denote by A_t^μ the PCAF corresponding to the smooth measure μ . For a signed Borel measure $\mu = \mu^+ - \mu^-$, let $|\mu| = \mu^+ + \mu^-$. When $|\mu|$ is a smooth measure, we define $A_t^\mu = A_t^{\mu^+} - A_t^{\mu^-}$ and $A_t^{|\mu|} = A_t^{\mu^+} + A_t^{\mu^-}$.

Following Chen [4], we introduce classes of potentials.

DEFINITION 2.1. (i) A signed Borel measure μ is said to be the *Kato measure* (in notation, $\mu \in \mathcal{K}$), if $|\mu| \in S_1$ and

$$\limsup_{t \rightarrow 0} \sup_{x \in X} \mathbb{E}_x(A_t^{|\mu|}) = 0.$$

(ii) A measure $\mu \in \mathcal{K}$ is said to be in the class \mathcal{K}_∞ , if for any $\epsilon > 0$ there exist a compact subset K and a positive constant $\delta > 0$ such that for all measurable set $B \subset K$ with $|\mu|(B) < \delta$,

$$\sup_{x \in X} \int_{K^c \cup B} R_1(x, y) |\mu|(dy) \leq \epsilon.$$

(iii) A signed Borel measure μ is said to be in the class \mathcal{S}_∞ , if for any $\epsilon > 0$ there exist a compact subset K and a positive constant $\delta > 0$ such that for all measurable set $B \subset K$ with $|\mu|(B) < \delta$,

$$\sup_{(x,z) \in X \times X \setminus d} \int_{K^c \cup B} \frac{R_1(x, y) R_1(y, z)}{R_1(x, z)} |\mu|(dy) \leq \epsilon.$$

It is known in [2] that μ belongs to \mathcal{K} if and only if

$$(2.3) \quad \lim_{\beta \rightarrow \infty} \sup_{x \in X} \int_X R_\beta(x, y) |\mu|(dy) = 0,$$

and in [4] that

$$(2.4) \quad \mathcal{S}_\infty \subset \mathcal{K}_\infty \subset \mathcal{K}.$$

We denote that $(N, H) = (N(x, dy), H_t)$ is the Lévy system of \mathbf{M} , that is, N is a kernel on $(X_\infty, \mathcal{B}(X_\infty))$ with $N(x, \{x\}) = 0$ and H is a positive continuous additive functional of \mathbf{M} such that for any non-negative measurable function F on $X \times X$ vanishing on the diagonal set and any $x \in X$,

$$\mathbb{E}_x \left(\sum_{0 < s \leq t} F(X_{s-}, X_s); t < \zeta \right) = \mathbb{E}_x \left(\int_0^t \int_X F(X_s, y) N(X_s, dy) dH_s \right).$$

We denote by μ^H be the smooth measure corresponding to H_t .

DEFINITION 2.2. Let F be a bounded measurable function on $X \times X$ vanishing on the diagonal set.

(i) F is said to be in the class \mathcal{A}_∞ , if for any $\epsilon > 0$ there exist a compact subset K and a positive constant $\delta > 0$ such that for all measurable set $B \subset K$ with $|\mu|(B) < \delta$,

$$\sup_{(x,z) \in X \times X \setminus d} \int_{((K \setminus B) \times (K \setminus B))^c} \frac{R_1(x, y) |F(y, z)| R_1(z, w)}{R_1(x, w)} N(y, dz) \mu^H(dy) \leq \epsilon.$$

(ii) F is said to be in the class \mathcal{A}_2 , if $F \in \mathcal{A}_\infty$ and

$$\mu_{|F|}(dx) = \left(\int_X |F(x, y)| N(x, dy) \right) \mu^H(dx) \in \mathcal{S}_\infty.$$

For properties and examples of \mathcal{A}_∞ and \mathcal{A}_2 , see [4], [5]. In the remainder of this paper, we assume that F is symmetric, $F(x, y) = F(y, x)$. We write $\mu + F \in \mathcal{K}_\infty + \mathcal{A}_2$ if $\mu \in \mathcal{K}_\infty$ and $F \in \mathcal{A}_2$.

For $\mu + F \in \mathcal{K}_\infty + \mathcal{A}_2$ define the AF $A_t^{\mu+F}$ by

$$A_t^{\mu+F} = A_t^\mu + \sum_{0 < s \leq t} F(X_{s-}, X_s),$$

and the generalized Feynman–Kac semigroup $\{p_t^{\mu+F}\}_{t \geq 0}$ by

$$p_t^{\mu+F} f(x) = \mathbb{E}_x \left(e^{A_t^{\mu+F}} f(X_t); t < \zeta \right), \quad f \in \mathcal{B}_b(X).$$

For $F \in \mathcal{A}_2$, we define the symmetric Dirichlet form $(\mathcal{E}_F, \mathcal{D}(\mathcal{E}))$ as follows: for $u, v \in \mathcal{D}(\mathcal{E})$

$$(2.5) \quad \begin{aligned} \mathcal{E}_F(u, v) &= \mathcal{E}^{(c)}(u, v) + \mathcal{E}^{(k)}(u, v) \\ &+ \frac{1}{2} \int_{X \times X} (u(x) - u(y))(v(x) - v(y)) e^{F(x, y)} N(x, dy) \mu^H(dx), \end{aligned}$$

where $\mathcal{E}^{(c)}$ and $\mathcal{E}^{(k)}$ are the local part and the killing part of the Dirichlet form $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ in Beurling–Deny formula ([14, Theorem 3.2.1]). Fundamental properties of non-local Feynman–Kac transforms were earlier studied by J. Ying [31], [32]. It is known in [8] that $\{p_t^{\mu+F}\}_{t \geq 0}$ is the semigroup generated by the Schrödinger form $(\mathcal{E}^{\mu+F}, \mathcal{D}(\mathcal{E}))$:

$$(2.6) \quad \mathcal{E}^{\mu+F}(u, v) = \mathcal{E}_F(u, v) - \int_X u(x)v(x) d\mu_{F_1}(x) - \int_X u(x)v(y) d\mu(x),$$

where $F_1 = \exp(F) - 1$. The form $\mathcal{E}^{\mu+F}$ is also written as

$$\begin{aligned} \mathcal{E}^{\mu+F}(u, v) &= \mathcal{E}(u, v) - \int_{X \times X} u(x)v(y) F_1(x, y) N(x, dy) d\mu^H(y) \\ &- \int_X u(x)v(x) d\mu(x), \quad u, v \in \mathcal{D}(\mathcal{E}). \end{aligned}$$

Let \mathcal{P} be the set of probability measures on X equipped with the weak topology. We define the function $I^{\mu+F}$ on \mathcal{P} by

$$I^{\mu+F}(\nu) = \begin{cases} \mathcal{E}^{\mu+F}(\sqrt{f}, \sqrt{f}) & \text{if } \nu = f \cdot m, \sqrt{f} \in \mathcal{D}(\mathcal{E}), \\ \infty & \text{otherwise.} \end{cases}$$

Let $\mu + F \in \mathcal{K}_\infty + \mathcal{A}_2$ and define $\kappa(\mu + F)$ by

$$\kappa(\mu + F) = \lim_{t \rightarrow \infty} \frac{1}{t} \log \|p_t^{\mu+F}\|_{\infty, \infty}.$$

We see from [1] that $\kappa(\mu + F)$ is finite. If $\alpha > \kappa(\mu + F)$ and $f \in \mathcal{B}_b(X)$, we define the resolvent $R_\alpha^{\mu+F}$ by

$$R_\alpha^{\mu+F} f(x) = \mathbb{E}_x \left(\int_0^\infty e^{-\alpha t + A_t^{\mu+F}} f(X_t) dt \right).$$

We set

$$\mathcal{D}_+(\mathcal{H}^{\mu+F}) = \{R_\alpha^{\mu+F} f : \alpha > \kappa(\mu + F), f \in L^2(X; m) \cap C_b(X), f \geq 0 \text{ and } f \not\equiv 0\}.$$

Each function $\phi = R_\alpha^{\mu+F} f \in \mathcal{D}_+(\mathcal{H}^{\mu+F})$ is strictly positive because $\mathbb{P}_x(\sigma_O < \zeta) > 0$ for any $x \in X$ by the assumption (I). Here O is a non-empty open set $\{x \in X : f(x) > 0\}$ and $\sigma_O = \inf\{t > 0 : X_t \in O\}$. We define the generator $\mathcal{H}^{\mu+F}$ by

$$\mathcal{H}^{\mu+F} u = \alpha u - f, \quad u = R_\alpha^{\mu+F} f \in \mathcal{D}_+(\mathcal{H}^{\mu+F}).$$

Let h be the function defined by $h(x) = \mathbb{E}_x(\exp(A_\zeta^{\mu+F}))$. We may assume that $\mu + F$ is gaugeable, that is, $\sup_{x \in X} h(x) < \infty$. In fact, it is enough to prove Theorem 2.1 and Theorem 4.1 below for the β -subprocess, $\mathbb{P}_x^{(\beta)} = e^{-\beta t} \mathbb{P}_x$. Moreover, we see that every $\mu + F \in \mathcal{K}_\infty + \mathcal{A}_2$ becomes gaugeable with respect to the β -subprocess of \mathbf{M} for a large enough β . In fact, we see from [5, Theorem 3.4] that $\mu + F \in \mathcal{K}_\infty + \mathcal{A}_2$ is gaugeable with respect to the β -subprocess if and only if

$$(2.7) \quad \inf \left\{ \mathcal{E}_F(u, u) + \int_X u(x)^2 (\mu^- + \mu_{F_1^-})(dx) + \beta \int_X u(x)^2 m(dx) : \int_X u(x)^2 (\mu^+ + \mu_{F_1^+})(dx) = 1 \right\} > 1,$$

where F_1^+ and F_1^- is the positive and negative part of F_1 . Since by (3.1)

$$\begin{aligned} & \mathcal{E}_F(u, u) + \int_X u(x)^2 (\mu^- + \mu_{F_1^-})(dx) + \beta \int_X u(x)^2 m(dx) \\ & \geq e^{-\|F^-\|_\infty} \left(\mathcal{E}(u, u) + \beta \int_X u(x)^2 m(dx) \right) \geq \frac{e^{-\|F^-\|_\infty}}{\|R_\beta(\mu^+ + \mu_{F_1^+})\|_\infty}, \end{aligned}$$

and the right hand side tends to ∞ as $\beta \rightarrow \infty$ because of $\mu^+ + \mu_{F_1^+} \in \mathcal{K}$, (2.7) holds for a large β .

We define the function I_h on \mathcal{P} by

$$(2.8) \quad I_h(v) = - \inf_{\substack{\phi \in \mathcal{D}_+(\mathcal{H}^{\mu+F}) \\ \epsilon > 0}} \int_X \frac{\mathcal{H}^{\mu+F}\phi}{\phi + \epsilon h} d\nu.$$

The gauge function $h(x)$ satisfies $0 < c \leq h(x) \leq C < \infty$. Indeed, it follows from Proposition 2.2 in [4] and (2.4) that for $\mu \in \mathcal{K}_\infty$ and $F \in \mathcal{A}_2$, $\sup_{x \in X} \mathbb{E}_x(A_\zeta^{|\mu|+|F|}) < \infty$. Hence, by Jensen's inequality,

$$\inf_{x \in X} \mathbb{E}_x(\exp(A_\zeta^{\mu+F})) \geq \exp\left(-\sup_{x \in X} \mathbb{E}_x(A_\zeta^{|\mu|+|F|})\right) > 0.$$

Let us define the function I_α on \mathcal{P} by

$$I_\alpha(v) = - \inf_{\substack{u \in \mathcal{B}_b^+(X) \\ \epsilon > 0}} \int_X \log\left(\frac{\alpha R_\alpha^{\mu+F}u + \epsilon h}{u + \epsilon h}\right) d\nu$$

Lemma 2.1. *It holds that*

$$I_\alpha(v) \leq \frac{I_h(v)}{\alpha}, \quad v \in \mathcal{P}.$$

Proof. For $u = R_\alpha^{\mu+F}f \in \mathcal{D}_+(\mathcal{H}^{\mu+F})$ and $\epsilon > 0$, set

$$\phi(\alpha) = - \int_X \log\left(\frac{\alpha R_\alpha^{\mu+F}u + \epsilon h}{u + \epsilon h}\right) d\nu.$$

Then, noting that $(d/d\alpha)(R_\alpha^{\mu+F}u) = -R_\alpha^{\mu+F}(R_\alpha^{\mu+F}u) = -(R_\alpha^{\mu+F})^2u$, we have

$$\frac{d\phi}{d\alpha}(\alpha) = - \int_X \frac{R_\alpha^{\mu+F}u - \alpha(R_\alpha^{\mu+F})^2u}{\alpha R_\alpha^{\mu+F}u + \epsilon h} d\nu = \int_X \frac{\mathcal{H}^{\mu+F}(R_\alpha^{\mu+F})^2u}{\alpha R_\alpha^{\mu+F}u + \epsilon h} d\nu.$$

Since

$$\begin{aligned} & (\alpha(R_\alpha^{\mu+F})^2u - R_\alpha^{\mu+F}u)(\alpha^2(R_\alpha^{\mu+F})^2u + \epsilon h) \\ & - (\alpha(R_\alpha^{\mu+F})^2u - R_\alpha^{\mu+F}u)(\alpha R_\alpha^{\mu+F}u + \epsilon h) \end{aligned}$$

equals $\alpha(\alpha(R_\alpha^{\mu+F})^2u - R_\alpha^{\mu+F}u)^2 \geq 0$, we have

$$\frac{\alpha(R_\alpha^{\mu+F})^2u - R_\alpha^{\mu+F}u}{\alpha R_\alpha^{\mu+F}u + \epsilon h} \geq \frac{\alpha(R_\alpha^{\mu+F})^2u - R_\alpha^{\mu+F}u}{\alpha^2(R_\alpha^{\mu+F})^2u + \epsilon h},$$

and thus

$$\begin{aligned}
\int_X \frac{\mathcal{H}^{\mu+F}(R_\alpha^{\mu+F})^2 u}{\alpha R_\alpha^{\mu+F} u + \epsilon h} d\nu &\geq \int_X \frac{\mathcal{H}^{\mu+F}(R_\alpha^{\mu+F})^2 u}{\alpha^2 (R_\alpha^{\mu+F})^2 u + \epsilon h} d\nu \\
&= -\frac{1}{\alpha^2} \left(- \int_X \frac{\mathcal{H}^{\mu+F}(R_\alpha^{\mu+F})^2 u}{(R_\alpha^{\mu+F})^2 u + \epsilon h/\alpha^2} d\nu \right) \\
&\geq -\frac{1}{\alpha^2} I_h(\nu).
\end{aligned}$$

Therefore

$$\phi(\infty) - \phi(\alpha) = \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} u + \epsilon h}{u + \epsilon h} \right) d\nu \geq -\frac{I_h(\nu)}{\alpha},$$

which implies

$$-\inf_{\substack{u \in \mathcal{D}_+(\mathcal{H}^{\mu+F}) \\ \epsilon > 0}} \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} u + \epsilon h}{u + \epsilon h} \right) d\nu \leq \frac{I_h(\nu)}{\alpha}.$$

Since $\|\beta R_\beta^{\mu+F} f\|_\infty \leq C \|f\|_\infty$, $\beta > 0$, and $\beta R_\beta^{\mu+F} f(x) \rightarrow f(x)$ as $\beta \rightarrow \infty$,

$$(2.9) \quad \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F}(\beta R_\beta^{\mu+F} f) + \epsilon h}{\beta R_\beta^{\mu+F} f + \epsilon h} \right) d\nu \xrightarrow{\beta \rightarrow \infty} \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} f + \epsilon h}{f + \epsilon h} \right) d\nu.$$

Define the measure ν_α by

$$\nu_\alpha(A) = \int_X \alpha R_\alpha^{\mu+F}(x, A) d\nu(x), \quad A \in \mathcal{B}(X).$$

Given $\nu \in \mathcal{B}_b^+(X)$, take a sequence $\{g_n\}_{n=1}^\infty \subset C_b^+(X) \cap L^2(X; m)$ such that

$$\int_X |v - g_n| d(\nu_\alpha + \nu) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

We then have

$$\int_X |\alpha R_\alpha^{\mu+F} v - \alpha R_\alpha^{\mu+F} g_n| d\nu \leq \int_X \alpha R_\alpha^{\mu+F}(|v - g_n|) d\nu = \int_X |v - g_n| d\nu_\alpha \rightarrow 0$$

as $n \rightarrow \infty$, and so

$$(2.10) \quad \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} g_n + \epsilon h}{g_n + \epsilon h} \right) d\nu \xrightarrow{n \rightarrow \infty} \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} v + \epsilon h}{v + \epsilon h} \right) d\nu.$$

Hence, combining (2.9) and (2.10)

$$\inf_{u \in \mathcal{D}_+(\mathcal{H}^{\mu+F})} \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} u + \epsilon h}{u + \epsilon h} \right) d\nu = \inf_{u \in \mathcal{B}_b^+(X)} \int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} u + \epsilon h}{u + \epsilon h} \right) d\nu,$$

which implies the lemma. \square

Lemma 2.2. *If $I_h(\nu) < \infty$, then ν is absolutely continuous with respect to m .*

Proof. By a similar argument in the proof of [12, Lemma 4.1], we obtain this lemma. Indeed, for $a > 0$ and $A \in \mathcal{B}(X)$, set $u(x) = a1_A(x) + 1 \in \mathcal{B}_b^+(X)$. Then

$$\int_X \log \left(\frac{\alpha R_\alpha^{\mu+F} u + \epsilon h}{u + \epsilon h} \right) d\nu = \int_X \log \left(\frac{a\alpha R_\alpha^{\mu+F}(x, A) + \alpha R_\alpha^{\mu+F}(x, X) + \epsilon h}{a1_A(x) + 1 + \epsilon h} \right) d\nu.$$

Define the measure ν_α as in the proof of Lemma 2.1. Put

$$c_\alpha = \int_X \alpha R_\alpha^{\mu+F}(x, X) d\nu(x) \quad (= \nu_\alpha(X)).$$

We see from Lemma 2.1 and Jensen's inequality that

$$\log(a\nu_\alpha(A) + c_\alpha + \epsilon h) \geq \nu(A) \log(a + 1 + \epsilon h) + \nu(A^c)(1 + \epsilon h) - \frac{I_h(\nu)}{\alpha},$$

and by letting $\epsilon \rightarrow 0$

$$\log(a\nu_\alpha(A) + c_\alpha) \geq \nu(A) \log(a + 1) - \frac{I_h(\nu)}{\alpha}.$$

Since $\log x \leq x - 1$ for $x > 0$, we have

$$a\nu_\alpha(A) + c_\alpha - 1 \geq \nu(A) \log(a + 1) - \frac{I_h(\nu)}{\alpha},$$

and so

$$\nu_\alpha(A) - \nu(A) \geq \frac{-I_h(\nu)/\alpha + \nu(A)(\log(a + 1) - a) + 1 - c_\alpha}{a}.$$

Noting that $\log(a + 1) - a < 0$, we have

$$\nu_\alpha(A) - \nu(A) \geq \frac{-I_h(\nu)/\alpha + (\log(a + 1) - a) + 1 - c_\alpha}{a}$$

for all $A \in \mathcal{B}(X)$ and

$$\begin{aligned} \nu(A) - \nu_\alpha(A) &= 1 - c_\alpha + (\nu_\alpha(A^c) - \nu(A^c)) \\ &\geq \frac{-I_h(\nu)/\alpha + (\log(a + 1) - a) + (1 - c_\alpha)(a + 1)}{a} \end{aligned}$$

for all $A \in \mathcal{B}(X)$. Therefore we can conclude that

$$\sup_{A \in \mathcal{B}(X)} |\nu(A) - \nu_\alpha(A)| \leq \frac{a - \log(a+1) + I_h(\nu)/\alpha + (1-c_\alpha)(a+1)}{a}.$$

Note that $c_\alpha \rightarrow 1$ as $\alpha \rightarrow \infty$. Then since

$$\limsup_{\alpha \rightarrow \infty} \sup_{A \in \mathcal{B}(X)} |\nu(A) - \nu_\alpha(A)| \leq \frac{a - \log(a+1)}{a}$$

and the right-hand side converges to 0 as $a \rightarrow 0$, the lemma follows. \square

Proposition 2.1. *It holds that for $\nu \in \mathcal{P}$*

$$I_h(\nu) = I^{\mu+F}(\nu).$$

Proof. We follow the argument of the proof of [12, Theorem 5]. Suppose that $I_h(\nu) = l < \infty$. By Lemma 2.2, ν is absolutely continuous with respect to m . Let us denote by f its density and let $f^n = \sqrt{f} \wedge n$. Since $\log(1-x) \leq -x$ for $-\infty < x < 1$ and

$$-\infty < \frac{f^n - \alpha R_\alpha^{\mu+F} f^n}{f^n + \epsilon h} < 1,$$

we have

$$\begin{aligned} \int_X \log\left(\frac{\alpha R_\alpha^{\mu+F} f^n + \epsilon h}{f^n + \epsilon h}\right) f \, dm &= \int_X \log\left(1 - \frac{f^n - \alpha R_\alpha^{\mu+F} f^n}{f^n + \epsilon h}\right) f \, dm \\ &\leq - \int_X \frac{f^n - \alpha R_\alpha^{\mu+F} f^n}{f^n + \epsilon h} f \, dm, \end{aligned}$$

and then

$$\int_X \frac{f^n - \alpha R_\alpha^{\mu+F} f^n}{f^n + \epsilon h} f \, dm \leq I_\alpha(f \cdot m).$$

By letting $n \rightarrow \infty$ and $\epsilon \rightarrow 0$, we have

$$\int_X \sqrt{f}(\sqrt{f} - \alpha R_\alpha^{\mu+F} \sqrt{f}) \, dm \leq I_\alpha(f \cdot m) \leq \frac{I_h(f \cdot m)}{\alpha},$$

which implies that $\sqrt{f} \in \mathcal{D}(\mathcal{E})$ and $\mathcal{E}^{\mu+F}(\sqrt{f}, \sqrt{f}) \leq I_h(f \cdot m)$.

Let $\phi \in \mathcal{D}_+(\mathcal{H}^{\mu+F})$ and define the semigroup P_t^ϕ by

$$P_t^\phi f(x) = \mathbb{E}_x \left(e^{A_t^{\mu+F}} \frac{(\phi + \epsilon h)(X_t)}{(\phi + \epsilon h)(X_0)} \exp\left(-\int_0^t \frac{\mathcal{H}^{\mu+F}\phi}{\phi + \epsilon h}(X_s) ds\right) f(X_t) \right).$$

Then, P_t^ϕ is $(\phi + \epsilon h)^2 m$ -symmetric and satisfies $P_t^\phi 1 \leq 1$. Given $v = f \cdot m \in \mathcal{P}$ with $\sqrt{f} \in \mathcal{D}(\mathcal{E})$, set

$$S_t^\phi \sqrt{f}(x) = \mathbb{E}_x \left(e^{A_t^{\mu+F}} \exp \left(- \int_0^t \frac{\mathcal{H}^{\mu+F} \phi}{\phi + \epsilon h}(X_s) ds \right) \sqrt{f}(X_t) \right).$$

Then

$$\begin{aligned} \int_X (S_t^\phi \sqrt{f})^2 dm &= \int_X (\phi + \epsilon h)^2 \left(P_t^\phi \left(\frac{\sqrt{f}}{\phi + \epsilon h} \right) \right)^2 dm \\ &\leq \int_X (\phi + \epsilon h)^2 P_t^\phi \left(\left(\frac{\sqrt{f}}{\phi + \epsilon h} \right)^2 \right) dm \\ &\leq \int_X (\phi + \epsilon h)^2 \left(\frac{\sqrt{f}}{\phi + \epsilon h} \right)^2 dm \\ &= \int_X f dm. \end{aligned}$$

Hence

$$0 \leq \lim_{t \rightarrow 0} \frac{1}{t} (\sqrt{f} - S_t^\phi \sqrt{f}, \sqrt{f})_m = \mathcal{E}^{\mu+F}(\sqrt{f}, \sqrt{f}) + \int_X \frac{\mathcal{H}^{\mu+F} \phi}{\phi + \epsilon h} f dm,$$

and thus $\mathcal{E}^{\mu+F}(\sqrt{f}, \sqrt{f}) \geq I_h(f \cdot m)$. \square

We now obtain a generalization of Theorem 1.1 in exactly the same way as the proof of it (cf. [10], [28]):

Theorem 2.1 ([24]). *Assume (I), (II) and (III). Suppose that $\mu + F \in \mathcal{K}_\infty + \mathcal{A}_2$.*

(i) *For each open set $G \subset \mathcal{P}$*

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_x (e^{A_t^{\mu+F}}; L_t \in G, t < \zeta) \geq - \inf_{v \in G} I^{\mu+F}(v).$$

(ii) *For each closed set $K \subset \mathcal{P}$*

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \sup_{x \in X} \mathbb{E}_x (e^{A_t^{\mu+F}}; L_t \in K, t < \zeta) \leq - \inf_{v \in K} I^{\mu+F}(v).$$

3. The existence of ground states

We first recall an inequality ([19]): for $\mu \in \mathcal{K}$,

$$(3.1) \quad \int_X \tilde{u}^2 d\mu \leq \|R_\alpha \mu\|_\infty (\mathcal{E}(u, u) + \alpha(u, u)_m), \quad u \in \mathcal{D}(\mathcal{E}).$$

Let $\lambda_2(\mu + F)$ be the bottom of the spectrum of $\mathcal{H}^{\mu+F}$:

$$(3.2) \quad \lambda_2(\mu + F) = \inf\{\mathcal{E}^{\mu+F}(u, u) : u \in \mathcal{D}(\mathcal{E}), \|u\|_2 = 1\}.$$

Proposition 3.1. *Assume (I), (II) and (III). There exists a unique ground state $\phi_0 \in \mathcal{D}(\mathcal{E})$: $\lambda_2(\mu + F) = \mathcal{E}^{\mu+F}(\phi_0, \phi_0)$.*

Proof. Let $\{u_n\}$ be a minimizing sequence of the right-hand side of (3.2), i.e., $\|u_n\|_2 = 1$ and $\lambda_2(\mu + F) = \lim_{n \rightarrow \infty} \mathcal{E}^{\mu+F}(u_n, u_n)$. Put $\mu' = |\mu| + |\mu_{F_1}|$. Since $\mathcal{E}(u_n, u_n) \leq c \cdot \mathcal{E}_F(u_n, u_n)$ ($c = \exp(-\|F\|_\infty)$) and $\int_X u_n^2 d\mu' \leq \|R_\alpha \mu'\|_\infty \cdot (\mathcal{E}(u_n, u_n) + \alpha)$,

$$\begin{aligned} \mathcal{E}^{\mu+F}(u_n, u_n) &= \mathcal{E}_F(u, u) - \int_X u_n^2 d\mu' \\ &\geq \frac{1}{c} \mathcal{E}(u_n, u_n) - \|R_\alpha \mu'\|_\infty (\mathcal{E}(u_n, u_n) + \alpha) \\ &= \left(\frac{1}{c} - \|R_\alpha \mu'\|_\infty \right) \mathcal{E}(u_n, u_n) - \alpha \|R_\alpha \mu'\|_\infty. \end{aligned}$$

Taking α large enough so that $c \|R_\alpha \mu'\|_\infty < 1$ on account of (2.3), we have

$$\sup_n \mathcal{E}(u_n, u_n) \leq \frac{c(\sup_n \mathcal{E}^{\mu+F}(u_n, u_n) + \alpha \|R_\alpha \mu'\|_\infty)}{1 - c \|R_\alpha \mu'\|_\infty} < \infty.$$

We see from the assumption (III) that for any $\epsilon > 0$ there exists a compact set K such that

$$\sup_n \int_{K^c} u_n^2 dm \leq \|R_1 1_{K^c}\|_\infty \cdot \left(\sup_n \mathcal{E}(u_n, u_n) + 1 \right) < \epsilon.$$

As a result, the subset $\{u_n^2 \cdot m\}$ of \mathcal{P} is tight. Hence there exists a subsequence $u_{n_k}^2 \cdot m$ which converges to a probability measure ν weakly. Since the function $I^{\mu+F}$ is lower semi-continuous by Proposition 2.1,

$$I^{\mu+F}(\nu) \leq \liminf_{k \rightarrow \infty} I^{\mu+F}(u_{n_k}^2 \cdot m) = \liminf_{k \rightarrow \infty} \mathcal{E}^{\mu+F}(u_{n_k}, u_{n_k}) < \infty.$$

Therefore ν can be written as $\nu = \phi_0^2 m$, $\phi_0 \in \mathcal{D}(\mathcal{E})$ by Proposition 2.1 and $\lambda_2(\mu + F) = \mathcal{E}^{\mu+F}(\phi_0, \phi_0)$, that is, ϕ_0 is the ground state. The uniqueness of the ground state follows from the irreducibility (I) (e.g. [9, Proposition 1.4.3]). \square

We also know from the proof above that the level set $\{\nu \in \mathcal{P} : I^{\nu+F}(\nu) \leq l\}$ is compact.

4. Large deviations from ground states

Given $\omega \in \Omega$ with $0 < t < \zeta(\omega)$, we define the occupation distribution $L_t(\omega) \in \mathcal{P}$ by

$$L_t(\omega)(A) = \frac{1}{t} \int_0^t 1_A(X_s(\omega)) ds$$

for a Borel set A of X , where 1_A is the indicator function of the set A .

Define the probability measure $Q_{x,t}$ on \mathcal{P} by

$$(4.1) \quad Q_{x,t}(B) = \frac{\mathbb{E}_x(e^{A_t^{\mu+F}}; L_t \in B, t < \zeta)}{\mathbb{E}_x(e^{A_t^{\mu+F}}; t < \zeta)}, \quad B \in \mathcal{B}(\mathcal{P}).$$

We define the function J on \mathcal{P} by

$$(4.2) \quad J(v) = I^{\mu+F}(v) - \lambda_2(\mu + F).$$

We then have the next lemma by Proposition 2.1 and Proposition 3.1.

Lemma 4.1. *The function J satisfies:*

- (i) $0 \leq J(v) \leq \infty$.
- (ii) J is lower semicontinuous.
- (iii) For each $l < \infty$, the set $\{v \in \mathcal{P}: J(v) \leq l\}$ is compact.
- (iv) $J(\phi_0^2 \cdot m) = 0$ and $J(v) > 0$ for $v \neq \phi_0^2 \cdot m$.

REMARK 4.1. Let $(\mathcal{E}^{\phi_0}, \mathcal{D}(\mathcal{E}^{\phi_0}))$ the bilinear form on $L^2(X; \phi_0^2 m)$ defined by

$$\begin{cases} \mathcal{E}^{\phi_0}(u, v) = \mathcal{E}^{\mu+F}(u\phi_0, u\phi_0) - \lambda_2(\mu + F)(u\phi_0, u\phi_0)_m, \\ \mathcal{D}(\mathcal{E}^{\phi_0}) = \{u \in L^2(X; \phi_0^2 m): u\phi_0 \in \mathcal{D}(\mathcal{E})\}. \end{cases}$$

We then see that $(\mathcal{E}^{\phi_0}, \mathcal{D}(\mathcal{E}^{\phi_0}))$ is a Dirichlet form and \mathcal{E}^{ϕ_0} is expressed by

$$\mathcal{E}^{\phi_0}(u, v) = \int_X \phi_0^2 d\mu_{\langle u, v \rangle}^c + \int_{X \times X \setminus \Delta} (u(x) - u(y))(v(x) - v(y))\phi_0(x)\phi_0(y)J(dx, dy).$$

Here $\mu_{\langle u, v \rangle}^c$ is the local part of energy measure ([6]). We then see that

$$J(v) = I_{\mathcal{E}^{\phi_0}}(v),$$

where $I_{\mathcal{E}^{\phi_0}}$ is defined by

$$(4.3) \quad I_{\mathcal{E}^{\phi_0}}(v) = \begin{cases} \mathcal{E}^{\phi_0}(\sqrt{f}, \sqrt{f}) & \text{if } v = f \cdot \phi_0^2 m, \sqrt{f} \in \mathcal{D}(\mathcal{E}^{\phi_0}), \\ \infty & \text{otherwise.} \end{cases}$$

We then have the main theorem:

Theorem 4.1. *Assume (I), (II) and (III). Let $\mu + F \in \mathcal{K}_\infty + \mathcal{A}_2$. Let $\{Q_{x,t}\}_{t>0}$ be a family of probability measures defined in (4.1). Then $\{Q_{x,t}\}_{t>0}$ obeys a large deviation principle with rate function J :*

(1) *For each open set $G \subset \mathcal{P}$*

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \log Q_{x,t}(G) \geq - \inf_{v \in G} J(v).$$

(2) *For each closed set $K \subset \mathcal{P}$*

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log Q_{x,t}(K) \leq - \inf_{v \in K} J(v).$$

Corollary 4.1. *The measure $Q_{x,t}$ converges to $\delta_{\phi_0^2 \cdot m}$ weakly.*

Proof. If a closed set K does not contain $\phi_0^2 \cdot m$, then $\inf_{x \in K} J(x) > 0$ by Lemma 4.1 (iv). Hence Theorem 4.1 (ii) says that $\lim_{t \rightarrow \infty} Q_{x,t}(K) = 0$ and $\lim_{t \rightarrow \infty} Q_{x,t}(K^c) = 1$. For a positive constant δ and a bounded continuous function f on the set of \mathcal{P} , define the closed set $K \subset \mathcal{P}$ by $K = \{v \in \mathcal{P} : |f(v) - f(\phi_0^2 \cdot m)| \geq \delta\}$. Then we have

$$\begin{aligned} \left| \int_{\mathcal{P}} f(v) Q_{x,t}(dv) - f(\phi_0^2 \cdot m) \right| &\leq \int_{\mathcal{P}} |f(v) - f(\phi_0^2 \cdot m)| Q_{x,t}(dv) \\ &= \int_K |f(v) - f(\phi_0^2 \cdot m)| Q_{x,t}(dv) + \int_{K^c} |f(v) - f(\phi_0^2 \cdot m)| Q_{x,t}(dv) \\ &\leq \delta Q_{x,t}(K^c) + 2 \|f\|_{\infty} Q_{x,t}(K) \rightarrow \delta \end{aligned}$$

as $t \rightarrow \infty$. Since δ is arbitrary, the weak convergence follows. \square

On account of Corollary 4.1, we can regard Theorem 4.1 as a genuine large deviation principle from the ground state.

5. Quasistationary distribution

In this section, we consider the existence of quasi-stationary distributions as an application of the existence of ground states. We continue with the setting of the preceding section.

Define the semigroup $\{p_t^{\phi_0}\}_{t \geq 0}$ on $L^2(X; \phi_0^2 m)$ generated by $(\mathcal{E}^{\phi_0}, \mathcal{D}(\mathcal{E}^{\phi_0}))$, that is

$$(5.1) \quad p_t^{\phi_0} f(x) = e^{\lambda_2(\mu+F)t} \frac{1}{\phi_0(x)} \mathbb{E}_x \left(e^{A_t^{\mu+F}} \phi_0(X_t) f(X_t) \right).$$

Let $\mathbf{M}^{\phi_0} = (\Omega, X_t, \mathbb{P}_x^{\phi_0})$ be the $\phi_0^2 m$ -symmetric Markov process generated by the Markov semigroup $p_t^{\phi_0}$ in (5.1).

Set

$$\mathcal{P}_0 = \left\{ \nu \in \mathcal{P} : \int_X \sqrt{p_1^{\mu+F}(x, x)} d\nu(x) < \infty, \int_X \phi_0(x) d\nu(x) < \infty \right\}.$$

We then have

Theorem 5.1. *Assume that $m(X) < \infty$. Then for $\nu \in \mathcal{P}_0$ and $B \in \mathcal{B}(X)$*

$$\lim_{t \rightarrow \infty} e^{\lambda_2(\mu+F)t} \mathbb{E}_\nu(e^{A_t^{\mu+F}}; X_t \in B) = \int_X \phi_0 d\nu \int_B \phi_0 dm.$$

Proof. Note that

$$e^{\lambda_2(\mu+F)t} \mathbb{E}_\nu(e^{A_t^{\mu+F}}; X_t \in B) = \int_X \phi_0(x) \mathbb{E}_x^{\phi_0} \left(\frac{1_B}{\phi_0}(X_t) \right) d\nu(x).$$

Let $\{E_\lambda, 0 \leq \lambda < \infty\}$ be the spectral family of $(\mathcal{E}^{\phi_0}, \mathcal{F}^{\phi_0})$. Then $\lim_{t \rightarrow \infty} p_t^{\phi_0} f = E_0 f$ in $L^2(X; \phi_0^2 m)$. Since $\mathcal{E}^{\phi_0}(E_0 f, E_0 f) = 0$, $E_0 f$ equals $\int_X f \phi_0^2 dm$, m -a.e. by the irreducibility of $(\mathcal{E}^{\phi_0}, \mathcal{F}^{\phi_0})$ (cf. [7, Theorem 5.2.13]). Note that $p_t^{\phi_0}(x, \cdot) \in L^2(X; \phi_0^2 m)$ because $\int_X p_t^{\phi_0}(x, y)^2 \phi_0^2(y) dm(y) = p_{2t}^{\phi_0}(x, x) < \infty$. Put $c = \int_B \phi_0 dm$. We then have

$$(5.2) \quad \begin{aligned} & \left| \int_X \phi_0(x) \mathbb{E}_x^{\phi_0} \left(\frac{1_B}{\phi_0}(X_t) \right) d\nu(x) - \int_X \phi_0 d\nu \int_B \phi_0 dm \right| \\ &= \left| \int_X \phi_0(x) \left(\int_X p_{1/2}^{\phi_0}(x, y) \left(\mathbb{E}_y^{\phi_0} \left(\frac{1_B}{\phi_0}(X_{t-1/2}) \right) - c \right) \phi_0(y)^2 dm(y) \right) d\nu(x) \right|. \end{aligned}$$

The right-hand side is dominated by

$$\int_X \phi_0(x) \sqrt{\int_X p_{1/2}^{\phi_0}(x, y)^2 \phi_0^2(y) dm(y)} d\nu(x) \cdot \sqrt{\int_X \left(\mathbb{E}_y^{\phi_0} \left(\frac{1_B}{\phi_0}(X_{t-1/2}) \right) - c \right)^2 \phi_0^2(y) dm(y)}.$$

Since

$$p_t^{\phi_0}(x, y) = e^{\lambda_2(\mu+F)t} \frac{p_t^{\mu+F}(x, y)}{\phi_0(x)\phi_0(y)},$$

the first factor is equal to

$$\int_X \phi_0(x) \sqrt{p_1^{\phi_0}(x, x)} d\nu(x) = e^{(1/2)\lambda_2(\mu+F)} \int_X \sqrt{p_1^{\mu+F}(x, x)} d\nu(x)$$

and is finite by the assumption that $\nu \in \mathcal{P}_0$. Hence the right-hand side of (5.2) converges to zero as $t \rightarrow \infty$ because $1_B/\phi_0 \in L^2(X; \phi_0^2 m)$. \square

Let η and $R_{\nu,t}$ be probability measures on X defined by

$$(5.3) \quad \eta(B) = \frac{\int_B \phi_0(x) dm(x)}{\int_X \phi_0(x) dm(x)}, \quad R_{\nu,t}(B) = \frac{\mathbb{E}_\nu(e^{A_t^{\mu+F}}; X_t \in B)}{\mathbb{E}_\nu(e^{A_t^{\mu+F}}; t < \zeta)} \quad \text{for } B \in \mathcal{B}(X).$$

Corollary 5.1. *For $\nu \in \mathcal{P}_0$ and $B \in \mathcal{B}(X)$*

$$(5.4) \quad \lim_{t \rightarrow \infty} R_{\nu,t}(B) = \eta(B).$$

Note that the Dirac measure δ_x belongs to \mathcal{P}_0 and so the distribution $R_{\delta_x,t}$ converges to η for all $x \in X$. Hence Corollary 5.1 says that the semigroup $\{p_t^{\mu+F}\}_{t \geq 0}$ is *conditionally ergodic* and η is a *quasi-stationary distribution* of the semigroup $\{p_t^{\mu+F}\}_{t \geq 0}$: for any $t > 0$

$$(5.5) \quad R_{\eta,t} = \eta$$

(e.g. [16]). If the semigroup $\{p_t^{\mu+F}\}_{t \geq 0}$ is ultracontractive, $p_t^{\mu+F}(x, y) \leq c_t$, then $p_t^{\mu+F}(x, x)$ and $\phi_0(x)$ are bounded and \mathcal{P}_0 equals \mathcal{P} . Consequently, for any $\nu \in \mathcal{P}$, the distribution $R_{\nu,t}$ converges to η .

When the measure m is not finite, we assume the *intrinsic ultracontractivity* of $\{p_t^{\mu+F}\}_{t \geq 0}$, that is,

$$(5.6) \quad p_t^{\mu+F}(x, y) \leq C_t \phi_0(x) \phi_0(y).$$

In [16], they proved that for a (not necessary symmetric) Markov process, the intrinsic ultracontractivity is a sufficient condition for the measure η being a unique quasi-stationary distribution, and the equation (5.4) holds for any initial distribution. We would like to give another proof of this fact by using the next theorem due to Fukushima [13].

Theorem 5.2. *Assume that $m(X) < \infty$ and \mathbf{M} is conservative, $p_t 1 = 1$, $t > 0$. Then for $f \in L^1(X; m)$,*

$$\lim_{t \rightarrow \infty} p_t f(x) = \frac{1}{m(X)} \int_X f(x) dm(x), \quad m\text{-a.e. and in } L^1(X; m).$$

Note that \mathbf{M}^{ϕ_0} satisfies the assumptions in Theorem 5.2.

Theorem 5.3. *Assume that $\{p_t^{\mu+F}\}_{t \geq 0}$ is intrinsically ultracontractive. Then for any $\nu \in \mathcal{P}$ and any $B \in \mathcal{B}(X)$*

$$\lim_{t \rightarrow \infty} e^{\lambda_2(\mu+F)t} \mathbb{E}_\nu(e^{A_t^{\mu+F}}; X_t \in B) = \int_X \phi_0 d\nu \int_B \phi_0 dm.$$

Consequently, the equation (5.4) follows.

Proof. First note that the upper bound (5.6) implies the lower bound ([9, Theorem 4.2.5]):

$$(5.7) \quad c_t \phi_0(x) \phi_0(y) \leq p_t^{\mu+F}(x, y).$$

As a result,

$$\sup_{x \in X} \phi_0(x) \int_X \phi_0(y) dm(y) \leq \frac{1}{c_t} \|p_t^{\mu+F} 1\|_\infty < \infty.$$

Hence ϕ_0 belongs to $L^1(X; m) \cap L^\infty(X; m)$ and $1_B/\phi_0 \in L^1(X; \phi_0^2 m)$. Applying Theorem 5.2 to \mathbf{M}^{ϕ_0} , we have

$$\mathbb{E}_y^{\phi_0} \left(\frac{1_B}{\phi_0}(X_t) \right) \rightarrow \int_B \phi_0 dm, \quad m\text{-a.e. } y \quad \text{and} \quad L^1(X; \phi_0^2 m)$$

as $t \rightarrow \infty$. Since $p_{1/2}^{\phi_0}(x, \cdot)$ is bounded by the ultracontractivity, it follows from the equation (5.2) that

$$\lim_{t \rightarrow \infty} \int_X \phi_0(x) \mathbb{E}_x^{\phi_0} \left(\frac{1_B}{\phi_0}(X_t) \right) d\nu(x) = \int_X \phi_0 d\nu \int_B \phi_0 dm. \quad \square$$

We finally consider the exponential integrability of hitting times of compact sets. Let $K \subset X$ be a compact set and D the complement of K , $D = X \setminus K$. We define the part (or absorbing) process X^D on D by

$$X_t^D = \begin{cases} X_t & t < \tau_D, \\ \Delta & t \geq \tau_D, \end{cases} \quad \tau_D = \inf\{t \geq 0: X_t \notin D\}.$$

Define the regular Dirichlet form $(\mathcal{E}^D, \mathcal{D}(\mathcal{E}^D))$ on D by

$$\begin{cases} \mathcal{E}^D = \mathcal{E}, \\ \mathcal{D}(\mathcal{E}^D) = \{u \in \mathcal{D}(\mathcal{E}): u = 0 \text{ q.e. on } K\}. \end{cases}$$

By [14, Theorem 4.4.3] the part process X^D is regarded as a Hunt process generated by $(\mathcal{E}^D, \mathcal{D}(\mathcal{E}^D))$. We see from [4, Theorem 4.2] that m is in \mathcal{K}_∞ . We write $\mathcal{K}_\infty(R_1)$ for \mathcal{K}_∞ to show the dependence. Let R_1^D be the 1-resolvent of X^D . The restriction m^D of m on D is in $\mathcal{K}_\infty(R_1^D)$. Indeed, let a compact set \tilde{K} and a positive constant δ in the definition of \mathcal{K}_∞ (Definition 2.1). We can suppose $K \subset \tilde{K}$. Let G be a relatively compact open set such that $K \subset G \subset \tilde{G} \subset \tilde{K}$ and $m(G \setminus K) < \delta$. Then $\tilde{K} \cap G^c$ is a compact subset of D and

$$R_1^D 1_{(\tilde{K} \cap G^c)^c} = R_1^D 1_{\tilde{K}^c \cup (G \setminus K)} \leq R_1 1_{\tilde{K}^c} + R_1 1_{G \setminus K} \leq 2\epsilon.$$

Moreover, $R_1^D 1_B \leq R_1 1_B$ for any Borel set $B \subset \tilde{K} \cap G^c$. Hence we have $m^D \in \mathcal{K}_\infty(R_1^D)$.

If X^D satisfies the irreducibility (I), it follows from [4, Theorem 4.1] that

$$\sup_{x \in D} \mathbb{E}_x(e^{\lambda \tau_D}) < \infty \iff \lambda < \lambda^D,$$

where λ^D is the bottom of the spectrum of $(\mathcal{E}^D, \mathcal{D}(\mathcal{E}^D))$.

Noting that by (3.1)

$$1 \leq \|R_1 1_D\|_\infty(\lambda^D + 1),$$

we see from (III) that

$$(5.8) \quad \lambda_D \uparrow \infty \quad \text{as} \quad K \uparrow X.$$

We can conclude that if for any compact set K , the part process X^D ($D = X \setminus K$) is irreducible, then for any $\lambda > 0$ there exists a compact set K such that

$$(5.9) \quad \sup_{x \in X} \mathbb{E}_x(e^{\lambda \tau_D}) < \infty.$$

If \mathbf{M} is conservative, τ_D equals the first hitting time σ_K of K , $\sigma_K = \inf\{t > 0 : X_t \in K\}$. Then the property (5.9) is called the *uniform hyper-exponential recurrence* in [30].

EXAMPLE 5.1 (One-dimensional diffusion processes). Let us consider a one-dimensional diffusion process $\mathbf{M} = (X_t, \mathbb{P}_x, \zeta)$ on an open interval $I = (r_1, r_2)$ such that $\mathbb{P}_x(X_{\zeta^-} = r_1 \text{ or } r_2, \zeta < \infty) = \mathbb{P}_x(\zeta < \infty)$, $x \in I$, and $\mathbb{P}_a(\sigma_b < \infty) > 0$ for any $a, b \in I$. The diffusion \mathbf{M} is symmetric with respect to its canonical measure m and it satisfies (I) and (II). The boundary point r_i of I is classified into four classes: *regular boundary*, *exit boundary*, *entrance boundary* and *natural boundary* ([15, Chapter 5]):

- (a) If r_2 is a regular or exit boundary, then $\lim_{x \rightarrow r_2} R_1 1(x) = 0$.
- (b) If r_2 is an entrance boundary, then $\lim_{r \rightarrow r_2} \sup_{x \in (r_1, r_2)} R_1 1_{(r_1, r_2)}(x) = 0$.
- (c) r_2 is a natural boundary, then $\lim_{x \rightarrow r_2} R_1 1_{(r_1, r_2)}(x) = 1$ and thus $\sup_{x \in (r_1, r_2)} R_1 1_{(r_1, r_2)}(x) = 1$.

Therefore, (III) is satisfied if and only if no natural boundaries are present. As a corollary of the equation (5.8), If r_2 is entrance, for any $\lambda > 0$ there exists $r_1 < r < r_2$ such that

$$\sup_{x > r} \mathbb{E}_x(\exp(\lambda \sigma_r)) < \infty,$$

where σ_r is the first hitting time of $\{r\}$. The statement above implies a uniqueness of quasi-stationary distributions ([3]).

References

- [1] S. Albeverio, P. Blanchard and Z.M. Ma: *Feynman–Kac semigroups in terms of signed smooth measures*; in Random Partial Differential Equations (Oberwolfach, 1989), Birkhäuser, Basel, 1991, 1–31.
- [2] S. Albeverio and Z.M. Ma: *Additive functionals, nowhere Radon and Kato class smooth measures associated with Dirichlet forms*, Osaka J. Math. **29** (1992), 247–265.
- [3] P. Cattiaux, P. Collet, A. Lambert, S. Martinez, S. Meleard and J. San Martin: *Quasi-stationary distributions and diffusion models in population dynamics*, Ann. Probab. **37** (2009), 1926–1969.
- [4] Z.-Q. Chen: *Gaugeability and conditional gaugeability*, Trans. Amer. Math. Soc. **354** (2002), 4639–4679.
- [5] Z.-Q. Chen: *Analytic characterization of conditional gaugeability for non-local Feynman–Kac transforms*, J. Funct. Anal. **202** (2003), 226–246.
- [6] Z.-Q. Chen, P.J. Fitzsimmons, M. Takeda, J. Ying and T.S. Zhang: *Absolute continuity of symmetric Markov processes*, Ann. Probab. **32** (2004), 2067–2098.
- [7] Z.-Q. Chen and M. Fukushima: *Symmetric Markov Processes, Time Change and Boundary Theory*, book manuscript, 2009.
- [8] Z.-Q. Chen and R. Song: *Conditional gauge theorem for non-local Feynman–Kac transforms*, Probab. Theory Related Fields **125** (2003), 45–72.
- [9] E.B. Davies: *Heat Kernels and Spectral Theory*, Cambridge Univ. Press, Cambridge, 1989.
- [10] G. De Leva, D. Kim and K. Kuwae: *L^p -independence of spectral bounds of Feynman–Kac semigroups by continuous additive functionals*, J. Funct. Anal. **259** (2010), 690–730.
- [11] J.-D. Deuschel and D.W. Stroock: *Large Deviations*, Academic Press, Boston, MA, 1989.
- [12] M.D. Donsker and S.R.S. Varadhan: *Asymptotic evaluation of certain Markov process expectations for large time*, I, Comm. Pure Appl. Math. **28** (1975), 1–47.
- [13] M. Fukushima: *A note on irreducibility and ergodicity of symmetric Markov processes*; in Stochastic Processes in Quantum Theory and Statistical Physics (Marseille, 1981), Lecture Notes in Phys. **173**, Springer, Berlin, 1982, 200–207.
- [14] M. Fukushima, Y. Oshima and M. Takeda: *Dirichlet Forms and Symmetric Markov Processes*, second rev. and ext. ed., de Gruyter, Berlin, 2010.
- [15] K. Itô: *Essentials of Stochastic Processes*, Amer. Math. Soc., Providence, RI, 2006.
- [16] R. Knobloch and L. Partzsch: *Uniform conditional ergodicity and intrinsic ultracontractivity*, Potential Anal. **33** (2010), 107–136.
- [17] E.H. Lieb and M. Loss: *Analysis*, second edition, Graduate Studies in Mathematics **14**, Amer. Math. Soc., Providence, RI, 2001.
- [18] Z.M. Ma and M. Röckner: *Introduction to the Theory of (Nonsymmetric) Dirichlet Forms*, Universitext, Springer, Berlin, 1992.
- [19] P. Stollmann and J. Voigt: *Perturbation of Dirichlet forms by measures*, Potential Anal. **5** (1996), 109–138.
- [20] M. Takeda: *Asymptotic properties of generalized Feynman–Kac functionals*, Potential Anal. **9** (1998), 261–291.
- [21] M. Takeda: *Conditional gaugeability and subcriticality of generalized Schrödinger operators*, J. Funct. Anal. **191** (2002), 343–376.
- [22] M. Takeda: *Large deviation principle for additive functionals of Brownian motion corresponding to Kato measures*, Potential Anal. **19** (2003), 51–67.
- [23] M. Takeda: *Large deviations for additive functionals of symmetric stable processes*, J. Theoret. Probab. **21** (2008), 336–355.
- [24] M. Takeda: *A large deviation principle for symmetric Markov processes with Feynman–Kac functional*, to appear in J. Theoret. Probab.
- [25] M. Takeda and Y. Tawara: *L^p -independence of spectral bounds of non-local Feynman–Kac semigroups*, Forum Math. **21** (2009), 1067–1080.
- [26] M. Takeda and K. Tsuchida: *Differentiability of spectral functions for symmetric α -stable processes*, Trans. Amer. Math. Soc. **359** (2007), 4031–4054.
- [27] M. Takeda and T. Uemura: *Subcriticality and gaugeability for symmetric α -stable processes*, Forum Math. **16** (2004), 505–517.

- [28] Y. Tawara: *L^p -independence of spectral bounds of Schrödinger-type operators with non-local potentials*, J. Math. Soc. Japan **62** (2010), 767–788.
- [29] S.R.S. Varadhan: *Asymptotic probabilities and differential equations*, Comm. Pure Appl. Math. **19** (1966), 261–286.
- [30] L. Wu: *Some notes on large deviations of Markov processes*, Acta Math. Sin. (Engl. Ser.) **16** (2000), 369–394.
- [31] J. Ying: *Bivariate Revuz measures and the Feynman–Kac formula*, Ann. Inst. H. Poincaré Probab. Statist. **32** (1996), 251–287.
- [32] J. Ying: *Dirichlet forms perturbated by additive functionals of extended Kato class*, Osaka J. Math. **34** (1997), 933–952.

Masayoshi Takeda
Mathematical Institute
Tohoku University
Aoba, Sendai, 980-8578
Japan
e-mail: takeda@math.tohoku.ac.jp

Yoshihiro Tawara
Division of General Education
Nagaoka National College of Technology
888 Nishikatakai, Nagaoka, Niigata 940-8532
Japan
e-mail: tawara@nagaoka-ct.ac.jp